A gene expression ratio-based diagnostic test for bladder cancer

Lingsheng Dong¹
Andrew J Bard¹
William G Richards¹
Matthew D Nitz²
Dan Theodorescu²
Raphael Bueno¹
Gavin J Gordon¹

¹The Thoracic Surgery Oncology laboratory and the Division of Thoracic Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA;

²Departments of Urology and Molecular Physiology, University of Virginia, Charlottesville, VA, USA

Purpose: Bladder cancer is relatively common but early detection techniques such as cystoscopy and cytology are somewhat limited. We developed a broadly applicable, platform-independent and clinically relevant method based on simple ratios of gene expression to diagnose human cancers. In this study, we sought to determine whether this technique could be applied to the diagnosis of bladder cancer.

Experimental design: We developed a model for the diagnosis of bladder cancer using expression profiling data from 80 normal and tumor bladder tissues to identify statistically significant discriminating genes with reciprocal average expression levels in each tissue type. The expression levels of select genes were used to calculate individual gene pair expression ratios in order to assign diagnosis. The optimal model was examined in two additional published microarray data sets and using quantitative RT-PCR in a cohort of 13 frozen benign bladder urothelium samples and 13 bladder cancer samples from our institution.

Results: A five-ratio test utilizing six genes proved to be 100% accurate (26 of 26 samples) for distinguishing benign from malignant bladder tissue samples ($P < 10^{-6}$).

Conclusions: We have provided a proof of principle study for the use of gene expression ratios in the diagnosis of bladder cancer. This technique may ultimately prove to be a useful adjunct to cytopathology in screening urine specimens for bladder cancer.

Keywords: bladder cancer, gene expression profiling, and diagnosis

Introduction

The American Cancer Society (www.cancer.org) estimates there were 67,160 new cases of bladder cancer in the United States during 2007 with approximately 13,750 bladder cancer-related deaths during the same period. The risk of developing bladder cancer is higher in men (~1 in 30) than in women (~1 in 90). Hematuria or changes in bladder habits are the most common symptoms associated with bladder cancer. However, fewer than 10% of patients with these symptoms actually have bladder cancer. Furthermore, many patients with early disease are asymptomatic and are only diagnosed at advanced stage when the therapeutic options are limited. Cystoscopy is currently the most accurate and sensitive method for bladder cancer diagnosis. However, this endoscopic procedure is relatively expensive and is associated with discomfort, inconvenience, and other risk. Therefore, it is only indicated in cases where clinical suspicion for bladder tumors exists or for surveillance in patients with known bladder cancer.

We have developed a simple, effective gene expression-based algorithm to develop specific tests for cancer detection, diagnosis, and prognosis.¹⁻⁶ This algorithm is based

Correspondence: Gavin J Gordon
Brigham and Women's Hospital, Division
of Thoracic Surgery, 75 Francis Street,
Boston, MA 02115, USA
Tel +1 617 732 6526
Fax +1 617 582 6085
Email ggordon@partners.org

on an initial supervised comparison of gene expression data between two groups that differ in a chosen cancer-related characteristic, such as benign versus malignant. Molecular biomarkers discovered are then used to calculate expression level ratios of select gene pairs that numerically assign tested samples to a given diagnostic or prognostic category. In this report, we utilized this ‘gene ratio’ algorithm in a proof of principle study using gene expression profiling data and discarded surgical specimens of malignant and benign bladder tissues.

Materials and methods

Human tissues

Thirteen sets of matched adjacent benign bladder and bladder cancer samples were obtained from the Tumor Bank at Brigham and Women’s Hospital. These frozen samples were discarded surgical tissues obtained after open surgical resection for bladder cancer and confirmed histologically by the tumor bank pathologist. Specimen identities were rendered anonymous and linked to select clinical and pathologic data. Eleven specimens were urothelial carcinoma and two were transitional cell carcinoma. Ten of the patients were male. Median age was 67 (range 40–79) years. Studies utilizing human tissues were approved by and conducted in accordance with the policies of the Institutional Review Board at Brigham and Women’s Hospital.

Gene expression profiling data

A total of four global gene expression datasets of bladder tissues were used. “Training Set A” consisted of six macrodissected normal bladder mucosa samples from patients undergoing bladder surgery for benign causes and 19 malignant bladder samples.⁷ “Training Set B” consisted of nine benign biopsies of normal bladder mucosa and 46 malignant bladder tissues samples (accessible at NCBI GEO database with accession number GSE3167).⁸ Training sets A and B were obtained using Affymetrix high-density oligonucleotide microarrays with probe sets representing approximately 22,000 genes. “Test Set A” (ie, combination of GSE88 and GSE89) consisted of 66 unique bladder tumors without matched normal tissue.⁹ These data were obtained using Affymetrix high-density oligonucleotide microarrays with probe sets representing approximately 5,600 genes. For these three datasets, Affymetrix .cel files were used to obtain gene hybridization intensities, which were scaled to a “target intensity” of 100 using Affymetrix Microarray Suite v.5.0. “Test Set B” was obtained from the supplemental

files of a recently published manuscript.¹⁰ This data set consisted of 157 samples composed of 48 histological normal urothelium samples obtained at distant sites from the bladder tumors resected by cystectomy or cystoprostatectomy and 109 bladder tumor samples, taken from 105 patients. There were several redundant samples and more than one microarray performed on each tissue, which reduced the number of unique samples to 39 benign bladder and 91 bladder tumors. Because Affymetrix .cel files were not available for this dataset, we used the values obtained by the authors analysis via Affymetrix Microarray Suite v.5.0 with a scaled “target intensity” of 500.

Real-time quantitative RT-PCR

Real-time quantitative reverse transcription-polymerase chain reaction (RT-PCR) was performed as described using 2 µg of total RNA.³ Forward and reverse RT-PCR primers (synthesized by Invitrogen Life Technologies, Carlsbad, CA) were designed to span large introns (>800 bp if possible) or to be positioned on the exon/exon junction. Primers used at a final concentration of 800 nM in the PCR reaction mixture were as follows (forward and reverse, respectively): SPARCL1 (5’-AGGAGTGTGACCCCAACAAG-3’ and 5’-CAGAGGAGGATGCTGGAAAG-3’), MYH11 (5’-CACAGGAACTTCGCAGTGA-3’ and 5’-GAGTGTCGTTTCCTCCTCA-3’), FHL1 (5’-GAAGTGTGCTGGATGCAAGA-3’ and 5’-CCAGATTCACGGAGCATT-3’), MIF (5’-CCGGACAGGGTCTACATCA-3’ and 5’-GGCGGGCCTAGAACACAG-3’), H2AFZ (5’-GCTGGTGGTGGTGTCATTC-3’ and 5’-CTGGAATCACCAACACTGGA-3’), and RPN2 (5’-CACTTTTGCTCCTAGCACGA-3’ and 5’-CAGCCAGAAACGTCACACTG-3’). PCR amplification of cDNA was performed using a Stratagene MX3000P machine and default thermal cycling parameters. No-template (ie, negative) controls that contained water instead of template were run in multiple wells on every reaction plate. An automatically calculated melting point dissociation curve generated after every assay was examined to ensure the presence of a single PCR species and the lack of primer-dimer formation in each well. The comparative CT equation (Applied Biosystems) describes the exponential nature of PCR-based amplification and was used, with minor modifications, to obtain quantitative values for gene expression ratios in all samples. The “CT” term stands for the fractional PCR cycle at which the quantity of the amplified product reaches a pre-determined threshold. The comparative CT equation states that the expression level of a gene in a given sample, normalized within the sample to an endogenous reference

gene, and relative to the expression level of the same gene in another sample (ie, an arbitrarily chosen “calibrator sample”) can be represented as: $2^{-\Delta\Delta CT}$ where $\Delta\Delta CT = [\Delta CT_{(\text{sample “x”})}] - [\Delta CT_{(\text{calibrator sample})}]$ and $\Delta CT = [CT_{(\text{target gene})}] - [CT_{(\text{reference gene})}]$. Calculation of an expression ratio using data from two rationally selected genes in any single sample negates the need for a calibrator sample and a reference gene to standardize when using different amounts of starting template. Therefore, to form expression ratios of two genes in a single sample, we simply presented the expression level of one gene relative to the expression level of the other gene. In this case, the $\Delta\Delta CT$ value in the comparative CT equation was expressed as $[CT_{(\text{gene 1})} - CT_{(\text{gene 2})}]$.

Data and statistical analysis

The selection of predictor genes was performed essentially as previously described using standard statistical techniques.^{1–5} Essentially, we used a two-sided Student’s (parametric) *t*-test for pair-wise comparisons of average gene expression levels of all genes represented on the microarrays of both training sets to identify those differentially expressed in a statistically significant manner ($P < 0.01$) between benign and malignant bladder. (Although Training Sets A and B were from the same platform, these data were considered separately for gene discovery purposes to minimize variability and account for potential site-specific confounders.) To further reduce the risk of false positives, we chose for additional consideration those genes with a ≥ 2 -fold difference in average expression levels and an average gene expression level > 600 in at least one of the two subsets in both Training Set A and Training Set B. The SAM algorithm¹¹ was used to estimate the false discovery rate.

Data from multiple highly accurate gene expression ratios were combined as described previously^{3,6} by calculating the geometric mean (ie, “combined score”), $(R_1 R_2 R_3)^{1/3}$, where R_i represents a single ratio value. This is equivalent to the average of $[\log_2(R_1), \log_2(R_2), \log_2(R_3)]$, and has the effect of giving equal weight to ratio fold-changes of identical magnitude but opposite direction. The classification accuracy of the model was assessed using an exact one-sample binomial test. The *P* values are reported under the null hypothesis of diagnosis randomly assigned with equal probability of 0.5 based on one-sided tests in order to reject lower levels of accuracy. The 95% confidence interval (CI) for proportions is based on the exact binomial distribution. All calculations and statistical comparisons were generated using S-PLUS,¹² except the exact binomial procedures that were computed using Stata 7 (StataCorp, College Station, TX).

Results

Identification of molecular markers in bladder cancer and prediction of diagnosis using gene expression ratios

The expression profiles of bladder tumors and normal urothelial tissues from Training Set A were compared to identify 22 differentially expressed candidate diagnostic genes that fit the filtering criteria (Table 1). The false discovery rate (FDR) associated with these genes was 13%, as calculated using the SAM algorithm.¹¹ All of these genes were also statistically significantly differentially expressed in Training Set B with a similar minimum two-fold difference in means (Table 1). Six of these genes with the lowest *P* values were chosen for further analysis: SPARCL1, MYH11, and FHL1 (overexpressed in adjacent nontumorous bladder specimens) and MIF, H2AFZ, and RPN2 (overexpressed in bladder tumor specimens).

Next, multiple expression ratios of these genes were evaluated as to whether they could accurately classify the 80 samples Training Sets A and B combined. Nine possible expression ratios were calculated per sample by dividing the expression value of each of the three genes expressed at relatively higher levels in the benign bladder samples by the expression value of each of the three genes expressed at relatively higher levels in the bladder tumors. Samples with ratio values > 1 were predicted to be “benign” and those with ratio values < 1 were predicted to be “malignant”. These nine individual ratios were found to be accurate (range 84%–94%; Table 2) in distinguishing benign from malignant bladder. The geometric mean of the combination of the five individual pair ratios that were at least 90% accurate from Table 2 (SPARCL1/MIF, SPARCL1/RPN2, MYH11/RPN2, FHL1/MIF, FHL1/RPN2) was calculated. This test was highly accurate (73/80, 91%; 95% CI 86%–100%, $P < 10^{-6}$) in the combined training sets. Importantly, this test was highly accurate in diagnosing tumors (61/65, 94%) but was slightly less accurate in the benign bladder subset (12/15, 80%) perhaps due to small sample size considerations and/or contamination by adjacent microscopic tumor elements.

This five-ratio test was then examined in the independent test sets. High diagnostic sensitivity was maintained in Test Set A (98%, 65/66, 95% CI 92%–100%, $P < 10^{-6}$). For Test Set B, when using all the available data (include duplicated samples), the sensitivity (85%, 86/109, 95% CI 78%–90%, 133/157) was significantly ($P < 10^{-6}$) high with nearly all benign samples called correctly (98%, 47/48). When the duplicated samples and possible histologically

Table 1 Bladder cancer diagnostic genes

Probe set	Accession #	Fold-change training set A ^a	P value training set A ^a	Fold-change training set B ^a	P value training set B ^a	Gene symbol	Description
200795_at	NM_004684	6.08	2.15 × 10 ⁻⁸	2.37	6.33 × 10 ⁻⁵	SPARCLI	Expressed at relatively higher levels in benign bladder
201497_x_at	NM_022844	24.08	4.18 × 10 ⁻⁸	6.36	2.01 × 10 ⁻⁷	MYH11	SPARC-like I (mast9, hev1n) Myosin, heavy polypeptide 11, smooth muscle
201540_at	NM_001449	11.07	3.15 × 10 ⁻⁷	2.97	1.10 × 10 ⁻³	FHL1	Four and a half LIM domains 1
200974_at	NM_001613	8.79	5.20 × 10 ⁻⁷	2.93	2.93 × 10 ⁻⁴	ACTA2	Actin, alpha 2, smooth muscle, aorta
202555_s_at	NM_005965	18.36	2.02 × 10 ⁻⁶	4.29	1.83 × 10 ⁻⁵	MYLK	Myosin, light polypeptide kinase
210987_x_at	M19267	7.30	2.52 × 10 ⁻⁶	2.85	3.52 × 10 ⁻⁵	TPM1	Tropomyosin
203951_at	NM_001299	44.07	2.53 × 10 ⁻⁶	7.30	4.44 × 10 ⁻⁶	CNN1	Calponin 1, basic, smooth muscle
202274_at	NM_001615	23.47	3.18 × 10 ⁻⁶	4.39	4.29 × 10 ⁻⁵	ACTG2	Actin, gamma 2, smooth muscle, enteric
201058_s_at	NM_006097	16.97	3.52 × 10 ⁻⁶	6.00	9.46 × 10 ⁻⁶	MYRL2	Myosin regulatory light chain 2, smooth muscle isoform
202222_s_at	NM_001927	29.79	4.19 × 10 ⁻⁶	11.36	1.73 × 10 ⁻⁶	DES	Desmin
205547_s_at	NM_003186	10.32	4.24 × 10 ⁻⁶	3.94	7.96 × 10 ⁻⁵	TAGLN	Transgelin
207961_x_at	NM_022870	24.82	1.53 × 10 ⁻⁵	6.28	1.74 × 10 ⁻⁶	MYH11	Myosin, heavy polypeptide 11, smooth muscle
214752_x_at	AI625550	5.27	1.78 × 10 ⁻⁵	2.95	4.53 × 10 ⁻⁵	FLNA	Filamin A, alpha (actin-binding protein-280)
200621_at	NM_004078	6.75	2.94 × 10 ⁻⁵	2.25	3.46 × 10 ⁻⁴	CSRPI	Cysteine and glycine-rich protein 1
200859_x_at	NM_001456	6.53	3.18 × 10 ⁻⁵	3.02	1.03 × 10 ⁻⁴	FLNA	Filamin A, alpha (actin-binding protein-280)
205382_s_at	NM_001928	27.85	1.36 × 10 ⁻⁴	2.51	3.48 × 10 ⁻³	DF	D component of complement (adipsin)
202822_at	BF221852	2.68	5.55 × 10 ⁻⁴	2.10	5.86 × 10 ⁻⁵	LPP	LIM domain-containing preferred translocation partner in lipoma
205132_at	NM_005159	23.63	9.13 × 10 ⁻⁴	5.34	4.77 × 10 ⁻⁶	ACTC	Actin, alpha, cardiac muscle
217871_s_at	NM_002415	0.39	5.18 × 10 ⁻⁴	0.45	5.10 × 10 ⁻³	MIF	Expressed at relatively higher levels in bladder cancer Macrophage migration inhibitory factor
213911_s_at	BF718636	0.44	5.49 × 10 ⁻³	0.45	4.01 × 10 ⁻⁵	H2AFZ	H2A histone family, member Z
213491_x_at	AL514285	0.45	7.62 × 10 ⁻³	0.38	1.75 × 10 ⁻⁵	RPN2	Ribophorin II
202428_x_at	NM_020548	0.48	8.09 × 10 ⁻³	0.47	4.72 × 10 ⁻³	DBI	Diazepam-binding inhibitor

Notes: ^aAverage expression level in benign bladder specimens/average expression level in malignant bladder specimens; Genes in bold identified for further analysis.

Table 2 Accuracy of all ratio combinations in predicting tumor diagnosis in training sets A and B combined

	SPARCL1	MYH11	FHL1
MIF	91% (73/80)	89% (71/80)	90% (72/80)
H2AFZ	84% (67/80)	84% (67/80)	89% (71/80)
RPN2	91% (73/80)	94% (75/80)	90% (72/80)

Notes: Six diagnostic genes were identified in a training set of samples as described in the text. A total of nine possible expression ratios (column/row intersection) were calculated where both genes used to form the ratio possessed inversely correlated expression levels in both tissue types. Predictions are stated as the fraction diagnosed correctly.

misdiagnosed samples (ie, samples that were clustered with the opposite histologic classification by unsupervised clustering analysis in the original analysis¹⁰) were excluded from analysis, the five-ratio test could diagnose the remaining 36 benign bladder and 81 bladder tumor samples with moderately higher (91%, 84%–95%, 106/117, $P < 10^{-6}$) accuracy (or 70/81 for tumor and 36/36 for normal).

Validation of expression level ratios as a diagnostic tool in bladder cancer

The five-ratio test was then examined in an independent set of 13 matched benign bladder urothelium samples and bladder cancer samples ($n = 26$ samples total) using quantitative RT-PCR (Figure 1). The five-ratio diagnostic test was 100% accurate (26/26, 95% CI 87%–100%, $P < 10^{-6}$) in classifying

Figure 1 Verification of expression level ratios as a diagnostic tool in bladder cancer. Quantitative RT-PCR was used to obtain gene expression levels for the six bladder cancer diagnostic genes and calculate the geometric mean of the five-ratio diagnostic test (SPARCL1/MIF, SPARCL1/RPN2, MYH11/RPN2, FHL1/MIF, FHL1/RPN2) in frozen samples of benign bladder urothelium ($n = 13$) and bladder cancer ($n = 13$). This test identified 26 of 26 samples and did not result in any false-negatives. Note that the y-axis is in log scale.

these samples (13/13, 95% CI 75%–100%, $P = 0.00012$ for each subset).

Discussion

To improve the diagnostic accuracy of cytology, we explored the feasibility of developing a genomic-based diagnostic test for bladder cancer that would use quantitative RT-PCR for data acquisition. The high accuracy (100%) of our test in surgical samples is encouraging in this limited proof of principle study, but a prospective collection of additional larger numbers of tissue specimens will be necessary before determining whether this approach should be used in a clinical setting. Because of the relatively small quantity of RNA needed, it is interesting to speculate that this approach would support sample attainment using urine samples, considering that sufficient mRNA quantity and quality can be isolated from bladder washing even for stringent applications such as microarray analysis.¹³ One possible confounder is the relatively small number of malignant cells in voided urine from bladder cancer patients and the presence of other non-tumor cell types. Encouragingly, these issues have not proved insurmountable in similar applications such as the use of gene expression levels of relatively rare cells to diagnosis malignant pleural effusions.¹⁴

The mechanistic role in carcinogenesis of the diagnostic genes we discovered (if any) is not immediately clear by examining their known functions. To our knowledge, only two of these six genes have been reported to be differentially expressed in human cancer tissues. SPARCL1 is widely expressed in human normal or nonneoplastic diseased tissues.¹⁵ In contrast, its expression is strongly downregulated in most neoplastic cells and tissues¹⁵ consistent with our findings. MIF is upregulated in multiple human cancers including breast cancer,¹⁶ prostate cancer,¹⁷ and nonsmall cell lung cancer.¹⁸ The remaining genes (eg, MYH11, FHL1, H2AFZ, and RPN2) are not well studied. None of the diagnostic genes we identified in this study were found in the original analysis of the training set samples, perhaps due to the fact that these investigators were primarily interested in elucidating genes associated with metastatic potential.

Large-scale expression profiling using microarrays is ideally suited for use as a gene discovery tool and can also aid in the development and validation of predictive models as in our analysis of the test set of samples. However, microarray technology has limited use in clinical scenarios due to the complexity of current bioinformatics tools¹⁹ and the reproducibility, or lack thereof, of measured gene

expression levels.²⁰ On the other hand, quantitative RT-PCR is widely considered to be more accurate (and subject to less variability) than microarray analysis at the current time. Since gene-ratio based tests can utilize quantitative RT-PCR for data acquisition, where as other equally accurate bioinformatics models are unproven on this platform, it logically follows that the gene ratio technique will provide a clear advantage to clinical use.

In summary, we provide evidence that gene expression ratio-based cancer classification can be used to distinguish bladder cancer from normal bladder tissues in this pilot study similar to other cancers²⁻⁶ including prostate cancer.¹ Specifically, we found that a five-ratio combination (SPARCL1/MIF, SPARCL1/RPN2, MYH11/RPN2, FHL1/MIF, FHL1/RPN2) consisting of six genes was capable of distinguishing malignant from nonmalignant bladder tissues with 100% accuracy in surgical specimens. These data provide evidence to warrant additional study by multiple investigators to assess more thoroughly the efficacy of this technique in the diagnosis of bladder cancer.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Bueno R, Loughlin KR, Powell MH, Gordon GJ. A diagnostic test for prostate cancer from gene expression profiling data. *J Urol.* 2004;171:903–906.
2. Gordon GJ, Deters LA, Nitz MD, Lieberman BC, Yeap BY, Bueno R. Differential diagnosis of solitary lung nodules using gene expression ratios. *J Thorac Cardiovasc Surg.* 2006;132:621–627.
3. Gordon GJ, Jensen RV, Hsiao LL, et al. Using gene expression ratios to predict outcome among patients with mesothelioma. *J Natl Cancer Inst.* 16 2003;95:598–605.
4. Gordon GJ, Jensen RV, Hsiao L-L, et al. Translation of microarray data into clinically relevant cancer diagnostic tests using gene expression ratios in lung cancer and mesothelioma. *Cancer Res.* 2002;62:4963–4967.
5. Gordon GJ, Richards WG, Sugarbaker DJ, Jaklitsch MT, Bueno R. A prognostic test for adenocarcinoma of the lung from gene expression profiling data. *Cancer Epidemiol Biomarkers Prev.* 2003;12:905–910.
6. Gordon GJ, Rockwell GN, Godfrey PA, et al. Validation of genomics-based prognostic tests in malignant pleural mesothelioma. *Clin Cancer Res.* 2005;11:4406–4414.
7. Nicholson BE, Frierson HF, Conaway MR, et al. Profiling the evolution of human metastatic bladder cancer. *Cancer Res.* 2004;64:7813–7821.
8. Dyrskjot L, Kruhoffer M, Thykjaer T, et al. Gene expression in the urinary bladder: a common carcinoma in situ gene expression signature exists disregarding histopathological classification. *Cancer Res.* 2004;64:4040–4048.
9. Dyrskjot L, Thykjaer T, Kruhoffer M, et al. Identifying distinct classes of bladder carcinoma using microarrays. *Nat Genet.* 2003;33:90–96.
10. Sanchez-Carbajo M, Socci ND, Lozano J, Saint F, Cordon-Cardo C. Defining molecular profiles of poor outcome in patients with invasive bladder cancer using oligonucleotide microarrays. *J Clin Oncol.* 2006;24:778–789.
11. Tusher VG, Tibshirani R, Chu G. Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci U S A.* 2001;98:5116–5121.
12. Venables WN, Ripley BD. *Modern Applied Statistics with S-Plus.* New York, NY: Springer; 1997.
13. Mengual L, Burset M, Ars E, et al. Partially degraded RNA from bladder washing is a suitable sample for studying gene expression profiles in bladder cancer. *Eur Urol.* 2006;50:1347–1355.
14. Holloway AJ, Diyagama DS, Opeskin K, et al. A molecular diagnostic test for distinguishing lung adenocarcinoma from malignant mesothelioma using cells collected from pleural effusions. *Clin Cancer Res.* 2006;12:5129–5135.
15. Claeskens A, Ongenaet N, Neefs JM, et al. Hevin is down-regulated in many cancers and is a negative regulator of cell growth and proliferation. *Br J Cancer.* 2000;82:1123–1130.
16. Bando H, Matsumoto G, Bando M, et al. Expression of macrophage migration inhibitory factor in human breast cancer: association with nodal spread. *Jpn J Cancer Res.* 2002;93:389–396.
17. Meyer-Siegler KL, Iczkowski KA, Vera PL. Further evidence for increased macrophage migration inhibitory factor expression in prostate cancer. *BMC Cancer.* 2005;5:73.
18. White ES, Flaherty KR, Carskadon S, et al. Macrophage migration inhibitory factor and CXC chemokine expression in non-small cell lung cancer: role in angiogenesis and prognosis. *Clin Cancer Res.* 2003;9:853–860.
19. Quackenbush J. Computational analysis of microarray data. *Nat Rev Genet.* 2001;2:418–427.
20. Gullans SR. Of microarrays and meandering time points. *Nat Genet.* 2000;26:4–5.

Advances and Applications in Bioinformatics and Chemistry

Publish your work in this journal

Advances and Applications in Bioinformatics and Chemistry is an international, peer-reviewed open-access journal that publishes articles in the following fields: Computational biomodelling; Bioinformatics; Computational genomics; Molecular modelling; Protein structure modelling and structural genomics; Systems Biology; Computational

Submit your manuscript here: <http://www.dovepress.com/advances-and-applications-in-bioinformatics-and-chemistry-journal>

Dovepress

Biochemistry; Computational Biophysics; Chemoinformatics and Drug Design; In silico ADME/Tox prediction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.