

# PD-I immune checkpoint blockade may be viable for the prevention and treatment of elderly patients with POCD [Letter]

This article was published in the following Dove Press journal:  
*Clinical Interventions in Aging*

Penghui Wei  
Fan Yang  
Jianjun Li

Department of Anesthesiology, Qilu  
Hospital of Shandong University  
(Qingdao), Qingdao 266035, People's  
Republic of China

## Dear Editor

Recently, we read with great interest the well-written review article published in the November 8, 2018 edition of the *Clinical Interventions in Aging* by Kotekar and colleagues titled “Postoperative cognitive dysfunction – current preventive strategies”.<sup>1</sup> The article elegantly addressed the current preventive strategies for post-operative cognitive dysfunction (POCD). The authors concluded that the best preventive strategies for POCD may involve early recognition and management of potential perioperative risk factors, such as intraoperative anesthetic monitoring of the depth of anesthesia, Enhanced Recovery After Surgery and a multidisciplinary collaboration. Furthermore, the authors proposed that pharmacological interventions may be effective preventative and therapeutic strategies to attenuate POCD. We appreciate the efforts of the authors and would like to present our opinion on the review.

POCD is a highly prevalent condition with significant impact on the prognosis of elderly patients undergoing an operation, experiencing problems with memory, concentration, information processing, language comprehension, and social integration that can last for months or may even be permanent. Chronic neuroinflammation and accompanying accumulation of amyloid- $\beta$  (A $\beta$ ) and phosphorylation of tau protein secondary to surgery and anesthesia are strongly implicated in the pathophysiology of POCD.<sup>2</sup> The exact cascade by which surgical stress and anesthesia induce neuroinflammation in association with POCD remains unknown. Boosting degenerative systemic immune response in elderly patients may be a potential pharmacological strategy in treating POCD. Although previous studies have highlighted the role of the systemic immune response in inducing neuroinflammation, these results still did not show a significant association between systemic immune response and neuroinflammation.<sup>3,4</sup> A famous study revealed that surgery-induced neuroinflammation was associated with infiltration of peripheral macrophages.<sup>5</sup> POCD is commonly observed in perioperative care in elderly patients; therefore, a limitation of this study is that they did not use aged mice as an experimental animal model of POCD.<sup>2</sup> Furthermore, elderly individuals more frequently present a degenerative systemic immune system and resident microglial priming in the aging brain, rather than the migration of peripheral macrophages, may have an important role in cognitive deficits after surgery.<sup>6</sup> Normal microglia respond to and degrade A $\beta$  by phagocytosis; however, primed microglia become more proinflammatory and less phagocytic.<sup>7</sup> Current anti-

Correspondence: Jianjun Li  
Department of Anesthesiology, Qilu  
Hospital of Shandong University  
(Qingdao), No.758 Hefei Road, Qingdao,  
People's Republic of China  
Tel +86 532 6685 0629  
Fax +86 532 6685 0629  
Email Ljj9573@163.com

inflammatory and immunosuppressive agents have demonstrated neuroprotection in POCD animal models, but there is no agreement on the efficiency of these treatments in clinical practice.<sup>2</sup> Given that most anti-inflammatory and immunosuppressive agents, such as ibuprofen,<sup>8</sup> can readily penetrate the blood-brain barrier damaged by surgical stress or anesthesia, current studies have not confirmed that these agents exert neuroprotective effects directly by targeting the systemic rather than the central immune response. Given the above, the systemic immune response induced by surgical stress and anesthesia may be the compensatory mechanism that maintains the stability of the immune-microenvironment in CNS. We, therefore, suggest that in POCD, degenerative systemic immunity should be boosted in elderly patients to drive an immune-dependent cascade needed for injured neuron repair.

Adaptive immune cell populations play an important role in restraining the accumulation of A $\beta$  and neuroinflammation.<sup>7</sup> Baruch et al<sup>9</sup> found that disrupting immune tolerance by targeting regulatory T (Treg) cells induced the accumulation of monocyte-derived macrophages and Treg cells at sites of CNS pathology as well as A $\beta$  clearance, contributing to a reversal of cognitive decline. While considering immune checkpoints to be regulatory pathways for maintaining the homeostasis of the systemic immune system, they further showed that immune checkpoint blockade directed against the programmed death-1 (PD-1) pathway, thereby evoking an IFN- $\gamma$ -associated systemic immune response and inducing CNS recruitment of myeloid cell, caused clearance of cerebral A $\beta$ , and improved cognitive performance.<sup>10</sup> In contrast to the continuous or bolus administration of anti-inflammatory and immunosuppressive agents in previous studies, immunotherapy conducted by PD-1 may maintain long-lasting beneficial effects. We, therefore, hypothesize that immunotherapy may be a potential therapeutic strategy for POCD, and PD-1 immune checkpoint blockade may be viable for the prevention and treatment of elderly patients with POCD in the future.

## Acknowledgments

This work was supported by grants from People's Benefit Project of Science and Technology in Qingdao (18-6-1-74-nsh) and the Scientific Research Foundation of Qilu Hospital of Shandong University (Qingdao) (Grant Nos. QDKY2016QN01 and QDKY2016ZD05).

## Disclosure

The authors report no conflicts of interest in this communication.

## References

1. Kotekar N, Shenkar A, Nagaraj R. Postoperative cognitive dysfunction – current preventive strategies. *Clin Interv Aging*. 2018;13:2267–2273. doi:10.2147/CIA.S133896
2. Skvarc DR, Berk M, Byrne LK, et al. Post-operative cognitive dysfunction: an exploration of the inflammatory hypothesis and novel therapies. *Neurosci Biobehav Rev*. 2018;84:116–133. doi:10.1016/j.neubiorev.2017.11.011
3. Leslie M. The post-op brain. *Science*. 2017;356(6341):898–900. doi:10.1126/science.356.6341.898
4. Zhu H, Liu W, Fang H. Inflammation caused by peripheral immune cells across into injured mouse blood brain barrier can worsen post-operative cognitive dysfunction induced by isoflurane. *BMC Cell Biol*. 2018;19(1):23. doi:10.1186/s12860-018-0172-1
5. Terrando N, Eriksson LI, Ryu JK, et al. Resolving postoperative neuroinflammation and cognitive decline. *Ann Neurol*. 2011;70(6):986–995. doi:10.1002/ana.22664
6. Murray C, Sanderson DJ, Barkus C, et al. Systemic inflammation induces acute working memory deficits in the primed brain: relevance for delirium. *Neurobiol Aging*. 2012;33(3):603–616.e3. doi:10.1016/j.neurobiolaging.2010.04.002
7. Marsh SE, Abud EM, Lakatos A, et al. The adaptive immune system restrains Alzheimer's disease pathogenesis by modulating microglial function. *Proc Natl Acad Sci U S A*. 2016;113(9):E1316–E1325. doi:10.1073/pnas.1525466113
8. Huang C, Irwin MG, Wong GTC, Chang RCC. Evidence of the impact of systemic inflammation on neuroinflammation from a non-bacterial endotoxin animal model. *J Neuroinflammation*. 2018;15(1):147. doi:10.1186/s12974-018-1220-7
9. Baruch K, Rosenzweig N, Kertser A, et al. Breaking immune tolerance by targeting Foxp3(+) regulatory T cells mitigates Alzheimer's disease pathology. *Nat Commun*. 2015;6:7967. doi:10.1038/ncomms8967
10. Baruch K, Deczkowska A, Rosenzweig N, et al. PD-1 immune checkpoint blockade reduces pathology and improves memory in mouse models of Alzheimer's disease. *Nat Med*. 2016;22(2):135–137. doi:10.1038/nm.4022

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Interventions in Aging 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Interventions in Aging editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

### Clinical Interventions in Aging

Dovepress

## Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier

Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>