Proportion of Japanese outpatients filling prescriptions for long-term medication regimens

This article was published in the following Dove Press journal:
Patient Preference and Adherence

Kaori Kurata¹
Michi Onuki²
Kazuki Yoshizumi³
Eitaro Taniai³
Akira Dobashi¹

¹Education and Research Institute of Information Science, Tokyo University of Pharmacy and Life Sciences, Hachioji, Tokyo, Japan; ²Insurance Pharmacy Division, Yakuju Co. Ltd, Minato-ku, Tokyo, Japan; ³Information Headquarters, Yakuju Co. Ltd., Yamato-shi, Kanagawa, Japan

Aim: Our study aimed to clarify how long outpatients with chronic diseases such as hypertension continuously fill prescriptions for the same medications as those prescribed initially and how many medications they take over the long term.

Methods: Medication records from April 1, 2016 to March 31, 2017 with total days between initial and final dispensation date ≥ 330 days and total days of medication supplies ≥ 180 days were extracted from an electronic database in a Japanese community pharmacy chain. The continuity of refilling medications over 1 year (ie, medication fill adherence) was measured using the medication possession ratio (MPR).

Results: A total of 34,549 outpatients received long-term medications under the above conditions (4.4% of all patients in the database). Mean age was 66.0 ± 17.4 years; 63.1% were ≥ 65 years. The mean number of medications prescribed per patient was 3.2 ± 2.3 . More than one-fifth of patients (22.6%) were taking ≥ 5 medications. The mean MPR for patients overall was $93.6 \pm 11.2\%$; 87.2% of patients had an MPR $\geq 80\%$ but $< 110\%$. Amlodipine besylate, an antihypertensive, was the most commonly prescribed drug ($n=5,537$ patients).

Conclusion: Outpatients that received long-term medications with no change in prescription had an MPR $> 90\%$ for around 3 medications. It can be reasonably assumed that these patients could receive a longer-term medication supply with the partial fills based on a physician's instruction ("Bunkatu Chozai" in Japanese). This longer-term supply would be similar to a basic prescription refill, but would require a physician's signature allowing for partial refills rather than a new prescription for each refill.

Keywords: outpatient, partial fill, long-term medication, medication possession ratio, chronic disease, refill prescription

Introduction

Outpatients with chronic diseases such as hypertension, diabetes mellitus, dyslipidemia, and hyperuricemia usually take medications over long periods. Prescriptions for these medications generally do not change as long as the patient's condition is stable and/or adverse events do not occur.

In Japan, almost all outpatients receive their medications on the day they are prescribed because, under the national health insurance program, prescriptions expire in 4 days. Furthermore, the Japanese national health system has not adopted a refill (or a repeat) prescription program, and the number of total days of medication supplies per single prescription is lower than that in other countries. For example, according to a Survey of Medical Care Activities in Public Health Insurance,¹ distribution of the number of prescription days for all medications prescribed in 2016 was as follows: ≤ 14 days, 87.6%; ≥ 15 days to ≤ 30 days,

Correspondence: Akira Dobashi
Education and Research Institute of Information Science, Tokyo University of Pharmacy and Life Sciences, 1432-1 Horinouchi, Hachioji, Tokyo 192-0392, Japan
Tel +81 42 676 3082
Fax +81 42 670 7067
Email dobashi@toyaku.ac.jp

8.7%, and ≥ 31 days, 3.7%. Prescription days per one single prescription are generally unlimited to the range that a physician in Japan foresees current medical condition lasting, and the proportion of prescriptions written for ≥ 31 days was very small.

However, outpatients with chronic diseases often take the same medications as those prescribed initially over the long term, so that actual number of days of medication supplies are longer than the prescription days reported in the survey above. Our study aimed to clarify how long they fill prescriptions for the same medications and how many medications they continuously take over 1 year. Based on our findings, we will discuss the possibility that these patients should receive a longer-term medication supply with partial fills by physician's instructions ("Bunkatu Chozai" in Japanese), similar to a basic prescription refill.

Methods

Medication records, including medicines prescribed repeatedly over 1 year, were extracted from an electronic database in a community pharmacy chain (Yakuju Co. Ltd.), consisting of 155 stores in the Japanese Kanto region (eastern half of Japan, including Tokyo). Medication records were collected from April 1, 2016, to March 31, 2017, and were anonymized to protect personal information. Outpatient records that showed total days between initial and final dispensation date ≥ 330 days and total days of medication ≥ 180 days for individual medications were included in this analysis. In addition, sex, age, and the date that medications prescribed for patients were filled in community pharmacies (ie, the dispensation date) were collected. The medicines were identified by individual medicine codes of Japan (referred to as YJ codes), brand names, and generic names.

The medication possession ratio (MPR) was used to assess medication fill adherence.²⁻⁸ The MPR was defined as the total days of medication supplies by prescription divided by the observation period that starts at the first dispensation date and ends at the last dispensation date for individual medications, and expressed as a percentage.⁴ Days' supply beyond the end of the observation period was excluded. When MPR is $>100\%$, it indicates that there may be overlaps in days between taking medications; this may leave medicines not taken regularly. When MPR is $<100\%$, it indicates that there may be gaps in the days between taking medications. As has been reported extensively in the literature describing adherence rates for

refilled prescriptions, medication refill adherence is considered satisfactory if MPR is $\geq 80\%$.^{5,6} Although MPR cannot ensure that patients are taking their medications, it provides an estimate of the highest possible level of medication consumption over a particular time frame.

Medication records of patients with the same prescription ≥ 2 times, each for ≥ 14 days, were collected by individual medications. Among these records, those having an observation period ≥ 330 days and total days of medication supplies ≥ 180 days were considered to be taking long-term medications. These records were extracted from the collected record and summed for each patient. Both mean MPR and the proportion of patients taking medications with an MPR $\geq 80\%$ but $<110\%$ were calculated; these values were compared between males and females, patients ≥ 65 years and <65 years, and between those taking ≥ 5 medications and <5 medications.

The ethics committee of the Social University (the General Incorporated Association, 8-5-26 Akasaka, Minato-ku, Tokyo, Japan) approved this study (approval day: March 16, 2017; approval number: SU-0005). Informed consent for each outpatient corresponding to the medication record stored in the database was not needed because the medication records analyzed in our study were anonymized in a linkable fashion according to the requirement for protection of personal information.

Statistical analysis

Statistical tests were performed using JMP Pro 14 (SAS, 6-10-1 Roppongi, Minato-ku, Tokyo, Japan). Chi-square test (P , 0.001) was used to determine differences in the proportion of patients with an MPR $\geq 80\%$ but $<110\%$ between males and females, between the two age groups (≥ 65 years and <65 years), and between two medication groups (≥ 5 medications and <5 medications).

Results

Medication records for the 1-year survey included 787,451 outpatients. Among these patients, 158,938 (20.2%) had medications prescribed more than twice and with each medication prescribed for ≥ 14 days. Of these patients, 34,549 (4.4% of all patients in the database) received medications for ≥ 330 total days between initial and final dispensation date and ≥ 180 total days of medication supplies. Records for these patients were included in this analysis.

The mean age of patients was 66.0 ± 17.4 years, 49.2% of male, and 21,789 (63.1%) were ≥ 65 years (Table 1).

Table 1 Characteristics of patients receiving long-term and repeat medications and those taking ≥ 5 medicines versus < 5 medicines

	Total 34,549 (100%)	<5 medica- tions 26,720 (77.3%)	≥ 5 medica- tions 7,829 (22.7%)
Age (SD), years	66.0 (17.4)	64.6 (17.8)	71.0 (15.0)
<65 years	12,760 (36.9%)	10,638 (39.8%)	2,122 (27.1%)
≥ 65 years	21,789 (63.1%)	16,082 (60.2%)	5,707 (72.9%)
Male	16,980 (49.2%)	12,973 (48.6%)	4,007 (51.2%)
Prescription days (SD)	33.2 (15.6)	33.8 (15.9)	31.1 (14.4)
<10 days	307 (0.9%)	238 (0.9%)	69 (0.9%)
10–40 days	27,279 (79.0%)	17,816 (66.7%)	6,460 (82.5%)
≥ 40 days	6,963 (20.2%)	8,666 (32.4%)	1,300 (16.6%)
Medications (SD)	3.2 (2.3)	2.2 (1.1)	6.7 (1.9)
Total days of med- ication supplies (SD)	320.5 (39.1)	317.2 (40.9)	332.0 (29.1)
Medication pos- session ratio (MPR) (SD), %	93.6 (11.2)	92.6 (11.8)	96.8 (8.2)
<80%	3,936 (11.4%)	3,633 (13.6%)	303 (3.9%)
80–110%	30,142 (87.2%)	22,745 (85.1%)	7,397 (94.5%)
$\geq 110\%$	471 (1.4%)	342 (1.3%)	129 (1.6%)

Figure 1 Frequency distribution of mean prescription days per one single patient for long-term medications. Minimum value of mean prescription days was 4.5-day ($n=1$) and maximum value 180-day ($n=2$).

had the prescription days ≥ 40 days, and only 0.9% had the prescription days < 10 days. The frequency distribution of the mean prescription days ranged from 4.5 to 180 days (Figure 1).

As can be seen in Figure 1, the proportion of the prescription days ≥ 30 days was considerably higher than that reported by a Survey of Medical Care Activities in Public Health Insurance (3.73% for ≥ 31 days).¹ Around one-half (49.4%) of patients in this study received long-term and repeat medications with the prescription days ≥ 30 days.

Prescription days and total days of medication supplies

The mean prescription days per single patient was 33.2 ± 15.6 days (Table 1). The mean total days of medication supplies was 320.5 ± 39.1 days within a mean observation period of 342.6 ± 8.1 days. A total of 80.5% received medications for ≥ 300 days. Thus, outpatients generally received, on average, medications around 10 times with a 33-day supply of the prescription from a community pharmacy per year.

A total of 79.0% of patients had a mean number of prescription days ≥ 10 days but < 40 days. A total of 20.2%

Number of medications

The number of medications prescribed per patient ranged from 1 to 22 for long-term and repeat medications (Figure 2). The mean number of medications prescribed was 3.2 ± 2.3 (Table 1). More than one-fifth of patients (22.7%) were taking ≥ 5 medications. There was a significant difference between males and females in those taking ≥ 5 medications (male, 51.2%; female, 48.8%; $p < 0.0001$).

Among all patients, 7,829 (51.2% male) were taking ≥ 5 medications and 26,720 patients were taking < 5 medications (48.6% male). In addition, patients taking ≥ 5

Figure 2 Number of long-term medications prescribed per patient.

medications were significantly more likely to be elderly (≥ 65 years vs < 65 years, 72.9% vs 60.2%, respectively; $P < 0.0001$).

Medication possession ratio (MPR)

The mean MPR for all patients was $93.6 \pm 11.2\%$ (Table 1). A total of 87.2% of patients had an MPR $\geq 80\%$ but $< 110\%$. A total of 11.4% had an MPR $< 80\%$, and only 1.4% had an MPR $\geq 110\%$. The frequency distribution of MPR ranged from 49.5% to 302.7% (Figure 3).

Patients taking ≥ 5 medications were significantly more likely to have an MPR $\geq 80\%$ but $< 110\%$ compared with those taking < 5 medications (94.5% vs 85.1%, respectively; $P < 0.0001$) (Table 1).

Representative medications prescribed for the long term

A total of 881 medications were prescribed repeatedly over 1 year. Table 2 shows the top 20 medications along with their anatomical therapeutic chemical (ATC) classification. These 20 medications had a mean MPR ranging from 92.7% to 96.4%, a mean of prescription days per patient ranging from 25.2 to 38.1, and a total days' supply (accumulated from all values except the last prescription day) ranging from 317.4 to 331.3.

Figure 3 Frequency distribution of medication possession ratio (MPR) for long-term medications. Minimum value of MPR was 49.5% ($n=1$) and maximum value 302.7% ($n=1$).

Amlodipine besylate, an antihypertensive, was prescribed for the highest number of patients. This drug showed a mean MPR of $95.5\% \pm 9.2\%$, a mean of 33.8 ± 14.6 prescription days per patient, and a mean of 326.9 ± 32.4 total medication days.

Among the 20 medications, eleven medications were included in four therapeutic categories related to chronic diseases shown in Table 3.

MPR of medications based on therapeutic category

Hypertensive agents were the most commonly prescribed drugs, followed by dyslipidemic agents, antidiabetic agents (limited to oral and non-insulin agents), and hyperuricemia agents (Table 3). The mean MPR for all four agents was $\geq 90\%$ and the proportion of patients with an MPR range of 80–110% was also $\geq 90\%$.

Discussion

Our study found that around 4% of Japanese outpatients generally received 3 medications with a 33-day filling period over 10 pharmacy visits during 1-year survey. The medication fill adherence was, on average, estimated at over 90%. Of these patients, those taking ≥ 5 medications

Table 2 Number of days and medication possession ratio (MPR) for top 20 medicines prescribed overall

Generic drug name	ATC classification ^a	n	Average prescription days per patient (SD)	Total days with prescription supplies (SD)	MPR (SD), %
Amlodipine besylate	C08	5,537	33.8 (14.6)	326.9 (32.4)	95.5 (9.2)
Rosuvastatin calcium	C10	2,436	34.5 (14.5)	322.3 (35.0)	94.2 (10.0)
Aspirin	B01	2,258	34.9 (16.9)	329.4 (29.7)	96.1 (8.4)
Magnesium oxide	A02, A06, A12	2,172	30.3 (15.8)	325.1 (42.2)	94.6 (12.0)
Atorvastatin calcium hydrate	C10	2,044	34.3 (14.8)	325.9 (31.9)	95.2 (9.1)
Lansoprazole	A02	1,852	33.2 (15.2)	328.8 (32.0)	96.0 (9.0)
Etizolam	N05	1,784	27.0 (8.9)	331.3 (60.7)	96.4 (17.3)
Nifedipine	C08	1,657	33.9 (14.9)	328.6 (32.2)	96.0 (9.2)
Pitavastatin calcium hydrate	C10	1,586	34.4 (14.1)	326.3 (31.0)	95.3 (8.9)
Candesartan cilexetil	C09	1,464	35.0 (15.5)	327.0 (31.5)	95.5 (9.0)
Telmisartan	C09	1,442	34.4 (14)	329.4 (29.2)	96.2 (8.2)
Brotizolam	N05	1,345	25.2 (6.0)	324.0 (52.7)	94.5 (15.1)
Allopurinol	M04	1,327	34.7 (16.0)	324.4 (33.6)	94.8 (9.6)
Zolpidem tartrate	N05	1,305	25.6 (5.7)	317.4 (52.0)	92.6 (14.9)
Febuxostat	M04	1,259	34.8 (15.2)	324.4 (33.6)	94.8 (9.5)
Esomeprazole magnesium hydrate	A02	1,230	34.3 (15.5)	324.6 (38.5)	94.8 (11.1)
Sitagliptin phosphate hydrate	A10	1,205	36.7(15.5)	328.9 (27.3)	96.2 (7.7)
Rebamipide	A02	1,195	30.4 (14.3)	318.0 (46.0)	92.7 (13.3)
Famotidine	A02	1,188	32.4 (16.3)	329.1 (31.4)	95.9 (8.9)
Metformin hydrochloride	A10	1,101	38.1 (15.0)	323.7 (33.4)	94.7 (9.7)

Note: ^aACT classification of being bold indicates that these medications are included in four therapeutic categories related to chronic diseases shown in Table 3.

Abbreviation: ATC, anatomical therapeutic chemical.

were significantly more likely to be ≥ 65 years higher and in the proportion with an MPR $\geq 80\%$ but $< 110\%$ than those taking < 5 medications.

MPR estimates the proportion of times that medications are filled over a particular observation period and the highest possible level of medication consumption.²⁻⁴ Based on the MPR values in our study, patients received medications prescribed by their physicians regularly. These findings are particularly important for patients undergoing long-term medical treatment and indicate that patients are making regular physician visits, as medications are not automatically refilled in Japan.

In a medication refill adherence study using a pharmacy claims database among patients with chronic diseases in the United State,⁷ MPR $\geq 80\%$ was detected for 72.3% of patients with hypertension and 65.4% with type 2 diabetes mellitus.

In our study, the MPR was $\geq 90\%$ for the topmost frequently prescribed medications, which is higher than that shown in the previous study. An adherence study of patients with chronic diseases receiving Medicaid in the United States found a mean MPR for a 30-day refill of

medications of 71% in patients with hypertension and 69% for patients with type 2 diabetes mellitus, and a mean MPR for a 90-day refill of 83% and 80%, respectively.⁸ The 33-day refill period in our study is similar to the 30-day period in the Medicaid study, although MPR findings were higher in our study.

The one way that prescriptions in Japan can be filled partially (“Bunkatu Chozai” in Japanese) is when it is difficult to preserve the medication for a long period; in these cases, medications are partially filled and the remaining portion is filled at a later time. Another way for partial filling of prescriptions is according to the physicians’ indications, which is similar to refilling prescriptions based on a specified number of refills before a certain expiration date. Thus, patients can obtain more of the same medications from pharmacists without seeing the prescribers within a total of the prescription days and 4 days of the prescription expiration date.⁹ This novel prescription format introduced in 2018 allows physicians to prescribe a partial days’ supply, and the number of partial fills of prescriptions in the remarks column of prescription sheets.

Table 3 Mean medication possession ratio (MPR) for four therapeutic categories

Therapeutic classification	ATC classification	Patients taking medications by therapeutic classification					
		n	Age (SD), years	Male, %	MPR (SD), %	n	
						MPR range of 80–110%	MPR range of ≥110%
Hypertension agents	C02-C09	14,824 (42.9%)	71.7 (11.8)	51.8	95.9 (8.9)	20,396 (92.9%)	200 (0.9%)
Dyslipidemia agents	C10	8,838 (25.6%)	70.2 (11.4)	45.2	95.1 (9.3)	8,507 (91.9%)	65 (0.7%)
Antidiabetic agents	A10	4,252 (12.3%)	68.5 (11.5)	60.1	95.6 (8.6)	6,841 (92.9%)	53 (0.8%)
Hyperuricemia agents	M04	2,883 (8.3%)	67.8 (13.4)	85.1	94.5 (9.8)	2,744 (90.7%)	21 (0.7%)

Abbreviation: ATC, anatomical therapeutic chemical.

The process for partial filling of a prescription in Japan differs substantially from that in the United States, where partial filling of a prescription is permitted when the pharmacist is unable to supply the prescribed amount or the prescriber requests less than the full quantity called for in a prescription.

Our study indicated that outpatients who repeatedly received the same medications as prescribed initially and who had a medication supply ≥ 180 days during this 1-year survey showed high adherence to both visiting their physicians to obtain a prescription. It is reasonable to assume that outpatients should be able to receive a long-term supply of medication (eg, 90 days), with an order for partial filling on the part of the physician, and without the need for a new prescription for each refill.

When this partial filling of prescriptions is expanded and extended, community pharmacists should regularly consult with patients about taking their medications as prescribed, not only to help them understand the need to continue treatment, but also to enhance their medication adherence. These consultations may also help to reduce polypharmacy, which is common in people aged ≥ 65 years. These elderly patients may have a greater risk for polypharmacy, that is, inappropriate use of multiple drugs, and should be considered in efforts to reduce unnecessary polypharmacy.¹⁰

Limitations

First, our study used MPR to estimate medication fill adherence. Although systematic reviews have shown that MPR is a reliable measure of adherence,² our study cannot report actual adherence to taking medications. However, we can discuss continuity and regularity of Japanese outpatients in terms of behavior regarding

receiving their medicines from a community pharmacy. Regular receipt of medications may mean that medications are being used. However, this could not be proven in our study.

Second, there are pharmacies within community pharmacy chains and individual pharmacies in Japan. Our study was based on medication records stored in a community pharmacy chain. This chain is controlled by a management headquarters, which may influence which medications are used. However, as medications are filled as prescribed by physicians in terms of both brand and generic medicines, the headquarters does not control the medication used except when a generic medicine is prescribed by the physician. Thus, all branches within Yakuju are similar to individual pharmacies, limiting the possibility for differences between these pharmacies.

Conclusion

Our study found that Japanese outpatients who repeatedly received the same medications as prescribed initially and who had a medication supply ≥ 180 days over 1 year were around 4% of all and showed high medication fill adherence over 90%. It is reasonably assumed that these patients could receive a longer-term medication supply with the novel partial fills based on a physician's instruction ("Bunkatu Chozai" in Japanese). This prescription format introduced in 2018 would be similar to a basic prescription refill and allows physicians to prescribe a partial days' supply, and the number of partial fills of prescriptions.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Survey of Medical Care Activities in Public Health Insurance 2016, Use of drugs, Table 6. [Publication day; July 25, 2017]. Available from: <https://www.e-stat.go.jp/stat-search/files?page=1&layout=data list&toukei=00450048&tstat=000001029602&cycle=7&tclass1=000001103215&tclass2=000001103256&tclass3=000001103259&second2=1>. Accessed July 27, 2018. Japanese.
2. Hess LM, Raebel MA, Conner DA, Malone DC. Measurement of adherence in pharmacy administrative databases: a proposal for standard definitions and preferred measures. *Ann Pharmacother*. 2006;40(7–8):1280–1288. doi:10.1345/aph.1H018
3. Sattler EL, Lee JS, Perri M. 3rd, Medication (re)fill adherence measures derived from pharmacy claims data in older Americans: a review of the literature. *Drugs Aging*. 2013;30(6):383–399. doi:10.1007/s40266-013-0074-z
4. Arnet I, Abraham I, Messerli M, Hersberger KE. A method for calculating adherence to polypharmacy from dispensing data records. *Int J Clin Pharm*. 2014;36:192–201. doi:10.1007/s11096-013-9891-8
5. Feehan M, Ranker L, Durante R, et al. Adherence to controller asthma medications: 6-month prevalence across a US community pharmacy chain. *J Clin Pharm Ther*. 2015;40(5):590–593. doi:10.1111/jcpt.12316
6. Feehan M, Munger MA, Cooper DK, et al. Adherence to glaucoma medications over 12 months in two US community pharmacy chains. *J Clin Med*. 2016;5(9):79–86. doi:10.3390/jcm5090079
7. Briesacher BA, Andrade SE, Fouayzi H, Chan KA. Comparison of drug adherence rates among patients with seven different medical conditions. *Pharmacotherapy*. 2008;28(4):437–443. doi:10.1592/phco.28.4.437
8. Taitel M, Fensterheim L, Kirkham H, Sekula R, Duncan I. Medication days' supply, adherence, wastage, and cost among chronic patients in Medicaid. *Medicare Medicaid Res Rev*. 2012;2(3):E1–E13. doi:10.5600/mmrr.002.03.A04
9. Ministry of Health, Labour and Welfare, Minister's Secretariat, Statistics and Information Department, Social Statistics Division. *Rules for health insurance-covered dispensing pharmacies and pharmacists*. [-Publication day; April 1, 2018]. Available from: http://elaws.e-gov.go.jp/search/elawsSearch/elaws_search/lsg0500/detail?lawId=332M50000100016&openerCode=1. Accessed August 23, 2018. Japanese.
10. Golchin N, Frank SH, Vince A, Isham L, Meropol SB. Polypharmacy in the elderly. *J Res Pharm Pract*. 2015;4(2):85–88. doi:10.4103/2279-042X.155755

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease

states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>