World Mental Health Day 2018, but how aware are medical students?

This article was published in the following Dove Press journal:
Advances in Medical Education and Practice

Godwin Tong

College of Medical and Dental Sciences,
University of Birmingham,
Birmingham, UK

Abstract: With the recent global interest in mental health, there has been an increase in the number of people coming forward to discuss the once frowned upon illness. It is true that some progress has been made in terms of establishing an equal footing within the health and social care landscape with the help of various campaigns. However, there is still much work to be done with regards to removing the stigma of mental health within the healthcare sector itself. It is no surprise that mental illness statistics are significantly higher amongst medical practitioners due to the high-stress nature of their jobs, yet it is believed to be under reported. Interestingly, there is emerging evidence that physician burn out stems early in their careers whilst in medical school, suggesting areas that need improvement in medical education with regards to how medical students view and handle mental illness. As such, two issues have been identified and discussed here - the stigmatisation of mental illness whilst in medical school and the underemphasis of mental illness as compared to other physical health problems in medical education. It is hoped that discussing these issues can help stimulate subsequent positive changes.

Keywords: mental health awareness, stigma, medical education, medical student attitudes

The 10th of October marks the World Federation of Mental Health's 22nd mental health awareness day. This day was created with a three-pronged approach in mind – highlighting the prevalence of mental illness amongst us, removing the stigma associated with mental illness and celebrating the recovery of those who were once victims of mental illness.¹ Depression is the second most common cause of disability worldwide, creating an increasing socio-economic burden.² With more emphasis being placed on mental health screening, it is no surprise that the prevalence will only increase with time. Indeed, it is currently estimated that one in six people in the past week have experienced a common mental health problem.³ These are just numbers that do not consider the number of individuals who have yet to come forth due to fear of stigmatization. Medical professionals without a doubt have a higher chance of developing mental illness due to a myriad of factors such as chronic stress, large workloads and compassion fatigue. Interestingly, there is a growing body of evidence in the past 5 years suggesting that physician burnout may have its origins in medical school.⁴ With national statistics showing more than 400 doctors in the UK taking their own lives between 2011 and 2015,⁵ perhaps one should look toward the root of the problem – medical education. Many questions remain but importantly, how are the medical students coping? How do medical students feel about mental illness?

Correspondence: Godwin Tong
College of Medical and Dental Sciences,
University of Birmingham, Birmingham
B15 2TT, UK
Email GXT545@bham.ac.uk

Mental health amongst medical students has been increasingly discussed in recent years. While not considered a mental illness, burnout can be considered a mental health issue. Recent studies suggest that students commonly suffer from burnout. One systematic review reported data from nine studies examining US medical students, with burnout prevalence ranging between 45% and 71%. Furthermore, symptoms of depression and suicidal ideologies were more rampant during medical school as compared to practicing physicians.⁴ The use of dysfunctional coping strategies, such as the use of alcohol and illicit drugs, has been significantly reported amongst UK and US medical students as well, along with high depression scores.^{6,7} This suggests that medical students have low levels of help-seeking when it comes to personal psychiatric problems, often only when a crisis arises. One reason for this is the fear of stigmatization. In an American study, it was found that depressed medical students more frequently endorsed several depression stigma attitudes than nondepressed students. This meant that compared with students with low self-identified depression, students with high depression scores more frequently agreed that the opinions of depressed medical students would be less respected, that the coping skills of depressed medical students would be viewed as less adequate, that they would be viewed as less able to handle their responsibilities by faculty members and that telling a counselor about depression would be risky. This is dangerous as it reflects how inaccurately distorted perceptions are amongst depressed students. Furthermore, students with high scores would be less likely to seek treatment if depressed than would students with low scores. These feelings were echoed by their UK counterparts where avoidance of appropriate help-seeking behavior is observed to start early in medical school for it is seen as a form of weakness.⁸ This distortion in perception is evident amongst doctors as well. In a survey of 6,000 American physicians, a whopping 40% of them revealed that they would be reluctant to seek treatment for mental health problems in fear of facing problems with obtaining their medical licensure.⁹ This is due to one-third of states having a mental health section on their medical licensure application questions. It is rather unexpected to see the medical community reinforce this stigma by their own attitudes, creating a vicious cycle by deterring doctors seeking treatment for themselves.

Underlying this problem of stigma amongst medical students is another issue – indifference. It is perhaps ironic

that despite being future health care professionals, medical students might be lacking empathy toward individuals with mental illness. Depersonalization and emotional exhaustion are core symptoms of burnout and thus could potentially explain the progressive lack of empathy as doctors progress through their training. However, this does not seem to be the case. A study of 760 UK medical students examining their attitudes to mental illness, as compared with attitudes toward other medical illness, showed that students showed the highest regard for patients with pneumonia as compared to other conditions such as abdominal pain and mental illness. What was shocking was that these attitudes were worse in the more experienced senior medical students.¹⁰ Another qualitative study in China, although with a small sample size, recorded the attitudes of fresh medical graduates at a local university.¹¹ They showed that of 241 coded responses concerning negative attitudes toward mental illness, 51% reflected fear of violent behavior and 23% reflected shame from interpersonal associations. From these studies, we can see that there are differing attitudes when comparing mental illness to other types of pathologies. It is easy to develop empathy towards patients with visible pathologies within a hospital, but if you replace what we can see with an invisible illness, do we still feel the same for these individuals?

Such low levels of social acceptance of individuals with mental illness should not be expected from future doctors. Current research on interventions to fight mental health stigma has been equivocal. Interventions that involve social contact are more effective in reducing stigmatizing attitudes than brief education sessions but only for the short term. There is yet concrete evidence for the medium-long term. Perhaps future studies should look to further understand the attitudes of medical students or even before they enter university. By tackling the root of the problem, we can then shape initiatives that promote psychiatric education, combat stereotypes and promote more positive attitudes amongst our future doctors. Hence, I ask all clinicians and medical educators, on this Mental Health Awareness Day, when you see your medical students under your charge, do not follow through with the mental health curriculum by showing plain statistics and clinical signs. Rather, take it to the next level by placing an emphasis on empathy toward individuals with mental illness. Perhaps medical students can learn to empathize more by emulating their role models. As for the medical students, I implore you to look inward before you speak to

a patient with mental illness – Would you treat the patient any differently if the patient was family? Would you still take the same elaborate history as with a complex cardiovascular pathology? Would you offer the same amount of compassion as with a patient with terminal stage breast cancer?

Acknowledgment

The author would like to thank the University of Birmingham Library Services division for supporting this work and Mr Ethan Khoo and Miss Inez Dale-Jones for their thoughtful insights.

Disclosure

The author reports no conflicts of interest in this work.

References

1. World Federation for Mental Health [Internet]. Available from: <https://wfmh.global/about-us/>. Accessed October 08, 2018.
2. Whiteford HA, Degenhardt L, Rehm J, et al. Global burden of disease attributable to mental and substance use disorders: findings from the Global Burden of Disease Study 2010. *Lancet*. 2013;382:1575–1586. doi:10.1016/S0140-6736(13)61611-6
3. McManus S, Bebbington P, Jenkins R, Brugha T. Mental health and Wellbeing in England: adult psychiatric morbidity survey 2014. *Leeds NHS Digit*. 2016. Available from: https://files.digital.nhs.uk/pdf/q/3/mental_health_and_wellbeing_in_england_full_report.pdf
4. Ishak W, Nikraves R, Lederer S, Perry R, Ogunyemi D, Bernstein C. Burnout in medical students: a systematic review. *Clin Teach*. 2013. doi:10.1111/tct.12014
5. Suicide by occupation, England: 2011 to 2015 [Internet]. *Off Natl Stat*; 2017. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/suicidebyoccupation/england2011to2015#references>
6. Vaysse B, Gignon M, Zerkly S, Ganry O. [Alcohol, tobacco, cannabis, anxiety and depression among second-year medical students. Identify in order to act]. *Sante Publique*. 2014;26(5):613–620.
7. Ayala EE, Roseman D, Winseman JS, Mason HRC. Prevalence, perceptions, and consequences of substance use in medical students. *Med Educ Online*. 2017;22:1392824. doi:10.1080/10872981.2017.1392824
8. Chew-Graham CA, Rogers A, Yassin N. “I wouldn’t want it on my CV or their records”: medical students’ experiences of help-seeking for mental health problems. *Med Educ*. 2003;37:873–880. doi:10.1046/j.1365-2923.2003.01627.x
9. Dyrbye LN, West CP, Satele D, et al. Burnout among U.S. medical students, residents, and early career physicians relative to the general U.S. population. *Acad Med*. 2014;89:443–451. doi:10.1097/ACM.0000000000000134
10. Korszun A, Dinos S, Ahmed K, Bhui K. Medical student attitudes about mental illness: does medical-school education reduce stigma? *Acad Psychiatry*. 2012;36:197. doi:10.1176/appi.ap.10110159
11. Luo A, He H, Mohamed S, Rosenheck R. Medical student attitudes towards people with mental illness in China: a qualitative study. *Cult Med Psychiatry*. 2018;42:535–551. doi:10.1007/s11013-018-9568-9

Advances in Medical Education and Practice

Dovepress

Publish your work in this journal

Advances in Medical Education and Practice is an international, peer-reviewed, open access journal that aims to present and publish research on Medical Education covering medical, dental, nursing and allied health care professional education. The journal covers undergraduate education, postgraduate training and continuing medical education

including emerging trends and innovative models linking education, research, and health care services. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/advances-in-medical-education-and-practice-journal>