The obstacles to diabetes self-management education and support from healthcare professionals' perspectives: a nationwide survey

This article was published in the following Dove Press journal:
Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Lukana Preechasuk¹
Porn Sriussadaporn¹
Supawadee Likitmaskul^{1,2}

¹Siriraj Diabetes Center of Excellence,
Faculty of Medicine Siriraj Hospital,
Mahidol University, Bangkok, Thailand;

²Division of Endocrinology and
Metabolism, Department of Pediatrics,
Faculty of Medicine Siriraj Hospital,
Mahidol University, Bangkok, Thailand

Purpose: This study identified the obstacles to diabetes self-management education and support (DSMES) from healthcare professionals' perspectives in Thailand.

Patients and methods: A cross-sectional survey was conducted at 613 hospitals throughout Thailand from December 2016 to March 2017. A self-completion questionnaire on the characteristics of, and obstacles to, diabetes education was sent to administrators, doctors, and diabetes educators (DEs). The views of DEs at public and private hospitals on the characteristics of diabetes education were compared. The obstructing factors reported by administrators, doctors, and DEs were also analyzed.

Results: The overall response was 76.6%; the response rates of DEs, doctors, and administrators were 85.6%, 58.9%, and 46.5%, respectively. Although the respondents reported that 75% of patients received diabetes education, only 30% of the DEs believed it was successful. An individual-patient education approach was used by 95.1% of public and 81.4% of private hospitals. The median durations of the individual education sessions were 15 (IQR 10, 28) and 30 (IQR 15, 30) minutes for public and private hospitals, respectively. The DE workload at public hospitals was 3 times heavier than at private hospitals (60 [IQR 30, 140] vs 20 [IQR 10, 33] patients per week; $p < 0.001$). Obstacles to diabetes education were a lack of time due to other duties, a lack of skills in assisting patients with behavior change, inadequate DE numbers, patient disinterest in diabetes education, and patient reluctance to change unhealthy behaviors.

Conclusions: High workloads, unclear DE roles, and a lack of DE skills to support patients with behavior change are the primary obstacles to diabetes education.

Keywords: diabetes education, diabetes self-management education, obstacle

Introduction

The 5th Thai National Health Examination Survey reported an increased prevalence of diabetes in people aged ≥ 15 years, with a rise from 6.9% in 2009 to 8.9% in 2014.¹ Data from the Medical Research Network of the Consortium of the Thai Medical Schools (MedResNet) indicated that there was a low level of glycemic control in Thailand, with only 36.3% of patients with type 2 diabetes having an HbA1c below 7% in 2015. Effective diabetes management support from health care providers and appropriate diabetes self-management by patients are fundamental to achieving diabetes goals. Diabetes self-management education (DSME) is the process of facilitating the development of the knowledge, skills, and abilities necessary for diabetes self-care, whereas diabetes self-management support (DSMS) refers to the activities that assist patients with diabetes to implement and

Correspondence: Supawadee Likitmaskul
Siriraj Diabetes Center of Excellence,
Faculty of Medicine Siriraj Hospital,
Mahidol University, 2 Wanglang Road,
Bangkok Noi, Bangkok 10700, Thailand
Tel +662 419 9568
Fax +662 419 9569
Email supawadee.lik@mahidol.ac.th

sustain the behaviors needed to manage diabetes on an ongoing basis.² Diabetes self-management education and support (DSMES) improves HbA1c,³ increases diabetes-complication screening,⁴ improves quality of life,² and is cost-effective in the long term.⁵ It has been recommended that all patients with diabetes receive DSMES, and that adequate resources are made available in local communities to support DSMES services.² In many developed countries, DSMES is usually provided by certified diabetes educators (CDEs),^{6–8} by contrast, there were no CDEs in Thailand either before or during the present survey. The majority of health professionals providing diabetes education to patients in Thailand attended short-course training in diabetes care for a few days at university hospitals, the Thai Association of Diabetes Educators, or the Diabetes Association of Thailand. Moreover, the costs of DSMES services in Thailand are not reimbursed by either the government or private organizations. All of these factors represent key differences between the approach to diabetes education in Thailand and other countries. Therefore, this study aimed to describe the characteristics of diabetes education and identify the perceptions of Thai health professionals regarding the obstacles to diabetes education.

Materials and methods

Questionnaire

A self-completion questionnaire (Supplementary 1) was created by an adult endocrinologist, a pediatric endocrinologist, and an experienced nurse working at Siriraj Diabetes Center of Excellence, Faculty of Medicine Siriraj Hospital. It comprised 7 questions on the characteristics of diabetes education, 4 on the workload associated with diabetes education, and a list of commonly perceived obstacles to diabetes education. This study listed the obstacles to diabetes education that had been identified in the national standards for DSMES in the USA⁹ and by a previous study;¹⁰ however, those obstacles were first modified to suit the diabetes education situation in Thailand. The questionnaire was validated by 4 lecturers from the Division of Endocrinology and Metabolism, Department of Medicine and Division of Endocrinology and Metabolism, Department of Pediatrics, Faculty of Medicine Siriraj Hospital, Mahidol University, and 1 lecturer from the Department of Medical Nursing, Faculty of Nursing, Mahidol University. Two of the four validators also worked as administrators, and three were expert in diabetes education for adult and pediatric patients. Using the averaging calculation method, the scale-level content validity index was 0.99.¹¹

Study design and data collection

A cross-sectional survey was conducted at 613 hospitals (508 public and 105 private) across Thailand from December 2016 to March 2017. It was conducted in every region in Thailand (68 in the north, 195 in the northeast, 169 in the central region, 35 in the west, 44 in the east, and 102 in the south) and at different-sized hospitals (277 public and 18 private hospitals with <60 beds; 141 public and 42 private hospitals with 60–119 beds; and 90 public and 45 private hospitals with ≥120 beds). Participant information sheets and self-completion questionnaires were sent to 3 groups: administrators, diabetes clinic doctors, and health professionals responsible for providing diabetes education (this third group is termed “diabetes educators” [DEs] in this paper). The survey collected data from all participants on the perceived obstacles to diabetes education, but data from only the DEs on the characteristics of diabetes education. Introductory letters were also sent to the administrators to inform them of the study’s purpose and to request their permission to conduct it at their institution. Reminder postcards were sent to the 3 groups of health professionals one month after the questionnaire had been sent. The questionnaires were remailed to hospitals that had not replied to the first approach within 3 months.

Sample size calculation and statistical analysis

The sample size was calculated using the Taro Yamane formula;¹² it was based on a population size (N) of 1,300 hospitals (the estimated number of hospitals in Thailand) and a margin of error of 0.05. The requisite sample size was calculated to be at least 306 hospitals. However, given that the response rate to mail surveys of medical professionals has been established to be around 50%,¹³ the information sheets and questionnaires were dispatched to 613 hospitals.

The statistical analyses were performed using IBM SPSS Statistics for Windows, version 18.0 (IBM Corp., Armonk, NY, USA). Continuous data were reported as means (SD) or medians (IQR), and categorical data as numbers or percentages. The continuous data were analyzed using the independent *t*-test, Mann–Whitney *U* test, one-way analysis of variance (ANOVA), or Kruskal–Wallis test, as appropriate, and the categorical data were evaluated with the Chi-squared test. A *p*-value of <0.05 was considered statistically significant.

Human research ethics

The Institutional Review Board of the Faculty of Medicine Siriraj Hospital, approved this study and waived the need for written informed consent, given the questionnaire-based nature of the study.

Results

Response rate

The overall response was 76.6% (470 hospitals). The response from the public hospitals was 81.5% (414 hospitals) and from the private hospitals was 53.3% (56 hospitals). The response rates from <60-bed, 60–120-bed, and ≥120-bed hospitals were 72.2% (213 hospitals), 75.9% (139 hospitals) and 87.4% (118 hospitals), respectively. The response rates of the administrators, doctors, and DEs were 46.5%, 58.9%, and 85.6%, respectively. The primary occupations of the DEs responding to the questionnaire were outpatient department nurse (85.2%), nutritionist (9.6%), health educator (3.1%), inpatient nurse (1.8%), pharmacist (0.2%), and other (0.1%). Sixty-seven percent of the respondents were female, and the mean ± SD age of the respondents was 41±9.9 years. In some hospitals, there was more than one answer from the same group of respondents. The authors analyzed all data from all of the respondents.

DE-reported characteristics of diabetes education in public and private hospitals

The DEs at both the public and private hospitals reported that nearly 75% of patients with diabetes received diabetes education, but that the larger hospitals tended to have a lower percentage of patients receiving diabetes education (Table 1). Diabetes education was provided to patients with newly diagnosed diabetes, uncontrolled diabetes, diabetes complications, and other individual problems. Of the hospital types, the private hospitals tended to have a lower glycemic control threshold for the provision of diabetes education, and they tended to prefer individual to group diabetes education classes. Although the diabetes education sessions provided by the public hospitals were shorter in duration, the public hospitals conducted more sessions per year than the private hospitals. Interestingly, only 30% of the DEs at the public and private hospitals reported that the diabetes education was successful. In contrast, the large majority (nearly 70%) stated that the results of the diabetes education programs were either uncertain or no evaluations were performed. However, because the success of diabetes education was assessed by DEs from each hospital, the

assessment process and definitions of diabetes educational outcomes might have differed between respondents.

Workload of diabetes care and diabetes education

Ninety-three percent of the hospitals reported that they had DEs. Their occupations were wide-ranging: outpatient department nurses (at 89.7% of the hospitals), nutritionists (at 61%), pharmacists (at 54.3%), doctors (at 44.4%), inpatient nurses (at 34.1%), physical therapists (at 16.6%), health educators (at 15%), Thai traditional-medicine staff (at 7.2%), and dentists (at 3%). Although the public hospitals reported having more patients with diabetes than private hospitals, the number of DEs did not differ between the types of hospital. The DEs in the public hospitals provided diabetes education to a greater median number of patients per week than the DEs in private hospitals (60 [IQR 30, 140] vs 20 [IQR 10, 33]; $p<0.001$), as shown in Figure 1. As the response rate of the DEs located at <60-bed private hospitals was very low ($n=2$), a statistical analysis was not performed for that subgroup.

Self-reported factors perceived as obstacles to diabetes education

The three most commonly perceived obstacles to diabetes education reported by staff at both public and private hospitals were: (1) a reluctance by patients to change their unhealthy behaviors, (2) a lack of time for DEs to deliver diabetes education due to other duties, and (3) patient disinterest in diabetes education. In the case of public hospitals, DEs reported higher percentages for nearly all of the queried obstructing factors than the administrators and doctors (Figure 2). A number of additional problems were reported by DEs at public hospitals, such as a lack of dietitians, communication problems with ethnic minorities, communication problems with older people, and a lack of evaluation after diabetes education programs.

Demand for diabetes educators

A need for more DEs was reported by 74.3% of the participating public hospitals and 62.3% of the private hospitals. The median number of additional DEs required by the public and private hospitals was 2 (IQR 2, 4) positions per hospital.

Discussion

This is the first nationwide survey of diabetes education in Thailand. The overall response was 76.6%. In total, around 75% of patients were reported to receive diabetes education.

Table 1 Characteristics of diabetes education programs in public and private hospitals, as reported by DEs

Characteristics	Public hospitals	Private hospitals
Patients receiving diabetes education, median (IQR), %		
<60-bed hospital	80 (50, 80)	75 (63, 88)
60–119-bed hospital	80 (60, 90)	80 (65, 85)
≥120-bed hospital	54 (38, 80)	70 (45, 85)
Total	75 (50, 80)	73 (58, 83)
Criteria to provide diabetes education		
Patients with newly diagnosed diabetes, n (%)	455 (97.6)	50 (87.7)
Patients with uncontrolled diabetes, n (%)	456 (97.9)	49 (83.1)
Definition of uncontrolled diabetes, median (IQR)		
Fasting plasma glucose (mg/dL)	182 (180, 200)	150 (130, 180)
Fasting plasma glucose (mmol/L)	10.1 (9.99, 11.1)	8.33 (7.22, 9.99)
HbA1c (%)	8 (7, 8)	7 (7, 8)
HbA1c (mmol/mol)	63.9 (53, 63.9)	53 (53, 63.9)
Patients with diabetes complications, n (%)	366 (78.5)	48 (81.4)
Method of diabetes education, n (%)		
Individual	443 (95.1)	48 (81.4)
Group	381 (81.8)	19 (32.2)
Duration of education session, median (IQR), min/session		
Individual	15 (10, 28)	30 (15, 30)
Group	30 (20, 30)	30 (20, 60)
Education sessions, median (IQR), n of sessions/y	3 (2, 4)	2 (1, 3.75)
Educational material, n (%)		
Brochures	391 (83.9)	44 (74.6)
Flip chart	309 (66.3)	20 (33.9)
Video	197 (42.3)	12 (20.3)
Computer slides	141 (30.3)	17 (28.8)
Conversation map	58 (12.5)	5 (8.5)
Food model	63 (13.5)	6 (10.2)
Poster	6 (1.3)	–
Evaluation of diabetes education, n (%)		
Successful	145 (31.1)	18 (30.5)
Uncertain outcome	199 (42.7)	22 (37.3)
Not evaluated	113 (24.2)	19 (32.2)

Note: Data are presented as n (%) or median (IQR).

Abbreviation: DEs, diabetes educators.

This proportion was higher than the corresponding figures that have been presented by other studies. For instance, in a self-report survey from the USA,¹⁴ physicians estimated that 37% of their patients received DSME in their service, while data drawn from Medicare and private-insurance reimbursements in the USA indicated that only 5%–6.8% of patients with newly diagnosed diabetes used a diabetes self-management training service.^{15,16} Unfortunately, there were no data on the number of patients receiving diabetes education through the health care system in Thailand. The major, government-provided, health insurance scheme in Thailand,

the Universal Coverage Scheme, records process indicators and outcome indicators related to diabetes care; unfortunately, it only records the percentage of patients receiving foot care education.¹⁷ Although our survey reported a high percentage of patients receiving diabetes education, 70% of DEs were uncertain about the educational outcomes. The lack of formal evaluation systems after the conduct of diabetes education programs is one of the diabetes-education weaknesses in Thailand.

Either individual or group approaches to DSMES is effective.³ Individual diabetes education is the approach

Figure 1 Number and workload of DEs at different-sized, public and private hospitals, as reported by DEs. **(A)** median (IQR) number of DEs per hospital; **(B)** median (IQR) number of patients receiving diabetes education per week per DE, * p -value <0.05, compared between public and private hospitals.

Abbreviation: DEs, diabetes educators.

predominantly utilized by public and private hospitals in Thailand. The public hospitals had a higher proportion of group education programs, a higher glycemic threshold for the provision of DSMES programs, and a shorter duration for individual programs than the private hospitals. The 2017 Diabetes Educator and the DSME National Practice Survey in the USA established that the majority of DEs (55%) saw ≤ 5 patients with diabetes per day, while 34% of DEs saw 6–10 patients with diabetes daily.¹⁸ In contrast, our study demonstrated that public hospital DEs in Thailand saw a median of 60 (IQR 30, 140) patients with diabetes weekly. Assuming that Thai public hospital DEs work a 5-day week, 22% of them advised ≤ 5 patients per day, 24.6% advised 6–10 patients daily, while 53.5% advised >10 patients each

working day. The higher workload of public hospital DEs might be one of the factors related to the low level of glycemic control achievement in Thailand.

Some of the reported perceived obstacles appear to relate to the absence of CDEs in Thailand. Because there is no official CDE training program and no formal position of “diabetes educator” in the Thai health care system, health professionals engaging in the duties of a DE are also obliged to work in their primary disciplines (such as nursing, clinical nutrition, and pharmacology). As a result, the most consequential obstacle to diabetes education that was reported was a lack of time to undertake diabetes education by the DEs due to their need to carry out their primary duties. In addition, DEs require a comprehensive knowledge of diabetes, effective teaching

Figure 2 Perceptions of obstacles to diabetes education, by respondent group. (A) public hospitals; (B) private hospitals. * p -value < 0.05 , compared between three groups of respondents.

Abbreviation: DEs, diabetes educators.

and learning skills, and a sound understanding of various psychosocial and behavior approaches.¹⁹ Interestingly, 60% of the public hospital DEs in the present study stated that their perceived lack of skills in assisting patients with behavior changes was an obstacle to diabetes education. Although the sustainability of behavior changes depends on many factors (such as self-efficacy, family support, social policy, and hierarchy of needs), developing behavior change skills is one of the strategies that can be employed by DEs to facilitate diabetes self-care by patients. Therefore, formally establishing the position of DE in the health care system, promoting DE as a career choice, and creating specific DE programs for health professionals may remove the identified obstacles.

Furthermore, healthcare professionals perceived some obstacles from patients in that they were often unwilling to change their unhealthy behaviors and/or showed a lack of interest in diabetes education. However, this study did not survey patients with diabetes. Therefore, we do not know whether the patients were really not interested in diabetes education or whether they had other reasons for not following the advice of the DEs. In a previous study, the reasons cited by people with diabetes for not attending DSME programs were an aversion to group classes, a lack of interest, transportation difficulties, and an inconvenient schedule.¹⁰ Furthermore, the current study demonstrated that administrators and doctors tended to report

a lower percentage of obstacles than DEs. Because administrators and doctors have a major role in determining hospital policies, an underestimation of the obstacles to diabetes education might result in less support by way of human, financial and facility resources.

This study has some limitations. Firstly, it did not specify the type of diabetes. However, because the prevalence of type 1 diabetes in Thailand is very low, the authors believe that the results represent the characteristics and obstacles of diabetes education for patients with type 2 diabetes.²⁰ In addition, while the study included the common obstacles to diabetes education listed in the literature, it did not specifically cover all existing obstacles in the Thai health care system (such as communication problems with ethnic minorities). Finally, the results of the current study were only based on the responses of health care professionals. In other words, the obstacles to diabetes education from the perspective of patients were not identified.

Conclusions

Both individual and group approaches to education are used in Thailand. Despite the reported percentage of patients receiving diabetes education being high, the large majority of DEs thought the programs were ineffective. For health-care professionals undertaking the defined roles of DEs, the chief obstacles to diabetes education were reported to be high workloads, unclear roles, and a lack of skills to assist patients to achieve behavior change. Lastly, a compounding factor is that hospital administrators and doctors often perceived a lower prevalence of obstacles than the nominal DEs, which may give rise to negative resource implications.

Acknowledgments

This study was supported by Siriraj Research Development Fund (Managed by Routine to Research: R2R). The authors give special thanks to Clinical Adjunct Professor Anne Belton for advice on the manuscript preparation, and thanks to Nujeejan Bunyapattanapong, Varisara Boonyuang, Bandit Pidokrath, and the Siriraj Diabetes Center staff for the data collection and administrative work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Aekplakorn W. The 5th Thai national health examination survey 2014. Nonthaburi Health systems research institute; 2014.
2. Powers MA, Bardsley J, Cypress M, et al. Diabetes self-management education and support in type 2 diabetes: a joint position statement of the American Diabetes Association, the American Association of Diabetes Educators, and the Academy of Nutrition and Dietetics. *Diabetes Care*. 2015;38(7):1372–1382. doi:10.2337/dc14-1100
3. Chrvala CA, Sherr D, Lipman RD. Diabetes self-management education for adults with type 2 diabetes mellitus: A systematic review of the effect on glycemic control. *Patient Educ Couns*. 2016;99(6):926–943. doi:10.1016/j.pec.2016.07.004
4. Kent D, D'Eramo Melkus G, Stuart PM, et al. Reducing the risks of diabetes complications through diabetes self-management education and support. *Popul Health Manag*. 2013;16(2):74–81. doi:10.1089/pop.2012.0020
5. Lian JX, McGhee SM, Chau J, Wong CKH, Lam CLK, Wong WCW. Systematic review on the cost-effectiveness of self-management education programme for type 2 diabetes mellitus. *Diabetes Res Clin Pract*. 2017;127:21–34. doi:10.1016/j.diabres.2017.02.021
6. The Canadian Diabetes Certification Board. Available from: <http://www.cdec.ca/>. Accessed April 24, 2018.
7. The National Certification Board for Diabetes Educators. Available from: <https://www.ncbde.org/>. Accessed April 24, 2018.
8. The Australian Diabetes Educators Association. Available from: <https://www.adea.com.au/>. Accessed April 24, 2018.
9. Haas L, Maryniuk M, Beck J, et al. National standards for diabetes self-management education and support. *Diabetes Care*. 2014;37(Suppl 1):S144–S153. doi:10.2337/dc14-S144
10. Maine Center for Disease Control and Prevention. Diabetes self-management education barrier study. 2006:1–38. Available from: <https://www.maine.gov/dhhs/meede/phdata/chronic-disease/reports-diabetes.htm>
11. Polit DF, Beck CT. The content validity index: are you sure you know what's being reported? Critique and recommendations. *Res Nurs Health*. 2006;29(5):489–497. doi:10.1002/(ISSN)1098-240X
12. Yamane T. *Statistics, an Introductory Analysis*. 2nd ed. New York: Horper and Row; 1967.
13. Asch DA, Jedrzejewski MK, Christakis NA. Response rates to mail surveys published in medical journals. *J Clin Epidemiol*. 1997;50(10):1129–1136. doi:10.1016/S0895-4356(97)00126-1
14. Peyrot M, Rubin RR, Funnell MM, Siminerio LM. Access to diabetes self-management education: results of national surveys of patients, educators, and physicians. *Diabetes Educ*. 2009;35(2):246–248, 252–246, 258–263. doi:10.1177/0145721708329546
15. Strawbridge LM, Lloyd JT, Meadow A, Riley GF, Howell BL. Use of medicare's diabetes self-management training benefit. *Health Educ Behav*. 2015;42(4):530–538. doi:10.1177/1090198114566271
16. Li R, Shrestha SS, Lipman R, Burrows NR, Kolb LE, Rutledge S. Diabetes self-management education and training among privately insured persons with newly diagnosed diabetes—United States, 2011–2012. *MMWR Morb Mortal Wkly Rep*. 2014;63(46):1045–1049.
17. National Health Security Office. คู่มือบริหาร กองทุน หลักประกันสุขภาพแห่งชาติ ปีงบประมาณ 2560 การบริหารงบ บริการ ควคม ป้องกัน และ รักษา โรค เรือรัง (*Manual of National Health Security Fund 2017*). Bangkok: แสงจันทร์การพิมพ์ (Sang Chan Publisher); 2016:38–56.
18. Rinker J, Dickinson JK, Litchman ML, et al. The 2017 diabetes educator and the diabetes self-management education national practice survey. *Diabetes Educ*. 2018;44(3):1–9.
19. Consultative Section on Diabetes Education International Diabetes Federation. *International Curriculum for Diabetes Health Professional Education*. 2008:15–106. Available from: <http://www.idf.org/e-library/education/64-the-international-curriculum-for-diabetes-health-professional-education,en.html>
20. International Diabetes Federation. *IDF Diabetes Atlas*. 8 ed. Brussels, Belgium: International Diabetes Federation; 2017.

Supplementary materials

Survey of characteristics and obstacles of diabetes education in Thailand

Administrators and doctors, please answer parts 1, 3, and 4.

Diabetes educators, please answer all parts of the survey.

Part 1: General information

- 1.1 Your hospital is located in:
- ☐ Northern Thailand   ☐ Northeastern Thailand   ☐ Central Thailand
- ☐ Eastern Thailand   ☐ Western Thailand   ☐ Southern Thailand
- 1.2 Your hospital is operated by:
- ☐ Office of the Permanent Secretary, Ministry of Public Health:
- ☐ Community hospital
- ☐ General Hospital
- ☐ Regional hospital
- ☐ Medical Service Department
- ☐ Ministry of Defence
- ☐ Royal Thai Police
- ☐ Private Sector
- 1.3 Your hospital has a capacity of beds
- 1.4 Your hospital provides treatment to around outpatients with diabetes per week
- 1.5 General information of respondent:
- 1.5.1 Gender: ☐ male   ☐ female
- 1.5.2 Age: years
- 1.5.3 Educational level: ☐ Bachelor   ☐ Master's degree   ☐ PhD
- ☐ Other _____
- 1.5.4 Status of respondent:
- ☐ Administrator: ☐ Director/Vice Director
- ☐ Chief Physician
- ☐ Other _____

Figure S1 (Continued).

☐ Physician responsible for diabetes clinic or non-communicable disease clinic:

☐ General Practitioner   ☐ Internist   ☐ Pediatrician

☐ Adult endocrinologist

☐ Pediatric endocrinologist

☐ Other _____

☐ Health professional responsible for providing diabetes education:

☐ Physician   ☐ Nurse (OPD)   ☐ Nurse (IPD)

☐ Nutritionist   ☐ Health educator   ☐ Pharmacist

☐ Other _____

Part 2: Characteristics of diabetes education (only to be completed by health professionals responsible for providing diabetes education)

2.1 In your opinion, what percentage of patients with diabetes in your hospital receive diabetes education?   %

2.2 Criteria to provide diabetes education in your hospital (you can choose more than 1 item):

☐ Patients with newly diagnosed diabetes

☐ Patients with uncontrolled diabetes; please specify the glycemic criteria for the provision of diabetes education:

Fasting plasma glucose over mg/dL

HbA1c over %

☐ Patients with diabetes complications, n (%)

☐ Other _____

2.3 Method of diabetes education in your hospital (you can choose more than 1 item):

☐ Individual education

☐ Group education; the typical number of patients per group is

2.4 Duration of education sessions (you can choose more than 1 item):

☐ Individual education; minutes/session

☐ Group education; minutes/session

2.5 The typical number of education sessions per year is

Figure S1 (Continued).

2.6 Educational material (you can choose more than 1 item):

- ☐ Brochures
- ☐ Flip charts
- ☐ Conversation map
- ☐ Video
- ☐ Computer slides
- ☐ Other _____

2.7 What is the outcome of diabetes education in your hospital?

- ☐ Successful
- ☐ Uncertain outcomes
- ☐ No evaluation conducted

Part 3: Workload of diabetes education

3.1 Does your hospital have health professionals responsible for providing diabetes education using the term “diabetes educator”?

- ☐ No
- ☐ Yes; number of diabetes educators ☐☐

What is the profession of the diabetes educators (you can choose more than 1 item):

- ☐ Physician   ☐ Nurse (OPD)   ☐ Nurse (IPD)
- ☐ Nutritionist   ☐ Health educator   ☐ Pharmacist
- ☐ Other _____

3.2 Does your hospital need more diabetes educators?

- ☐ Yes   ☐ by recruiting new staff, at around ☐☐ positions
- ☐ by training current staff to become diabetes educators, at around ☐☐ positions
- ☐ Uncertain
- ☐ No

3.3 **For diabetes educators**, how many patients with diabetes do you provide diabetes education to per week?

around ☐☐☐ patients/week

3.4 **For physicians**, how many outpatients with diabetes do you treat per week?

around ☐☐☐ patients/week

Figure S1 (Continued).

Part 4: Obstacles to diabetes education

In your opinion, what are the obstacles to diabetes education in your hospital?

- ☐ Inadequate number of diabetes educators
- ☐ Lack of time by diabetes educators due to other duties
- ☐ Lack of skills in assisting behavior change
- ☐ Lack of interest from patients
- ☐ Patients unwilling to change unhealthy behaviors
- ☐ Lack of budget
- ☐ Lack of suitable place for diabetes education
- ☐ Lack of suitable educational materials
- ☐ Other _____

Figure S1 Survey of characteristics and obstacles of diabetes education in Thailand.

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion

and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>