Association of *pvl* gene with incomplete hemolytic phenotype in clinical *Staphylococcus aureus*

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Menglu Gao¹
Ruirui Sang¹
Gang Wang¹
Yuanhong Xu^{1,2}

¹Department of Clinical Laboratory, The First Affiliated Hospital of Anhui Medical University, Hefei, 230000, People's Republic of China; ²Provincial Laboratories of Pathogen Biology and Zoonoses, Anhui Medical University, Anhui, Hefei 230000, People's Republic of China

Purpose: This experiment aimed to evaluate the correlation between the hemolytic phenotype of *Staphylococcus aureus* and *pvl* gene in terms of characteristics of antibiotic resistance.

Materials and methods: Two-hundred and eleven strains of hospital-acquired *S. aureus* and their bacterial susceptibility to 20 antibiotics were determined by MicroScan WalkAway96. All strains were cultured on Columbia sheep blood agar plates for 24 hours and then underwent ten passages for investigation of their hemolytic phenotypes. *S. aureus* produced incomplete β -hemolytic phenotype, termed as *S. aureus* strains with incomplete hemolytic phenotype (SIHP). The *pvl* gene was identified by PCR amplification followed by DNA sequencing. Statistical analyses of the data were performed using SPSS version 16.0 software.

Results: Fifty-two (24.64%) strains were confirmed to maintain the incomplete hemolytic phenotype of *S. aureus* (SIHP). Meanwhile, 15 (7.11%) of 211 strains were found to carry the *pvl* gene, and eight of the 15 strains were SIHP. Compared with *S. aureus* strains with complete hemolytic phenotype (SCHP), SIHP showed higher susceptibility to seven of the 20 antibiotics (oxacillin, ciprofloxacin, gentamicin, ceftriaxone, ceftiofur, levofloxacin, and moxifloxacin) ($P < 0.05$). The *pvl*-positive bacteria had a higher rate of resistance to four antibiotics (rifampin, ciprofloxacin, levofloxacin, and moxifloxacin) in comparison with the *pvl*-negative strains ($P < 0.05$).

Conclusion: SIHP had a high frequency of *pvl* gene. The *pvl*-positive isolates showed less resistance to rifampin, ciprofloxacin, levofloxacin, and moxifloxacin. Additionally, the majority of SIHP isolates (61.54%) were methicillin-resistant *S. aureus*. SIHP strains had significantly higher antibiotic resistance to ceftiofur when compared with SCHP, while SCHP strains had a high rate of antibiotic resistance to ciprofloxacin, gentamicin, ceftriaxone, levofloxacin, and moxifloxacin. The results may help to provide medical advice for selection of antibiotics for patients with SIHP-associated infections.

Keywords: *S. aureus*, incomplete hemolytic phenotype, *pvl*, antimicrobial drug resistance, virulence factor

Introduction

Staphylococcus aureus is one of the most common pathogens isolated from both healthy individuals and patients worldwide, which can produce a variety of secreted toxins and cause various infections in humans.^{1–3} PVL is one of the toxins of synergohymenotropic exotoxin produced by *S. aureus* and belongs to the pore-forming toxin family.^{4–6} PVL shows strong lytic activity against host defense cells such as human polymorphonuclear neutrophils,⁷ monocytes, macrophages, and rabbit neutrophils but not murine neutrophils in vitro.^{8–10} Pore formation requires

Correspondence: Yuanhong Xu
Department of Clinical Laboratory, The First Affiliated Hospital of Anhui Medical University, 218 Jixi Road, Shushan District, Anhui, Hefei 230022, People's Republic of China
Tel +86 135 0569 4447
Fax +86 551 6292 2125
Email xyhong1964@ahmu.edu.cn

the presence of the two components of the toxin, LukS-PV and LukF-PV. Previous studies demonstrated that PVL aggravated many infections,¹¹ such as skin and soft tissue infection,¹ necrotizing pneumonia, bone joint infections, and even bacteremia.^{12–14}

It has been found that the *pvl* gene can spread from strain to strain by bacteriophages and plasmids.^{15,16} As a result of the *pvl* gene's transmission mechanism, the prevalence of *pvl*-positive strains has gradually increased.¹⁷

The prevalence of the *pvl* gene has been less common in methicillin-susceptible *S. aureus* isolates than in methicillin-resistant *S. aureus* (MRSA).¹⁸ The *pvl* gene locus represents a stable genetic marker of community-acquired MRSA (CA-MRSA) strains.¹⁹ Studies have shown that PVL had a high proportion in CA-MRSA.²⁰ The PVL genes have also been reported to be present in 77% of the CA-MRSA isolates, but only 4% of HA-MRSA isolates were *pvl*-positive.²¹ Although the toxin may be a highly linked epidemiological marker for CA-MRSA strains, PVL is not the major virulence determinant of CA-MRSA.²²

Incomplete hemolytic phenotype has an opaque, darker hemolytic ring in contrast with the transparent hemolytic phenotype of *S. aureus* strains with complete hemolytic phenotype (SCHP).²³ Recently, Zhang et al proposed that a number of *S. aureus* strains with incomplete hemolytic phenotype (SIHP) strains belong to the class of *S. aureus*. The authors demonstrated that SIHP strains were MRSA which highly expressed β -hemolysin and carried the *tst* gene. Because toxic shock syndrome toxin (expressed by *tst*) causes toxic shock syndrome,^{24,25} SIHP possesses high virulence potential.²³ Here we determined the prevalence of SIHP, antimicrobial resistance, and PVL genes carried in *S. aureus*, and compared the diversity of these factors between SIHP and SCHP strains.

Materials and methods

Bacterial strains

After excluding strains from repeated sources, a total of 211 strains were selected randomly and isolated from patients in the First Affiliated Hospital of Anhui Medical University from January 2016 to December 2017. Constituent ratios of departments where strains were isolated are shown in Table 1. The specimens included 30 samples of blood (14.22%), 70 of secretions (33.18%), 45 of sputa (21.33%), nine of shunt fluids (4.27%), 25 from

wounds (11.85%), eight from interstitial fluids (3.79%), and 24 others (11.37%). The Medical Ethics Committee of the First Affiliated Hospital of Anhui Medical University approved this study (no Quick-PJ2018-07–29) and all isolates were collected with written informed consent from the patients, conducted in accordance with the Declaration of Helsinki.

Antimicrobial sensitivity testing

The bacteria were cultured on Columbia sheep blood agar plates at 35°C in an atmosphere containing 5% CO₂ (v/v) for 24 hours. According to the manufacturer's instructions, 1–2 single colonies were added to Mueller-Hinton broth (MHB), followed by proper adjustment of bacteria concentration. Based on the broth microdilution method and the Clinical and Laboratory Standards Institute criteria, bacterial identification and antimicrobial susceptibility testing were performed using MicroScan WalkAway 96 (Beckman Coulter, Brea, CA, USA).²⁶ The bacterial isolates were subjected to 20 antibiotics for sensitivity testing. Antibiotics included: tetracycline, beta-lactams (oxacillin, ceftriaxone, cefoxitin), fluoroquinolones (levofloxacin, ciprofloxacin, moxifloxacin), gentamicin,

Table 1 Constituent ratios of departments where strains were isolated

Department	Strains, n (%)
Neurology	9 (4.27)
Orthopedic	21 (9.95)
Pediatrics	5 (2.37)
ENT	3 (1.42)
Out-patient	11 (5.21)
Infectious diseases	11 (5.21)
Oncology	2 (0.95)
Intensive Care Unit	24 (11.37)
General surgery	15 (7.11)
Respiration	3 (1.42)
Plastic surgery	6 (2.84)
Cerebral surgery	3 (1.42)
Urology	5 (2.37)
Endocrinology	14 (6.64)
Dermatology	24 (11.37)
Burns	12 (5.69)
Nephrology	18 (8.53)
Hematology	6 (2.84)
Obstetrics and gynecology	1 (0.47)
Plastic surgery	9 (4.27)
Rheumatism and Immunological Disease	9 (4.27)
Total	211 (100.00)

ampicillin, amoxicillin, daptomycin, clindamycin, nitrofurantoin, linezolid, penicillin, rifampin, sulfamethoxazole–trimethoprim, quinupristin-dalfopristin, erythromycin, and vancomycin.

Detection of hemolytic phenotype

All the strains were identified as *S. aureus* by matrix-assisted laser desorption ionization time-of-flight mass spectrometry (MALDI-TOF MS VITEK V3.0, BioMérieux, Craponne, France). The strain ATCC25923 with the complete hemolytic phenotype was taken as SCHP reference. All the isolates were cultured on Columbia sheep blood agar plates from different companies (OXOID, UK; BioMérieux; BD Biosciences, San Jose, CA, USA; TianDa, People's Republic of China) at 35°C in an atmosphere containing 5% CO₂ (v/v) for 24 hours, and 1–2 single colonies from one agar plate were taken to the next Columbia sheep blood agar plates from the same company and cultured for 24 hours, and then underwent ten serial passages. The hemolytic phenomenon was observed. Strains with incomplete hemolytic phenotype on all four blood agar plates and their phenotypes stable to ten passages were identified as SIHP and otherwise, as SCHP. Several SIHP colonies gathered and formed a grid-like hemolytic phenotype.

Detection of *pvl* genes

The strains were inoculated on Columbia blood agar plates and cultured at 35°C in an atmosphere containing 5% CO₂ for 24 hours. Four bacterial colonies were selected and suspended in 8 mL of MHB and cultured at 37°C with shaking at 200 rpm for 12 hours. One mL of MHB bacterial suspension was moved to a sterile centrifuge tube with 200 µL lysostaphin (Sangon, Shanghai, People's Republic of China), boiled at 37°C for 1 hour. After that, DNA were extracted and purified using DNA Extraction Kit (TianGen, Beijing, People's Republic of China) according to the manufacturer's instructions. The DNA were stored at –20°C and prepared for PCR detection.

PCR was run as follows: pre-denaturation at 93°C for 3 minutes, denaturation at 93°C for 30 seconds, annealing at 55°C for 30 seconds, extension at 72°C for 1 minute. PCR products were sequenced and analyzed by BLAST for the expected sequences of peptides. Primers for *pvl* gene (PVL-F, 5'-ATCATTAGGTAAAATGTCTGGACATGATCCA-3'; and PVL-R, 5'-GCATCAAGTGTATTGGATAGCAAAAGC-3') were used to generate an internal control with an amplicon size of 433 bp.²⁷

Statistical analysis

Chi-squared test and Fisher's exact test were used for the statistical analysis and $P < 0.05$ was considered statistically significant. All the data were analyzed by using SPSS version 16.0 software (SPSS Inc., Chicago, IL, USA) and Figures 3–7 were made by GraphPad Prism version 5.0 (GraphPad Software, Inc., La Jolla, CA, USA).

Results

Hemolytic phenotype of SIHP strains on sheep agar blood plates

One-hundred and fifty-nine strains cultured on blood agar plates for 24 hours presented complete hemolytic rings (β-hemolytic phenotype) (Figure 1B). The incomplete hemolytic phenotype formed by a single colony of bacteria looked like a donut in which a small, lighter circular hemolytic ring located at the center and the outside was wrapped in an opaque, darker hemolytic ring (Figure 1A). When the two colonies grew close, a transparent linear hemolysis zone was formed at the junction of the two colonies on the hemolytic ring. Several SIHP colonies aggregated, forming a grid-like hemolytic phenotype (Figure 1C). In addition, after ten serial passages, the incomplete hemolytic phenotype still remained (Figure 1D). Fifty-two of 211 hospital-acquired *S. aureus* were identified as SIHP strains.

Testing for the *pvl* gene

Among the 211 clinical isolates, 15 (7.11%) yielded positive *pvl* gene amplification (Figure 2). Eight (53.33%) of the 15 *pvl*-positive isolates belonged to SIHP strains and seven (3.57%) of 196 *pvl*-negative isolates were SCHP strains. The prevalence of *pvl* gene in SIHP was significantly higher than that in SCHP ($P < 0.05$) (Figure 3).

Drug resistance of *S. aureus*

Antimicrobial characteristics of 211 *S. aureus* isolates are demonstrated in Figure 4. It shows that *S. aureus* strains presented high resistance to oxacillin (45.02%), cefoxitin (46.45%), ampicillin (76.30%), penicillin (82.46%), and erythromycin (65.88%) when compared with the other antibiotics tested such as tetracycline (31.28%), ceftriaxone (21.80%), levofloxacin (27.01%), ciprofloxacin (32.70%), moxifloxacin (17.06%), gentamicin (28.44%), amoxicillin (24.64%), daptomycin (0.47%), clindamycin (37.91%), nitrofurantoin (0.95%), linezolid (0.95%), rifampin (13.74%), sulfamethoxazole–trimethoprim (9.00%), and quinupristin-dalfopristin (2.84%). Among 211 *S. aureus*

Figure 1 Hemolytic phenotype comparative analyses of *Staphylococcus aureus* on blood agar plates.

Notes: (A) *S. aureus* strains with incomplete hemolytic phenotype (SIHP) strains growing in the microaerophilic condition. (B) *S. aureus* strains with complete hemolytic phenotype (SCHP) control strain of *S. aureus* ATCC25923. (C) Several SIHP colonies aggregated, forming a grid-like hemolytic phenotype. (D) SIHP strains after ten serial passages.

Figure 2 Detection of the *pvl* gene.

Notes: ATCC49775. Due to limitations of instrument, only 14 *pvl*-positive strains are shown here.

Abbreviations: M, marker; P, *pvl*-positive strain; N, *pvl*-negative strain.

isolates, the minimum inhibitory concentrations (MICs) of vancomycin were found to be MIC =0.5 (two strains), MIC =1.0 (109 strains), MIC 2.0 (99 strains), and MIC =4.0 (one strain). The MICs of *S. aureus* are shown in Figure 5. No statistically significant difference in MICs was noted between SIHP and SCHP strains. Similarly, no difference was seen between *pvl*-positive and -negative isolates.

Antimicrobial characteristics of *pvl*-positive and -negative strains

In comparison with the sensitivity of *pvl*-negative strains to quinolones (levofloxacin 29.08%, ciprofloxacin 35.20%, and moxifloxacin 18.37%) and rifampin (14.29%), the sensitivity of *pvl*-positive *S. aureus* to quinolones (levofloxacin 0%, ciprofloxacin 0%, and moxifloxacin 0%) and rifampin

Figure 3 Comparison of *pvl* gene between *Staphylococcus aureus* strains with incomplete hemolytic phenotype (SIHP) strains and *S. aureus* strains with complete hemolytic phenotype (SCHP) strains.

Notes: ** $P < 0.01$.

Abbreviations: pvl, pvl-positive *S. aureus*; Npvl, pvl-negative *S. aureus*.

Figure 5 Comparison of MIC to vancomycin between *Staphylococcus aureus* strains with incomplete hemolytic phenotype (SIHP) strains and *S. aureus* strains with complete hemolytic phenotype (SCHP) strains as well as pvl-positive and -negative strains.

Notes: (NS), $P > 0.05$, difference was not statistically significant.

Abbreviations: MIC, minimum inhibitory concentration; VAN, vancomycin; R, resistant; I, intermediate; S, susceptible; pvl, pvl-positive *S. aureus*; Npvl, pvl-negative *S. aureus*; NS, not significant.

(6.67%) was significantly elevated ($P < 0.05$) (Figure 6). No statistically significant difference in antibacterial sensitivity was noted in the remaining 16 antibiotics between pvl-positive and -negative strains ($P > 0.05$).

Antimicrobial characteristics of SIHP and SCHP strains

Compared with SCHP strains, SIHP strains had a remarkably higher susceptibility to levofloxacin, ciprofloxacin, and moxifloxacin. Additionally, SIHP strains were more susceptible to gentamicin and ceftriaxone than SCHP strains, but more resistant to ceftioxin than SCHP strains. There were 61.54% of SIHP strains but only 39.62% of SCHP strains found to be MRSA ($P > 0.05$). All statistical differences between SIHP and SCHP strains are shown in Figure 7.

Discussion

Bacteriophages are responsible for spreading *pvl* gene among bacterial strains.¹⁵ Recently, phage-therapy as a therapeutic strategy in some infections has been proposed.²⁸ Bacteriophages are able to completely prevent biofilm formation through this strain and eliminate the staphylococcal biofilms within 4 hours.²⁹ In in vitro inhibitory assays, phage κ lysed a range of clinically isolated MRSA strains successfully.³⁰ We speculate that bacteriophages might inhibit the resistance of pvl-positive strains to quinolones and rifampin.

The *pvl* gene prevalence of SIHP strains is significantly higher than that of SCHP strains. Moreover, SIHP strains have high susceptibility to quinolones and the susceptibility pattern is similar to pvl-positive strains. We believe that bacteriophages might inhibit the resistance of pvl-positive

Figure 4 Antimicrobial characteristics of 211 *Staphylococcus aureus* in terms of 19 antibiotics.

Abbreviations: AMC, amoxicillin; AMP, ampicillin; CIP, ciprofloxacin; CLI, clindamycin; CRO, ceftriaxone; DAP, daptomycin; ERY, erythromycin; FOX, ceftioxin; GEN, gentamicin; I, intermediate; LNZ, linezolid; LVX, levofloxacin; MPX, moxifloxacin; NIT, nitrofurantoin; OXA, oxacillin; PEN, penicillin; QDA, quinupristin-dalfopristin; R, resistant; RIF, rifampin; S, susceptible; SXT, sulfamethoxazole-trimethoprim; TCY, tetracycline.

Figure 6 Comparison of drug resistance to four antibiotics in *pvl*-positive and -negative strains.

Notes: * $P < 0.05$. pvl, *pvl*-positive *Staphylococcus aureus*; Npvl, *pvl*-negative *S. aureus*. **Abbreviations:** CIP, ciprofloxacin; I, intermediate; LVX, levofloxacin; MFX, moxifloxacin; R, resistant; RIF, rifampin; S, susceptible; pvl, *pvl*-positive *Staphylococcus aureus*; Npvl, *pvl*-negative *S. aureus*.

strains to quinolones, and the correlation between *pvl* gene and incomplete hemolytic phenotype may be responsible for quinolones sensitivity of SIHP strains.

MRSA occurs more frequently in SIHP strains than SCHP ones, our results are similar to those in the study of Zhang et al.²³ They demonstrated that SIHP isolates carried the *tst* gene which expressed toxic shock syndrome toxin and caused toxic shock syndrome. In the present study, we noted that the *pvl* gene was frequently carried by SIHP strains and they were more resistant to cefoxitin. Therefore, we suggest that SIHP might be a class of *S. aureus* with potentially high virulence, that commonly

carries virulence factors and presents relatively high resistance to oxacillin and cefoxitin.

SIHP strains are more susceptible to gentamicin and ceftriaxone, and more resistant to oxacillin and cefoxitin than SCHP strains. There may be a series of common regulatory genes that regulate antibiotic sensitivity and hemolytic phenotype. The mechanism of susceptibility, however, remains unclear.

As described previously, SIHP strains with *tst* and *pvl* gene cause toxic shock syndrome and aggravate many infections.^{11,24} They have high susceptibility to quinolones, gentamicin, and ceftriaxone as well. Unfortunately, identification of antibiotic susceptibility from the isolation and culture of bacteria is time-consuming. Therefore, for patients with severe *S. aureus* infection, selection of antibiotics will be preferentially based on clinical experience. These results may assist when providing medical advice on the appropriate use of antibiotics when the *S. aureus* isolated from patients has been defined as incomplete hemolytic phenotype. The relationship between the *pvl* gene and drug resistance provides microbiologists with a way to explore how the PVL gene reduces drug resistance in *S. aureus*. Our study, however, may not reflect a general situation of hospital-acquired *S. aureus* in central China, due to the limitation of district and the number of *S. aureus* strains collected. Further approaches would be needed to clarify the significance of PVL gene in formation of incomplete hemolytic ring and increased antibiotic resistance of *S. aureus*.

Figure 7 Comparison of drug resistance to seven antibiotics between *Staphylococcus aureus* strains with incomplete hemolytic phenotype (SIHP) strains and *S. aureus* strains with complete hemolytic phenotype (SCHP) strains.

Notes: ** $P < 0.005$; * $P < 0.05$.

Abbreviations: CIP, ciprofloxacin; CRO, ceftriaxone; FOX, cefoxitin; GEN, gentamicin; I, intermediate; LVX, levofloxacin; MFX, moxifloxacin; OXA, oxacillin; R, resistant; S, susceptible.

Conclusion

Taken together, the frequency of *pvl* gene is likely associated with the incomplete hemolytic ring of *S. aureus* strains. *pvl*-positive *S. aureus* shows a high sensitivity to quinolones and rifampin. SIHP strains are susceptible to quinolones, ceftriaxone, and gentamicin but have a higher resistance to oxacillin and ceftiofloxacin when compared with SCHP strains, which may be helpful for correct selection of antibiotics in patients with SIHP-associated infections.

Ethical approval

The Medical Ethics Committee of the First Affiliated Hospital of Anhui Medical University approved this study (no Quick-PJ2018-07-29) and all isolates were collected with written informed consent from the patients, in accordance with the Declaration of Helsinki.

Data availability

All data generated or analyzed during this study are included in this published article.

Acknowledgments

We are grateful to the Department of Anhui Provincial Hospital for the materials used in this study. We especially thank Dr JL Shen from the Key Laboratory of Parasitology and Zoonoses, Anhui for his technical assistance and for access to the laboratory facilities. This work was supported by Anhui Natural Science Foundation (9021138203).

Author contributions

All authors contributed to data analysis, drafting and revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- Cribier B, Prevost G, Couppie P, Finck-Barbançon V, Grosshans E, Piémont Y. Staphylococcus aureus leukocidin: a new virulence factor in cutaneous infections? An epidemiological and experimental study. *Dermatology*. 1992;185:175–180. doi:10.1159/000247443
- Lina G, Piémont Y, Godail-Gamot F, et al. Involvement of Pantone-Valentine leukocidin-producing Staphylococcus aureus in primary skin infections and pneumonia. *Clin Infect Dis*. 1999;29:1128–1132. doi:10.1086/313461
- Diep BA, Gillet Y, Etienne J, Lina G, Vandenesch F. Pantone-Valentine leukocidin and pneumonia. *Lancet Infect Dis*. 2013;13:566. doi:10.1016/S1473-3099(13)70102-6
- Prevost G, Cribier B, Couppie P, et al. Pantone-Valentine leukocidin and gamma-hemolysin from Staphylococcus aureus ATCC 49775 are encoded by distinct genetic loci and have different biological activities. *Infect Immun*. 1995;63:4121–4129.
- Woodin AM. Fractionation of a leukocidin from Staphylococcus aureus. *Biochem J*. 1959;73:225–237. doi:10.1042/bj0730225
- Pedelacq JD, Maveyraud L, Prevost G, et al. The structure of a Staphylococcus aureus leukocidin component (LukF-PV) reveals the fold of the water-soluble species of a family of transmembrane pore-forming toxins. *Structure*. 1999;7:277–287.
- Finck-Barbançon V, Duportail G, Meunier O, Colin DA. Pore formation by a two-component leukocidin from Staphylococcus aureus within the membrane of human polymorphonuclear leukocytes. *Biochim Biophys Acta*. 1993;1182:275–282.
- Gladstone GP, Van Heyningen WE. Staphylococcal leukocidins. *Br J Exp Pathol*. 1957;38:123–137.
- Diep BA, Chan L, Tattevin P, et al. Polymorphonuclear leukocytes mediate Staphylococcus aureus Pantone-Valentine leukocidin-induced lung inflammation and injury. *Proc Natl Acad Sci U S A*. 2010;107:5587–5592. doi:10.1073/pnas.0912403107
- Hongo I, Baba T, Oishi K, Morimoto Y, Ito T, Hiramatsu K. Phenol-soluble modulin alpha 3 enhances the human neutrophil lysis mediated by Pantone-Valentine leukocidin. *J Infect Dis*. 2009;200:715–723. doi:10.1086/605332
- Tong SY, Davis JS, Eichenberger E, Holland TL, Fowler VG. Staphylococcus aureus infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev*. 2015;28:603–661. doi:10.1128/CMR.00134-14
- Kreienbuehl L, Charbonney E, Eggimann P. Community-acquired necrotizing pneumonia due to methicillin-sensitive Staphylococcus aureus secreting Pantone-Valentine leukocidin: a review of case reports. *Ann Intensive Care*. 2011;1:52. doi:10.1186/2110-5820-1-52
- Flammier S, Rasigade JP, Badiou C, et al. Human monocyte-derived osteoclasts are targeted by Staphylococcal pore-forming toxins and superantigens. *PLoS One*. 2016;11:e0150693. doi:10.1371/journal.pone.0150693
- Shallcross LJ, Fragaszy E, Johnson AM, Hayward AC. The role of the Pantone-Valentine leukocidin toxin in staphylococcal disease: a systematic review and meta-analysis. *Lancet Infect Dis*. 2013;13:43–54. doi:10.1016/S1473-3099(12)70238-4
- Xia G, Wolz C. Phages of Staphylococcus aureus and their impact on host evolution. *Infect Genet Evol*. 2014;21:593–601. doi:10.1016/j.meegid.2013.04.022
- Boyle-Vavra S, Daum RS. Community-acquired methicillin-resistant Staphylococcus aureus: the role of Pantone-Valentine leukocidin. *Lab Invest*. 2007;87:3–9. doi:10.1038/labinvest.3700501
- Klein S, Menz MD, Zanger P, Heeg K, Nurjadi D. Increase in the prevalence of Pantone-Valentine leukocidin and clonal shift in community-onset methicillin-resistant Staphylococcus aureus causing skin and soft-tissue infections in the Rhine-Neckar Region, Germany, 2012-2016. *Int J Antimicrob Agents*. 2019;53:261–267. doi:10.1016/j.ijantimicag.2018.10.026
- David MZ, Daum RS. Community-associated methicillin-resistant Staphylococcus aureus: epidemiology and clinical consequences of an emerging epidemic. *Clin Microbiol Rev*. 2010;23:616–687. doi:10.1128/CMR.00081-09
- Vandenesch F, Naimi T, Enright MC, et al. Community-acquired methicillin-resistant Staphylococcus aureus carrying Pantone-Valentine leukocidin genes: worldwide emergence. *Emerg Infect Dis*. 2003;9:978–984. doi:10.3201/eid0908.030089
- Nakagawa S, Taneike I, Mimura D, et al. Gene sequences and specific detection for Pantone-Valentine leukocidin. *Biochem Biophys Res Commun*. 2005;328:995–1002. doi:10.1016/j.bbrc.2005.01.054

21. Naimi TS, LeDell KH, Como-Sabetti K, et al. Comparison of community- and health care-associated methicillin-resistant *Staphylococcus aureus* infection. *Jama*. 2003;290:2976–2984. doi:10.1001/jama.290.22.2976
22. Voyich JM, Otto M, Mathema B, et al. Is Pantone-Valentine leukocidin the major virulence determinant in community-associated methicillin-resistant *Staphylococcus aureus* disease? *J Infect Dis*. 2006;194:1761–1770. doi:10.1086/509506
23. Zhang H, Zheng Y, Gao H, et al. Identification and characterization of *Staphylococcus aureus* strains with an incomplete hemolytic phenotype. *Front Cell Infect Microbiol*. 2016;6:146. doi:10.3389/fcimb.2016.00146
24. Saha B, Harlan DM, Lee KP, June CH, Abe R. Protection against lethal toxic shock by targeted disruption of the CD28 gene. *J Exp Med*. 1996;183:2675–2680. doi:10.1084/jem.183.6.2675
25. Thomas D, Chou S, Dauwalder O, Lina G. Diversity in *Staphylococcus aureus* enterotoxins. *Chem Immunol Allergy*. 2007;93:24–41. doi:10.1159/000100856
26. Song KH, Kim ES, Park KH, et al. Clinical and molecular characterization of pantone-valentine leukocidin-positive invasive *Staphylococcus aureus* infections in Korea. *Microb Drug Resist*. 2019;25:450–456. doi:10.1089/mdr.2018.0238
27. McClure JA, Conly JM, Lau V, et al. Novel multiplex PCR assay for detection of the staphylococcal virulence marker Pantone-Valentine leukocidin genes and simultaneous discrimination of methicillin-susceptible from -resistant staphylococci. *J Clin Microbiol*. 2006;44:1141–1144. doi:10.1128/JCM.44.3.1141-1144.2006
28. Capparelli R, Parlato M, Borriello G, Salvatore P, Iannelli D. Experimental phage therapy against *Staphylococcus aureus* in mice. *Antimicrob Agents Chemother*. 2007;51:2765–2773. doi:10.1128/AAC.01513-06
29. Fenton M, Keary R, McAuliffe O, Ross RP, O'Mahony J, Coffey A. Bacteriophage-derived peptidase CHAP(K) eliminates and prevents *Staphylococcal* biofilms. *Int J Microbiol*. 2013;2013:625341. doi:10.1155/2013/625341
30. O'Flaherty S, Ross RP, Meaney W, Fitzgerald GF, Elbreki MF, Coffey A. Potential of the polyvalent anti-*Staphylococcus* bacteriophage K for control of antibiotic-resistant staphylococci from hospitals. *Appl Environ Microbiol*. 2005;71:1836–1842. doi:10.1128/AEM.71.4.1836-1842.2005

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>