Effect of a mindfulness-based pilot intervention on maternal-fetal bonding

This article was published in the following Dove Press journal:
International Journal of Women's Health

Karina M Shreffler¹
Stacy Tiemeyer¹
Lucia Ciciolla²
Julie M Croff³

¹Department of Human Development and Family Science, Oklahoma State University, Stillwater, OK, USA;

²Department of Psychology, Oklahoma State University, Stillwater, OK, USA;

³Department of Rural Health, Center for Health Sciences, Oklahoma State University, Tulsa, OK, USA

Abstract: This pilot study explored the effects of a two-week mindfulness-based intervention designed to enhance maternal-fetal bonding among pregnant women. Participants who listened to their baby's heartbeat every day for two weeks and received four texts per week with mindfulness-based activities to do from home reported a significant increase in reported maternal-fetal attachment scores across the study period. Enhancing maternal-fetal bonding through this low-cost intervention has the potential for reducing adverse birth outcomes.

Keywords: pregnancy, mindfulness, intervention, attachment, bonding, MFAS

Introduction

Maternal-fetal bonding refers to the affectionate feelings and perceptions a mother develops for her fetus during pregnancy¹ and includes behavioral acts during pregnancy that demonstrate care and commitment to the fetus.² Maternal-fetal bonding lays the foundation for post-birth mother-infant bonding and early caregiving, including maternal sensitivity, nurturing behaviors, and infant attachment security,^{3,4} which are necessary for infants to reach their developmental potential.^{5,6}

Maternal-fetal bonding is also critical for healthy fetal development because it is associated with healthier maternal behaviors during pregnancy.⁷ For example, women with high levels of prenatal bonding are more likely to stop or reduce smoking cigarettes, alcohol, and illicit drug use.^{7,8} Fortunately, maternal-fetal bonding appears to be modifiable; prior research indicates ultrasounds can increase prenatal attachment and improve maternal health behaviors during pregnancy.⁹ Yet the high costs (eg, equipment, medical personnel, patient time off of work, transportation, etc.) may be prohibitive for the women who would benefit most from intervention.

Methods

The data for this project, Babies and Moms connected by Love, Openness, and Opportunity (BLOOM) come from a larger study following 177 pregnant girls and women (ages 15–40) from two prenatal clinics serving high proportions of Medicaid patients. The Oklahoma State University Institutional Review Board approved the study protocol before data collection began. Participants provided written informed consent for the overall study at the time of enrollment; participants in BLOOM electronically reviewed and signed additional informed consent forms before participating in the additional BLOOM intervention. Participants completed online surveys about their pregnancy, including prenatal

Correspondence: Karina M Shreffler
Department of Human Development and Family Science, Oklahoma State University, 233 Human Sciences Building, Stillwater, OK 74078, USA
Tel +1 918 594 8389
Email karina.shreffler@okstate.edu

bonding and sociodemographic information. All women enrolled in the larger study were invited to participate in BLOOM if they were still pregnant, in their second trimester, and were planning to take the baby home from the hospital. The sample for the present study consists of 34 pregnant girls and women (15–40 years of age) in their second trimester who volunteered for the BLOOM intervention.

The study was conducted in 2018. Participants were randomly assigned to a control group ($n=9$) or to one of three intervention groups that received either (1) fetal Doppler heartrate monitors ($n=8$); (2) four texts per week with mindfulness tasks designed to enhance feelings of maternal attachment ($n=11$); or (3) both Doppler monitors and mindfulness exercises ($n=6$). The groups did not significantly differ by sociodemographic, health, or pregnancy characteristics. The participants who received fetal Doppler monitors were trained on appropriate use and instructed to listen to their baby's heartbeat for at least 1 min per day over the two-week period. We examined compliance with the protocol by asking participants during the follow-up how many times per day that actually used the Doppler during the study period. On average, participants with the Dopplers reported using the monitors 1.5 times per day.

The mindfulness exercises were sent via text messages with instructions for the specific exercise assigned that day. All participants in the mindfulness groups received the same text message. Text messages were sent at the same time of day for all groups (mid-evening). Women were asked to “take a few moments alone and complete the following task”, which were different and included: deep breathing, meditation, prenatal massage, responding to kicks, nursery rhymes, telling the baby about a cherished person in their life, planning an activity with the baby, and reading a story to the baby.

All participants completed the Maternal-Fetal Attachment Scale (MFAS)¹⁰ prior to and immediately following the intervention. The MFAS is a 23-item scale measuring maternal interaction and feelings toward the fetus.¹⁰ For example, participants were asked whether they think or do the following things: talk to their baby, refer to the baby by a nickname, decided on a name, wonder what the baby thinks and feels inside me, gave up doing certain things because they want to help their baby. Responses to the questions ranged from “definitely yes (1) to uncertain (3) to definitely no (5)”. We reverse coded one item, “I feel my body is ugly.” The potential

MFAS score range is between 23 and 115. Cronbach's alpha for our sample is 0.87.

A change score analysis was conducted on participants' pre- and post-test MFAS scores to determine the effect of the intervention. We first calculated the change score by subtracting the pre-MFAS score from the post-MFAS score. To compare differences in pre- and post-test MFAS scores we used one-way analysis of variance. A one-way analysis of covariance was used to estimate the association between the intervention groups and MFAS change score, controlling for the pre-MFAS score.

Results

Table 1 presents pre- and post-test results for each group. The average pre-MFAS score was 93.86 ($SD=9.14$). The intervention groups did not differ significantly from each other on pre-MFAS scores. For the post-MFAS scores, the average score was 94.67 ($SD=9.57$), and all intervention groups reported higher levels of post-MFAS relative to the control group, but the difference was not statistically significant. Comparing change scores before and after the two-week period, the control group was the only one to report a decrease in feelings of maternal-fetal bonding ($\Delta S = -1.78$). Using the change score as the outcome variable, we fit an ANCOVA model to determine if changes in MFAS scores between baseline and time 2 were significant. After adjusting for baseline MFAS, women in the Doppler + Mindfulness group had significantly higher levels of MFAS, with $F(1, 28) = 5.39$, $p < 0.05$. Controlling for baseline scores, results indicate a significant increase in MFAS scores for the Doppler + Mindfulness group ($\Delta S = 4.83$, $p < 0.05$, Cohen's $d = 0.59$). The other three groups all reported an increase in scores post-intervention, but their change scores did not differ significantly from the control group.

Discussion

Although maternal-fetal bonding typically increases throughout pregnancy, there is considerable variation among pregnant women.¹ Our findings suggest that mindfulness skill-building in combination with the use of inexpensive fetal Doppler monitors at home by pregnant women quickly achieves an increase in maternal feelings of bonding. Future research is needed to determine whether the increase in maternal-fetal bonding scores due to the BLOOM intervention is associated with subsequent improvements in maternal health behaviors during pregnancy.

Table I BLOOM intervention pilot results (N=34)

	Condition									
	Control (n=9)		Doppler only (n=8)		Doppler+ Mindfulness (n=6)		Mindfulness only (n=11)		Total (n=34)	
	M	SD	M	SD	M	SD	M	SD	M	SD
Pre-MFAS	90.40	11.32	94.88	8.46	93.00	6.10	96.73	8.81	93.86	9.14
Post-MFAS	88.22	11.46	96.57	7.21	97.83	9.83	97.00	7.51	94.67	9.57
Difference MFAS	-1.78	7.31	2.57	5.50	4.83	6.68	0.27	5.90	1.03	6.52
Change Score Analysis ^a										
β			0.34		0.44*		0.28			
Adj. R^2									0.15	
Analysis of covariance										
	Sum of squares				df		Mean square		F	p
Model	353.13				4		88.28		2.45	0.069
Control	111.55				1		111.55		3.10	0.089
Doppler only	194.14				1		194.14		5.39	0.028
Doppler + Mindfulness	65.62				1		65.62		1.82	0.188
Mindfulness only	172.44				1		172.44		4.79	0.037
Error	1007.84				28		35.99			
Total	1360.97				32		42.53			

Notes: ^aChange score analysis of difference in MFAS between time 1 and time 2 controlling for baseline MFAS. * $p < 0.05$.

Abbreviations: BLOOM, Babies and Moms connected by Love, Openness, and Opportunity; MFAS, Maternal-Fetal Attachment Scale.

Acknowledgments

This research was supported in part by the National Institute of General Medical Sciences of the National Institutes of Health (P20GM109097; Jennifer Hays-Grudo, PI). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The study sponsor had no role in study design, collection, analysis, interpretation of the data, writing the report, or the decision to submit the report for publication. An earlier version of this paper was presented at the 2019 annual conference of the American Academy of Health Behavior in Greenville, SC.

Disclosure

Dr. Karina M Shreffler reports grants from National Institute of General Medical Sciences, during the conduct of the study. The authors report no other conflicts of interest in this work.

References

- Brandon A, Pitts S, Denton W, Stringer A, Evans H. A history of the theory of prenatal attachment. *J Prenat Perinat Psychol Health*. 2009;23:201–222.
- Salisbury A, Law K, LaGasse L, Lester B. Maternal-fetal attachment. *Jama*. 2003;289:1701. doi:10.1001/jama.289.13.1701
- Shin H, Park YJ, Kim MJ. Predictors of maternal sensitivity during the early postpartum period. *J Adv Nurs*. 2006;55(425–434). doi:10.1111/j.1365-2648.2006.03943.x
- Siddiqui A, Hagglof B. Does maternal prenatal attachment predict postnatal mother-infant interaction? *Early Hum Dev*. 2000;59:13–25.
- Black MM, Walker SP, Fernald LC, et al. Early childhood development coming of age: science through the life course. *Lancet*. 2017;389:77–90. doi:10.1016/S0140-6736(16)31389-7
- Shaw DS, Vondra JJ. Infant attachment security and maternal predictors of early behavior problems: a longitudinal study of low-income families. *J Abnorm Child Psychol*. 1995;23:335–357. doi:10.1007/BF01447561
- Lindgren K. Relationships among maternal-fetal attachment, prenatal depression, and health practices in pregnancy. *Res Nurs Health*. 2001;24:203–217. doi:10.1002/(ISSN)1098-240X
- Massey SH, Bublitz MH, Magee SR, et al. Maternal-fetal attachment differentiates patterns of prenatal smoking and exposure. *Addict Behav*. 2015;45:51–56. doi:10.1016/j.addbeh.2015.01.028

9. Sedgmen B, McMahon C, Cairns D, Benzie RJ, Woodfield RL. The impact of two-dimensional versus three-dimensional ultrasound exposure on maternal-fetal attachment and maternal health behavior in pregnancy. *Ultrasound Obstet Gynecol.* 2006;27:245–251. doi:10.1002/uog.2703
10. Cranley MS. Development of a tool for the measurement of maternal attachment during pregnancy. *Nurs Res.* 1981;30:281–284. doi:10.1097/00006199-198109000-00008

International Journal of Women's Health

Dovepress

Publish your work in this journal

The International Journal of Women's Health is an international, peer-reviewed open-access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of women's healthcare including gynecology, obstetrics, and breast cancer. The

manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-womens-health-journal>