Should embolectomy be performed in late acute lower extremity arterial occlusions?

Hikmet Iyem
M Nesimi Eren

Department of Cardiovascular
Surgery, Dicle University, Diyarbakir,
Turkey

Background: We analyzed the embolectomy results and complications of patients who were operated on after a diagnosis of late acute arterial occlusion of lower extremities.

Methods: A total of 122 patients operated on in our clinic between 2004 and 2009 for late acute arterial occlusion were included in the study. Late arterial occlusion was defined as occlusion occurring 72 hours after initial manifestation of the patient complaints related to the affected lower extremity.

Results: Average age of the 122 patients (71 male, 51 female) was 54.2 ± 16.8 years. In this cohort, 64.75% of patients had cardiac pathologies, while 28.68% had extracardiac causes; 1.64% patients had catheterization, 0.81% patient had malignancy, and 2.46% patients had a history of trauma. In 1.64% of the cases, no reason for thromboembolysis could be found. Thirty-one patients (25.40%) had additional surgical operations, 14 (11.47%) had fasciotomy, and 9 (7.37%) had amputation. Re-embolectomy was performed on 37 patients (30.32%) who had ongoing ischemia after an operation. Additional surgical operations were performed on 31 patients (25.40%) with ongoing ischemia. In 14 of these cases (11.47%), patients were treated with fasciotomy due to development of compartment syndrome. Amputation was performed on a total of 9 patients. Early in the postoperative phase, mortality was observed in 11 patients (9.01%).

Conclusion: We believe that late embolectomies of acute late leg ischemia increases blood flow in the extremity and reduces the number of amputations required.

Keywords: lower extremity, embolism, prognosis, treatment outcome

Introduction

Critical leg ischemia can be observed as an acute event that is either the initial finding of peripheral arterial disease for a patient or the reason for symptomatic deterioration in a patient with peripheral arterial disease of the lower extremities (intermittent claudication). Since some cells are more sensitive to anoxia, it is very difficult to determine the tolerance of extremities to ischemia. After 4 to 6 hours of ischemia, irreversible changes arise in skeletal muscles and peripheral nerves.

In acute arterial occlusive disease, rapid diagnosis and performing the appropriate treatment with a multidisciplinary approach are critically important. Nevertheless, despite the progress in treatment and techniques, amputation and mortality rates have remained high, even in special/private vascular surgery centers.¹ Late use of embolectomy on an extremity with ischemia remains controversial, but has been gaining acceptance.

It is known that a delay of more than 8 hours increases ischemic complications in patients with arterial emboli in an extremity. On the other hand, the literature

Correspondence: Hikmet Iyem
Dicle Üniversitesi Kalp Damar Cerrahisi
AD/Diyarbakir/Turkey
Tel +90 412 2488001
Fax +90 412 2488021
Email hikmetiyem@gmail.com

shows a decrease in appearance of complications when the period of ischemia exceeds 7 days.² In light of this observation, it can be inferred that performance of a surgical operation after the patient has survived the first week of ischemia with little extremity or tissue loss could be more successful. In this clinical study, our aim was to analyze retrospectively the results of arterial embolectomies and complications in patients, who are admitted to surgery with diagnosis of late acute lower extremity arterial occlusion.

Materials and methods

A total of 122 patients who were admitted to our clinic with diagnosis of late acute lower extremity arterial occlusion and who were operated on between January 2004 and February 2009 were included in the study. Computer records of these patients were retrospectively analyzed and recorded.

Late arterial occlusion is defined as occlusion occurring 72 hours after initial manifestation of patient complaints related to the effected lower extremity. Only the patients who met this criterion were included in this study. We excluded the patients with polytrauma because we thought that it may negatively affect mortality and morbidity. In all cases, diagnosis was made by the use of anamnesis, physical examination, and doppler ultrasonography; we employed no additional tests, such as conventional angiography or CT angiography, to any patients prior to surgery. No intraoperative angiography was performed on any patient, but all patients duplex examination was performed on all patients in the preoperative period.

When examining the patients, we sought answers to three questions: is the extremity still viable? Is the viability of the extremity acutely threatened? Are there irreversible changes that hinder preservation of the extremity? All of the patients were analyzed according to the Turkish Society of Vascular Surgery classification scheme (Table 1).³ Nonetheless, we performed advanced tests on those patients whose ischemia were not treated with embolectomy and who had planned reoperations. After embolectomy, all the

patients were examined with abdominal ultrasonography and cardiac ecocardiography to investigate the origin of the arterial occlusion.

Surgical method

Femoral arterial embolectomies of the lower extremities were performed under local anesthesia by a longitudinal incision from the femoral region. Spinal anesthesia was used for patients that had popliteal embolectomy and additional surgical operation. Standard 3F, 4F and 5F Fogarty embolectomy catheters were used. 3F and 4F Fogarty catheters were used for distal embolectomy while 5F Fogarty catheters were used for proximal embolectomy.

The embolectomy was stopped when the thrombus was completely removed from distal and proximal approaches (when the thrombus could not be contacted distally or proximally by a Fogarty catheter), and distal backflow and proximal flow was deemed adequate. In all cases, arteriotomy was closed by washing the distal bed with 5000 U of heparin diluted in 100 mL of 0.9% physiological saline solution. For patients with ongoing ischemia despite embolectomy, additional surgery was performed after CT angiography or conventional angiography (Table 2).

Preoperative and postoperative medical treatment protocol

For all patients diagnosed with arterial occlusion, heparin-dextran 40-pentoxifyllin infusion was administered and continued for 3 days postoperatively. Conventional heparin was started at a dose of 100 U/kg/h and dextran 40 at 0.5 mL/kg/h. Heparin was assessed with ACT (activated coagulation time). ACT level was maintained within the range of 200 to 300 seconds. Following this, patients were administered anticoagulants.

Statistical analysis

SPSS Windows 10.0 statistical software package was used for analyzing patient records. Continuous variables are presented as means $\pm$ standard deviation. Student's test, Chi-square

Table 1 Assessment of viability of the extremity

	Description/ Prognosis	Sensory loss	Functional failure in muscle	Arterial doppler signals	Venous doppler signals
Stage 1: Viable	No urgent threat	No	No	Yes	Yes
Stage 2a: Marginal threat	Can be saved	Minimal (fingers)	No	Yes	Yes
Stage 2b: Serious threat	Can be saved urgently	Extensive	Mild-moderate	No	Yes
Stage 3: Irreversible	Inevitable tissue loss	Anesthesia	Paralysis	No	No

Table 2 Surgical operations and occlusion levels

	N	%
Surgical operations		
Femoral embolectomy	88	72.13
Popliteal embolectomy	21	17.21
Femoral + popliteal embolectomy	13	10.65
Additional surgical operations		
Aortafemoral bypass	7	5.73
Iliofemoral bypass	5	4.09
Femorofemoral bypass	7	5.73
Femoropopliteal bypass	12	9.83
Occlusion level		
Aortic bifurcation	7	5.73
Iliac	12	9.83
Femoral	78	63.93
Popliteal	25	20.49

and Fisher's exact tests were used. A *P* value <0.05 was considered as statistically significant.

Results

Average age of the 122 patients (71 male, 51 female) operated on was 54.2 ± 16.8 years. Preoperative demographic data of the patients are presented in Table 3. We observed that peripheral arterial occlusions were frequently caused by cardiac pathologies. Cardiac-related reasons were present in 79 cases (64.75%) (atrial fibrillation in 46 cases, mitral stenosis in 5 cases, mitral stenosis and atrial fibrillation in 14 cases, prosthetic cardiac valves in 4 cases, myocardial infarction history in 9 cases, endocarditis in 1 case); extracardiac reasons were present in 35 cases (28.68%) (atherosclerosis in 24 cases, vascular trauma in 4 cases, abdominal aort aneurism in 7 cases); catheterization in 2 cases (1.64%); malignancy in 1 case (0.81%); and history of trauma in 3 cases (2.46%). In 2 cases (1.64%) no reason could be found to explain the thromboembolysis.

Re-embolectomy was performed on 37 patients (30.32%) that had ongoing ischemia after an operation. Additional surgical operations were performed on 31 patients (25.40%) with ongoing ischemia (Table 2). In 14 of these cases (11.47%), patients were treated with fasciotomy due to development of compartment syndrome. Amputation was performed on a total of 9 patients (7.37%) who had life-threatening risks; these included 7 patients who had ongoing ischemia despite fasciotomy and had developed a demarcation line, and 2 patients with graft infections. Seven of the amputations were above and 2 were below the knee.

Table 3 Preoperative demographic data

	N	%
Age (average)	54.2 ± 16.8	
Sex (male/female)	71/51	58.19/41.80
Accompanying diseases		
Diabetes mellitus	28	22.95
Hypertension	31	25.40
Atrial fibrillation	60	49.18
Congestive heart failure	9	7.37
Rheumatismal heart disease	23	18.85
Cerebrovascular disease	3	2.45
Ischemic heart disease	13	10.65
Onset of complaints		
3–7 days	91	74.59
>7 days	31	25.40
Admittance complaints		
Pain	114	93.44
Coldness	88	72.13
Paleness	94	77.04
Sensory loss	12	9.83
Motor deficit	4	3.27
Cyanosis	13	10.65

Incision site infections were observed in 11 of the patients (9.01%). Infections developed at the amputation site in 2 patients, in the popliteal region below the knee in 2 patients, at abdominal incision site in 1 patient, and 3 patients had graft infections. Two patients with graft infections had amputations performed and the other patients were treated with antibiotherapy. Among the less common complications, myonephropathic metabolic syndrome (reperfusion damage) occurred in 5 patients (4.09%), and venous thrombosis in 3 patients (2.45%).

Mortality was observed early in the postoperative period in 11 patients (9.01%). The cases of mortality comprised patients who were notably older (64.3 ± 5.26 years), mostly with cardiac problems. However, the age difference between cases with and without mortality was not statistically significant (*P* > 0.05). When patients were admitted to the hospital after 72 hours, mortality was not affected by whether they were admitted between 3 and 7 days or after 7 days (*P* > 0.05). However, morbidity was significantly higher for patients admitted after 7 days (*P* = 0.027). Demographic features related to morbidity and mortality are presented on Table 4.

Table 4 Demographic data related to morbidity and mortality

	N	%
Second time embolectomy	37	30.32
Third time embolectomy	9	7.37
Complications		
Fasciotomy	14	11.47
Amputation	9	7.37
Infection	11	9.01
Myoneuropathic metabolic syndrome	5	4.09
Venous thrombosis	3	2.45
Mortality	11	9.01
Causes of mortality		
Myocardial infarction	3	2.45
Congestive heart failure	2	1.63
Cerebral emboli	1	0.81
Kidney failure	1	0.81
Multiorgan failure	2	1.63
Arrhythmia	1	0.81
Gastrointestinal bleeding	1	0.81

Discussion

We analyzed patients admitted to our clinic with acute arterial occlusion who have been operated on in the last 5 years. Delayed use of embolectomy on the extremity with ischemia is still controversial. However, it has been gaining acceptance. Studies in which clinical and pathological data of late arterial embolectomies are investigated have shown that there are four major factors for successful treatment: 1) relatively less damage on arterial intima; 2) thrombus not adhering to intima and no secondary thrombus on intima; 3) patent distal arterial tree despite embolization; and 4) presurgical treatment with anticoagulants. When these factors are maintained, adequate circulation can be achieved in the extremities of patients with the use of delayed embolectomy.⁴

Critical leg ischemia is defined as a sudden decrease in leg perfusion, which threatens viability of the extremity. This condition not only results in local ischemia, but also to a pathology that can lead to systemic complications. It is the reason for 10% to 15% of surgeries performed in vascular surgery clinics and involves 5% to 7% of patients admitted to these clinics with vascular symptoms.⁵

Despite the improved intensive care and surgical conditions, mortality rates remain high.⁶ Mortality and morbidity risks persist even when the reason for acute ischemic condition is eliminated and reperfusion is attained. When surgical intervention is delayed, these risks increase.⁷ Approximately a hundred years have passed since the first

successful embolectomy performed by Georges Labey in 1911.⁸ The embolectomy catheters devised by Fogarty have facilitated this operation and increased its success.⁹

Our classical knowledge states that the earlier the embolectomy (especially when it is within the first 8 hours), the better the result. However, cases of successful embolectomies performed a few days after an acute occlusion have also been reported.^{2,10–12} Eliot et al² showed that for patients with arterial emboli, a delay of more than 8 hours increases ischemic complications, but when the delay exceeds 7 days, the occurrence of complications decreases.

On the other hand, Blaisdell et al¹³ findings contradict those of Eliot and suggests that a single high-dose heparin therapy course be employed (starting with a 20,000 U bolus and continuous perfusion of 2000 to 4000 U/h). According to his view, embolectomy should be performed only on patients with less than 8 hours of ischemia, who are low risk. Cardiac diseases, most commonly mitral narrowness and atrial fibrillation, are the reason for 80% to 90% of acute arterial emboli. Embolus is found on the left ventricle apical thrombus, which forms after anterior transmural infarction, at a rate of 5%. Cardiac mixoma, mechanical heart valves, and grafts are other possible reasons for emboli.^{14,15} Farshad et al¹⁶ stated that 10% to 15% of emboli as not cardiac-related. Cranley et al¹⁷ could not find the cause for emboli in 10% of the cases. However, in our study, the causes for the emboli were 64.75% cardiac, 28.68% extracardiac, and 9.04% other reasons; and, unlike Cranley et al, we could not find the reason for emboli in only 2 cases (1.63%).

Compartment syndrome caused by reperfusion after embolectomy must be carefully assessed. Early fasciotomy is required to prevent ischemia and complications that may occur. The time between the onset of ischemia and treatment, the cause of embolization, localization of the emboli, and accompanying pathologies are reported to be important for therapeutic success.¹⁸

Free oxygen radicals, which emerge after re-establishment of blood flow in an extremity that has remained ischemic for a long time, interact with endothelium and neutrophils, and rapidly increase lipid peroxidation, resulting in many local and systemic affects. Cellular swelling, edema, toxin and myoglobin release, accompanied by the effects of free oxygen radicals may cause systemic damages such as acute kidney failure, lung edema, sudden onset of adult respiratory distress syndrome and liver shock. Even when extremity reperfusion has been completely restored by removal of the underlying cause of the acute ischemic condition, a chain of events may result in loss of the extremity, acute kidney

and respiratory failure, or functional deterioration in tissues such as heart, intestine, brain or spleen.¹⁹ This “reperfusion damage”, or “myonephrotic metabolic syndrome” as suggested by Haimovici, is the main cause of mortality and morbidity especially in acute cases treated late.²⁰

In various studies, systemic organ damage has been shown to appear as a result of ischemia in the skeletal muscle after 6 to 8 hours of ischemia and as a result of the production of ischemic metabolites, myoglobulinemia and myoglobulinuria, after revascularization.^{19–22} Aune et al²³ reported that both serious cardiac disease and reperfusion damage increase mortality in cases of embolism. In our study, 5 patients developed myonephrotic metabolic syndrome by production of ischemic metabolites, and by myoglobinemia and myoglobulinuria, after revascularization. However, we were prepared for this complication and had taken the necessary precautions and immediately started dialysis. Among these 5 patients, only 1 was lost.

The amputation rate following operations undertaken during the first 12 hours is reported in the range of 2.1% to 5.9%; but the rate increases to 39.2% after 12 hours.^{16,17} Mortality rates have reported as 12.5% within 12 hours and 37.7% thereafter.²⁴ Hight et al²⁵ analyzed 11 different series between 1954 and 1974 and determined an amputation rate of 4% to 48% and a mortality rate of 14% to 50%. In a study by Yangni et al²⁶ that included 24 patients, the amputation rate was reported as 29.2% and the mortality rate as 29.2%. In another study, the amputation rate of arterial embolectomies beyond 24 hours was reported to be 11.4%.¹⁵

In our study, fasciotomy was performed for 11.47% of the patients, those who had developed compartment syndrome. For 7 patients who had ongoing ischemia despite fasciotomy and who had developed a demarcation line and for 2 patients with graft infections, amputation was performed (altogether 7.37%) because of life-threatening risks. Although mortality rates of patients who have received embolectomy 12 hours later is reported to be high in the literature, only 11 patients died (9.01%) in our study, mostly older (average age, 64.3 ± 5.26 years) patients, mostly with cardiac problems. In addition, we think that postoperative heparin and dextran 40 infusion for at least 3 days is the reason for the lower morbidity and mortality rates observed in our study cohort.

Early diagnosis and treatment of acute arterial emboli play an important role in morbidity and mortality. Considering our morbidity and mortality rates, we think that performing an additional surgical operation in acute leg ischemia would be appropriate even if a week has passed.

Conclusion

We believe that late embolectomies of acute leg ischemia increase blood flow in the extremity and reduce the number of amputations required. We also think it is advantageous to give the patient a chance by performing embolectomy, even as late as a week after diagnosis of ischemia.

Disclosure

The authors declare no conflicts of interest.

References

1. Braithwaite BD, Davies B, Birch PA, Heather BP, Earnshaw JJ. Management of acute leg ischemia in the elderly. *Br J Surg*. 1988; 85:217–220.
2. Elliot JP, Hageman JH, Szilagyi DE, Ramakrishnan V, Bravo JJ, Smith RF. Arterial embolization: Problems of source, multiplicity, recurrence and delayed treatment. *Surgery*. 1980;88:833–845.
3. Rutherford RB, Baker JD, Ernst C, et al. Recommended standards for reports dealing with lower extremity ischemia. *J Vasc Surg*. 1997;26:517–538.
4. Haimovici H, Ascer E, Holier HL, Strandness DE, Towne JB. Peripheral arterial aneurysms. In: Haimovici's Vascular Surgery. Principal and Techniques (eds: Haimovici H), 4th ed. Blackwell Science Cambridge 1996:893–910.
5. Ouriel K. Acute limb ischemia. In: Vascular surgery. Rutherford RB ed. 6th ed. Philadelphia: Elsevier; 2005:959–986.
6. Patel N, Sacks D, Patel RI, et al. SCVIR reporting standards for the treatment of acute limb ischemia with use of transluminal removal of arterial thrombus. *J Vasc Interv Radiol*. 2001;12:559–570.
7. Hobson RW II, Milazzo VJ, Duran WN. Pathophysiology of skeletal muscle ischemia-reperfusion injury. In: Haimovici's Vascular Surgery Principal and Techniques. Haimovici H ed. 4th ed. Cambridge: Blackwell Science; 1996:497–508.
8. Cranley JJ, Krause RJ, Strasser ES, Hafner CD, Fogarty TJ. Peripheral arterial embolism: Changing concepts. *Surgery*. 1964;55: 57–63.
9. Fogarty TJ. Management of arterial emboli. *Surg Clin North Am*. 1979;59:749–753.
10. Shifrin EG, Anner H, Eid A, Romanoff H. Practice and theory of ‘delayed’ embolectomy. *J Cardiovasc Surg*. 1986;27:553–556.
11. Cambria RP, Ridge BA, Brewster DC, Moncure AC, Darling RC, Abbott WM. Delayed presentation and treatment of popliteal artery embolism. *Ann Surg*. 1991;214:50–55.
12. Jarrett GC, Dacumos GC, Crummy AB, et al. Late appearance of arterial emboli: diagnosis and management. *Surgery*. 1979;86: 898–905.
13. Blaisdell FW, Steele M, Allen RE. Management of acute lower extremity arterial ischemia due to embolism and thrombosis. *Surgery*. 1978;84:822–825.
14. Erentug V, Mansuroglu D, Bozbuga NU, et al. Treatment of acute arterial occlusions. [in Turkish]. *Turkish J Thorac Cardiovasc Surg*. 2003;11:236–239.
15. Karapolat S, Dag O, Abanoz M, Aslan M. Arterial embolectomy: a retrospective evaluation of 730 cases over 20 years. *Surg Today*. 2006;36:416–419.
16. Abir F, Barkhordorian B, Sumpio BE. Non cardiac vascular complications of coronary bypass procedures: A review. *Int J Angiol*. 2004;13:1–6.
17. Cranley JJ, Krause RJ, Strasser ES, Hafner CD, Fogarty TJ. Peripheral arterial embolism: Changing concepts. *Surgery*. 1964;55:57–63.
18. Panetta T, Thompson JE, Talkington CM, Garrett WV, Smith BL. Arterial embolectomy: A 34-year experience with 400 cases. *Surg Clin North Am*. 1986;66:339–353.

19. Haimovici H. Metabolic complications of acute arterial occlusions and skeletal muscle ischemia: Myonephropathic-metabolic syndrome. In: *Haimovici's Vascular Surgery. Principal and Techniques*. Haimovici H ed. 4th ed. Cambridge: Blackwell Science; 1996: 509–530.
20. Parrino PE, Laubach VE, Gaughen JR Jr, et al. Inhibition of inducible nitric oxide synthase after myocardial ischemia increases coronary flow. *Ann Thorac Surg*. 1998;66:733–739.
21. Duprez D. Natural history and evolution of peripheral obstructive arterial disease. *Int Angiol*. 1992;11:165–168.
22. Stoney RJ, Thompson RW, Nelken NA. Surgical recanalization of occluded peripheral arteries. *Surg Clin North Am*. 1992;72:749–756.
23. Annue S, Trippestad A. Operative mortality and Long term survival of patients operated on for acute lower limb ischemia. *Eur J Endovasc Surg*. 1998;15:143–146.
24. Taviloğlu K, Günay K, Asaoğlu O, Güloğlu R, Kurtoglu M. Is primary amputation required in peripheral arterial occlusion cases that are admitted to the clinic with 12-hour delay? [in Turkish]. *Turkish J Vasc Surg*. 1995;4:91–96.
25. Hight DW, Tilney NL, Couch NP: Changing clinical trends with arterial emboli. *Surgery*. 1976;79:172–176.
26. Yangni AH, Adoubi A, Adoh M, Yapobi Y, Coulibaly AO. Acute nontraumatic limb ischemia. *West Afr J Med*. 2006;25:101–104.

Vascular Health and Risk Management

Publish your work in this journal

Vascular Health and Risk Management is an international, peer-reviewed journal of therapeutics and risk management, focusing on concise rapid reporting of clinical studies on the processes involved in the maintenance of vascular health; the monitoring, prevention and treatment of vascular disease and its sequelae; and the involvement of

metabolic disorders, particularly diabetes. This journal is indexed on PubMed Central and MedLine. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/vascular-health-and-risk-management-journal>

Dovepress