Antimicrobial resistance pattern and molecular genetic distribution of metallo- β -lactamases producing *Pseudomonas aeruginosa* isolated from hospitals in Minia, Egypt

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Sara M Farhan¹
Reham A Ibrahim²
Khaled M Mahran³
Helal F Hetta^{4,5}
Rehab M Abd El-Baky^{1,2}

¹Department of Microbiology and Immunology, Faculty of Pharmacy, Deraya University, Minia 11566, Egypt;

²Department of Microbiology and Immunology, Faculty of Pharmacy, Minia University, Minia 61519 Egypt; ³General Surgery and Laparoscopic Surgery, Faculty of Medicine, Minia University, Minia 61519 Egypt; ⁴Department of Medical Microbiology and Immunology, Faculty of Medicine, Assiut University, Assiut 71515, Egypt; ⁵Department of Internal Medicine, University of Cincinnati College of Medicine, Cincinnati, OH, USA

Background: *Pseudomonas aeruginosa* (*P. aeruginosa*) represents a great threat to public health worldwide, due to its high ability to acquire resistance to different antibiotic classes. Carbapenems are effective against multidrug resistant (MDR) *P. aeruginosa*, but their widespread use has resulted in the emergence of carbapenem-resistant strains, which is considered a major global concern. This study aimed to determine the prevalence of carbapenem resistance among *P. aeruginosa* strains isolated from different sites of infection.

Methods: Between October 2016 and February 2018, a total of 530 clinical specimens were collected from patients suffering from different infections, then processed and cultured. Isolates were tested for extended spectrum β -lactamase (ESBL) and metallo- β -lactamase (MBL) production using double-disk synergy test, modified Hodge tests, and disc potentiation test. PCR was used for the detection of selected OXA carbapenemases encoding genes.

Results: Of 530 samples, 150 (28.3%) *P. aeruginosa* isolates were obtained. MDR strains were found in 66.6% (100 of 150) of isolates. Of 100 MDR *P. aeruginosa* isolates, 54 (54%) were ESBL producers and 21 (21%) carbapenem resistant *P. aeruginosa*. MBL production was found in 52.3% (eleven) carbapenem-resistant isolates. CTX-M15 was found among 55.5% of ESBL-producing *P. aeruginosa*. Carbapenemase genes detected were *bla*_{IMP} (42.8%, nine of 21), *bla*_{VIM} (52.3%, eleven of 21), *bla*_{GIM} (52.3%, eleven of 21), *bla*_{SPM} (38%, 8/21). In addition, isolates that were positive for the tested genes showed high resistance to other antimicrobials, such as colistin sulfate and tigecycline.

Conclusion: Our study indicates that *P. aeruginosa* harboring ESBL and MBL with limited sensitivity to antibiotics are common among the isolated strains, which indicates the great problem facing the treatment of serious infectious diseases. As such, there is a need to study the resistance patterns of isolates and carry out screening for the presence of ESBL and MBL enzymes, in order to choose the proper antibiotic.

Keywords: MDR; *P. aeruginosa*, ESBL, MBL, antimicrobial resistance

Introduction

Pseudomonas aeruginosa is an opportunistic pathogen that can cause outbreaks of hospital-acquired and life-threatening infections, especially among immunocompromised and critically ill patients.¹ *P. aeruginosa* can cause respiratory tract, burn, wound infections and otitis media.² *P. aeruginosa* infections are commonly associated with high mortality, attributed to its intrinsic resistance to many classes of antimicrobial agents and ability to acquire resistance by mutation and horizontal

Correspondence: Helal F Hetta
Department of Internal Medicine,
University of Cincinnati College of
Medicine, 231 Albert B. Sabin Way, PO
Box 670595, Cincinnati, OH 45267-0595,
USA
Email helal.hetta@uc.edu

transfer of resistance determinants.³ The rapid emergence of penicillin and cephalosporin resistance among *P. aeruginosa* strains has become a serious clinical problem worldwide. Carbapenems (imipenem and meropenem), potent antipseudomonal drugs, have been used as the last resort for the treatment of infections associated with multi-drug resistant (MDR) *P. aeruginosa* isolates.⁴ Resistance to carbapenems has developed through decreased permeability, overexpression of the efflux-pump system, alterations in penicillin-binding protein and carbapenem-hydrolyzing enzymes (carbapenemases).⁵

Carbapenemases represent three classes of β -lactamase (BL). Ambler class A and D (serine carbapenemases) and class B (zinc-dependent). These enzymes require zinc for their catalytic activity and are inhibited by metal chelators, such as EDTA and thiol-based compounds, and are called metallo-BLs (MBLs). MBL enzymes are able to hydrolyze all β -lactam antibiotics, with the exception of monobactams. The genes encoding these enzymes have found to be carried on highly mobile elements, which is the main cause of their dissemination in the hospital environment. MBLs are mainly plasmid-mediated and in some cases chromosomally mediated. The most common MBLs enzymes belong to the Verona integron-encoded MBL (VIM), imipenemase (IMP), São Paulo MBL (SPM), German imipenemase MBL (GIM), Seoul imipenemase MBL, and New Delhi MBL families.⁶

Infections caused by MBL-producing organisms are associated with high morbidity and mortality rate, especially in hospitalized and immunosuppressed patients.⁷ Recently, many studies reported the prevalence of *P. aeruginosa* strains harboring both extended-spectrum BL (ESBL) and MBL genes, which is considered a great challenge for antimicrobial therapy.⁸ In addition, it is difficult to detect ESBLs phenotypically.⁹ As such, molecular techniques are required to analyze the coexistence of carbapenemases and ESBLs in the same strain. The aim of this study was to study the prevalence and DR profile of carbapenem-resistant *P. aeruginosa* (CRPA) isolates obtained from hospitalized patients with various infections

Methods

Bacterial isolates

A total of 150 (28.3%) *P. aeruginosa* isolates were isolated from 530 samples collected from hospitalized patients with various infections as part of routine hospital-laboratory procedures. Samples were processed and cultured on

blood agar at 37°C and 42°C for 24 hours. One colony was picked and subcultured on MacConkey agar plates and cetrimide agar. Isolated colonies were further identified according to colony morphology, lactose fermentation, and biochemical characteristics (oxidase, triple sugar iron, urease tests, sulphide–indole–motility). Colonies were able to grow on cetrimide agar, show positive reactions on catalase and oxidase tests, grow at 42°C (used to distinguish *P. aeruginosa* from other lactose nonfermenting Gram-negative rods), and show negative results in triple-sugar iron and glucose-fermentation tests.^{10,11} *P. aeruginosa* colonies were purified by streaking, and pure colonies were stored at 4°C.

Antimicrobial-susceptibility testing

Antimicrobial susceptibility was determined by Kirby–Bauer disk diffusion test.¹² Results were assessed on the basis of Clinical and Laboratory Standards Institute criteria. The following antimicrobial disks (Oxoid, Basinstoke, UK) were used: azlocillin (75 μ g), ciprofloxacin (5 μ g), ampicillin–sulbactam (20 μ g), levofloxacin (5 μ g), cefepime (30 μ g), meropenem (10 μ g), aztreonam (30 μ g), imipenem (10 μ g), polymyxin B (300 μ g), colistin sulfate (10 μ g), tigecycline (15 μ g), tobramycin (10 μ g), ceftazidime (30 μ g), amoxicillin–clavulanic (20/10 μ g), carbenicillin (100 μ g), amikacin (30 μ g), gentamicin (10 μ g), piperacillin (100 μ g), and cefoperazone (75 μ g).

Phenotypic detection of ESBL production

Detection of ESBL production by *P. aeruginosa* strains was performed by double-disk synergy test (DDST).¹³ Disks of ceftazidime, cefotaxime, aztreonam, and cefepime (30 μ g each) were placed at a distance of 30 or 20 mm (center to center) from an amoxicillin 20 μ g–clavulanic acid 10 μ g disk. Increase in zones of inhibition toward amoxicillin–clavulanic acid antibiotic disks is indicative of the presence of ESBL.

Phenotypic detection of MBL production¹⁴

Imipenem–EDTA combined disk synergy testing was used for identification of MBL-producing isolates according to Lee et al.^{14,15} A solution of 0.5 M EDTA (pH 8) was prepared by dissolving 18.61 g of EDTA in 100 mL distilled water and adjusting its pH to 8 using NaOH, then, autoclaving. The tested organisms were cultured on the surface of Müller–Hinton agar plates. Two 10 μ g imipenem disks or two 10 μ g meropenem disks were placed on the surface of agar plates and 5 μ L EDTA solution added

to one imipenem and one meropenem disks. Zones of inhibitions around discs with EDTA were examined after 16–18 hours' incubation at 35°C and compared to those without EDTA. An increase in zone diameter of at least 7 mm around the imipenem–EDTA disc and meropenem–EDTA disks were considered positive results.

Amplification of ESBL-CTX-M15 and MBL genes

Boiling was used to prepare DNA templates of genes. Specific primers — cefotaximase (*bla*_{CTX-M15}), *bla*_{VIM}, *bla*_{IMP}, *bla*_{GIM}, and *bla*_{SPM} (Table 1) — were used for PCR amplification of the genes. PCR amplification was done using 25 µL reaction mixture containing 0.2 µL Taq polymerase 5 U/µL 1 pmol of each forward and reverse primer, 2.5 µL dNTP mix (2 Mm), 3 µL DNA template, and 14.8 µL DNase-free and RNase-free water. PCR reactions were performed using a Mastercycler personal 5332 (Eppendorf, Hamburg, Germany). Amplified products were analyzed by electrophoresis in 2% agarose gel at 80 V for 45 minutes in Tris–Borate–EDTA buffer containing ethidium bromide under ultraviolet irradiation.^{16,17}

Results and discussion

P. aeruginosa is commonly associated with hospital-acquired infections. With regard to the specimen site, of 530 samples, 150 (28.3%) were positive for *P. aeruginosa*, which was similar to results reported by Al-Haik et al¹⁸ and Mansour et al¹⁹ and fewer than Gad et al.²⁰ *P. aeruginosa* isolates were isolated from 65 of 332 (19.5%) wound swabs, 39 of 57 (68.4%) ear swabs, five of 26

(19.2%) burn swabs, six of 30 (20%) urine samples, eight of 12 (66.6%) sputum samples, eight of 35 (22.8%) stool samples, 19 of 38 (50%) of patients admitted to the intensive-care unit (ICU). Our results showed high incidence (68.4%) of *P. aeruginosa* among samples collected from patients suffering from otitis media, which was higher than reported by Umar et al,²¹ who found that 23.2% of samples of otitis media were positive for *P. aeruginosa*. The distribution of isolates across major hospitals in Minia Governorate was analyzed. High incidence of *P. aeruginosa* was observed among samples collected from the chest hospital, while all samples obtained from Minia General Hospital were negative for *P. aeruginosa* (Figure 1).

P. aeruginosa possesses MDR against a wide variety of antibiotics. Resistance of *P. aeruginosa* is usually accompanied by the production of many BLs, active expulsion of antibiotics by efflux pump, and alteration of outer-membrane protein expression.^{9,22} Resistance to variety of β-lactam antibiotics is a growing problem, due to their continuous mutation, which makes BLs production the most common cause of DR and antimicrobial therapy failure.²³ Among BLs, ESBLs are widely distributed among Enterobacteriaceae members. They are also found in *Acinetobacter baumannii* and *P. aeruginosa*. At first, TEM-type ESBLs and SHV-type ESBLs were the most dominant among Gram negative isolates in Europe and other regions. Since the last decade, CTX-M type ESBL has become the most prevalent.

ESBL production is widely spread among Enterobacteriaceae, especially *P. aeruginosa*. Our study showed that all *P. aeruginosa* isolates were completely resistant to azlocillin and amoxicillin–clavulanic acid. Of 150

Table 1 PCR primers used for detection of ESBL-CTX-M15 and MBL genes in *Pseudomonas aeruginosa*

Gene	Primers	Sequence	Product size
ESBL-CTXM15	CTX-M15-F CTX-M15-R	5'-CGTCACGCTGTTGTTAGGAA-3' 5'-ACGGCTTTCTGCCTTAGGTT-3'	780 bp
<i>bla</i> _{VIM}	VIM-F VIM-R	5'-GATGGTGTG TGG TCG CAT A-3' 5'-CGA ATG CGC AGC ACC AG-3'	390 bp
<i>bla</i> _{IMP}	IMP-F IMP-R	5'-GGAATAGAGTGGCTTAATTCTC3' 5'-CCAAACCACTACGTTATCT-3'	188 bp
<i>bla</i> _{GIM}	GIM-F GIM-R	5'-TCG ACA CAC CTT GGT CTG AA 3' 5'-AAC TTC CAA CTT TGC CAT GC-3'	477 bp 271 bp
<i>bla</i> _{SPM}	SPM-F SPM-R	5'-AAA ATC TGG GTA CGC AAA CG-3' 5'-ACA TTA TCC GCT GGA ACA GG-3'	

Abbreviations: ESBL, extended-spectrum β-lactamase; MBL, metallo-β-lactamase; IMP, imipenemase; VIM, Verona integron–encoded MBL; GIM, German imipenemase MBL; SPM, São Paulo MBL.

Figure 1 Prevalence of *Pseudomonas aeruginosa* isolated from different hospitals in Minia.

P. aeruginosa isolates, 100 (66.6%) were MDR and 21 (21%) of these were CRPA (eleven isolates were imipenem-resistant and ten meropenem-resistant). Figure 2 shows that 46%, 28.7%, and 28% of *P. aeruginosa* were resistant to polymyxin B, colistin sulfate, and tigecycline, respectively.

In this study, it was found that 54 (54%) isolates of MDR *P. aeruginosa* were ESBL producers. Similarly high production of ESBL was reported by Ahmad et al,²⁴ who reported that ESBL production by *P. aeruginosa* isolates was 61.6%, while lower incidence (27.33%) was reported by Dutta et al.²⁵ In addition, our results showed that eleven (11%) isolates were MBL-producing *P. aeruginosa*. Furthermore, MBL-producing strains represented 52.3% (eleven of 21) of CRPA isolates. Coexistence of ESBL and MBL was found among 5% of MDR *P. aeruginosa* and five of 21 (23.8%) CRPA isolates. Antibiotic-resistance patterns of ESBL-producing strains revealed that all ESBL producers were completely resistant to azlocillin, amoxicillin-clavulanic acid, ampicillin/sulbactam and cefipime. Co-resistance

with other antibiotics was observed including colistine sulfate, tigecycline, and polymyxin B (Table 2). Also, MBL-producing strains showed high resistance to cefipime and carbenicillin (72.7% each), but lower resistance was observed against ciprofloxacin, colistin sulfate, and levofloxacin (36.3% each). Ilyas et al²⁶ showed higher incidence of antibiotic resistance exhibited by MBL- and ESBL-producing *P. aeruginosa*. They reported that ESBL- and MBL-producing *P. aeruginosa* isolates were completely resistant to amoxicillin-clavulanic acid, ceftriaxone, ciprofloxacin, and cefipime. Also, they showed higher incidence of MBL-producing *P. aeruginosa* (25.7%) and lower incidence of ESBL production (8.5%). Mirsalehian et al²⁷ found that all MBL-producing *P. aeruginosa* were colistin-sensitive and 37.5% were resistant to aztreonam, while in the present study low incidence of resistance to colistin, ciprofloxacin, and levofloxacin (36.4%) and a resistance rate of 54.5% were reported against aztreonam. Bashir et al²⁸ reported that all MBL-producing *P. aeruginosa* isolates were resistant to gentamicin, ceftazidime, carbenicillin, tobramycin, ceftriaxone, ofloxacin, cefoperazone, cefoperazone-sulbactam, and ceftazidime-clavulanic acid and low resistance to polymyxin B.

The rapid spread and the emergence of MBL- and ESBL-producing *P. aeruginosa* isolated from hospitals is of great concern and threat. In addition, differences in resistance patterns among strains isolated from different countries may be attributed to antibiotic use, horizontal gene transfer, and environmental conditions. Therefore, it is important to test isolates for MBL and ESBL production and to test for antibiotic susceptibility before antimicrobial therapy.

Figure 2 Resistance pattern of *Pseudomonas aeruginosa* isolates to different antimicrobial agents.

Table 2 Antibiotic-resistance patterns of ESBL- and MBL-producing strains

	% Resistance	
	ESBL-producing <i>P. aeruginosa</i> (n=54)	MBL-producing <i>P. aeruginosa</i> (n=11)
Azlocillin	100	100
Amoxicillin-clavulanic acid	100	100
Ampicillin/sulbactam	100	100
Tobramycin	81.5	45.5
Aztreonam	83.3	54.5
Cefepime	100	72.7
Polymyxin B	74.1	54.5
Cefoperazone	90.8	54.5
Colistin sulfate	37	36.4
Levofloxacin	55.6	36.4
Amikacin	55.6	63.6
Ciprofloxacin	35.2	36.4
Piperacillin	51.9	54.5
Gentamicin	46.3	63.6
Imipenem	5.6	100
Meropenem	7.4	100
Carbenicillin	46.3	72.7
Tigecycline	42.6	45.5

Abbreviations: ESBL, extended-spectrum β -lactamase; MBL, metallo- β -lactamase; *P. aeruginosa*, *Pseudomonas aeruginosa*.

Table 3 shows that the highest incidence of ESBL production was observed among MDR *P. aeruginosa* samples isolated from ear infections (80%), followed by those isolated from chest infections (75%), and ICU patients (70%). The highest incidence of MBL production was observed among MDR *P. aeruginosa* samples isolated

from wound infections (19%) followed by those isolated from ear infections (14.3%). Nithyalakshmi et al²⁹ reported that the frequency of occurrence of ESBL among *P. aeruginosa* isolates was 21.96%, and most ESBL producers were obtained from urine samples (27.7%), followed by respiratory infection (23.68%), and wound infection (22.95%).

All MDR *P. aeruginosa* isolates were tested for CTX-M15 and carbapenem-resistance genes: *bla*_{IMP}, *bla*_{VIM}, *bla*_{GIM}, and *bla*_{SPM}. It was found that 55.5% (30 of 54) of ESBL-producing *P. aeruginosa* isolates were harboring CTX-M15, which was higher than another study¹⁷ reporting that out of 200 MDR *P. aeruginosa* isolates, 19 were positive for CTX-M15, of which 64.28% were ESBL-positive. Although carbapenem resistance was found among 21 *P. aeruginosa* isolates, only eleven were found to harbor MBL genes. Of 21 carbapenem-resistant strains, 42.8% (nine of 21) were positive for *bla*_{IMP}, 52.3% (eleven of 21) positive for *bla*_{VIM}, 52.3% (eleven of 21) positive for *bla*_{GIM}, and 38% (eight of 21) positive for *bla*_{SPM}.

The distribution of carbapenem-resistance genes and *bla*_{CTX-M15} among MDR *P. aeruginosa*-producing ESBL and/or MBL isolates were tested (Table 4). Of eleven MBL-producing MDR *P. aeruginosa*, three (27.2%) were CTX-M15, nine (81.8%) positive for *bla*_{IMP}, four (36.3%) for *bla*_{VIM}, five (45.4%) for *bla*_{SPM} and six (54.5%) for *bla*_{GIM}. Lower incidence was found by Zubair et al³⁰ who reported that among 22 isolates positive for MBL production phenotypically, only five were harboring MBL genes. Furthermore, they reported that *bla*_{VIM} was the predominant gene, and none of the other genes were detected.

Also, It was found that 55.1% of ESBL/non-MBL-producing MDR *P. aeruginosa* isolates were positive for

Table 3 Distribution of ESBL- and MBL-producing isolates among MDR *Pseudomonas aeruginosa* isolates from different clinical specimens

Type of infection	MDR <i>P. aeruginosa</i>	ESBLs		MBL	
		n	%	n	%
Wound	59	24	40%	4	19%
Ear	20	16	80%	3	14.3%
Burns	2	1	50%	—	—
Urinary tract	2	1	50%	—	—
Chest	4	3	75%	1	4.8%
Gastroenteritis	3	2	66.7%	1	4.8%
Patients admitted to ICU (from buccal cavity, skin swab, and eye swab)	10	7	70%	2	9.5%
Total	100	54	54%	11	11%

Note: Percentages correlated with number of MDR *P. aeruginosa* isolates from each type of infection.

Abbreviations: ICU, intensive-care unit; MDR, multidrug-resistant; ESBL, extended-spectrum β -lactamase; MBL, metallo- β -lactamase.

Table 4 Distribution of different groups of carbapenem-resistance genes in phenotypically positive ESBL- and MBL-producing *Pseudomonas aeruginosa* isolates

	<i>bla</i> _{CTX-M-15} positive isolates	<i>bla</i> _{IMP}	<i>bla</i> _{VIM}	<i>bla</i> _{GIM}	<i>bla</i> _{SPM}
	n (%)	n (%)	n (%)	n (%)	n (%)
ESBL ⁺ /MBL ⁻ , n=49	27 (55.1)	0 (0)	7 (14.2%)	3 (6.1)	2(4)
ESBL ⁻ /MBL ⁺ , n=6	0 (0)	6 (100)	2 (33.3)	3 (50)	2 (33.3)
MBL ⁺ /ESBL ⁻ , n=5	3 (60)	3 (60)	2 (40)	3 (60)	3 (60)

Note: *Percentages were correlated with number of isolates positive for ESBL, MBL, or both.

Abbreviations: ESBL, extended-spectrum β -lactamase; MBL, metallo- β -lactamase; IMP, imipenemase; VIM, Verona integron-encoded MBL; GIM, German imipenemase MBL; SPM, S>o Paulo MBL.

CTX-M15, while none of these strains was found to harbor *bla*_{IMP}. On the other hand, *bla*_{VIM} was the most common carbapenem-resistance gene (14.2%). Rafiee et al³¹ and Laudy et al³² showed that all ESBL-producing isolates were negative for CTX-M gene, while Ahmed et al³³ reported a lower incidence of *bla*_{CTX-M} production (10.7%) among *P. aeruginosa* strains isolated from Makkah hospitals. All MBL/non-ESBL-producing *P. aeruginosa* harbored *bla*_{IMP}-like genes and 50% were positive for *bla*_{GIM}, while 33.3% only were positive for both *bla*_{VIM} and *bla*_{SPM} (Table 4). Similar findings were shown by Abiri et al³⁴. In contrast, Mirsalehian et al²⁷ reported that *bla*_{VIM} was the most prevalent carbapenemase gene among MBL-producing *P. aeruginosa*, while 25% of MBL isolates were positive for *bla*_{IMP} and all MBL isolates negative for *bla*_{GIM} and *bla*_{SPM}. Our results showed that five isolates of MDR *P. aeruginosa* were ESBL and MBL coproducers. Three isolates (60%) were found to have *bla*_{CTX-M15}, *bla*_{IMP}, *bla*_{GIM}, and *bla*_{SPM}, and two (40%) were positive for *bla*_{VIM}.

MDR *P. aeruginosa* samples were classified into seven groups according to the number of carbapenem-resistant genes harbored by MBL-producing *P. aeruginosa* isolates, in order to study their demographic, phenotypic, and genotypic features: group A comprised MBL-producing *P. aeruginosa* isolates harboring two genes (*bla*_{IMP} and *bla*_{GIM}), group B isolates positive for *bla*_{IMP}, *bla*_{VIM}, and *bla*_{SPM}, group C including those which were positive for *bla*_{IMP}, group D isolates positive for *bla*_{IMP} and *bla*_{SPM}, group E isolates positive for *bla*_{IMP}, *bla*_{GIM}, and *bla*_{SPM}, Group F isolates positive for *bla*_{IMP} and *bla*_{VIM}, group G MBL-producing *P. aeruginosa* isolates positive for *bla*_{VIM}, *bla*_{GIM}, and *bla*_{SPM}, and group H including isolates positive for *bla*_{VIM} and *bla*_{GIM} (Table 5).

Our study showed that all MBL-producing *P. aeruginosa* isolates in groups A–H were obtained from Minia University Hospital except one isolate that had been obtained from a chest hospital. Of eleven MBL-producing *P. aeruginosa*, five

were ESBL producers and obtained from surgery and ICU units of Minia University Hospital. Of these, three (two from surgery unit and one from ICU) were positive for CTX-M15 gene. The isolate obtained from the ICU unit showed resistance to meropenem, polymyxin B, tigecycline, gentamicin, amikacin, and ceftazidime, which represents a great challenge for antimicrobial therapy patients. The other two isolates (surgery unit) showed resistance to gentamicin, ceftazidime, meropenem, imipenem, tigecycline, and colistin sulfate. Furthermore, the isolate obtained from the chest hospital belonged to group H, was positive for ESBL but negative for CTX-M15, and showed resistance to ceftazidime, cefoperazone, gentamicin, amikacin, tigecycline, polymyxin B, and meropenem. Chaudhary et al³⁵ found that the frequency of *bla*_{IMP} and *bla*_{VIM} among MBL-producing strains was 28.73% and 47.12%, respectively. Coexistence of MBL and ESBL was found among 14.3% of isolates, of which 17.5% were positive for TEM and IMP genes and 14.8% positive for AMP-C and VIM. Also, they found that isolates coproducing ESBL and MBL were highly resistant to cefepim, piperacillin–tazobactam, ceftazidime, meropenem, and imipenem.

Our study showed the prevalence of ESBL- and MBL-producing *P. aeruginosa* with limited sensitivity to antibiotics among the isolated strains, which indicates the great problem in the treatment of serious infectious diseases. In addition, there is a need to study resistance pattern of isolates and carry out screening for the presence of ESBL and MBL enzymes, in order to choose the proper antibiotic.

Study limitations and future recommendations

We detected the distribution of genes only among resistant strains. Quantitative PCR assays are recommended for future studies, and should be performed to verify expression differences of different resistance genes in MDR *P. aeruginosa*.

Table 5 Demographic, phenotypic and genotypic features of MBL-producing *Pseudomonas aeruginosa*

Group	Carbapenem-resistance genes				Sample	Isolate source	Hospital	ESBL production	bla _{CTXM15}	Resistance pattern
	bla _{IMP}	bla _{VIM}	bla _{GIM}	bla _{SPM}						
A	+	–	+	–	PA1 PA2 PA8	ICU Wound swab ICU	Minia University	–, +, –	–, –	CAZ, CN, AK, PB, MEM CAZ, CEP, TGC, MEM CAZ, CN, CT, CIP, IPM
B	+	+	–	+	PA7	Stool	Minia University	–	–	CAZ, CEP, AK, CT, PB, MEM
C	+	–	–	–	PA9	Wound swab	Minia University	+	+	CAZ, CN, TGC, CIP, IPM
D	+	–	–	+	PA11 PA33	Ear discharge Wound swab	Minia University	–, +	–, +	CAZ, CEP, AK, PB, CIP, IPM, CAZ, CEP, CN, CT, MEM
E	+	–	+	+	PA13	ICU	Minia University	+	+	CAZ, CN, AK, TGC, PB, MEM
F	+	+	–	–	PA16	Stool	Minia University	–	–	CAZ, CEP, AK, TGC, CIP, MEM
G	–	+	+	+	PA19	Wound swab	Minia University	–	–	CAZ, CN, AK, CT, PB, IPM
H	–	+	+	–	PA27	Sputum	Chest Hospital	+	–	CAZ, CEP, CN, AK, TGC, PB, MEM

Notes: A, MBL-producing *P. aeruginosa* isolates positive for bla_{IMP} and bla_{GIM}; B, MBL-producing *P. aeruginosa* isolates positive for bla_{IMP}, bla_{VIM}, and bla_{SPM}; C, MBL-producing *P. aeruginosa* isolates positive for bla_{IMP}, bla_{VIM}, and bla_{SPM}; D, MBL-producing *P. aeruginosa* isolates positive for bla_{IMP} and bla_{SPM}; E, MBL-producing *P. aeruginosa* isolates positive for bla_{IMP}, bla_{GIM}, and bla_{SPM}; F, MBL-producing *P. aeruginosa* isolates positive for bla_{IMP} and bla_{VIM}; G, MBL-producing *P. aeruginosa* isolates positive for bla_{VIM}, bla_{GIM}, and bla_{SPM}; H, MBL-producing *P. aeruginosa* isolates positive for bla_{VIM} and bla_{GIM}.

Abbreviations: ESBL, extended-spectrum β-lactamase; MBL, metallo-β-lactamase; ICU, intensive-care unit; CAZ, Cefazolin, CN, Gentamicin, AK, Amikacin, PB, polymyxin B, MEM, meropenem, TGC, Tigecycline, CIP, cefepime, CT, colistin, CIP, ciprofloxacin, IPM, imipenem; IMP, imipenem; VIM, Verona integron-encoded MBL; GIM, German imipenemase MBL; SPM, S^o Paulo MBL.

Conclusion

Using carbapenems in clinical practice was initially the solution to treatment of serious bacterial infections caused by β -lactam-resistant bacteria. Due to their widespread use, the emergence of MBL-producing strains and strains coproduce both ESBL and MBL was observed. As found in our study, strains showed high resistance to the commonly used antibiotics, which emphasizes the need to know the resistance patterns and testing for the coexistence of these enzymes, in order to design newer policies for antimicrobial chemotherapy.

Abbreviations list

MHT, modified Hodge Test; MDR, multidrug-resistant; ESBL, extended-spectrum β -lactamase; MBL, metallo- β -lactamase.

Disclosure

The authors report no conflicts of interest in this work.

References

- Santajit S, Indrawattana N. Mechanisms of antimicrobial resistance in ESKAPE pathogens. *Biomed Res Int*. 2016;2016:2475067. doi:10.1155/2016/2475067
- Lister PD, Wolter DJ, Hanson ND. Antibacterial-resistant *Pseudomonas aeruginosa*: clinical impact and complex regulation of chromosomally encoded resistance mechanisms. *Clin Microbiol Rev*. 2009;22(4):582–610. doi:10.1128/CMR.00040-09
- Potron A, Poirel L, Nordmann P. Emerging broad-spectrum resistance in *Pseudomonas aeruginosa* and *Acinetobacter baumannii*: mechanisms and epidemiology. *Int J Antimicrob Agents*. 2015;45(6):568–585. doi:10.1016/j.ijantimicag.2015.03.001
- Pitout JD, Gregson DB, Poirel L, McClure JA, Le P, Church DL. Detection of *Pseudomonas aeruginosa* producing metallo-beta-lactamases in a large centralized laboratory. *J Clin Microbiol*. 2005;43(7):3129–3135. doi:10.1128/JCM.43.7.3129-3135.2005
- Kaleem F, Usman J, Hassan A, Khan A. Frequency and susceptibility pattern of metallo-beta-lactamase producers in a hospital in Pakistan. *J Infect Dev Ctries*. 2010;4(12):810–813.
- Owlia P, Saderi H, Karimi Z, Rad A, Bagher SM, Bahar MA. Phenotypic detection of Metallo-beta-Lactamase producing *Pseudomonas aeruginosa* strains isolated from burned patients. *Iran J Pathol*. 2008;3(1):20–25.
- Hong DJ, Bae IK, Jang IH, Jeong SH, Kang HK, Lee K. Epidemiology and characteristics of metallo-beta-lactamase-producing *Pseudomonas aeruginosa*. *Infect Chemother*. 2015;47(2):81–97. doi:10.3947/ic.2015.47.2.81
- Bae IK, Suh B, Jeong SH, et al. Molecular epidemiology of *Pseudomonas aeruginosa* clinical isolates from Korea producing beta-lactamases with extended-spectrum activity. *Diagn Microbiol Infect Dis*. 2014;79(3):373–377. doi:10.1016/j.diagmicrobio.2014.03.007
- Polotto M, Casella T, de Lucca Oliveira MG, et al. Detection of *P. aeruginosa* harboring bla CTX-M-2, bla GES-1 and bla GES-5, bla IMP-1 and bla SPM-1 causing infections in Brazilian tertiary-care hospital. *BMC Infect Dis*. 2012;12. doi:10.1186/1471-2334-12-166
- Fazeli H, Akbari R, Moghim S, Narimani T, Arabestani MR, Ghoddousi AR. *Pseudomonas aeruginosa* infections in patients, hospital means, and personnel's specimens. *J Res Med Sci*. 2012;17(4):332–337.
- Tille P. *Bailey & Scott's Diagnostic Microbiology-E-Book*. Philadelphia: Elsevier Health Sciences; 2015.
- Wayne P. Clinical and Laboratory Standards Institute: performance standards for antimicrobial susceptibility testing: 20th informational supplement. *CLSI document M100-S20*. 2010.
- Tzelepi E, Giakkoupi P, Sofianou D, Loukova V, Kemeroglou A, Tsakris A. Detection of extended-spectrum beta-lactamases in clinical isolates of *Enterobacter cloacae* and *Enterobacter aerogenes*. *J Clin Microbiol*. 2000;38(2):542–546.
- Clinical and Laboratory Standards Institute (2007) *Performance Standards for Antimicrobial Susceptibility Testing; Seventeenth Informational Supplement. CLSI Document M100-S17*. Wayne, PA: Clinical and Laboratory Standards Institute; 2007.
- Lee K, Lim YS, Yong D, Yum JH, Chong Y. Evaluation of the Hodge test and the imipenem-EDTA double-disk synergy test for differentiating metallo-beta-lactamase-producing isolates of *Pseudomonas spp.* and *Acinetobacter spp.* *J Clin Microbiol*. 2003;41(10):4623–4629. doi:10.1128/jcm.41.10.4623-4629.2003
- Ellington MJ, Kistler J, Livermore DM, Woodford N. Multiplex PCR for rapid detection of genes encoding acquired metallo-beta-lactamases. *J Antimicrob Chemother*. 2007;59(2):321–322. doi:10.1093/jac/dkl481
- Ullah W, Qasim M, Rahman H, et al. CTX-M-15 and OXA-10 beta lactamases in multi drug resistant *Pseudomonas aeruginosa*: first report from Pakistan. *Microb Pathog*. 2017;105:240–244. doi:10.1016/j.micpath.2017.02.039
- Al-Haik WM, Al-Mahbush AA, Al-Mahdi AY, Mohamed MM, Al-Haddad A. Genotypic characteristics of clinical and non-clinical isolates of *Pseudomonas aeruginosa*: distribution of different antibiogram profiles and molecular typing. *Jordan J Biol Sci*. 2016;9:3.
- Mansour SA, Eldaly O, Jiman-Fatani A, Mohamed ML, Ibrahim EM. Epidemiological characterization of *P. aeruginosa* isolates of intensive care units in Egypt and Saudi Arabia. *East Mediterr Health J*. 2013;19(1):71–80.
- Gad GF, El-Domany RA, Zaki S, Ashour HM. Characterization of *Pseudomonas aeruginosa* isolated from clinical and environmental samples in Minia, Egypt: prevalence, antibiogram and resistance mechanisms. *J Antimicrob Chemother*. 2007;60(5):1010–1017. doi:10.1093/jac/dkm348
- Umar JB, Ibrahim MM, Tom IM, Umore AM, Isa T. *Pseudomonas aeruginosa* in otitis media. *Int J Med*. 2016;4(2):55–57. doi:10.14419/ijm.v4i2
- Bonfiglio G, Laksai Y, Franchino L, Amicosante G, Nicoletti G. Mechanisms of beta-lactam resistance amongst *Pseudomonas aeruginosa* isolated in an Italian survey. *J Antimicrob Chemother*. 1998;42(6):697–702. doi:10.1093/jac/42.6.697
- Dallenne C, Da Costa A, Decré D, Favier C, Arlet G. Development of a set of multiplex PCR assays for the detection of genes encoding important beta-lactamases in Enterobacteriaceae. *J Antimicrob Chemother*. 2010;65(3):490–495. doi:10.1093/jac/dkp498
- Ahmad SS, Ali FA. Detection of ESBL, AmpC and Metallo Beta-Lactamase mediated resistance in Gram-negative bacteria isolated from women with genital tract infection. *Eur Sci J*. 2014;10(9):193–209.
- Dutta H, Nath R, Saikia L. Multi-drug resistance in clinical isolates of Gram-negative bacilli in a tertiary care hospital of Assam. *Indian J Med Res*. 2014;139(4):643.
- Ilyas M, Khurram M, Ahmad S, Ahmad I. Frequency, susceptibility and co-existence of MBL, ESBL & AmpC positive *Pseudomonas aeruginosa* in Tertiary Care Hospitals of Peshawar, KPK, Pakistan. *J Pure Appl Microbiol*. 2015;9(2):981–988.

27. Mirsalehian A, Kalantar-Neyestanaki D, Taherikalani M, Jabalameli F, Emameini M. Determination of carbapenem resistance mechanism in clinical isolates of *Pseudomonas aeruginosa* isolated from burn patients, in Tehran, Iran. *J Epidemiol Glob Health*. 2017;7(3):155–159. doi:10.1016/j.jegh.2017.04.002
28. Bashir D, Thokar MA, Fomda BA, et al. Detection of metallo-beta-lactamase (MBL) producing *Pseudomonas aeruginosa* at a tertiary care hospital in Kashmir. *Afr J Microbiol Res*. 2011;5(2):164–172.
29. Nithyalakshmi J, Vidhyarani R, Mohanakrishnan K, Sumathi G. ESBL producing *Pseudomonas aeruginosa* in clinical specimens: is it a scary nightmare or paper tiger? *Indian J Microbiol Res*. 2016;3(3):287–291. doi:10.5958/2394-5478.2016.00062.5
30. Zubair K, Iregbu K. Resistance pattern and detection of metallo-beta-lactamase genes in clinical isolates of *Pseudomonas aeruginosa* in a Central Nigeria Tertiary Hospital. *Niger J Clin Pract*. 2018;21(2):176–182. doi:10.4103/njcp.njcp_229_17
31. Rafiee R, Eftekhari F, Tabatabaei SA, Minaee Tehrani D. Prevalence of extended-spectrum and metallo β -lactamase production in AmpC β -lactamase producing *Pseudomonas aeruginosa* Isolates From Burns. *Jundishapur J Microbiol*. 2014;7(9):e16436. doi:10.5812/jjm
32. Laudy AE, Róg P, Smolińska-Król K, et al. Prevalence of ESBL-producing *Pseudomonas aeruginosa* isolates in Warsaw, Poland, detected by various phenotypic and genotypic methods. *PLoS One*. 2017;12(6):e0180121. doi:10.1371/journal.pone.0180121
33. Ahmed OB, Asghar AH, Bahwerth FS. Prevalence of ESBL genes of *Pseudomonas aeruginosa* strains isolated from Makkah Hospitals, Saudi Arabia. *Euro J Biol Med Sci Res*. 2015;3(6):12–18.
34. Abiri R, Mohammadi P, Shavani N, Rezaei M. Detection and genetic characterization of metallo- β -lactamase IMP-1 and VIM-2 in *Pseudomonas aeruginosa* strains from different hospitals in Kermanshah, Iran. *Jundishapur J Microbiol*. 2015;8(9):e22582. doi:10.5812/jjm
35. Chaudhary M, Payasi A. Rising antimicrobial resistance of *Pseudomonas aeruginosa* isolated from clinical specimens in India. *J Proteomics Bioinform*. 2013;6(1):005–009.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>