Quinolones and fluoroquinolones are useless to counter uropathogenic *Escherichia coli* infections [Letter]

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Bhoj R Singh

Division of Epidemiology, The Indian
Council of Agricultural Research-Indian
Veterinary Research Institute, Izatnagar,
243122, India

In the study by Malekzadegan et al,¹ conducted on a limited number of cases, it is reported that 71.9% and 48.4% of isolates of uropathogenic *Escherichia coli* were resistant to nalidixic acid and ciprofloxacin, respectively. It appears of normal occurrence but certainly enriches the knowledge of clinicians towards the selection of appropriate antibiotic. In the study, *E. coli* isolates claimed to be uropathogenic were not confirmed for their uropathogenic potential with any other test than their isolation in pure culture from urinary tract infection (UTI) cases. Uropathogenic *E. coli* may also occur in mixed culture and sometimes may be just contaminants from perineal region.² My intention is not to criticise the report but I feel that people in Iran are certainly lucky not to have super-drug-resistant *E. coli* causing urinary tract infections. In Bareilly, one of the big city in India, on looking at bacteriological culture and antibiotic sensitivity data of recent 209 human UTI cases referred to Clinical Epidemiology Laboratory of the Institute, *E. coli* was at the top to be associated with urinary tract infections in 22% cases, followed staphylococci (20.6%), streptococci (10.5%), enterococci (8.1%), *Enterobacter* sp. (6.2%), *Proteus* sp. (4.3%), *Aerococcus* sp. (3.8%), *Erwinia* sp. (3.8%), klebsiellae (3.3%), pseudomonads (2.9%), *Raoultella* sp. (2.9%), *Acinetobacter* sp. (1.9%), *Alcaligenes* sp. (1.9%), aeromonads (1.4%), *Flavobacter* sp. (1%), *Micrococcus* sp. (1%) strains, and isolation of *Actinobacillus*, *Bacillus*, *Candia*, *Citrobacter*, *Edwardsiella*, *Moraxella*, *Morganella*, *Serratia* and *Vibrio* species from one case each was also recorded.

Of the 46 isolates of *E. coli*, isolated in our laboratory from UTI cases, all were resistant to nalidixic acid, only 13% were sensitive for ciprofloxacin and 7 of them were also resistant to one or more carbapenem drugs (producing Metallo- β -lactamase); however, nitrofurantoin, a drug of choice earlier in UTI cases, was found inhibitory to >90% strains of *E. coli* from UTI cases.

I feel that there is a continuous need for monitoring of *E. coli* and other bacteria too causing UTI and gastrointestinal tract infections to keep an eye on the emergence of drug-resistant strains as such *E. coli* strains have a high potential of spread beyond geographical boundaries through excretions of travellers either clinically ill or carriers of the potentially pathogenic strains.

Correspondence: Bhoj R Singh
438 MLB, Division of Epidemiology, ICAR-
Indian Veterinary Research Institute,
Izatnagar 243122, India
Email brs1762@gmail.com

Disclosure

The author reports no conflicts of interest in this communication.

References

1. Malekzadegan Y, Rastegar E, Moradi M, Heidari H, Sedigh Ebrahim-Saraie H. Prevalence of quinolone-resistant uropathogenic *Escherichia coli* in a tertiary care hospital in south Iran. *Infect Drug Resist.* 2019;12:1683–1689. doi:10.2147/IDR.S206966
2. Stork C, Kovács B, Rózsai B, et al. Characterization of asymptomatic bacteriuria *Escherichia coli* isolates in search of alternative strains for efficient bacterial interference against uropathogens. *Front Microbiol.* 2018;9:214. doi:10.3389/fmicb.2018.00214

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Infection and Drug Resistance 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Infection and Drug Resistance editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>