Genetic characterization of a novel sequence type of multidrug-resistant *Citrobacter freundii* strain recovered from wastewater treatment plant

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Xiawei Jiang¹
Xinjie Cui¹
Wenhong Liu¹
Hao Xu²
Beiwen Zheng²

¹College of Basic Medical Sciences, Zhejiang Chinese Medical University, Hangzhou, Zhejiang, People's Republic of China; ²Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou, Zhejiang, People's Republic of China

Abstract: A multidrug-resistant *Citrobacter freundii* strain R17 was isolated from a wastewater treatment plant in China. Whole-genome sequencing of strain R17 revealed a new sequence type (ST412) chromosome (length 5,124,258 bp) and an Inc FII (Yp) group plasmid pCFR17_1 (length 206,820 bp). A total of 13 antibiotic-resistance genes (ARGs) that confer resistance to eight different antibiotic groups were encoded by strain R17 and 12 of them were carried by plasmid pCFR17_1. These data and analysis suggest that the environment-derived *C. freundii* strains may serve as potential sources of ARGs and highlight the need of further surveillance of this bacteria in the future.

Keywords: *Citrobacter freundii*, multidrug resistant, antibiotic-resistance genes, whole-genome sequencing, wastewater treatment plant, insertion elements

Introduction

Citrobacter freundii is a Gram-negative and facultative anaerobic bacterium which belongs to family Enterobacteriaceae. After its first isolation from soil, *C. freundii* has been found in various natural habitats, as well as intestines of animals and humans.¹⁻⁴ Some environment-derived *C. freundii* strains have potential use in many fields. For example, *C. freundii* strain JPG1, which was isolated from a gold mining tailing in China, was resistant to heavy metals and capable of removing copper. Thus, it was suggested to have a great potential in the treatment of copper-rich industrial wastewater.⁵ *C. freundii* strain IFO 13545 has the ability to produce bioflocculant which can be used in fields of water supply, wastewater treatment, and food production.⁶ Although *C. freundii* was previously recognized as a bacterium of low virulence, it has been demonstrated to be the causative agent of a wide spectrum of infections including diarrhea, pneumonia and septicemia.^{7,8} Moreover, there has been a growing body of literature that reported the plasmid-mediated multidrug resistance in *C. freundii*, indicating its threat to human health.^{9,10}

Wastewater treatment plants (WWTPs) have been regarded as the major source of antibiotic-resistant bacteria (ARB) and antibiotic-resistance genes (ARGs).^{11,12} Previous studies have reported the isolation of several *C. freundii* strains from WWTPs.^{13,14} In our investigation of the ARB and ARGs in WWTPs, we obtained a multidrug-resistant strain of *C. freundii*. This strain (R17) was isolated from the sludge of a WWTP that treats wastewater of a pharmaceutical industry in Taizhou, China. Sludge samples were collected from this WWTP in April 2017. Samples were first

Correspondence: Beiwen Zheng
Collaborative Innovation Center for
Diagnosis and Treatment of Infectious
Diseases, State Key Laboratory for
Diagnosis and Treatment of Infectious
Diseases, The First Affiliated Hospital,
College of Medicine, Zhejiang University,
No. 79 Qingchun Road, Hangzhou
310000, People's Republic of China
Tel +86 5 718 723 6423
Fax +86 5 718 723 6423
Email zhengbw@zju.edu.cn

cultured overnight in Mueller-Hinton broth without antimicrobial agents, and then the enriched samples were spread on chromID® ESB� agar plates (bioMérieux, Marcy-l'Étoile, France). After incubating at 37°C for 48 hrs, colonies were picked up for identification using both Microflex MALDI-TOF mass spectrometer (Bruker Daltonics, Bremen, Germany) and 16S rRNA sequencing technology as described previously.⁴ In the present study, we reported the whole-genome sequencing of *C. freundii* strain R17. Further investigations were conducted on the genetic features and ARGs associated with its resistance phenotypes.

Antimicrobial susceptibility testing of strain R17 was performed using VITEK 2 system employing panel AST-GN-16 (bioMérieux) with *Escherichia coli* ATCC 25922 as control. The results were interpreted according to the standards of the Clinical and Laboratory Standards Institute.¹⁵ Minimum inhibitory concentration (MIC) of strain R17 indicated its resistance to various types of antibiotics, including ampicillin (MIC ≥32 µg/mL), amoxicillin (MIC ≥32 µg/mL), cefazolin (MIC ≥64 µg/mL), cefoxitin (MIC ≥64 µg/mL), gentamicin (MIC ≥16 µg/mL), ciprofloxacin (MIC ≥4 µg/mL), levofloxacin (MIC ≥8 µg/mL) and trimethoprim (MIC ≥320 µg/mL).

Genomic DNA of strain R17 was extracted using QIAamp DNA Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. Whole-genome sequencing was performed using Pacific Biosciences RSII platform (PacBio, Menlo Park, CA, USA) and Illumina Hiseq 2000 sequencer (Illumina, San Diego, CA, USA) as described before.¹⁶ A chromosome of 5,124,258 bp in length with a GC content of 51.5% and a plasmid (pCFR17_1) of 206,820 bp in length with a GC content of 53.8% were assembled from the sequencing reads using SPAdes software (<https://github.com/ablab/spades#sec5>).^{17,18}

Annotation with RAST server¹⁹ revealed that the chromosome of strain R17 contained 4,960 protein-coding genes, 42 repeat regions, 85 tRNAs and 25 rRNAs (eight complete 5S-23S-16S rRNA operons and one additional 5S rRNA). Two intact phage regions (one was 48 kb in length and another one was 44.8 kb in length) were predicted from the chromosome using PHASTER.²⁰ One AmpC-type beta-lactamase encoding gene, *bla*_{CMY-85}, was found in the chromosome of strain R17 using ResFinder 3.1²¹ (Table 1). AmpC enzymes were reported to be involved in expanded-spectrum cephalosporins resistance and were found in the chromosomes of several members of *Enterobacteriales* including *C. freundii*.²² Up

Table 1 Distribution of ARGs in *C. freundii* strain R17

Resistance gene	Phenotype	Number of ARGs
Chromosome		1
<i>bla</i> _{CMY-85}	Beta-lactam resistance	1
pCFR17_1		12
<i>aadA2</i>	Aminoglycoside resistance	1
<i>aac</i> (3)-IId	Aminoglycoside resistance	1
<i>bla</i> _{DHA-1}	Beta-lactam resistance	1
<i>bla</i> _{TEM-1B}	Beta-lactam resistance	1
<i>qnrB4</i>	Fluoroquinolone resistance	1
<i>mph</i> (A)	Macrolide resistance	1
<i>catA2</i>	Phenicol resistance	1
<i>sul1</i>	Sulphonamide resistance	2
<i>sul2</i>	Sulphonamide resistance	1
<i>tet</i> (D)	Tetracycline resistance	1
<i>dfpA12</i>	Trimethoprim	1

Abbreviation: ARGs, antibiotic-resistance genes.

to date, 76.62% (272 of 355) *C. freundii* strains in NCBI Pathogen Detection database (<https://www.ncbi.nlm.nih.gov/pathogens/isolates#/search/>) carried *bla*_{CMY} genes, indicating that *bla*_{CMY} genes were ubiquitous in *C. freundii*. To reveal the phylogenetic relationship between strain R17 and other *C. freundii* strains with published complete chromosome sequences, Average Nucleotide Identities (ANIs) were calculated based on MUMmer using JSpeciesWS.²³ Though strain R17 was environmental-derived, it showed high ANI values with the clinical or host-associated strains (Table S1), indicating its potential of being a pathogen. Notably, by searching against *C. freundii* locus/sequence definitions database (https://pubmlst.org/bigdb?db=pubmlst_cfreundii_seqdef), we found that strain R17 could not be classified to any known STs, so it was assigned to a new ST, ST412.

The plasmid typing using PlasmidFinder²⁴ indicated that pCFR17_1 was a Inc FII (Yp) type plasmid. A total of 12 ARGs were predicted in pCFR17_1 using ResFinder 3.1²¹ (Table 1) and these genes were associated with resistance phenotypes of strain R17. By Blastn against NCBI database, we found that pCFR17_1 shared the highest similarity (99.99% identity and 73% coverage) with pCFR-e790 (GenBank accession number CP026241). Plasmid pCFR-e790 was identified in *C. freundii* strain CFNIH9 which was collected from wastewater of hospital internal pipes in USA.²⁵ These two plasmids shared the core backbone gene loci (Figure 1).

lie downstream of *bla*_{TEM-1B} in both plasmids. This conserved sequence showed high similarity to TnA33 which belonged to Tn3 composite transposon family. However, a ~24 kb region in pCFR17_1 was absent in pCFR-e790. And, this region exhibited a high similarity to pQnrB4 (GenBank accession number CP028537), a plasmid of *Enterobacter hormaechei* strain SCEH020042 which was a human-related strain isolated from Sichuan, China (Figure 1). Two insertion elements, *IS6* and *IS110*, located at each ends of this multidrug-resistant

8. Anderson MT, Mitchell LA, Zhao L, Mobley HL. *Citrobacter freundii* fitness during bloodstream infection. *Sci Rep*. 2018;8(1):11792. doi:10.1038/s41598-018-30196-0
9. Ouyang J, Sun F, Zhou D, et al. Comparative genomics of five different resistance plasmids coexisting in a clinical multi-drug resistant *Citrobacter freundii* isolate. *Infect Drug Resist*. 2018;11:1447. doi:10.2147/IDR.S165818
10. Wu W, Espedido B, Feng Y, Zong Z. *Citrobacter freundii* carrying bla KPC-2 and bla NDM-1: characterization by whole genome sequencing. *Sci Rep*. 2016;6:30670. doi:10.1038/srep30670
11. Obayiwana A, Ogunjobi A, Yang M, Ibekwe M. Characterization of bacterial communities and their antibiotic resistance profiles in wastewaters obtained from pharmaceutical facilities in Lagos and Ogun States, Nigeria. *Int J Environ Res Public Health*. 2018;15:7. doi:10.3390/ijerph15061188
12. Zhai W, Yang F, Mao D, Luo Y. Fate and removal of various antibiotic resistance genes in typical pharmaceutical wastewater treatment systems. *Environ Sci Pollut Res*. 2016;23(12):12030–12038. doi:10.1007/s11356-016-6350-9
13. Duceppe M-O, Phipps-Todd B, Carrillo C, Huang H. Draft genome sequences of eight Canadian *Citrobacter braakii* and *Citrobacter freundii* strains. *Microbiol Resour Announcements*. 2019;8(27):e00273–e00219. doi:10.1128/MRA.00273-19
14. Aristizábal-Hoyos AM, Rodríguez EA, Arias L, Jiménez JN. High clonal diversity of multidrug-resistant and extended spectrum beta-lactamase-producing *Escherichia coli* in a wastewater treatment plant. *J Environ Manage*. 2019;245:37–47. doi:10.1016/j.jenvman.2019.05.073
15. Performance CLSI. *Standards for Antimicrobial Susceptibility Testing*. 27th ed. CLSI supplement M100. Wayne, PA: Clinical and Laboratory Standards Institute; 2017.
16. Jiang X, Liu W, Zheng B. Complete genome sequencing of *Comamonas kerstersii* 8943, a causative agent for peritonitis. *Sci Data*. 2018;5:180222. doi:10.1038/sdata.2018.187
17. Nurk S, Bankevich A, Antipov D, et al. Assembling genomes and mini-metagenomes from highly chimeric reads. In: Deng M, Jiang R, Sun F, Zhang X (editors). *Research in Computational Molecular Biology: RECOMB 2013. Lecture Notes in Computer Science*, vol 7821. Heidelberg: Springer, Berlin; 2013.
18. Antipov D, Korobeynikov A, McLean JS, Pevzner PA. hybridSPAdes: an algorithm for hybrid assembly of short and long reads. *Bioinformatics*. 2015;32(7):1009–1015. doi:10.1093/bioinformatics/btv688
19. Aziz RK, Bartels D, Best AA, et al. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics*. 2008;9(1):75. doi:10.1186/1471-2164-9-75
20. Arndt D, Marcu A, Liang Y, Wishart DS. PHAST, PHASTER and PHASTEST: tools for finding prophage in bacterial genomes. *Brief Bioinform*. 2017. doi:10.1093/bib/bbx121
21. Zankari E, Hasman H, Cosentino S, et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother*. 2012;67(11):2640–2644. doi:10.1093/jac/dks261
22. Meini S, Tascini C, Cei M, Sozio E, Rossolini GM. AmpC β -lactamase-producing Enterobacterales: what a clinician should know. *Infection*. 2019;47(3):363–375.
23. Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplies J. JSpeciesWS: a web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics*. 2015;32(6):929–931. doi:10.1093/bioinformatics/btv681
24. Carattoli A, Zankari E, Garcia-Fernandez A, et al. PlasmidFinder and pMLST: in silico detection and typing of plasmids. *Antimicrob Agents Chemother*. 2014;58(7):3895–3903.
25. Weingarten RA, Johnson RC, Conlan S, et al. Genomic analysis of hospital plumbing reveals diverse reservoir of bacterial plasmids conferring carbapenem resistance. *MBio*. 2018;9(1):e02011–e02017. doi:10.1128/mBio.02011-17

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

Dovepress