

Diversity and frequency of resistance and virulence genes in *bla*_{KPC} and *bla*_{NDM} co-producing *Klebsiella pneumoniae* strains from China

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Xin Liu^{1,*}
Jie Zhang^{1,*}
Yini Li¹
Qiuni Shen¹
Wenting Jiang¹
Kelei Zhao²
Yancheng He¹
Penggao Dai¹
Zhihao Nie¹
Xiyan Xu^{1,3}
Yingshun Zhou¹

¹Department of Pathogenic Biology, School of Basic Medicine, Southwest Medical University, Luzhou, Sichuan, People's Republic of China; ²Antibiotics Research and Re-evaluation Key Laboratory of Sichuan Province, Sichuan Industrial Institute of Antibiotics, Chengdu University, Chengdu, Sichuan 610052, People's Republic of China; ³Department of Histology and Embryology, School of Basic Medicine, Southwest Medical University, Luzhou, Sichuan, People's Republic of China

*These authors contributed equally to this work

Background: Emergence of *bla*_{KPC} and *bla*_{NDM} co-producing *Klebsiella pneumoniae* strains have led to the limited therapeutic options for clinical treatment. Understanding the diversity and frequency of resistance and virulence genes of these isolates is of great significance.

Purpose: The aim of this study is to research the diversity and frequency of resistance and virulence genes in the *bla*_{KPC} and *bla*_{NDM} co-producing *Klebsiella pneumoniae* strains.

Methods and Results: In this study, 117 *K. pneumoniae* strains were isolated from China, and among of which, 24 were found to be *bla*_{KPC} and *bla*_{NDM} co-producing with significant resistance against almost all the commonly used antibiotics. Additionally, 4 strains were hypermucoviscous and 8 showed high serum resistance. Overall, *bla*_{SHV}, *bla*_{CTX-M}, *tetA* and *sul1* resistance genes found in 100% of the isolates, followed by *bla*_{TEM} (95.8%), *oqxA/B* (91.7%), *qnrB* (87.5%), *aac(6')Ib-cr* (83.3%), *bla*_{DHA} (79.2%), *rmtB* (66.7%), *qnrS* (54.2%), *cat* (54.2%), *floR* (50.0%), *sul2* (45.8%) *cmlA* (20.8%) and *bla*_{CMY} (8.33%), respectively. What's more, seven *bla*_{CTX-M} subtypes [*bla*_{CTX-M-14} (n=18), *bla*_{CTX-M-3} (n=11), *bla*_{CTX-M-65} (n=4), *bla*_{CTX-M-15} (n=3), *bla*_{CTX-M-28} (n=2), *bla*_{CTX-M-55} (n=2), *bla*_{CTX-M-22} (n=1)] and six *bla*_{SHV} subtypes [*bla*_{SHV-12} (n=16), *bla*_{SHV-11} (n=4), *bla*_{SHV-2a} (n=1), *bla*_{SHV-1} (n=1), *bla*_{SHV-38} (n=1) and *bla*_{SHV-28} (n=1)] were detected. The frequency of virulence genes was as follows: 100% for *entB*, *ybtS* and *irp*, 95.8% for *mrkD*, 91.66% for *fimH*, 79.2% for *iutA*, 62.5% for *iroBCDE*, *aerobactin* and *kfu*, 66.7% for *allS*, 45.8% for *wcaG*, 37.5% for *rmpA*, 20.8% for *pagO* and 16.7% for *maga*.

Conclusion: From this study, we concluded that the *bla*_{KPC} and *bla*_{NDM} co-producing *Klebsiella pneumoniae* strains have a high diversity and frequency of resistance and virulence genes. This study may offer hospitals important information about the control of infections caused by *bla*_{KPC} and *bla*_{NDM} co-producing *Klebsiella pneumoniae*.

Keywords: *Klebsiella pneumoniae*, *bla*_{NDM}, *bla*_{KPC}, resistance genes, virulence factors

Introduction

Carbapenemase-producing bacteria can hydrolyse carbapenems and most other β -lactam antibiotics which pose significant challenges to clinical diagnosis and treatment. *Klebsiella pneumoniae* carbapenemase (KPC) and Metallo-B-Lactamases (*bla*_{NDM}) are the two major groups of carbapenemases that produced by the most of Carbapenemase-Resistant *Enterobacteriaceae* strains (CRE). The *bla*_{KPC} and *bla*_{NDM} genes are commonly found in CRE strains in recent years.¹⁻³ Those type of the

Correspondence: Xiyan Xu; Yingshun Zhou
Department of Pathogenic Biology, School of Basic Medicine, Southwest Medical University, No. 319, Zhongshan Road, Luzhou, Sichuan 646000, People's Republic of China
Tel +86 0 830 316 0073
Email yingshunzhou@swmu.edu.cn

carbapenem resistance genes and other resistance genes including the key Extended-Spectrum β -lactamases (ESBLs) genes (*bla*_{CTX-M}, *bla*_{SHV} and *bla*_{TEM}), the fluoroquinolone resistance genes (*qnrA*, *qnrB*, *qnrS*, *oqxA/B*), aminoglycoside resistance genes (*rmtA*, *rmtB* and *rmtC*), chloramphenicol resistance genes (*cat*, *floR*, *cmlA*, *cfr*) and tetracycline resistance genes (*tetA*, *tetB*, *tetC*) are carried by the same strain and resulting in high resistance to almost all kinds of antibiotics.⁴⁻⁷ The more worrisome is hypervirulent *K. pneumoniae* strains (hvKP) emergency sharply in recent years, especially the carbapenemase-producing hvKP related infections in immunocompromised patients which is a serious threat to the patients.⁸⁻¹¹

More and more researchers report that HvKP strains are characterized a number of virulence factors including *aerobactin* (encodes high-affinity iron chelators), *rmpA* (regulators of mucoid phenotype), *wcaG* (involved in the biosynthesis of the outer core lipopolysaccharide), *allS* (associated with allantoin metabolism), *kfu* (responsible for an iron uptake system), *yptS*, *irp* (yersiniabactin biosynthesis) and *iroBCDN* (salmocheilin biosynthesis), *entB* (catecholate siderophore), *fimH* and *mrkD* (fimbrial adhesin, which mediate binding to the extracellular matrix to form the biofilm), *pagO* (involved in liver abscess formation by liver abscess-Kp).^{9,12-15}

Understanding the diversity and frequency of resistance and virulence genes of these isolates is of great significance to disease prevention and control. For offer hospitals important information about the control of infections caused by *bla*_{KPC} and *bla*_{NDM} co-producing *K. pneumoniae*. In this study, we mainly present the diversity and frequency of resistance and virulence genes in the *bla*_{KPC} and *bla*_{NDM} co-producing *K. pneumoniae*.

Materials and methods

Isolates collection and screening of *bla*_{KPC} and *bla*_{NDM} genes

A total of 117 non-repetitive *K. pneumoniae* strains were isolated from sputum, cerebrospinal fluid, wound, and urine samples for routine examination between Aug. 2016 and Sept.2018 at several hospitals in Sichuan, Henan, Fujian province of China. These isolates were identified by VITEK2 Compact System (bioMérieux, France) and 16sRNA sequencing. *K. pneumoniae* ATCC700603 was used as the control strain for the species identification and antimicrobial susceptibility test. The

*bla*_{KPC} and *bla*_{NDM} detection were performed according to our previous work by PCR.^{9,16}

Antimicrobial susceptibility testing

Antimicrobial susceptibility testing of the *bla*_{KPC} and *bla*_{NDM} co-producing *K. pneumoniae* strains were performed according to the recommendations of the Clinical Laboratory Standards Institute (CLSI) guidelines (CLSI, 2017). Antimicrobial agents (Oxoid, England) used in this study included CXM (cefuroxime axetil), TZP (piperacillin-tazobactam), CAZ (ceftazidime), CRO (ceftriaxone), IPM (imipenem), MEM (meropenem), ATM (aztreonam), AMK (amikacin), CIP (ciprofloxacin), CHL (chloramphenicol), TMP-SMZ (trimethoprim/sulfamethoxazole). *E. coli* strain ATCC 25922 was used as quality control.¹⁷

Hypermucoviscosity, biofilm formation and serum killing assay

The hypermucoviscosity phenotype of 24 *K. pneumoniae* was detected by string test.¹⁸ The colonies were cultured on blood agar plate overnight at 37°C, stretched by a bacteriology inoculation loop. The strain formed a viscous string of >5 mm was designated as hypermucoviscous. Biofilm formation assay was performed by crystal violet staining assay.⁹ Biofilm formation in each well was measured by microplate reader (Bio-Rad, US) at optical density (OD) 595 nm. The susceptibility of the *K. pneumoniae* isolates to human serum was explored by an established method.¹⁹ Briefly, *K. pneumoniae* strains were inoculated into LB Broth Medium and incubated at 37°C with shaking until the logarithmic phase was reached (T=4 h, OD₆₀₀=0.6). 25 μ L of diluted culture (containing 10⁶ CFU of bacteria) and 75 μ L human serum were then added into a 10 \times 75 mm Falcon polypropylene tube and incubated at 37°C with shaking. Viable counts were checked at 0, 1, 2, and 3 h. The response to serum killing in terms of viable counts was scored on six grades as described previously method.²⁰

ERIC-PCR

Enterobacterial repetitive intergenic consensus PCR (ERIC-PCR) method was used to evaluate the genetic diversity of the 24 isolates, as previously described using the primers.²¹ The PCR products were loaded on a 1% agarose gel with the gelred at 90 V for 40 mins, and the banding patterns were analyzed by gel imaging and analysis system. To determine the similarity rate among the acquired outcomes, Genetic diversity were analyzed using

the unweighted pair-group method with arithmetic mean (UPGMA) and isolates with $\geq 80\%$ similarity were treated as a single cluster.²²

Detection of resistance and virulence genes

By using PCR, the carriage of carbapenemase-encoding genes (*bla*_{VIM}, *bla*_{GES}, *bla*_{DIM}, *bla*_{GIM}, *bla*_{SPM} and *bla*_{AIM}),²³ ESBL-encoding genes (*bla*_{TEM}, *bla*_{SHV}, *bla*_{CTX-M}, *bla*_{CTX-M-1}, *bla*_{CTX-M-2} and *bla*_{CTX-M-9}),⁷ AmpC β -lactamase genes (*bla*_{DHA}, *bla*_{CMY}),²⁴ 16 s rRNA methylase genes (*rmtA*, *rmtB* and *rmtC*),²⁵ sulfonamides resistance genes (*sul1*, *sul2* and *sul3*), chloramphenicol resistance genes (*cmlA*, *floR* and *catB*), multiresistance gene (*cfr*), tigecycline resistance gene (*tetA*, *tetB* and *tetC*)^{26,27} and quinolone resistance genes (*qnrA*, *qnrB*, *qnrS*, *aac(6')-Ib-cr*, *qepA* and *oqxAB*)^{28–30} were detected as described previously. PCR assays were also used to assess the capsular serotypes (*K1*, *K2*, *K5*, *K20*, *K54* and *K57*)³¹ and fourteen virulence genes (*magA*, *rmpA*, *allS*, *wcaG*, *ybtS*, *kfu*, *iroBCDE*, *entB*, *irp*, *iutA*, *aerobactin*, *mrkD*, *fimH* and *pagO*).^{12–14,31,32} PCR amplicons were sequenced by Shanghai Sangon Bioengineering Company. Sequences were analyzed by the BLAST programs (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). The primers used were shown in Table S1.

Results

Antimicrobial susceptibility, hypermucoviscosity, serotyping, biofilm, serum resistance assay and ERIC-PCR typing

A total of 24 *bla*_{KPC} and *bla*_{NDM} co-producing strains were screened from 117 non-repetitive *K. pneumoniae* strains. All the isolates were resistant to piperacillin-tazobactam, cefuroxime axetil, ceftazidime, ceftriaxone, imipenem, meropenem and aztreonam (Table 1). Among the 24 *bla*_{KPC} and *bla*_{NDM} co-producing strains, 16.7% (n=4) were the K1 type, while the K2, K5, K20, K57 and K54 serotype were not found (Figure 1). String test showed that 4 (KP103L, KP48L, KP97L, KP36L) *bla*_{KPC} and *bla*_{NDM} co-producing *K. pneumoniae* isolates were hypermucoviscous. Biofilm formation was observed in all the 24 strains, with values of OD₅₉₅ nm ranged from 0.33 to 2.70, whereas the mean value of the negative control wells is 0.168. Serum killing assay showed that 33.3% (n=8) of the strains were high serum resistance (Grade 5 or Grade 6). Analysis of genetic linkage among isolates by ERIC-PCR showed 34–100% similarity among 24 isolates (Table 2). Genetic diversity was established

among 24 *bla*_{KPC} and *bla*_{NDM} co-producing *K. pneumoniae* isolates by detecting 15 different ERIC fingerprints with the similarity cutoff of 80% (Table 2).

Diversity and frequency of resistance and virulence gene

As shown in Table 1, all isolates (100%, n=24) carried the resistance gene *bla*_{SHV}, *bla*_{CTX-M}, *tetA* and *sul1*, followed by *bla*_{TEM} (95.8%), *oqxA/B* (91.7%), *qnrB* (87.5%), *aac(6')Ib-cr* (83.3%), *bla*_{DHA} (79.2%), *rmtB* (66.7%), *qnrS* (54.2%), *cat* (54.2%), *floR* (50.0%), *sul2* (45.8%) *cmlA* (20.8%) and *bla*_{CMY} (8.33%), respectively. While the carbapenemase encoding genes *bla*_{GES}, *bla*_{VIM}, *bla*_{AIM}, *bla*_{GIM}, *bla*_{DIM} were not detected in any of those strains. Regarding the *bla*_{CTX-M} group (Table 2; Supplement Sequences), the most widespread subtype was *bla*_{CTX-M-14}, which was found in 75% (n=18) of the tested isolates, followed by *bla*_{CTX-M-3} in 45.8% (n=11), *bla*_{CTX-M-65} in 16.7% (n=4), *bla*_{CTX-M-15} in 12.5% (n=3), *bla*_{CTX-M-28} in 8.3% (n=2), *bla*_{CTX-M-55} in 8.3% (n=2), *bla*_{CTX-M-22} in 4.2% (n=1). In addition, there are 17 isolates carried two subtypes of *bla*_{CTX-M}. And the majority of the 8 isolates carried *bla*_{CTX-M-14} co-existing with *bla*_{CTX-M-3}, while 2 isolates co-carried *bla*_{CTX-M-14} and *bla*_{CTX-M-65} (Table 2). Regarding the *bla*_{SHV} group, *bla*_{SHV-12} (66.7%; n=16) was the most prevalent *bla*_{SHV} in those 24 *bla*_{KPC} and *bla*_{NDM} co-producing strains, followed by *bla*_{SHV-11} in 16.7% (n=4), *bla*_{SHV-2a}, *bla*_{SHV-1}, *bla*_{SHV-38} and *bla*_{SHV-28} in 4.2% (n=1) (Table 2).

Diversity and frequency of virulence genes

The prevalence and distribution of virulence factors are given in Table 2. All strains carried the *ybtS*, *entB* and *irp* gene. 95.8% (n=23) strains harbored *mrkD* gene, 91.6% (n=22) strains harbored *fimH* gene, 79.2% (n=19) strains contained *iutA* gene, 66.7% (n=16) strains carried *allS* gene, 62.5% (n=15) strains carried *iroBCDE*, *aerobactin* and *kfu* gene, 45.8% (n=11) strains contained *wcaG* gene, 37.5% (n=9) strains involved *rmpA* gene, 20.8% (n=5) strains involved *pagO* gene and 16.7% (n=4) carried *magA* gene.

Discussion

The prevalence of co-carried *bla*_{NDM} and *bla*_{KPC} in a single bacterial isolate in hospitals has led to heightened concerns because often makes the isolate an extremely drug-resistant variant.^{2,3} In this study, 117 non-repetitive *K. pneumoniae* strains were isolated from China, and among of which, 24

Table 1 The antibiotic resistance phenotype profile and positive rate of the resistance gene of the isolates

Antibiotic Resistance phenotype profile		Resistance gene																	
		<i>bla</i> _{NDM}	<i>bla</i> _{KPC}	<i>bla</i> _{CTX-M}	<i>bla</i> _{SHV}	<i>bla</i> _{TEM}	<i>bla</i> _{DHA}	<i>bla</i> _{CMY}	<i>sul</i> ₁	<i>sul</i> ₂	<i>rmt</i> _B	<i>cat</i> _B	<i>flor</i>	<i>cm</i> _I A	<i>qnr</i> _B	<i>qnr</i> _S	<i>oxy</i> _{A/B}	<i>aac</i> (6')- <i>lb-cr</i>	<i>tet</i> _A
Kp6L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/TMP-SMZ/TZP																		
Kp32L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TGC/TMP-SMZ																		
Kp5L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TGC																		
Kp50L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TMP-SMZ																		
Kp22L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/CHL/TMP-SMZ																		
Kp49L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TGC																		
Kp42L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/CIP																		
Kp93L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TMP-SMZ																		
Kp11L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TGC/CHL																		
Kp105L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/CIP/TGC/TMP-SMZ																		
Kp103L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TGC/TMP-SMZ																		
Kp31L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP																		
Kp48L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TGC																		
Kp87L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/CIP																		
Kp104L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/CIP/TGC/CHL																		
Kp20L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TGC/CHL																		
Kp116L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/CHL/TMP-SMZ																		
Kp29L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TGC/CHL																		
Kp12L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TMP-SMZ																		
Kp36L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/CIP/CHL/TMP-SMZ																		
Kp97L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/CHL/TMP-SMZ																		
Kp9L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/AMK/CIP/TGC/CHL																		
Kp40L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/CIP/TGC																		
Kp13L	TZP/CXM/CAZ/CRO/IMP/MEM/ATM/CIP/TGC/CHL																		
Positive rate		100%	100%	100%	100%	95.8%	79.17%	8.33%	100%	45.8%	66.7%	54.2%	50%	20.8%	87.5%	54.2%	91.7%	83.3%	100%

Note: The green check represent the positive while the blank is the negative.

Abbreviations: TZP, piperacillin-tazobactam; CXM, cefuroxime axetil; CAZ, ceftazidime; CRO, ceftriaxone; IMP, imipenem; MEM, meropenem; ATM, aztreonam; AMK, amikacin; CIP, ciprofloxacin; CHL, chloramphenicol; TMP-SMZ, trimethoprim/sulfamethoxazole.




Figure 1 The dendrogram of ERIC-PCR fingerprints and diversity of the ESBLs genotypes.

Notes: The dendrogram of ERIC-PCR fingerprints was constructed using the Dice coefficient and the unweighted pair-group method with arithmetic mean (UPGMA) and the diversity of the ESBLs (bla_{CTX-M} group and bla_{SHV} group) genotypes.

were found to be bla_{KPC} and bla_{NDM} co-producing with significant resistance against almost all the commonly used antibiotics. This results showed that the positive incidence of the bla_{NDM} and bla_{KPC} co-producing *K. pneumonia* is increasing. The results were expected that all 24 isolates resist almost the all test antibiotic and biofilm formation was observed in all the 24 strains. This is a dangerous situation for antibiotic treatment because the high biofilm formation pathogenic bacteria often involved in hospital infections and always lead to the failure of antibiotic treatments.⁹ Additionally, 4 strains were hypermucoviscous and 8 strains showed high serum resistance. To our knowledge, the phenotype of hypermucoviscous, biofilm formation ability and serum resistance were as the virulence evaluation criterion.^{18,20} Those results indicated that there are harboring hypervirulent variant of *Klebsiella pneumonia* (hvKp) among the 24 bla_{NDM} and bla_{KPC} co-

producing strains. This results suggest that urgent need to enhance clinical awareness and epidemiologic surveillance. Although the genetic diversity was established among 24 bla_{KPC} and bla_{NDM} co-producing *K. pneumoniae* isolates by detecting 15 different ERIC fingerprints with the similarity cutoff of 80%, we should pay more attention about this like strains clonal spread in the hospital.

In recent years, more and more researchers report that the co-carried bla_{NDM} and bla_{KPC} *K. pneumoniae* strains carried a large number of resistance genes, making this isolate highly resistance against almost all the commonly used antibiotics. For example, the bla_{KPC-2} and bla_{NDM-1} co-carriage strain *C. freundii* 112298 existence many resistance genes including the bla_{SHV-12}, bla_{CTX-M-14}, aac(6')-Ib-cr, bla_{OXA-1}, catB3, arr-3, fosA3 and sul1.¹ The bla_{KPC-2} and bla_{NDM-5} co-carriage strain ZSH6 carried

Table 2 The string test, serotyping, Serum killing and biofilm formation assay and diversity and frequency of the virulence factors of the *bla_{KPC}* and *bla_{NDM}* co-producing *Klebsiella pneumoniae*

	String test	Serotype	Serum resistance	Biofilm formation (OD value)	Virulence factor														
					rmpA	ybtS	mrkD	entB	kfu	wcaG	allS	iutA	aerobactin	magA	fimH	pagO	iroBCDE	irp	
Kp6L	—	ND	G1	Weak (0.33)															
Kp32L	—	ND	G1	Moderate (0.70)															
Kp5L	—	K1	G1	Strong (1.79)															
Kp50L	—	ND	G2	Strong (1.95)															
Kp22L	—	ND	G6	Strong (0.80)															
Kp49L	—	ND	G6	Strong (0.94)															
Kp42L	—	ND	G1	Strong (1.35)															
Kp93L	—	ND	G3	Weak (0.33)															
Kp11L	—	ND	G1	Strong (1.05)															
Kp105L	—	K1	G2	Strong (1.12)															
Kp103L	+	ND	G6	Strong (0.86)															
Kp31L	—	ND	G1	Strong (1.99)															
Kp48L	+	K1	G1	Strong (1.76)															
Kp87L	—	ND	G6	Strong (0.99)															
Kp104L	—	ND	G2	Strong (0.98)															
Kp20L	—	ND	G1	Strong (1.09)															
Kp116L	—	ND	G1	Strong (2.24)															
Kp29L	—	ND	G2	Strong (2.70)															
Kp12L	—	ND	G5	Strong(0.81)															
Kp36L	+	ND	G1	Strong (2.55)															
Kp97L	+	ND	G5	Strongp(1.04)															
Kp9L	—	K1	G5	Strong (1.04)															
Kp40L	—	ND	G6	Strong (0.83)															
Kp13L	—	ND	G1	Strong (0.74)															
Positive rate					37.5%	100%	95.8%	100%	62.5%	45.8%	66.7%	79.2%	62.5%	16.7%	91.66%	20.8%	62.5%		100%

Notes: “+”: positive, “—”: negative; The green check represent the positive while the blank is the negative. Biofilm formation expressed as crystal violet optical density value (OD at 595 nm).

Abbreviations: ND, Not Determination; OD, optical density; G, grade.

twenty resistance genes *bla*_{KPC-2}, *bla*_{NDM-5}, *bla*_{CTX-M-3}, *bla*_{CTX-M-65}, *bla*_{TEM-1}, *floR*, *tet(A)*, *tet(B)*, *dfrA17*, *aadA5*, *sul1*, *mdf(A)*, *mph(A)*, *erm(B)*, *aph(3')-Ia*, *aph(3')-Ib*, *aph(4)-Ia*, *aph(6)-Id*, *aac(3)-Iva*, *aac(3)-IId*.³ In this study, we also found that the high frequency and diversity of the resistance gene were emergency in the *bla*_{KPC-2} and *bla*_{NDM-1} co-carriage strains. All 24 isolates carried the *bla*_{SHV}, *bla*_{CTX-M}, *tetA* and *sul1*, followed by *bla*_{TEM} (95.8%), *oqxAB* (91.7%), *qnrB* (87.5%), *aac(6')Ib-cr* (83.3%), *bla*_{DHA} (79.2%), *rmtB* (66.7%), *qnrS* (54.2%), *cat* (54.2%), *floR* (50.0%), *sul2* (45.8%) and *cmlA* (20.8%). Particularly the high frequency and diversity of the ESBLs. (*bla*_{CTX-M} group and *bla*_{SHV} group) gene. For the *bla*_{CTX-M} group, there are seven *bla*_{CTX-M} subtypes including (*bla*_{CTX-M-14}, *bla*_{CTX-M-3}, *bla*_{CTX-M-65}, *bla*_{CTX-M-15}, *bla*_{CTX-M-28}, *bla*_{CTX-M-55} and *bla*_{CTX-M-22}) in all 24 strains. Our study showed that *bla*_{CTX-M-14} was the most frequent. In addition, there are 17 isolates carried two subtypes of *bla*_{CTX-M}. And the majority of the 8 isolates carried *bla*_{CTX-M-14} co-existing with *bla*_{CTX-M-3}, while 2 isolates co-carried *bla*_{CTX-M-14} and *bla*_{CTX-M-65} (Table 1). Regarding the *bla*_{SHV} group, *bla*_{SHV-12} (66.7%, n=16) was the most prevalent *bla*_{SHV} subtype in 24 *bla*_{KPC} and *bla*_{NDM} co-producing strains. The threat of the high frequency and diversity of the resistance gene emergency in the *bla*_{KPC-2} and *bla*_{NDM-1} co-carriage strains should be strict surveillance and management, although its resist almost all the commonly used antibiotics.²

Besides of the high frequency and diversity of the resistance gene, the virulence genes were also high emergency in 24 *K. pneumoniae* strains. In this study, we found that the frequency of virulence genes (*ybtS*, *entB*, *irp*, *mrkD*, *fimH*) was similar to most of others researcher's reports. However, the frequency of *wcaG* (45.8%), *allS* (66.7%) and *pagO* (20.8%) gene was slightly higher than our previous work. This results indicated the frequency of some virulence is rising. The high frequency of virulence factors found in these *bla*_{NDM} and *bla*_{KPC} bacteria is a problem for treatment. Some researchers suggested that molecular typing and virulence gene analysis are powerful tools that can shed light on *Klebsiella pneumoniae* infections.^{12,15,33,34} However, in this study, we found that some isolates were high serum resistance (Grade 5 or Grade 6) but the number of the virulence factors was less to some serum resistance strains. This results showed that how to identify the hvKP is still unknown. We suspect that the comprehensive analysis the frequency of the

virulence factors, phenotype (biofilm, sting test and serum killing assay) and clinical characteristics maybe a preferable method to identify the hvKP strains.

In conclusion, this study demonstrated that the high frequency and diversity of the resistance and virulence factors was in the *bla*_{NDM} and *bla*_{KPC} co-producing *K. pneumoniae* making this strain resistant to almost all antibiotics. This study may offer hospitals important information about the control of infections caused by *bla*_{KPC} and *bla*_{NDM} co-producing *Klebsiella pneumoniae*.

Acknowledgment

This research was funded by the National Natural Science Foundation of China (31500114) and by a grant from the Sichuan Province Science and Technology project (2016JY0223) and Luzhou and Southwest Medical University Natural Science Foundation [2018LZXNYD-ZK51] and Southwest Medical University Science Park funding [2019005].

Disclosure

The authors declare that there are no conflicts of interest in this work.

References

- Feng J, Qiu Y, Yin Z, et al. Coexistence of a novel KPC-2-encoding MDR plasmid and an NDM-1-encoding pNDM-HN380-like plasmid in a clinical isolate of *Citrobacter freundii*. *J Antimicrob Chemother.* 2015;70:2987. doi:10.1093/jac/dku445
- Zheng B, Xu H, Yu X, et al. Identification and genomic characterization of a KPC-2-, NDM-1- and NDM-5-producing *Klebsiella michiganensis* isolate. *J Antimicrob Chemother.* 2017;73:536–538.
- Fu L, Wang S, Zhang Z, et al. Co-carrying of KPC-2, NDM-5, CTX-M-3 and CTX-M-65 in three plasmids with serotype O89: H10 *Escherichia coli* strain belonging to the ST2 clone in China. *Microb Pathog.* 2019;128:1–6. doi:10.1016/j.micpath.2018.12.033
- Freire Martin I, AbuOun M, Reichel R, La Ragione RM, Woodward MJ. Sequence analysis of a CTX-M-1 IncI1 plasmid found in *Salmonella* 4,5,12: i:-, *Escherichia coli* and *Klebsiella pneumoniae* on a UK pig farm. *J Antimicrob Chemother.* 2014;69:2098–2101. doi:10.1093/jac/dku098
- Peirano G, Schreckenberger PC, Pitout JD. Characteristics of NDM-1-producing *Escherichia coli* isolates that belong to the successful and virulent clone ST131. *Antimicrob Agents Chemother.* 2011;55:2986. doi:10.1128/AAC.01763-10
- Pan YS, Zong ZY, Yuan L, et al. Complete sequence of pEC012, a multidrug-resistant IncI1 ST71 plasmid carrying *bla*_{CTX-M-65}, *rmtB*, *fosA3*, *floR*, and *oqxAB* in an Avian *Escherichia coli* ST117 strain. *Front Microbiol.* 2016;7:1117.
- Tian GB, Wang HN, Zou LK, et al. Detection of CTX-M-15, CTX-M-22, and SHV-2 extended-spectrum beta-lactamases (ESBLs) in *Escherichia coli* fecal-sample isolates from pig farms in China. *Foodborne Pathog Dis.* 2009;6:297. doi:10.1089/fpd.2008.0164

8. Chao L, Shi J, Guo J. High prevalence of hypervirulent *Klebsiella pneumoniae* infection in the genetic background of elderly patients in two teaching hospitals in China. *Infect Drug Resist*. 2018;11:1031–1041. doi:10.2147/IDR.S161075
9. Fu L, Huang M, Zhang X, et al. Frequency of virulence factors in high biofilm formation blaKPC-2 producing *Klebsiella pneumoniae* strains from hospitals. *Microb Pathog*. 2018;116:168–172. doi:10.1016/j.micpath.2018.01.030
10. Xu M, Fu Y, Fang Y, et al. High prevalence of KPC-2-producing hypervirulent *Klebsiella pneumoniae* causing meningitis in Eastern China. *Infect Drug Resist*. 2019;12:641–653. doi:10.2147/IDR.S191892
11. Struve C, Roe CC, Stegger M, et al. Mapping the evolution of Hypervirulent *Klebsiella pneumoniae*. *mBio*. 2015;6:e00630. doi:10.1128/mBio.00630-15
12. Yu WL, Ko WC, Cheng KC, Lee CC, Lai CC, Chuang YC. Comparison of prevalence of virulence factors for *Klebsiella pneumoniae* liver abscesses between isolates with capsular K1/K2 and non-K1/K2 serotypes. *Diagn Microbiol Infect Dis*. 2008;62:1. doi:10.1016/j.diagmicrobio.2008.04.007
13. Wasfi R, Elkhatib WF, Ashour HM. Molecular typing and virulence analysis of multidrug resistant *Klebsiella pneumoniae* clinical isolates recovered from Egyptian hospitals. *Sci Rep*. 2016;6:38929. doi:10.1038/srep38929
14. Ye M, Tu J, Jiang J, et al. Clinical and genomic analysis of liver abscess-causing *Klebsiella pneumoniae* identifies new liver abscess-associated virulence genes. *Front Cell Infect Microbiol*. 2016;6:165. doi:10.3389/fcimb.2016.00165
15. Russo TA, Marr CM. Hypervirulent *Klebsiella pneumoniae*. *Clin Microbiol Rev*. 2019; 32:e00001–19.
16. Liu Y, Zhang H, Zhang X, et al. Characterization of an NDM-19-producing *Klebsiella pneumoniae* strain harboring 2 resistance plasmids from China. *Diagn Microbiol Infect Dis*. 2019;93:355–361. doi:10.1016/j.diagmicrobio.2018.11.007
17. Fu L, Tang L, Wang S, et al. Co-location of the blaKPC-2, blaCTX-M-65, rmtB and virulence relevant factors in an IncFII plasmid from a hypermucoviscous *Klebsiella pneumoniae* isolate. *Microb Pathog*. 2018;124:301–304. doi:10.1016/j.micpath.2018.08.055
18. Shon AS, Bajwa RP, Russo TA. Hypervirulent (hypermucoviscous) *Klebsiella pneumoniae*: a new and dangerous breed. *Virulence*. 2013;4:107–118. doi:10.4161/viru.22718
19. Podschun R, Sievers D, Fischer A, Ullmann U. Serotypes, hemagglutinins, siderophore synthesis, and serum resistance of *Klebsiella* isolates causing human urinary tract infections. *J Infect Dis*. 1993;168:1415–1421. doi:10.1093/infdis/168.6.1415
20. Zhang Y, Zhao C, Wang Q, et al. High prevalence of hypervirulent *Klebsiella pneumoniae* infection in China: geographic distribution, clinical characteristics, and antimicrobial resistance. *Antimicrob Agents Chemother*. 2016;60:6115–6120. doi:10.1128/AAC.01127-16
21. Versalovic J, Koeuth T, Lupski JR. Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Res*. 1991;19:6823–6831. doi:10.1093/nar/19.24.6823
22. Duan H, Chai T, Liu J, et al. Source identification of airborne *Escherichia coli* of swine house surroundings using ERIC-PCR and REP-PCR. *Environ Res*. 2009;109:511–517. doi:10.1016/j.envres.2009.02.014
23. Poirel L, Walsh TR, Cuvillier V, Nordmann P. Multiplex PCR for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis*. 2011;70:119. doi:10.1016/j.diagmicrobio.2010.12.002
24. Pérezpérez FJ, Hanson ND. Detection of plasmid-mediated AmpC beta-lactamase genes in clinical isolates by using multiplex PCR. *J Clin Microbiol*. 2002;40:2153. doi:10.1128/JCM.40.6.2153-2162.2002
25. Liu Z, Ling B, Zhou L. Prevalence of 16S rRNA methylase, modifying enzyme, and extended-spectrum beta-lactamase genes among *Acinetobacter baumannii* isolates. *J Chemother*. 2015;27:207–212. doi:10.1179/1973947814Y.0000000190
26. Aminov RI, Chee-Sanford JC, Garrigues N, Mehboob A, Mackie RI. Detection of tetracycline resistance genes by PCR methods. *Methods Mol Biol*. 2004;268:3–13.
27. Zhang AY, Wang HN, Tian GB, et al. Phenotypic and genotypic characterisation of antimicrobial resistance in faecal bacteria from 30 Giant pandas. *Int J Antimicrob Agents*. 2009;33:456. doi:10.1016/j.ijantimicag.2008.10.030
28. Wu JJ, Ko WC, Tsai SH, Yan JJ. Prevalence of plasmid-mediated quinolone resistance determinants QnrA, QnrB, and QnrS among clinical isolates of *Enterobacter cloacae* in a Taiwanese hospital. *Antimicrob Agents Chemother*. 2007;51:1223–1227. doi:10.1128/AAC.01195-06
29. Andres P, Lucero C, Soler-Bistue A, et al. Differential distribution of plasmid-mediated quinolone resistance genes in clinical enterobacteria with unusual phenotypes of quinolone susceptibility from Argentina. *Antimicrob Agents Chemother*. 2013;57:2467–2475. doi:10.1128/AAC.01615-12
30. Kim HB, Wang M, Park CH, Kim EC, Jacoby GA, Hooper DC. oqxAB encoding a multidrug efflux pump in human clinical isolates of Enterobacteriaceae. *Antimicrob Agents Chemother*. 2009;53:3582–3584. doi:10.1128/AAC.01574-08
31. Turton JF, Perry C, Elgohari S, Hampton CV. PCR characterization and typing of *Klebsiella pneumoniae* using capsular type-specific, variable number tandem repeat and virulence gene targets. *J Med Microbiol*. 2010;59:541–547. doi:10.1099/jmm.0.015198-0
32. Compain F, Babosan A, Brisse S, et al. Multiplex PCR for detection of seven virulence factors and K1/K2 capsular serotypes of *Klebsiella pneumoniae*. *J Clin Microbiol*. 2014;52:4377–4380. doi:10.1128/JCM.02316-14
33. Min X, Fu Y, Kong H, et al. Bloodstream infections caused by *Klebsiella pneumoniae*: prevalence of bla KPC, virulence factors and their impacts on clinical outcome. *BMC Infect Dis*. 2018;18:358. doi:10.1186/s12879-018-3109-6
34. Wang X, Xie Y, Li G, et al. Whole-Genome-Sequencing characterization of bloodstream infection-causing hypervirulent *Klebsiella pneumoniae* of capsular serotype K2 and ST374. *Virulence*. 2018;9:510–521. doi:10.1080/21505594.2017.1421894

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>