Preliminary psychometric properties of the Arabic version of Sheu and colleagues Perceived Stress Scale among nursing students at Jordanian universities

This article was published in the following Dove Press journal:
Journal of Multidisciplinary Healthcare

Abdulnaser Algaralleh ¹

Diala Altwalbeh ²

Abdulkarim Alzayyat ³

¹Department of Counseling and Special Education, Faculty of Educational Sciences, Mutah University, Mutah, Jordan; ²Department of Allied Medical Sciences, Faculty of Karak, Al-Balqa Applied University, Salt, Jordan;

³Community Health Nursing Department, Faculty of Nursing, The University of Jordan, Amman, Jordan

Purpose: Nursing is known to be a stressful profession, as it necessitates constant interaction with different individuals in an environment that is described as highly stressed. The stress phenomenon mainly affects the students of nursing more than other health-related students. The Perceived Stress Scale (PSS), which is developed by Sheu and colleagues represents the most implemented instrument to examine stress among nursing students that belong to different cultures in different countries. Previous studies have not fully examined the psychometric properties of the PSS, especially in the Arab World. Therefore, this study aims to assess the psychometric properties of the Arabic version of Sheu and colleagues Perceived Stress Scale (PSS), which is used to measure the clinical stress among nursing students.

Patients and methods: This methodological study utilized a cross-sectional design to test the PSS preliminary psychometric properties among 320 nursing students at 5 Jordanian universities.

Results: The Arabic version of the PSS showed high internal consistency reliability (Cronbach's α was 0.90 in addition to an excellent content validity (CVI was 0.94). The exploratory factor analysis indicated that 54.54% of the total variance was accounted for by the five factors model, which confirmed the construct validity of the Arabic version of PSS. The findings of this study revealed that the Arabic version of PSS can be appropriately implemented among nursing students.

Conclusion: This study supports and recommends the use of the Arabic version of PSS to measure clinical stress among nursing students at Jordanian Universities.

Keywords: validity, reliability, exploratory factor analysis, clinical training, stress

Introduction

Stress has become a commonly used word and concept in both everyday language,¹ and is also a medical concern, for example, the world health organization (who) has described stress as the "21st century epidemic".² Many definitions have been given to this phenomenon; and the emphasis depends on the approach or framework being posited.³ Linguistically speaking, stress is defined as "uncountable or countable pressure or worry caused by the problems in someone's life".⁴ Similarly, Lazarus and Folkman (1984) and based on the transactional model of stress, defined psychological stress as "a particular relationship between the person and the environment that is appraised by the person as taxing or exceeding his or her resources and endangering his or her well-being".⁵

Correspondence: Diala Altwalbeh
Department of Allied Medical sciences,
Faculty of Karak, Al-Balqa Applied
University, Salt, Jordan
Tel +96 279 602 5066
Email diala.tawalbeh@bau.edu.jo

Nursing is a demanding profession requiring constant interaction with different individuals within a complex clinical environment. Stress has many effects on nurses, such as alteration in their professional skills, absenteeism, somatic illnesses, coronary artery diseases, and alcoholism.⁶⁻⁹ Nursing students have been reported as more affected by stress than other health-related students.¹⁰ Stress can build the student's knowledge background and enhance performance when it remains at tolerable levels. However, if stress escalates, students will be more prone to make mistakes, lose confidence in their ability to perform their assigned tasks, and potentially fail.¹¹ In addition, stress can result in health problems such as hypertension, heart diseases, immune deficiency, and depression.¹²

According to Alzayyat and Al-Gamal (2014), nursing students usually encounter stress that is generated from three key sources including academic sources, clinical sources, and personal or social sources of stress. The academic sources of stress comprise assessments and examinations, assignment workload, and the fear of failure. The clinical sources of stress comprise clinical settings, initial clinical experience, the fear of conducting mistakes, the death of patients, and relations with other health team members including superiors' attitudes that are negative and hostile. The personal/social sources involve finance, as well as issues concerning the home-college interface, e.g., lack of free time.¹³ Clinical sources of stress are the most stressful stressors for nursing students.¹⁴⁻¹⁶

To measure stress among nursing students, scholars of nursing used several tools. For example, the Student Stress Survey (SSS) was used to identify the source of stress among Iranian nursing students.¹⁷ The stress in nurse education questionnaire (SNEQ) was used to investigate nursing students' experiences of stress in the United Kingdom.¹⁸ The SNEQ was also used to detect sources of stress among nursing students across five different countries: Albania, Brunei, the Czech Republic, Malta, and Wales.¹⁹ The General Health Questionnaire (GHQ) was used to identify the source of stress among nursing students in Australia²⁰ and Japan.²¹ The Perceived Stress Questionnaire (PSQ) was used to describe stressors among nursing students in the USA.²² However, several systematic reviews reported that the Perceived Stress Scale (PSS), developed by Sheu and colleagues in 1997, has been the most utilized tool for assessing stress in nursing.¹³⁻¹⁵ The PSS was used to assess stressors among nursing students in different countries including Taiwan,²³⁻²⁵ Hong Kong,²⁶ China,²⁷ Philippines,

Greece, and Nigeria,²⁸ Spain,²⁹ Turkey,¹² Jordan,^{30,32-34} and Saudi Arabia.³⁵⁻⁴⁰ Based on the previous literature, nursing students, including Jordanian nursing students, suffer from several clinical stressors that can inflict a negative effect on their psychological well-being and deter their ability to provide optimal nursing care for their patients.

In summary and based on the existing literature, the PSS tool was utilized among nursing students from different cultures in different countries. However, despite extensive use among nursing students from different cultures in different countries, the PSS psychometric properties of the PSS have not been previously investigated thoroughly. Previous studies investigated mainly the consistency reliability but did not mention other details such as construct validity and content validity or the detailed translation process except for a previous study in Spain.²⁹ The PSS tool was used in 11 studies in two Arab countries (Jordan³⁰⁻³⁴ and Saudi Arabia.³⁵⁻⁴⁰) However, a validation of the Arabic language version of the PSS has not been investigated so far. The absence of a validated Arabic instrument to assess the clinical stress in the Arab World, where Arabic is the first language, is a real concern. Therefore, investigating a validated tool, which measures clinical stress among the nursing students, contributes to the body of literature as it provides insights into the clinical stress among nursing students. Moreover, this is essential for assessing the impact of interventions, which aim at decreasing the levels of stress among Arabic speaking nursing students in different countries.

This study aims to evaluate the psychometric properties of the Arabic version of the PSS tool among nursing students at Jordanian universities. This study also aims at achieving the following objectives: (i) to translate the PSS from English into Arabic, (ii) to determine whether the PSS is appropriate for the Jordanian nursing students, and (iii) to examine the reliability and validity of the translated Arabic version of the PSS tool.

Materials and methods

Perceived Stress Scale (PSS): development, reliability and validity

The Perceived Stress Scale (PSS) tool was originally developed by Sheu et al (1997) to assess the extent and type of stress as perceived by nursing students at Taiwanese universities. The PSS consists of 29 items rated on a 5-point Likert scale, where (0= never, 1= almost never, 2= sometimes, 3= fairly often, and 4= very often).

The total score ranges from 0 to 116. Higher score indicates a higher degree of stress. The items were clustered into six factors or subscales that related to the source of stress as follows: 1) the first factor consists of 3 items, and it measures “stress from lack of professional knowledge and skills”, 2) the second factor consists of 8 items that measure “stress from taking care of patients”, 3) the third factor comprises 5 items; this factor measures “stress from assignments and workload”, 4) the fourth factor consists of 6 items that measure “stress from teachers and nursing staff”, 5) the fifth factor comprises 3 items that measure “stress from the clinical environment”, and 6) the sixth factor, which consists of 4 items, measures “stress from peers and daily life”. The exploratory factor analysis (EFA) indicated that 50.7% of the total variance was accounted for by the six factors. Sheu et al (2002) reported good internal consistency reliability of 0.89, whereas the one-week test-retest stability reliability coefficient was $r=0.60$. The content validity index (CVI) was 0.94, which showed excellent levels.^{23,41}

Design, setting and participants

This methodological study utilized a cross-sectional design to test the preliminary psychometric properties of the PSS among nursing students at five Jordanian universities, including four public universities and one private university in the central province of Jordan. Based on the type and location of the university, a stratified random sampling technique was used to select the five universities from a total of 15 universities offering an undergraduate nursing program in Jordan. The nursing students in the selected universities come from diverse socioeconomic and cultural backgrounds. The classes were randomly selected from each of the five selected universities. The sample size for exploratory factor analysis (EFA) is usually determined based on the number of variables (items) on the scale. As a general rule, the minimum ratio is 10–15 participants for each item.⁴² Based on the total number of items in the PSS, which is 29 items, a minimum of 290 participants is required in the study.

Data collection procedures

The data collection procedure was carried out during the academic years from 2015 to 2016. Based on certain arrangements with the administrators of the nursing schools at each of the participating universities, only one elective theoretical class of the second, third, and fourth-year nursing students were randomly selected from each university.

The first-year nursing students do not have clinical courses at hospitals and, therefore, they were not included in the study. Accordingly, the researchers contacted the instructors of those selected classes; they explained the objectives of the study and arranged for the data collection procedure during class sessions. The students' registration list at each class was used to select students randomly. The students, who agreed to participate in the survey, completed two types of self-report questionnaires in 15 mins. The first questionnaire was about the students' demographic data that was designed for this study, whereas the second questionnaire was about the PSS tool.

Ethical considerations

An official permit was obtained to use the PSS English version in this study from the original author. The current study was approved by the Research Ethics Committee at Mutah University (reference number: REC-25/2015).

Scale translation

In this study, the PSS was translated from English into Arabic in accordance with Brislin's model. A well-known method, for constructing reliable and valid instruments for cross-cultural investigations.^{43–46} The English-Arabic translation process involved three steps: 1) the translation was conducted from English (the source Language) into Arabic (the target language) by a bilingual expert, 2) back-translation from Arabic into English was conducted by another bilingual expert, and 3) a comparison between the back-translated versions with the original translated version was carried out. The first two steps were carried out by two (different) clinical nurse specialists. These translators were fluent in English and Arabic, and they were familiar with the nature of this study as they worked as clinical instructors for the nursing students. The third step was undertaken by a bilingual expert (an Assistant Professor in clinical nursing), who compared the PSS back-translated version with the originally translated version item-by-item to transfer the semantic equivalence of the translated items. The translator is fluent in English and knowledgeable in instrument development. Moreover, an English native speaker double checked the PSS back-translated version with the originally translated version to achieve an adequate translation.

Pilot study

To assess the PSS, readability, clarity and reliability of the PSS, and to determine if modifications were required

before conducting the main study, a pilot study was conducted on a sample of 30 nursing students, who were enrolled in a clinical nursing course at one of the selected universities. Before embarking on filling out the questionnaires, the researcher directed the nursing students to add their own comments on whether the items are clear or difficult to grasp. Moreover, the students were given the opportunity to suggest new items for the study questionnaires. Therefore, spaces were provided for the qualitative comments after each questionnaire.

The pilot study revealed a few minor issues. First, regarding the addition of new items of the study tools, most of the students' suggestions were implicitly included in the existing items. For instance, it was commented by several students that excessive assignments constitute a stressor. Item No. 15 in PSS, i.e., (requirements of clinical practice exceed my physical and emotional endurance) implies the students' suggestions. Therefore, no new items were identified. Second, regarding the clarity of the tools, the students reported that PSS items were clear and easy to comprehend. Third, regarding the reliability of the pilot Arabic language PSS, Cronbach's alpha was 0.89 indicating good internal consistency reliability. The technique of "Alpha if item deleted" was used to examine the items separately to establish how every item affected the reliability of the scale.⁴⁷ However, none of the item's deletion resulted in improving the Cronbach's alpha of the total PSS. Therefore, all PSS items were used in this study.

Data analysis

SPSS Statistics 17.0 was used for data analysis. Descriptive statistics were used to describe the participants' demographics. The psychometric properties of the Arabic version of PSS (content validity, construct validity, internal consistency reliability) were measured using several statistical tests, described in detail in the following section.

Content validity

Content validity can be defined as "the degree to which an instrument has an appropriate sample of items for the construct being measured and adequately covers the construct domain".⁴⁸ (p 336) There exist possible differences in the parameters of clinical stress between Taiwanese and Jordanian nursing students. Therefore, the CVI of the PSS Arabic version was evaluated to ensure that the items were familiar to the Jordanian students and their clinical training environment. Therefore, five nursing experts were invited to assess the content validity. All the invited experts had

an adequate background about tools' and they had extensive experience in clinical nursing education. The experts were asked to rate each item of the PSS based on a four-point scale labeled as: "1: not relevant, 2: somewhat relevant, 3: quite relevant, or 4: highly relevant".

Moreover, the panel of experts was asked to evaluate each item separately in the Arabic version of PSS in addition to the overall PSS. The CVI of the total PSS is the proportion of the items that are rated 3 or 4. A CVI score of 0.80 or better indicates good content validity.⁴⁹

Factorial construct validity

The construct validity of the instrument is the degree to which it measures the construct under investigation.⁴⁸ The construct validity of the PSS was assessed through Exploratory Principal Component Factor Analysis (PCA) with the varimax rotation method.⁵⁰ To determine the number of factors to be retained for interpretation, the criterion of eigenvalues greater than 1.0 was adopted.^{50,51} The selection of items for a given factor was based on their factor loadings; items with a minimum factor loading of 0.40 were selected.⁴⁸ Assumptions for PCA (Kaiser–Meyer–Olkin (KMO) and Barlett's Test of Sphericity (BTS) were checked.⁴⁸

Internal consistency reliability

The reliability of an instrument is the consistency with which it measures the target attribute. Reliability can be equated with a measure's stability, consistency, or dependability. The internal consistency reliability is the most widely used reliability approach.⁴⁸ The internal consistency reliability of the total PSS and the identified subscales from EFA were determined by Cronbach's alpha coefficient. Cronbach's alpha is calculated based on the pairwise correlations between items. The normal range of values is in between .00 and 1.00 and higher values reflect higher internal consistency. The "alpha if item deleted" technique was used to identify whether an item's deletion enhanced the Cronbach's alpha coefficient of the total PSS or subscales substantially.⁴⁷

Results

Demographics of the sample

A total of 350 questionnaires were distributed to nursing students. Thirty questionnaires were excluded because of missing data. Table 1 illustrates the demographics of the 320 participants. The nursing students' age ranged from 20 to 25 years ($M=21.15$; $SD=1.42$). The students' academic

Table 1 Demographics characteristics of nursing students (N=320)

Variable	M (SD)	SD	Minimum	Maximum
Age (years)	21.15	1.42	20	25
Academic cumulative score average	2.97	0.33	2.26	3.65
Academic credit hours	16	1.2	12	20
Variable	n	Frequency		
Gender				
Female	272	85.0%		
Male	48	15.0%		
Marital status				
Single	276	86.25%		
Married	36	11.25%		
Divorced	8	2.50%		
Employment status				
Full time	44	13.75%		
Part Time	33	10.31%		
Unemployed	243	75.93%		
Current university year				
2nd year	48	15.0%		
3rd year	218	68.13%		
4th year	54	16.88%		

score ranged from 2.26 to 3.65 out of 4; academic score under 2 is considered weak while 3.65 and above is considered excellent ($M=2.97$; $SD=0.33$). Most of the participants were females ($n=272$; 85.0%), single ($n=276$; 86.25%), unemployed ($n=243$; 75.93%), and third-year students ($n=218$; 68.13%). The students registered between 12 and 20 academic credit hours during the semesters of data collection ($M=16$; $SD=1.2$).

PSS validity

Translation process

Translation of the PSS from English into Arabic was straightforward due to the simplicity and non-technical language of the original PSS. However, the English native speaker, who checked the back-translated version with the original one suggested that item 8, that is, (feeling anxious when trust and acceptance of patients and their families are lacking) and item 13 (fear of having bad grades) in the back-translated version was rewritten. The English native speaker believed that these items should measure the students' fear of having future stressors as reported in the original version. Accordingly, these items were reworded to reflect future stressors in the Arabic version. In the back-translated version, item 8 was reworded to become

(feeling anxious that I will not be gaining the trust and acceptance of patients and their families), and item 13 was reworded to become (fear of receiving future bad grades).

Content validity

Content validity: the findings of the content validity measurements were mainly positive. The computed item-CVI was 1.00 (the maximum) for 20 from a total of the total 29 items. The remaining nine items had an item-cvi of 0.80. According to the Polit and Beck's (2012) criteria, all the 29 items in the Arabic version of PSS were acceptable. With an overall scale-level CVI for the PSS of 0.94, content validity can be considered excellent.⁴⁸

Construct validity

The KOM test result was 0.88, which indicates sampling adequacy for the factor analysis. BTS findings were statistically significant ($X^2=1947.26$; $DF=406$; $P<0.0001$),

Indicating that the correlation matrix is suitable for EFA. Results of the PCA are shown in Table 2. There were 5 factors accounting for 54.54% of the variance. Factor 1 consists of seven items measuring "stress from teachers and nursing staff". Factor 2 consists of six items measuring "stress from taking care of patients". Factor 3 consists of five items measuring "stress from lack of

Table 2 Factor loadings of the Arabic version of PSS items after varimax rotation (N=320)

	Item	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5
25	Lack of care and guidance from teachers	0.75	0.11	0.23	0.15	0.20
14	Teachers do not give fair evaluation on students	0.71	0.13	-0.15	0.27	-0.22
20	Type and content of teacher's instruction do not meet my expectation	0.67	0.03	0.09	-0.16	0.29
21	Pressure from teachers who evaluate students' performance	0.61	0.09	0.17	0.11	0.18
1	Experience discrepancy between theory and practice	0.61	0.07	0.13	0.15	0.25
17	Medical personnel lack empathy and are not willing to help	0.59	0.15	-0.20	0.12	0.06
22	My performances do not meet teacher's expectation	0.54	-0.03	0.05	0.09	0.10
3	Unable to provide patients with good nursing care	0.03	0.77	0.04	0.01	0.02
2	Lack of experience and ability in providing nursing care and in making judgment	0.11	0.66	0.27	-0.05	0.08
11	Unable to reach one's expectation	0.15	0.65	0.13	0.27	0.01
10	Unable to provide appropriate responses to doctor's, teacher's and patients' questions	0.02	0.65	0.23	-0.06	0.15
4	Do not know how to communicate with patients	0.12	0.61	0.09	0.15	0.17
12	Do not know how to help patients with physio- psycho-social problems	0.05	0.61	0.11	-0.12	0.18
6	Unfamiliar with patients' diagnosis and treatments	0.04	-0.23	0.80	0.30	0.18
7	Unfamiliar with medical history and terms	0.11	0.03	0.79	0.05	0.20
28	Unfamiliar with ward's facilities	0.09	0.17	0.59	0.21	-0.28
26	Unfamiliar with professional nursing skills when dealing with patients	0.16	0.04	0.56	-0.15	0.21
18	Do not know how to discuss patients' illness with teachers, and medical and nursing personnel	0.09	0.12	0.57	0.27	0.13
19	Feel that dull and inflexible clinical practice affects one's family and social life	0.12	0.17	0.24	0.73	-0.03
15	Feel that the requirements of clinical practice exceed one's physical and emotional endurance	0.11	0.08	0.31	0.72	0.05
5	Feel that clinical practice affects one's involvement in extracurricular activities	0.01	0.03	0.50	0.64	0.12
27	Feel stressed from the rapid change in patient's condition	0.23	0.01	0.20	0.55	-0.12
29	Feel stressed in the hospital environment where clinical practice takes place	-0.14	0.07	0.22	0.54	0.15
16	Experience pressure from the nature and quality of clinical practice	0.03	0.14	0.17	0.47	-0.01
24	Experience competition from peers in school and clinical practice	0.10	0.21	0.08	0.28	0.61
8	Feeling anxious that I will not be gaining the trust and acceptance of patients and their families	0.40	0.30	0.01	0.02	0.55
9	Experience difficulties in changing from the role of a student to that of a nurse.	0.00	-0.21	0.08	0.11	0.53
13	Fear of receiving future bad grades	0.12	0.05	0.42	0.15	0.52
	Initial eigenvalues	8.80	2.52	1.68	1.56	1.26
	Percentages of variance explained	30.34	8.68	5.79	5.38	4.35
	Cumulative variance	30.34	39.02	44.81	50.19	54.54

Notes: Item 23 has been excluded since it did not have any significant load (≥ 0.4) in any of the five identified factors. Factor loadings in bold mean the items were assigned to the designated factor.

professional knowledge and skills". Factor 4 consists of six items measuring "stress from the nature of clinical practice". Factor 5 consists of four items measuring "stress from the student's role demands". From the original 29-item set, item 23 was removed since the factor loading of this item was lower than 0.40, and this item did not fit well into one of the five identified factors. Three items with substantial loadings on more than one factor (items 5, 8, and 13) were assigned to the factor with the highest loading. Table 3 illustrates a comparison between the English and Arabic versions of the PSS. The table showed that factors 1, 2 & 3 in the Arabic version correspond to factors 4, 2 and 1 in the English version respectively.

Internal consistency reliability

The total PSS scale alpha coefficient was 0.90, which indicates an excellent internal consistency. The Cronbach's alphas of the five subscales ranged from 0.65 to 0.83. All items were retained as "alpha if item deleted" showed that deletion of any item did not lead to an improvement in the Cronbach's alpha of the total PSS or its subscales. The reliability tests of the total PSS and the five factors in this study are summarized in Table 4.

Discussion

Clinical stress has frequently been frequently evaluated using the PSS, which was designed for nursing students with adequate English language skills. However,

applying the original, English language PSS in the Jordanian context potentially has limitations due to language and cultural variations. This study aimed to translate and evaluate the psychometric properties of the PSS among Jordanian nursing students. It was concluded that the translation process was successfully conducted and that the PSS Arabic version is suitable to be utilized for nursing students that are native speakers of Arabic. This result was further supported by the panel of experts, who examined the CVI of the Arabic version of PSS and reported that the items adequately measured all the dimensions of the clinical stress construct. The CVI of the Arabic version PSS was 0.94, which is similar to the original scale.²³ This reflects an excellent level.⁴⁸

The Arabic language PSS resolved in to a five-factors model reflecting different sources of stress perceived by students during clinical practice. Accounting for 54.54% of the total variance, in comparison with the six-factor model, which explained 50.7% and 56.11% of the total variance of the original version²³ and the Spanish version²⁹ respectively.

While the factors total number and order differed somewhat, similar factors emerged across the two versions and for three of the factors in the Arabic version, the names of the factors were retained, because the items' grouping of the factors in the Arabic version is somewhat like those of the English version. The fourth factor in the Arabic PSS "stress from the nature of clinical practice"

Table 3 Comparison between the English and Arabic versions of the PSS of 28 Items loaded on five factors

Factor rank	English PSS Subscales	Factor rank	Arabic PSS Subscales
	# of items, items loaded		# of items, items loaded
4	Stress from teachers and nursing staff (6 items), 1, 14, 17, 18, 20, 25	1	Stress from teachers and nursing staff (7 items) 1, 14, 17, 20, 21, 22, 25.
2	Stress from taking care of patients (8 items), 2, 3, 4, 8, 9, 10, 11, 12	2	Stress from taking care of patients (8 items) (6 items) 2, 3, 4, 10, 11, 12
1	Stress from lack of professional knowledge and skills (3 items), 6, 7, 26	3	Stress from lack of professional knowledge and skills (5 items) 6, 7, 18, 26, 28
3	Stress from assignments and workload (5 items), 11, 13, 15, 16, 19	4	Stress from the nature of clinical practice (6 items) 5, 15, 16, 19, 27, 29,
6	Stress from peers and daily life (4 items), 24, 21, 5, 23 (omitted from new scale)	5	Stress from student's role demands (4 items) 8, 9, 13, 24
5	Stress from clinical environment (3 items) 29, 28, 27		

Notes: Items in bold were matched in the subscales between the English and Arabic versions of the PSS.

Table 4 The factors structure of the Arabic PSS scales and its Cronbach's alpha (α)

Item number	28 Items/5 PSS Subscales	α
	Total perceived stress scale	0.90
Factor 1	Stress from teachers and nursing staff	0.83
25	Lack of care and guidance from teachers	
1	Experience discrepancy between theory and practice	
22	My performances do not meet teacher's expectation	
20	Type and content of teacher's instruction do not meet my expectations	
14	Teachers do not give fair evaluation on students	
21	Pressure from teachers who evaluate students' performance	
17	Medical personnel lack empathy and are not willing to help	
Factor 2	Stress from taking care of patients	0.78
11	Unable to reach one's expectation	
4	Do not know how to communicate with patients	
10	Unable to provide appropriate responses to doctor's, teacher's and patients' questions	
3	Unable to provide patients with good nursing care	
12	Do not know how to help patients with physio- psycho-social problems	
2	Lack of experience and ability in providing nursing care and in making judgment	
Factor 3	Stress from lack of professional knowledge and skills	0.83
28	Unfamiliar with ward's facilities	
18	Do not know how to discuss patients' illness with teachers, and medical and nursing personnel	
7	Unfamiliar with medical history and terms	
6	Unfamiliar with patients' diagnosis and treatment	
26	Unfamiliar with professional nursing skills when dealing with patients	
Factor 4	Stress from the nature of clinical practice	0.78
19	Feel that dull and inflexible clinical practice affects one's family and social life	
27	Feel stressed from the rapid change in patient's condition	
29	Feel stressed in the hospital environment where clinical practice takes place	
16	Experience pressure from the nature and quality of clinical practice	
15	Feel that the requirements of clinical practice exceed one's physical and emotional endurance.	
5	Feel that clinical practice affects one's involvement in extracurricular activities	
Factor 5	Stress from student's role demands	0.65
8	Feeling anxious that I will not be gaining the trust and acceptance of patients and their families	
9	Experience difficulties in changing from the role of a student to that of a nurse.	
13	Fear of receiving future bad grades	
24	Experience competition from peers in school and clinical practice	

comprises items from the third factor “stress from assignments and workload”, as well as the fifth factor of the English PSS. The remaining items were included under the fifth factor, which was labeled as “stress from the student's role demands. Fourteen from a total of 28 items in the Arabic version of the PSS were distributed across 3 similar factors to the original English version. These 3 factors in the Arabic version are therefore conceptually equivalent with the original instrument. Another Fourteen items were loaded onto factors that were different from those of the original English version. Such inconsistencies are most likely to happen because of dissimilarities that are inherent

to the samples utilized for analysis (such as sample size and sample recruitment method) or differences in the composition of the clinical educational programs rather than because of contradictory representations of the clinical stress construct.

The results of this study, however, indicated that the proposed PSS model needs some modifications, especially in terms of the included items. Given the obviously inappropriate loadings of item 23 remarked in EFA, it was determined to eliminate this item from the proposed PSS model as it should not be incorporated in the computation of the PSS scores. Item 23, that is, (cannot get along with

other peers in the group) may represent a significant stressor in the earlier context of the PSS development. In the Jordanian context, the cultural factors may normalize this stress in the perceptions of nursing students as there are strong and positive affiliations between students' classmates in Jordan. Furthermore, these affiliations are usually used by the students to buffer stress instead of being a source of stress.⁵² Therefore, a critical evaluation of item 23 wording, as well as replication of the study findings in different contexts (e.g., in another Arab world country) are recommended in future studies. Moreover, further research should aim to confirm the status of item 23 in the PSS model by confirmatory factor analysis procedures.

Regarding the PSS reliability, the Cronbach's alpha coefficient of the total PSS scale was 0.90 in this study compared with 0.89 of the original scale. This is in accordance with results reported in previous studies from Jordan³³ and Saudi Arabia,³⁸ but less than those reported by studies from Spain²⁹ and Turkey.¹² In general, The results indicated that PSS has excellent internal consistency reliability. According to Polit and Beck (2012), the first four identified factors of the PSS achieved reasonable internal consistency reliability ($\alpha \geq 0.7$). However, the last identified factor (i.e., student's role demands subscale) achieved relatively inadequate internal consistency reliability ($\alpha < 0.7$). This is partly because of the minimal number of items loading on the fifth factor.⁴⁸

In summary, the five-factor model of PSS was found to be valid, reliable, and empirically supported. The results of this study showed that the 28-item PSS and the five-factor solution achieved the necessary validity of the scale among the nursing students at Jordanian universities.

Limitations

The findings of this study are promising, and they contribute to the development and validation of the PSS model. However, a few limitations should be acknowledged. First, the utilized sample is comparatively homogeneous (female nursing students in the third year of their study). Therefore, generalizing the study results can be limited. Second, the current study has only examined the content and constructs validity in addition to consistency reliability. Therefore, the authors recommended further studies to explore other dimensions of psychometric properties such as the stability reliability, the criterion-related validity, and predictive validity of the PSS model. However, the confirmatory analysis of the five-factor model of the PSS is a necessary second step, which is

highly recommended before the validation of this tool to measure clinical education stress among Arab-native nursing students.⁴² These studies would be useful in evaluating stress resources during clinical training. At the same time, they would provide opportunities to compare the differences in the perceived level of stress between Eastern and Western nursing students.

Conclusion

This study has offered initial evidences for the validity and reliability of the Arabic version of the PSS. The Arabic version of the PSS appears to be an appropriate tool to measure nursing students' clinical stress. It showed high internal consistency reliability (Cronbach's alpha was 0.90), and excellent content validity (CVI 0.94). The exploratory factor analysis indicated that 54.54% of the total variance was accounted for by the five factors model and supported the construct validity of the Arabic version of PSS. Future work should validate the hypothesized PSS model in this study in a more diverse and larger sample of nursing students, using more advanced confirmatory analyses that boost the obtained results of this exploratory study. Moreover, the impact of deleting or re-development of the low-loading items on the whole PSS performance should be verified. To conclude, this study supports and recommends the use of the Arabic version of PSS to measure clinical stress among nursing students at Jordanian Universities.

Disclosure

The researchers declare that there are no conflicts of interest to disclose.

References

- Goodnite PM. Stress: a concept analysis. In: *Nursing Forum*. Vol. 49. Wiley Online Library; 2014: 71–74. doi: 10.1111/nuf.12044
- Fink G. *Stress: Concepts, Cognition, Emotion, and Behavior: Handbook of Stress Series*. Vol. 1. Academic Press; 2016.
- Carrieri-Kohlman V, Lindsey AM, West CM. *Pathophysiological Phenomena in Nursing: Human Responses to Illness*. Saunders; 2003.
- Oxford Advanced Learner's Dictionary; 2019. Available from: https://www.oxfordlearnersdictionaries.com/definition/english/stress_1?q=stress. Accessed August 26, 2019.
- Lazarus RS, Folkman S. *Stress, Appraisal, and Coping*. Springer publishing company; 1984.
- Beck CT. Secondary traumatic stress in nurses: a systematic review. *Arch Psychiatr Nurs*. 2011;25(1):1–10. doi:10.1016/j.apnu.2010.05.005
- Kane PP. Stress causing psychosomatic illness among nurses. *Indian J Occup Environ Med*. 2009;13(1):28. doi:10.4103/0019-5278.50721
- McVicar A. Workplace stress in nursing: a literature review. *J Adv Nurs*. 2003;44(6):633–642. doi:10.1046/j.0309-2402.2003.02853.x
- Wu H, Chi T, Chen L, Wang L, Jin Y. Occupational stress among hospital nurses: cross-sectional survey. *J Adv Nurs*. 2010;66(3):627–634. doi:10.1111/j.1365-2648.2009.05203.x

10. Beck DL, Hackett MB, Srivastava R, McKim E, Rockwell B. Perceived level and sources of stress in university professional schools. *J Nurs Educ*. 1997;36(4):180–186.
11. Wallace L, Bourke MP, Tormoehlen LJ, Poe-Greskamp MV. Perceptions of clinical stress in baccalaureate nursing students. *Int J Nurs Educ Scholarsh*. 2015;12(1):91–98. doi:10.1515/ijnes-2014-0056
12. Karaca A, Yildirim N, Ankarali H, AÇIKGÖZ F, AKKUŞ D. Nursing Students' perceived levels of clinical stress, stress responses and coping behaviors. *J Psychiatr Nursing/Psikiyatri Hemsireleri Dern*. 2017;8(1):32–39. doi:10.14744/phd.2017.22590
13. Alzayyat A, Al-Gamal E. A review of the literature regarding stress among nursing students during their clinical education. *Int Nurs Rev*. 2014;61(3):406–415. doi:10.1111/inr.12114
14. Labrague LJ, McEnroe-Petitte DM, Gloe D, Thomas L, Papathanasiou IV, Tsaras K. A literature review on stress and coping strategies in nursing students. *J Ment Heal*. 2017;26(5):471–480. doi:10.1080/09638237.2016.1244721
15. Labrague LJ, McEnroe-Petitte DM, Al Amri M, Fronda DC, Obeidat AA. An integrative review on coping skills in nursing students: implications for policymaking. *Int Nurs Rev*. 2018;65(2):279–291. doi:10.1111/inr.12393
16. Pulido-Martos M, Augusto-Landa JM, Lopez-Zafra E. Sources of stress in nursing students: a systematic review of quantitative studies. *Int Nurs Rev*. 2012;59(1):15–25. doi:10.1111/j.1466-7657.2011.00939.x
17. Seyedfatemi N, Tafreshi M, Hagani H. Experienced stressors and coping strategies among Iranian nursing students. *BMC Nurs*. 2007;6(1):11. doi:10.1186/1472-6955-6-11
18. Edwards D, Burnard P, Bennett K, Hebden U. A longitudinal study of stress and self-esteem in student nurses. *Nurse Educ Today*. 2010;30(1):78–84. doi:10.1016/j.nedt.2009.06.008
19. Burnard P, Edwards D, Bennett K, et al. A comparative, longitudinal study of stress in student nurses in five countries: Albania, Brunei, the Czech Republic, Malta and Wales. *Nurse Educ Today*. 2008;28(2):134–145. doi:10.1016/j.nedt.2007.04.002
20. Lo R. A longitudinal study of perceived level of stress, coping and self-esteem of undergraduate nursing students: an Australian case study. *J Adv Nurs*. 2002;39(2):119–126. doi:10.1046/j.1365-2648.2000.02251.x
21. Yamashita K, Saito M, Takao T. Stress and coping styles in Japanese nursing students. *Int J Nurs Pract*. 2012;18(5):489–496. doi:10.1111/j.1440-172X.2012.02056.x
22. Wolf L, Stidham AW, Ross R. Predictors of stress and coping strategies of US accelerated vs. generic baccalaureate nursing students: an embedded mixed methods study. *Nurse Educ Today*. 2015;35(1):201–205. doi:10.1016/j.nedt.2014.07.005
23. Sheu S, Lin H-S, Hwang S-L. Perceived stress and physio-psychosocial status of nursing students during their initial period of clinical practice: the effect of coping behaviors. *Int J Nurs Stud*. 2002;39(2):165–175. doi:10.1016/S0020-7489(01)00016-5
24. Chen Y-W, Hung C-H. Predictors of Taiwanese baccalaureate nursing students' physio-psychosocial responses during clinical practicum. *Nurse Educ Today*. 2014;34(1):73–77. doi:10.1016/j.nedt.2013.02.021
25. Li H, Wang LS, Lin Y, Lee I. The effect of a peer-mentoring strategy on student nurse stress reduction in clinical practice. *Int Nurs Rev*. 2011;58(2):203–210. doi:10.1111/j.1466-7657.2010.00839.x
26. Chan CKL, So WKW, Fong DYT. Hong Kong baccalaureate nursing students' stress and their coping strategies in clinical practice. *J Prof Nurs*. 2009;25(5):307–313. doi:10.1016/j.profnurs.2009.01.018
27. Zhao F, Lei X, He W, Gu Y, Li D. The study of perceived stress, coping strategy and self-efficacy of Chinese undergraduate nursing students in clinical practice. *Int J Nurs Pract*. 2015;21(4):401–409. doi:10.1111/ijn.12273
28. Labrague LJ, McEnroe-Petitte DM, Papathanasiou IV, et al. A cross-country comparative study on stress and quality of life in nursing students. *Perspect Psychiatr Care*. 2018;54(4):469–476. doi:10.1111/ppc.12248
29. Jimenez C, Navia-Osorio PM, Diaz CV. Stress and health in novice and experienced nursing students. *J Adv Nurs*. 2010;66(2):442–455. doi:10.1111/j.1365-2648.2009.05183.x
30. Al-Zayyat AS, Al-Gamal E. Perceived stress and coping strategies among Jordanian nursing students during clinical practice in psychiatric/mental health courses. *Int J Ment Health Nurs*. 2014;23(4):326–335. doi:10.1111/inm.12054
31. Shaban IA, Khater WA, Akhu-Zaheya LM. Undergraduate nursing students' stress sources and coping behaviours during their initial period of clinical training: a Jordanian perspective. *Nurse Educ Pract*. 2012;12(4):204–209. doi:10.1016/j.nepr.2012.01.005
32. Khater WA, Akhu-Zaheya LM, Shaban IA. Sources of stress and coping behaviours in clinical practice among baccalaureate nursing students wejdan. *Int J Humanit Soc Sci*. 2014;4(6):194–202.
33. Akhu-Zaheya LM, Shaban IA, Khater WA. Nursing students' perceived stress and influences in clinical performance. *Int J Adv Nurs Stud*. 2015;4(2):44. doi:10.14419/ijans.v4i2.4311
34. Alzayyat A, Al-Gamal E. Correlates of stress and coping among Jordanian nursing students during clinical practice in psychiatric/mental health course. *Stress Heal*. 2016;32(4):304–312. doi:10.1002/smi.2606
35. Hamaideh SH, Al-Omari H, Al-Modallal H. Nursing students' perceived stress and coping behaviors in clinical training in Saudi Arabia. *J Ment Heal*. 2017;26(3):197–203. doi:10.3109/09638237.2016.1139067
36. Alsaqri SH. Stressors and coping strategies of the Saudi nursing students in the clinical training: a cross-sectional study. *Educ Res Int*. 2017;2017. doi:10.1155/2017/4018470.
37. Aedh AI, Elfaki NK, Mohamed IA. Factors associated with stress among nursing students (Najran University-Saudi Arabia). *IOSR J Nurs Heal Sci*. 2015;4(6):33–38.
38. Al-Gamal E, Alhosain A, Alsunaye K. Stress and coping strategies among Saudi nursing students during clinical education. *Perspect Psychiatr Care*. 2018;54(2):198–205. doi:10.1111/ppc.12223
39. Ismaile S. Perceived clinical stressors among Saudi nursing students. *Open J Nurs*. 2017;7(04):463. doi:10.4236/ojn.2017.74036
40. Mohamed BM, Ahmed ES. Perception of nursing students towards clinical stressors in the faculty of applied medical sciences-Al Jouf University-Saudia Arabia. *J Am Sci*. 2012;8:608–617.
41. Sheu S, Lin HS, Hwang SL, Yu PJ, Hu WY, Lou MF. The development and testing of perceived stress scale of clinical practice. *Nurs Res (Republic China)*. 1997;5(4):341–351.
42. Plichta SB, Kelvin EA, Munro BH. *Munro S Statistical Methods for Health Care Research*. Wolters Kluwer Health/Lippincott Williams & Wilkins; 2013.
43. Brislin RW, Lonner WJ, Thorndike RM. *Cross-cultural Research Methods*. 1973.
44. Brislin RW. Back-translation for cross-cultural research. *J Cross Cult Psychol*. 1970;1(3):185–216. doi:10.1177/135910457000100301
45. Brislin RW. The wording and translation of research instruments. In: Lonner WJ, Berry JW, editors. *Cross-Cultural Research and Methodology Series. Field Methods in Cross-Cultural Research*. Sage Publications; 1986:137–164.
46. Jones PS, Lee JW, Phillips LR, Zhang XE, Jaceldo KB. An adaptation of Brislin's translation model for cross-cultural research. *Nurs Res*. 2001;50(5):300–304.
47. Raykov T. Alpha if item deleted: A note on loss of criterion validity in scale development if maximizing coefficient alpha. *Br J Math Stat Psychol*. 2008;61(2):275–285. doi:10.1348/000711007X188520
48. Polit DF, Beck CT. *Nursing Research: Generating and Assessing Evidence for Nursing Practice*. Lippincott Williams & Wilkins; 2012.
49. Polit DF, Beck CT. *Nursing Research: Principles and Methods*. Lippincott Williams & Wilkins; 2004.
50. Field A. *Discovering Statistics Using IBM SPSS Statistics*. Sage; 2013.

51. Fabrigar LR, Wegener DT, MacCallum RC, Strahan EJ. Evaluating the use of exploratory factor analysis in psychological research. *Psychol Methods*. 1999;4(3):272. doi:10.1037/1082-989X.4.3.272
52. Hamdan-Mansour AM, Dawani HA. Social support and stress among university students in Jordan. *Int J Ment Health Addict*. 2008;6(3):442–450. doi:10.1007/s11469-007-9112-6

Journal of Multidisciplinary Healthcare

Dovepress

Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal

covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>