Coproduction Of MCR-9 And NDM-I By Colistin-Resistant *Enterobacter hormaechei* Isolated From Bloodstream Infection

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Yi Yuan^{1,*}

Ying Li^{2,*}

Guangxi Wang³

Chengwen Li²

Li Xiang³

Junping She³

Yan Yang³

Fangcai Zhong¹

Luhua Zhang³

¹Department of Clinical Laboratory, The First People's Hospital of Neijiang, Sichuan, People's Republic of China;

²Department of Immunology, School of Basic Medical Sciences, Southwest Medical University, Luzhou, Sichuan, People's Republic of China; ³Department of Pathogenic Biology, School of Basic Medical Sciences, Southwest Medical University, Luzhou, Sichuan, People's Republic of China

*These authors contributed equally to this work

Background: Colistin acts as the last line of defense against severe infections caused by carbapenem-resistant *Enterobacteriaceae*. Infections caused by extensively drug-resistant isolates coproducing MCR and carbapenemases have posed a serious public health concern.

Purpose: In this study, we reported the first clinical colistin and carbapenem-resistant *Enterobacter hormaechei* isolate SCNJ07 coharboring *bla*_{NDM-1} and *mcr-9* from a patient with bloodstream infection in China.

Methods: Bacterial antimicrobial susceptibility testing was performed using the broth microdilution method. Conjugation assay was carried out to investigate the transferability of *mcr-9* and *bla*_{NDM-1}. Whole-genome sequencing of strain SCNJ07 was performed using an Illumina HiSeq system and the genetic characteristics of the *mcr-9*- and *bla*_{NDM-1}-harboring plasmids were analyzed.

Results: Conjugation assays revealed that both *bla*_{NDM-1} and *mcr-9* genes could successfully transfer their resistance phenotype to *Escherichia coli* strain J53. Whole genome sequencing showed that SCNJ07 possessed an FIB36:FIH4 type self-transmissible plasmid bearing *bla*_{NDM-1}, which possessed high similarity to previously reported pRJF866 in China. *mcr-9* was located on a ~28-kb self-transmissible plasmid pMCR-SCNJ07 with both IncHI2 and IncR replicons. Two copies of intact IS903 that bracketed a ~8-kb region containing the *mcr-9* gene were identified in pMCR-SCNJ07. BLASTn analysis revealed that a number of *mcr-9*-positive plasmids have been around for a while among *Enterobacteriaceae* worldwide.

Conclusion: This study reveals the likelihood of a wide dissemination of this newly identified colistin resistance gene *mcr-9* among *Enterobacteriaceae*. Further surveillance is urgently needed to understand the prevalence and dissemination of *mcr-9*, thereby facilitating establishment of measures to control its spread.

Keywords: colistin, *mcr-9*, CRE, IS903

Introduction

Carbapenem-resistant *Enterobacteriaceae* (CRE)-related infections are global public health issues.¹ The New Delhi metallo-β-lactamase (NDM) is one of the most common carbapenemases worldwide.² Polymyxins are among the last-line therapeutic options to treat serious infections caused by CRE.³ However, concerns were raised regarding the increasing prevalence of the plasmid-borne colistin resistance gene, *mcr-1*, which has been detected from the environment, food, animals and humans around the world since its first discovery in China in late 2015.⁴⁻⁶ More worrisomely, cases of infection caused by extensively drug-resistant isolates that

Correspondence: Luhua Zhang
Department of Pathogenic Biology,
School of Basic Medical Sciences,
Southwest Medical University, Xiang Lin
Road, Longmatan District, Luzhou City
646000, Sichuan Province, People's
Republic of China
Tel/Fax +86 830 3160073
Email zhluhua@swmu.edu.cn

coproduce MCR-1 and NDM have been reported by groups from different regions.^{7–13}

Following a recent study which identified an additional *mcr* homolog, *mcr-9*, in a *Salmonella enterica* serotype Typhimurium isolate,¹⁴ we isolated an extensively drug-resistant *Enterobacter hormaechei* strain, which coproduces NDM-1 and MCR-9, from a patient with bloodstream infection. The genetic characteristics of the *bla*_{NDM-1} and *mcr-9*-harboring plasmids were analyzed. In addition, comparative genetic analysis of *mcr-9* in pMCR-SCNJ07 and closely related plasmids were also carried out. To the best of our knowledge, this is the first report describing a clinical colistin and carbapenem-resistant *E. hormaechei* isolate coharboring *bla*_{NDM-1} and *mcr-9* in China.

Materials And Methods

Bacterial Isolation And Identification

The *E. hormaechei* strain SCNJ07 was recovered from the blood sample of a 50-year-old male patient with obstructive jaundice in a hospital in Sichuan, China, in September 2018. It was initially identified as *Enterobacter cloacae* using the Vitek-2 compact system (bioMérieux, Marcy-l'Étoile, France) and by species identification established by sequencing of the 16S rRNA gene amplified with the universal primers 27F and 1492R.¹⁵ The presence of the acquired carbapenemase genes (*bla*_{KPC}, *bla*_{NDM}, *bla*_{GES}, *bla*_{IMB}, *bla*_{OXA-48}, and *bla*_{VIM}) and *mcr* genes (*mcr-1* to *mcr-8*) in SCNJ07 was determined by PCR amplification as described previously.^{16–21} The detection of *mcr-9* was amplified using the primers MCR9-F (5'-CTTTCCATAACAGCGAGACA C-3') and MCR9-R (5'-GTATCCTTCCTGCCATCCTC-3').

Antimicrobial Susceptibility Testing

In vitro susceptibility tests of ceftazidime, ceftriaxone, cefepime, cefazolin, cefotetan, imipenem, aztreonam, ciprofloxacin, gentamycin, tobramycin, levofloxacin, amikacin, piperacillin-tazobactam and trimethoprim-sulfamethoxazole were performed using Vitek-2 system. The minimum inhibitory concentrations (MICs) of meropenem, colistin, polymyxin B, doxycycline, fosfomycin and tigecycline against the strain were determined using the microdilution broth method following recommendations of the Clinical Laboratory Standards Institute (CLSI) (CLSI, 2019). Breakpoints of colistin and tigecycline were interpreted according to the European Committee on Antimicrobial Susceptibility Testing (EUCAST) (<http://www.eucast.org/>); otherwise, we applied those defined by the CLSI.

Conjugation Assay

Conjugation experiments were carried out using broth-based methods with *Escherichia coli* J53 (azide-resistant) as the recipient strain. Transconjugants were selected on LB agar plates containing 150 µg/mL sodium azide plus 1 µg/mL meropenem or 4 µg/mL colistin. The presence of *bla*_{NDM-1} or *mcr-9* in transconjugants was confirmed by PCR and Sanger sequencing.

Genome Sequencing And Analysis

Total genomic DNA of *E. coli* isolates was extracted using Rapid Bacterial Genomic DNA Isolation Kit (Sangon Biotech, Shanghai, China) according to the manufacturer's protocol. Purified DNA was subjected to whole genomic sequencing on the Illumina HiSeq 2000 system with the 150-bp paired-end approach and 150× coverage. Reads were trimmed using Trimmomatic²² and were then assembled using the SOAP *de novo* program.²³ Annotation was carried out using Prokka.²⁴ The species identification was performed by average nucleotide identity (ANI) analysis with JSpeciesWS (<http://jspecies.ribohost.com/jspeciesws/#analyse>). Antimicrobial resistance genes were identified using ResFinder v3.1 of the Center for genomic epidemiology (<http://genomicepidemiology.org/>).

Plasmid Sequencing And Analysis

Plasmid DNA from transconjugants containing *bla*_{NDM-1} or *mcr-9* was extracted and sequenced using the Illumina HiSeq system. After filtering J53 chromosomal DNA data and assembling the remaining reads, the plasmid carrying *bla*_{NDM-1} or *mcr-9* was completely circularized using PCR and Sanger sequencing to fill in gaps between contigs. The plasmid replicon type and MLST were determined using the PlasmidFinder (<https://cge.cbs.dtu.dk/services/PlasmidFinder/>) and pMLST tools (<https://cge.cbs.dtu.dk/services/pMLST/>). The annotations of the plasmid sequences were conducted using the RAST tools and edited manually.²⁵ Sequence alignment of *mcr-9*-carrying plasmids was performed using BLAST and visualized with Easyfig v 2.2.3.²⁶ Alignments with highly homologous complete plasmid sequences of pMCR-SCNJ07 available in NCBI were performed by using the BRIG tool.²⁷

Nucleotide Sequence Accession Numbers

Draft genome sequence of the strain SCNJ07 has been deposited into GenBank under the accession no. VBSC000000. The complete sequences of pNDM-SCNJ07 and pMCR-SCNJ07 have been deposited into GenBank under accession no. MK933278 and MK933279, respectively.

Ethics Statement

The clinical isolate *E. hormaechei* SCNJ07 was generated as part of the routine hospital laboratory procedure. The Ethics Committee of the Southwest Medical University exempted this study from review because the present study only focused on bacteria and patient intervention was not required.

Results And Discussion

Antimicrobial Susceptibility Of The Strain *E. hormaechei* SCNJ07

Susceptibility testing by Vitek-2 system showed that the strain SCNJ07 was resistant to all tested drugs, including ceftazidime (MIC, ≥ 64 $\mu\text{g/mL}$), ceftriaxone (MIC, ≥ 64 $\mu\text{g/mL}$), cefepime (MIC, ≥ 64 $\mu\text{g/mL}$), cefazolin (MIC, ≥ 64 $\mu\text{g/mL}$), cefotetan (MIC, ≥ 64 $\mu\text{g/mL}$), imipenem (MIC, ≥ 16 $\mu\text{g/mL}$), aztreonam (MIC, ≥ 64 $\mu\text{g/mL}$), ciprofloxacin (MIC, ≥ 4 $\mu\text{g/mL}$), gentamycin (MIC, ≥ 16 $\mu\text{g/mL}$), tobramycin (MIC, ≥ 16 $\mu\text{g/mL}$), levofloxacin (MIC, ≥ 8 $\mu\text{g/mL}$), amikacin (MIC, ≥ 64 $\mu\text{g/mL}$), piperacillin-tazobactam (MIC, ≥ 128 $\mu\text{g/mL}$) and trimethoprim-sulfamethoxazole (MIC, ≥ 320 $\mu\text{g/mL}$). Additional broth microdilution testing²⁸ showed that it also exhibited resistance to meropenem (MIC, 256 $\mu\text{g/mL}$), doxycycline (MIC, 128 $\mu\text{g/mL}$), colistin (MIC, 16 $\mu\text{g/mL}$) and polymyxin B (MIC, 8 $\mu\text{g/mL}$) but was susceptible to fosfomycin (MIC, 64 $\mu\text{g/mL}$) and tigecycline (MIC, 2 $\mu\text{g/mL}$). PCR and sequencing analysis showed that the strain SCNJ07 harbored both a *bla*_{NDM-1} carbapenemase gene and the recently identified plasmid-borne colistin resistance gene *mcr-9*.

Location And Transferability Of *mcr-9* And *bla*_{NDM-1}

Conjugation assays showed that *bla*_{NDM-1} and *mcr-9* were successfully transferred to *E. coli* J53 at the frequency of $\sim 10^{-2}$ and 10^{-4} (transconjugant/recipient), respectively. Compared with the MICs for the recipient strain J53, transformants containing *bla*_{NDM-1} showed a 128-fold increase for meropenem (from 0.5 to 64 $\mu\text{g/mL}$) and those harboring *mcr-9* 16-fold for colistin (from 0.5 to 8 $\mu\text{g/mL}$). These findings revealed that *bla*_{NDM-1} and *mcr-9* were functional and carried by two self-transmissible plasmids.

Genome Characteristics Of The Strain SCNJ07

Draft genome sequence of SCNJ07 was assembled into 120 contigs (112 were >1000 bp in length), which comprises

5,321,397 bp, with a 54.77% GC content. Species identification based on ANI analysis confirmed that the strain SCNJ07 actually belongs to *E. hormaechei*, as it only had an 86.32% identity (75.45% query coverage) to *E. cloacae* subsp. *cloacae* ATCC 13,047, but a 98.41% identity (80.30% coverage) to *E. hormaechei* subsp. *steigerwaltii* strain DSM 16691, obviously above the 95–96% cut-off for defining a bacterial species.²⁹

Resistance Profile Of The Strain SCNJ07

Analysis of the whole genome sequence of SCNJ07 allowed us to more confidently conclude that *bla*_{NDM-1} and *mcr-9* co-existed in this strain. Besides, in consistency with its multidrug resistance phenotype, SCNJ07 also had multiple genes mediating resistance to β -lactams (*bla*_{CTX-M-3}, *bla*_{TEM-1B}, *bla*_{SHV-12} and *bla*_{ACT-7}), aminoglycosides (*aac*(6')-IIa, *aadA16*, *aac*(6')-Ib3 and *rmtC*), fluoroquinolones (*qnrS1* and *aac*(6')-Ib-cr), macrolide (*mph*(A)), rifampicin (*ARR-3*), tetracycline (*tet*(D)), sulfonamide (*sul1*) and trimethoprim (*dfrA16*). *E. hormaechei*, a common nosocomial pathogen of clinical significance, was reported in several outbreaks of sepsis in neonatal intensive care units in the USA and in Brazil,³⁰ while it was only sporadically reported in China.³¹ The identification of *E. hormaechei* SCNJ07 coproducing NDM-1 and MCR-9 in this study highlights the need to enhance the epidemiologic surveillance of this novel colistin resistance gene in CRE.

Analysis Of The *bla*_{NDM-1}-Harboring Plasmid pNDM-SCNJ07

Plasmid analysis revealed that the *bla*_{NDM-1} was carried by an FIB36:FIY4 replication type plasmid, designated pNDM-SCNJ07, which was 110,786 bp in length and had an average GC content of 54.84%. pNDM-SCNJ07 is almost identical (99% identity and 99.98% coverage) to the plasmid pRJF866 (GenBank accession no. KF732966) from *K. pneumoniae* from Shanghai, China, in 2015.³² Besides, similar FIB36:FIY4-type plasmids carrying *bla*_{NDM-1} have been widely found among *Enterobacteriaceae*, including pKOX_NDM1 (Accession no. JQ314407) from *Klebsiella michiganensis* from Taiwan,³³ pNDM_20ES (Accession no. MF042356) from *E. cloacae* and pNDM_4TM (Accession no. MF042352) from *Serratia marcescens* from Romania.³⁴ On these plasmids, the *bla*_{NDM-1} gene was embedded in the same genetic environment and its flanking miniature inverted-repeat transposable elements (MITes, positions

70834 to 71089 and 77035 to 77290 of pNDM-SCNJ07) were suggested to be responsible for the mobilization of *bla*_{NDM-1} onto the FIB36:FIY4 plasmids.^{33,35}

Genetic Characteristics Of The Plasmid-borne *mcr-9*

Complete sequence of the *mcr-9*-carrying plasmid, designated pMCR-SCNJ07, was circularized by PCR mapping using pT5282-mphA (Accession no. KY270852) as the template. pMCR-SCNJ07 was 285,587 bp in size with an average G+C content of 47.31%. The plasmid contained 327 predicted ORFs, and two replicons, IncHI2 (ST1) and IncR, and carried a number of additional resistance genes, including *bla*_{SHV-12}, *aadA16*, *aac(6')-IIa*, *mph(A)*, *sul1* and *tet(D)*. Sequence alignment showed that pMCR-SCNJ07 displayed 97% query coverage and 99.99% identity with the reference plasmid pT5282-mphA from an *E. cloacae* isolated in 2012 from a teaching hospital in Chongqing, China.³⁶ By BLAST,

pMCR-SCNJ07 was also similar to several previously sequenced IncHI2 plasmids (Figure 1), including pMRVIM0813 (86% query coverage and 99.97% identity, accession no. KP975077), pCTXM9_020038 (85% query coverage and 99.97% identity, accession no. CP031724), pC45-VIM4 (85% query coverage and 99.96% identity, accession no. LT991958) and pSE15-SA01028 (83% query coverage and 99.99% identity, accession no. CP026661), from a clinical *E. cloacae* isolate from USA in 2015, an *E. hormaechei* from China in 2016, an *E. cloacae* from France in 2018 and a *Salmonella enterica subsp. Enterica* isolate from Germany in 2015, respectively. It is noteworthy that all these plasmids were *mcr-9*-positive, highlighting an earlier presence of *mcr-9* among *Enterobacteriaceae* around the world than previously known and raising the likelihood of ongoing undetected transmission. Close surveillance is urgently needed to determine *mcr-9* prevalence and effective actions are required to control its further dissemination.

Figure 1 Circular comparison of *mcr-9*-carrying IncHI2 plasmids. The complete sequence of pMCR-SCNJ07 was used as the reference. The arrows indicate deduced ORFs and their orientations. The *mcr-9* gene is indicated in red. The circular maps were generated using BRIG²⁷ and plasmids were included in the following order (inner to outer circles): pMCR-SCNJ07 (this study, accession no. MK933279), pT5282-mphA (KY270852), pN1863-HI2 (MF344583), pMRVIM0813 (KP975077), pCTXM9_020038 (CP031724), pC45-VIM4 (LT991958), pSE15-SA01028 (CP026661), p505108-MDR (KY978628), pGW1 (CP028975), pRH-R27 (LN555650) and p707804-NDM (MH909331).

The genetic environment of *mcr-9* has not been described before. On pMCR-SCNJ07, *mcr-9* was located in a ~8kb region bracketed by two copies of intact IS903, an insertion sequence originating from *E. coli*. In this region, *wbuC* (encoding a cupin fold metalloprotein) was located downstream of *mcr-9*, followed by a two-component system encoding genes *qseC* and *qseB*, a truncated insertion sequence Δ IS1R, and a remnant of Δ silR (encoding a transcriptional regulatory protein) (Figure 2A). The two copies of intact IS903 could form a composite transposon or a circular intermediate, which has the potential to mobilize the *mcr-9* gene. However, no circular intermediate was detected despite repeated attempts via inverse PCR

in our work. Whether *mcr-9* could transfer through the formation of a circular intermediate requires further investigation. Of note, similar to the scenario in all known *mcr-1* cassettes, in which one or both ISAp11 sequences are consistently absent in the *mcr-1*-bearing region,³⁷ the downstream IS903 was absent on some *mcr-9*-carrying plasmids (Figure 2A).

A pairwise comparison of the *sil-cop* region of IncHI2 plasmids R478 and pRH-R27 enabled us to learn that *mcr-9* was likely to be transferred by the IS1R to IS903 region initially, and that insertion of this *mcr-9*-carrying region between *sil* and *cop* resulted in truncation of *cop* resistance gene clusters into Δ *copS-copE1* (Figure 2B).³⁸ BLASTn

Figure 2 Colinear analyses for genetic environment of *mcr-9*. **(A)** Comparison of the *mcr-9*-containing regions from different plasmid reservoirs and the chromosome of *E. hormaechei* strain S5. **(B)** Comparison of the *mcr-9*-neighboring genetic contexts on IncHI2-type plasmids. The corresponding region on non-*mcr-9*-carrying plasmid R478 (top) is shown for comparison. Arrows indicate the positions and directions of the genes and Δ indicates the truncated gene. *mcr-9* genes are indicated in dark blue and IS903 are highlighted in purple. Gray shades denote shared regions with a high degree of homology. Vertical green bars represent the direct repeats of IS10R. The accession numbers were: *E. hormaechei* strain S5 (Accession no. CP031571), pT5282-mphA (KY270852), pN1863-HI2 (MF344583), pGW1 (CP028975), R478 (NC_005211), pRH-R27 (LN555650), pN1863-HI2 (MF344583), and pMCR-SCNJ07 (this study, MK933279). The construction of sequence comparison was performed using BLAST (<http://blast.ncbi.nlm.nih.gov>) and Easyfig version 2.2.3.²⁶

analysis suggested that the *mcr-9*-carrying region of pMCR-SCNJ07 could be derived from pRH-R27 as a result of multiple genetic events (Figure 2B). No 9-bp target site duplication repeats, which are characteristic of the insertion of IS903, could be identified in the sequence adjacent to both IS903 elements. The detail of the evolutionary route of the *mcr-9*-carrying region, therefore, remains unclear.

To address the possible origin of *mcr-9*, a BLAST search against the NCBI's non-redundant protein sequence database (nr) was conducted by using the amino acid sequence of *mcr-9* as a probe. The result identified hits aligned to phosphoethanolamine transferases from multiple genera of *Enterobacteriaceae* (100% query coverage and 94–100% identity), followed by that from *Buttiauxella brennerae* (WP_064558897, 100% query coverage and 86.83% identity). It should be noted that of the flanking genes of *mcr-9* within the ~8kb region, none but the *wbuC* gene showed homologous to that from *Buttiauxella*, with 98% query coverage and 85.9% identity between their amino acid sequences. These findings suggested that the disseminated *mcr-9* gene in *Enterobacteriaceae* might have originated from *Buttiauxella* species and that the *wbuC* gene had been likely comobilized with *mcr-9* from its original genetic context. Additional studies are needed to fully characterize the origin of *mcr-9*.

Conclusion

In summary, we here report the discovery of a clinical *E. hormaechei* strain carrying both *bla*_{NDM-1} and *mcr-9*. Despite few reports characterizing *mcr-9*-harboring plasmids, this newly identified mobile colistin resistance gene is likely to have already been widely disseminated. Yet more worryingly, two copies of IS903 encompassing the *mcr-9* gene have the potential to mobilize *mcr-9* by hijacking more plasmids as the vehicle to disseminate this gene. Therefore, screening for the *mcr-9* gene should be urgently included in the surveillance of colistin-resistant *Enterobacteriaceae* from humans, animals, and the environment.

Acknowledgments

This work was supported by the Project of Education Department in Sichuan, China (18ZB0633), Natural Science Foundation of Southwest Medical University (No. 2017-ZRZD-022 and 2018-ZRZD-011), and National Undergraduate Innovation and Entrepreneurship Project (No.201816032021). The funders had no role in study

design, data collection and interpretation, or the decision to submit the work for publication.

Author Contributions

LZ designed the experiments. YY, GW, and CL performed the experiments. LX, JS and YY analyzed the data. YL and LZ analyzed the data and wrote the manuscript. FZ edited the original draft. All authors contributed to data analysis, drafting or revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- Logan LK, Weinstein RA. The epidemiology of carbapenem-resistant *Enterobacteriaceae*: the impact and evolution of a global menace. *J Infect Dis*. 2017;215(suppl_1):S28–S36. doi:10.1093/infdis/jiw282
- Wu WJ, Feng Y, Tang GM, Qiao F, McNally A, Zong ZY. NDM metallo- β -lactamases and their bacterial producers in health care settings. *Clin Microbiol Rev*. 2019;32(2):e00115–00118. doi:10.1128/CMR.00115-18
- Kaye KS, Pogue JM, Tran TB, Nation RL, Li J. Agents of last resort: polymyxin resistance. *Infect Dis Clin North Am*. 2016;30(2):391–414. doi:10.1016/j.idc.2016.02.005
- Liu YY, Wang Y, Walsh TR, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis*. 2016;16(2):161–168. doi:10.1016/S1473-3099(15)00424-7
- Sun J, Zhang HM, Liu YH, Feng YJ. Towards understanding MCR-like colistin resistance. *Trends Microbiol*. 2018;26(9):794–808. doi:10.1016/j.tim.2018.02.006
- Wang Q, Sun J, Li J, et al. Expanding landscapes of the diversified *mcr-I*-bearing plasmid reservoirs. *Microbiome*. 2017;5(1):70. doi:10.1186/s40168-017-0288-0
- Delgado-Blas JF, Ovejero CM, Abadia-Patino L, Gonzalez-Zorn B. Coexistence of *mcr-I* and *bla*_{NDM-1} in *Escherichia coli* from Venezuela. *Antimicrob Agents Chemother*. 2016;60(10):6356–6358. doi:10.1128/AAC.01319-16
- Liu BT, Song FJ, Zou M, Hao ZH, Shan H. Emergence of colistin resistance gene *mcr-I* in *Cronobacter sakazakii* producing NDM-9 and in *Escherichia coli* from the same animal. *Antimicrob Agents Chemother*. 2017;61(2):e01444–16.
- Feng S, Shen C, Chen H, et al. Co-production of MCR-1 and NDM-5 in *Escherichia coli* isolated from a colonization case of inpatient. *Infect Drug Resist*. 2018;11:1157–1161. doi:10.2147/IDR.S171164
- Li X, Mu X, Zhang P, et al. Detection and characterization of a clinical *Escherichia coli* ST3204 strain coproducing NDM-16 and MCR-1. *Infect Drug Resist*. 2018;11:1189–1195. doi:10.2147/IDR.S175041
- Xu L, Wang P, Cheng J, Qin S, Xie W. Characterization of a novel *bla*_{NDM-5}-harboring IncFII plasmid and an *mcr-I*-bearing IncI2 plasmid in a single *Escherichia coli* ST167 clinical isolate. *Infect Drug Resist*. 2019;12:511–519. doi:10.2147/IDR.S192998
- Lin YC, Kuroda M, Suzuki S, Mu JJ. Emergence of an *Escherichia coli* strain co-harboring *mcr-I* and *bla*_{NDM-9} from a urinary tract infection in Taiwan. *J Glob Antimicrob Resist*. 2019;16:286–290. doi:10.1016/j.jgar.2018.10.003

13. Mediavilla JR, Patrawalla A, Chen L, et al. Colistin- and carbapenem-resistant *Escherichia coli* harboring *mcr-1* and *bla_{NDM-5}*, causing a complicated urinary tract infection in a patient from the United States. *MBio*. 2016;7(4). doi:10.1128/mBio.01191-16.
14. Carroll LM, Gaballa A, Guldinmann C, Sullivan G, Henderson LO, Wiedmann M. Identification of novel mobilized colistin resistance gene *mcr-9* in a multidrug-resistant, colistin-susceptible *Salmonella enterica* serotype Typhimurium isolate using a combination of high-throughput, in silico screening and functional analysis. *mBio*. 2019;10(3): e00853-19. doi:10.1128/mBio.00853-19
15. Lane DJ. 16S/23S rRNA sequencing. *Nucleic Acid Tech Bacterial Sys*. 1991;115–175.
16. Szekely E, Damjanova I, Janvari L, et al. First description of *bla_(NDM-1)*, *bla_(OXA-48)*, *bla_(OXA-181)* producing *Enterobacteriaceae* strains in Romania. *Int J Med Microbiol*. 2013;303(8):697–700. doi:10.1016/j.ijmm.2013.10.001
17. Dallenne C, Da Costa A, Decre D, Favier C, Arlet G. Development of a set of multiplex PCR assays for the detection of genes encoding important beta-lactamases in Enterobacteriaceae. *J Antimicrob Chemother*. 2010;65(3):490–495. doi:10.1093/jac/dkp498
18. Ellington MJ, Kistler J, Livermore DM, Woodford N. Multiplex PCR for rapid detection of genes encoding acquired metallo-beta-lactamases. *J Antimicrob Chemother*. 2007;59(2):321–322. doi:10.1093/jac/dkl481
19. Poirel L, Le Thomas I, Naas T, Karim A, Nordmann P. Biochemical sequence analyses of GES-1, a novel class A extended-spectrum beta-lactamase, and the class 1 integron In52 from *Klebsiella pneumoniae*. *Antimicrob Agents Chemother*. 2000;44(3):622–632. doi:10.1128/AAC.44.3.622-632.2000
20. Wang X, Wang Y, Zhou Y, et al. Emergence of a novel mobile colistin resistance gene, *mcr-8*, in NDM-producing *Klebsiella pneumoniae*. *Emerg Microbes Infect*. 2018;7(1):122. doi:10.1038/s41426-018-0124-z
21. Yang YQ, Li YX, Lei CW, Zhang AY, Wang HN. Novel plasmid-mediated colistin resistance gene *mcr-7.1* in *Klebsiella pneumoniae*. *J Antimicrob Chemother*. 2018;73:1791–1795. doi:10.1093/jac/dky111
22. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 2014;30(15):2114–2120. doi:10.1093/bioinformatics/btu170
23. Li R, Li Y, Kristiansen K, Wang J. SOAP: short oligonucleotide alignment program. *Bioinformatics*. 2008;24(5):713–714. doi:10.1093/bioinformatics/btn025
24. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics*. 2014;30(14):2068–2069. doi:10.1093/bioinformatics/btu153
25. Aziz RK, Bartels D, Best AA, et al. The RAST server: rapid annotations using subsystems technology. *BMC Genomics*. 2008;9:75. doi:10.1186/1471-2164-9-75
26. Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer. *Bioinformatics*. 2011;27(7):1009–1010. doi:10.1093/bioinformatics/btr039
27. Alikhan NF, Petty NK, Ben Zakour NL, Beatson SA. BLAST ring image generator (BRIG): simple prokaryote genome comparisons. *BMC Genomics*. 2011;12:402. doi:10.1186/1471-2164-12-402
28. CLSI. *Performance Standards for Antimicrobial Susceptibility Testing*. 29th. CLSI supplement M100. Wayne, PA: Clinical and Laboratory Standards Institute; 2019
29. Richter M, Rossello-Mora R. Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A*. 2009;106(45):19126–19131. doi:10.1073/pnas.0906412106
30. Mezzatesta ML, Gona F, Stefani S. *Enterobacter cloacae* complex: clinical impact and emerging antibiotic resistance. *Future Microbiol*. 2012;7(7):887–902. doi:10.2217/fmb.12.61
31. Zheng F, Sun J, Cheng C, Rui Y. Molecular characteristics of carbapenem-resistant gram-negative bacteria in southern China. *Microb Drug Resist*. 2015;21(2):178–185. doi:10.1089/mdr.2014.0085
32. Qu H, Wang X, Ni Y, et al. NDM-1-producing *Enterobacteriaceae* in a teaching hospital in Shanghai, China: incX3-type plasmids may contribute to the dissemination of *bla_{NDM-1}*. *Int J Infect Dis*. 2015;34:8–13. doi:10.1016/j.ijid.2015.02.020
33. Huang TW, Wang JT, Lauderdale TL, et al. Complete sequences of two plasmids in a *bla_{NDM-1}*-positive *Klebsiella oxytoca* isolate from Taiwan. *Antimicrob Agents Chemother*. 2013;57(8):4072–4076. doi:10.1128/AAC.02266-12
34. Phan HTT, Stoesser N, Maciucă IE, et al. Illumina short-read and MinION long-read WGS to characterize the molecular epidemiology of an NDM-1 *Serratia marcescens* outbreak in Romania. *J Antimicrob Chemother*. 2018;73(3):672–679. doi:10.1093/jac/dkx456
35. Wailan AM, Sidjabat HE, Yam WK, et al. Mechanisms involved in acquisition of *bla_{NDM}* genes by IncA/C2 and IncFIIY plasmids. *Antimicrob Agents Chemother*. 2016;60(7):4082–4088. doi:10.1128/AAC.00368-16
36. Liang Q, Yin Z, Zhao Y, et al. Sequencing and comparative genomics analysis of the IncHI2 plasmids pT5282-mphA and p112298-catA and the IncHI5 plasmid pYNKP001-dfrA. *Int J Antimicrob Agents*. 2017;49(6):709–718. doi:10.1016/j.ijantimicag.2017.01.021
37. Li R, Xie M, Zhang J, et al. Genetic characterization of *mcr-1*-bearing plasmids to depict molecular mechanisms underlying dissemination of the colistin resistance determinant. *J Antimicrob Chemother*. 2017;72(2):393–401. doi:10.1093/jac/dkw411
38. Dolejska M, Villa L, Poirel L, Nordmann P, Carattoli A. Complete sequencing of an IncHI1 plasmid encoding the carbapenemase NDM-1, the ArmA 16S RNA methylase and a resistance-nodulation-cell division/multidrug efflux pump. *J Antimicrob Chemother*. 2013;68(1):34–39. doi:10.1093/jac/dks357

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

Dovepress