Postpartum Contraceptive Use In Parakou (A City In Northern Benin) In 2018: A Community Based Cross-Sectional Study

This article was published in the following Dove Press journal:
Open Access Journal of Contraception

Fanny Hounkponou¹
Yolaine Glèlè Ahanhanzo²
Chabi Olaniran Alphonse
Biaou²
Candide Russel Dos-Santos³
Aurelle Ahouingnan¹
Awadé Achille Obossou¹
Rachidi Sidi Imorou¹
Mahublo Vodouhe¹
Hubert Laourou³
Kabibou Salifou¹
René Xavier Perrin⁴

¹Department of Mother and Child, Faculty of Medicine, University of Parakou, Parakou, Benin; ²Epidemiology and Biostatistics Department, Public Health Regional Institute, Ouidah, Benin; ³Public Health Department, Claude Bernard Lyon I University, Lyon, France; ⁴Department of Mother and Child, Faculty of Health Science, University of Abomey Calavi, Cotonou, Benin

Background: Over 30% of maternal deaths and 10% of infant deaths are avoidable if couples space out births at intervals of 2 years or greater. This study aimed to identify factors associated with contraceptive use in the postpartum period.

Methods: This was a cross-sectional study among postpartum women living in Parakou. Participants were selected using a two-stage sampling method. Postpartum contraceptive use was defined as the use of at least one contraceptive method at the time of the survey. Independent variables included socio demographic and reproductive history. For the analysis we performed a multiple logistic regression and we calculated odds ratios with 95% confidence intervals.

Results: The study included 453 postpartum women. Among the participants, 59 (13%) were using a contraceptive method. Injectable contraceptives were the most common method used (25.4%). Factors associated with contraceptive use in the postpartum period included history of contraceptive use (AOR=9.4; $p<0.001$), resumption of sexual intercourse (AOR=5.7; $p<0.001$), discussion with partner about contraceptive method (AOR=5.7; $p=0.005$), need of partner's approval before adopting a contraceptive method (AOR=0.4; $p=0.014$) and counseling during pregnancy, in the delivery room, and during postnatal visits (AOR=2.7; $p=0.019$).

Conclusion: The prevalence of contraceptive use during the postpartum period remains low. Interventions should be designed with a focus on the associated factors in order to increase postpartum contraceptive use in Parakou.

Keywords: family planning, postpartum period, risk factors, Benin

Introduction

In 2015, approximately 303,000 women died during pregnancy, delivery, or postpartum, and the majority of these deaths occurred in low income countries.¹ A key strategy in reducing maternal mortality is family planning (FP) during the postpartum period, which can prevent unintended pregnancies and short interpregnancy intervals during the twelve month period after delivery.² We consider that spacing births at least every two years could lead to a more than 30% reduction in maternal deaths and a 10% reduction in infant deaths.³ An expansion of contraceptive use should help achieve this goal.

In developing countries, two-thirds of lactating women do not use any contraceptive method, while one-third will become pregnant again within fifteen months after their last delivery.⁴ About one fifth of pregnancies in Benin are unplanned. Among women in the Borgou Department, the prevalence of current modern family planning methods is 11%. An estimated 27.4% of women have an unmet need for family planning.⁵

Correspondence: Chabi Olaniran Alphonse Biaou
Epidemiology and Biostatistics Department, Public Health Regional Institute, Ouidah, Benin
Tel +229 21 97730142
Email alphonse.biaou@yahoo.fr

Although many studies have assessed the uptake of family planning and associated factors generally, few have specifically focused on contraceptive utilization in the postpartum period. And yet, women in the post-partum period are among those most in need of FP services. Indeed, the consequences of inadequate use of contraceptive methods during the post-partum period are an increase in the number of pregnancies and a reduction in the inter-reproductive interval. Such consequences put the mother and child at increased risk of maternal morbidity, low birth weight, intrauterine growth retardation, prematurity, and malnutrition.⁶⁻⁸ In addition, there are a range of psychological and social consequences that can affect the mother and the breastfeeding child, as well as the unborn child. In order to guide strategies aimed at increasing FP prevalence, this study aimed to identify the factors associated with contraceptive use among women during the postpartum period.

Methods

Study Design, Study Population And Sampling Procedure

The study was conducted in Parakou, a city in northern Benin (West Africa), specifically in the Borgou Department. This was a community-based cross-sectional study of women in the postpartum period. Inclusion criteria included all women living in Parakou who had given birth in the twelve months prior to the study. The data were collected from May 1–31, 2018. Women who had a tubal ligation or a hysterectomy, as well as those who did not consent to participate, were excluded from the study. Parakou consists of 58 villages and districts.⁹ Sampling was conducted in two stages. For the first stage, half (29) were selected by simple random sampling. In the second stage, we selected households with women who were eligible for the survey. In the center of each selected district, we used the direction indicated by the throwing of a ballpoint pen, and then proceeded to number all the households in the chosen direction and select one household to survey in that area by simple random draw. It was in this household that the selection of women began and the following was done according to the nearest door technique, until we obtained the number of eligible women needed per district. A maximum of two women per household were randomly selected. The sample size was calculated based on modern contraceptive prevalence in Borgou department (11%) in 2012,⁵ following the

Schwartz formula with an accuracy of 3%. The minimum sample size needed for the study was $n = 418$, however we increased the number of participants sampled by 10% to take into account non-response. In total, 460 women were surveyed. The number of eligible women surveyed by area was obtained by proportional allocation to the number of expected births by district/village of Parakou in 2017.

Techniques, Data Collection And Variables

Data were collected via a pre-tested questionnaire. The dependent variable was defined as the use of at least one contraceptive method during the postpartum period. This was a binary qualitative variable measured from the woman's statement. This variable was coded « yes » if the woman reported using a contraceptive method at the time of the survey and « No » if otherwise. In order to identify factors associated with PF in the postpartum period, we included sixteen (16) independent variables at the level of the respondents. These included i) socio demographic features such as age, level of education, occupation, religion, marital status, husband's education level, socio-economic level, ii) factors related to the woman's reproductive history such as gravidity, parity, menses resumption, knowledge of contraceptives methods, contraceptive use history, resumption of sexual activities post-pregnancy, last intended pregnancy, counseling during pregnancy/in the delivery room/in postpartum period, and iii) discussion around contraceptive use with husband/partner, need of husband's permission before adopting a contraceptive method.

The socio-economic level of the respondent's household was determined using the Demographic and Health Surveys' poverty index, which is a composite index of socio-economic status that assigns a weight or scores. Scores are obtained through a principal component analysis, as described by Filmer and al,¹⁰ which takes into account the number of properties the respondent owns and the household's characteristics. Each respondent is ranked according to the household asset score and is assigned to poverty tertiles as follows: poorest (Poor), average (Average), and wealthiest (Rich).

Knowledge of contraceptive methods was assessed in a dichotomized manner ("yes" vs "no") based on the answers given by the respondent; those categorized as "Yes" "could cite at least one role and were able to give at least

two examples of contraceptive methods” and those categorized as “No” did not meet such criteria.

Data Processing And Analysis

The data were processed using Epi Data 3.1 and Stata 11. We gathered descriptive statistics and the results are presented as a mean (standard deviation) or median (Interquartile range) for the quantitative variables. Categorical variables were expressed as a percentage. To identify the factors associated with contraceptive use in the postpartum period, logistic regression was used to estimate crude and adjusted odds ratios (OR) and their corresponding 95% confidence intervals (95% CI). Using a significance threshold of 20%, a univariate analysis was conducted to pre-select variables that were included in the multivariate model. We used a step down (descending) modeling strategy for the gradual elimination of variables which are less significantly related to postpartum contraceptive use. Variables were retained in the final model if they had a statistical significance level of less than 5%. We also searched for possible interactions and confounding factors. The fitness of the final model was verified using the Hosmer Lemeshow test.

Ethical Statement

This study was conducted in accordance with the Declaration of Helsinki. Prior to the data collection, ethical authorization (N° 0123/CLERB-UP/P/SP/R/SA) was obtained from the Faculty of Medicine (University of Parakou) ethical clearance committee, as well as from Parakou's city health authorities. When recruiting participants, researchers informed subjects of the purpose of the study, the fact that it was anonymous, and of their right to agree or refuse to participate. Participants were also told that they could withdraw from the study at any time if they wished to do so. The interviews took place in private to ensure confidentiality. Prior to data collection, the research team obtained written consent to participate from all respondents.

Results

Characteristics Of Participants

A total of 460 women in the post-partum period were recruited. 453 of participants responded fully to the questionnaire, making the response rate 98.47%. The average age was 27.1 years (SD=6). Participants ranged in age from 15 to 45 years and 36.2 % were under 25. The

majority were married (74.6%), and most were merchants (36.9%), Muslim (68.4%) and uneducated (39.3%).

The majority of respondents (85%) were familiar with birth control methods. One of every 2 women surveyed (50.3%), had discussions around contraception with their partner. About 3 out of 4 women (73.3%) reported needing their partner's permission before using any contraceptive method. 67.1% of women who did not discuss contraception with their partner said they needed their husband's permission before adopting a contraceptive method. Additional respondent characteristics are summarized in [Table 1](#).

Prevalence Of Postpartum Contraceptive Use

In total, 59 women were already using a contraceptive method at the time of the survey, making the prevalence rate at 13% (CI 95%: [9.9%–16.1%]). Among women who were familiar with contraceptive methods (n=385), 15.3% were using contraception at the time of the survey. Injectable contraceptives (25.4%) and natural methods (23.7%) were the most frequently used ([Figure 1](#)).

Factors Related To Postpartum Contraceptive Use

In the univariate analysis, factors associated with postpartum contraceptive use included: woman's and husband's educational level, religion ($p=0.028$), socioeconomic status, menses resumption, prior use of contraceptives, resumption of sexual activities, discussion of contraceptive use with husband/partner, the need for husband's permission prior to starting a contraceptive method, and counseling during the pregnancy, in the delivery room, and during postpartum visits ([Table 1](#)).

Results from the multiple logistic regression analysis ([Table 2](#)) demonstrated that the history of contraceptive use before the last pregnancy increases the probability of contraceptive use in the postpartum period (AOR=9.4; $p<0.001$). Women who had already resumed sexual activities were more likely to use contraceptives than those who had not (AOR=5.7; $p<0.001$). In addition, women who discussed FP with their partner were more likely to use contraceptive methods in the post-partum period than others (AOR=3.5; $p=0.005$). Women who reported that they received counseling during pregnancy, in the delivery room, and during postpartum visits were also more likely to be using contraceptive methods than others (AOR=2.7; $p=0.019$). On the other hand, those who needed their husband's permission were

Table 1 Univariate Logistic Regression Of Factors Associated With Postpartum Contraceptive Use In Parakou, Benin, 2018

	Total Population		Postpartum Contraceptive Use		
	Number	%	%	OR [95% CI]	p-value
Age (Years)					0.230
< 25	164	36.2	9.1	1	
25–29	132	29.1	16.7	2.0 [1.0–4.0]	
30–34	91	20.1	15.4	1.8 [0.8–3.9]	
≥ 35	66	14.6	12.1	1.4 [0.6–3.4]	
Educational level					< 0.001
No education	178	39.3	7.3	1	
Primary	106	23.4	10.4	1.5 [0.6–3.4]	
Secondary	147	32.4	17.7	2.7 [1.3–5.5]	
University	22	4.9	40.9	8.8 [3.2–24.4]	
Occupation					< 0.001
Student	21	4.6	19.1	2.6 [0.7–9.7]	
Employed	39	8.6	35.9	6.1 [2.2–16.8]	
Merchants	167	36.9	13.2	1.6 [0.7–4.0]	
Housewife	143	31.6	8.4	1.0 [0.4–2.6]	
Artisan	83	18.3	8.4	1	
Religion					0.028
Christian	143	31.6	18.2	1.86 [1.06–3.25]	
Other ^a	310	68.4	10.7	1	
Marital status					0.682
Single	19	4.2	10.5	1	
Concubinage	96	21.2	15.6	1.6 [0.3–7.5]	
Married	338	74.6	12.4	1.2 [0.3–5.4]	
Husband education level					< 0.001
No education	136	30.0	8.1	1	
Primary	101	22.3	4.9	0.6 [0.2–1.8]	
Secondary	172	38.0	16.9	2.3 [1.1–4.8]	
University	44	9.7	31.8	5.3 [2.2–12.8]	
Socio-economic status ^b					0.010
Poor	151	33.3	6.6	1	
Average	150	33.1	16.0	2.7 [1.2–5.8]	
Rich	152	33.6	16.5	2.8 [1.3–6.0]	
Gravidity					0.757
1	106	25.6	11.3	1	
2 – 3	96	21.2	14.9	1.4 [0.6–3.2]	
≥ 4	241	53.2	13.1	1.2 [0.6–2.4]	
Parity					0.985
1	116	25.6	12.9	1	
2 – 3	96	21.2	13.5	1.1 [0.5–2.3]	
≥ 4	241	53.2	12.9	1.0 [0.5–1.9]	
Menses resumption					< 0.001
No	282	62.3	8.2	1	
Yes	171	37.7	21.1	3.0 [1.7–5.3]	

(Continued)

Table 1 (Continued).

	Total Population		Postpartum Contraceptive Use		
	Number	%	%	OR [95% CI]	p-value
History of contraceptive use					
No	341	75.3	4.4	1	< 0.001
Yes	112	24.7	39.3	14.1 [7.4–26.7]	
Sexual intercourse resumption					
No	294	64.9	4.8	1	< 0.001
Yes	159	35.1	28.3	7.9 [4.2–14.9]	
Last pregnancy was planned ?					
No	33	7.3	9.1	1	0.489
Yes	420	92.7	13.3	1.5 [0.5–5.2]	
Discussion about contraceptive use with partner					
No	225	49.7	4.9	1	< 0.001
Yes	228	50.3	21.1	5.2 [2.6–10.3]	
Partner's permission needed ^c					
No	121	26.7	20.7	1	0.004
Yes	332	73.3	10.2	0.4 [0.2–0.8]	
Counseling Received ^d					
No	189	41.7	5.3	1	< 0.001
Yes	264	58.3	18.6	4.1 [2.0–8.3]	

Notes: ^aMuslim (n=301); (Animist n=9); ^bSocio-economic status was categorized using tertiles, with the lowest being "Poor" and the highest being "Rich"; ^cPartner's permission needed before adopting a contraceptive method; ^dCounseling during pregnancy/in the delivery room/in postpartum period

Abbreviations: OR, odds ratio; %, Percentage.

Figure 1 Contraceptives methods used at times of the survey in Parakou, Benin, 2018.

less likely to use a contraceptive method during the postpartum period (AOR=0.4; p=0.014).

Discussion

Prevalence Of Postpartum Contraceptive Use

The prevalence of postpartum contraceptive use was 13%, revealing a very low uptake of contraceptive methods for

postpartum women in Parakou. This finding aligns closely to results reported in the 2012 Demographic Health Survey (DHS),⁵ but it remains significantly lower than most of the results found in the literature. For example, Jalang'o and al¹¹ reported a prevalence of 86.3% in Kenya in 2017. Bwazi and al¹² found a prevalence of 74.6% in 2014 in Ntchisi (Malawi). Such differences could be explained by the fact that the target populations of these studies were women in the 18th–24th month¹¹ and 6th–12th

Table 2 Multivariate Analysis Of Factors Associated With Postpartum Contraceptive Use In Parakou, Benin, 2018

	OR [95% CI]	P value
Contraceptive method use history		
No	1	
Yes	9.4 [4.6–18.8]	< 0.001
Resumption of sexual relationships		
No	1	
Yes	5.7 [2.8–11.7]	< 0.001
Discussion about contraceptive use with husband/partner		
No	1	
Yes	3.5 [1.5–8.4]	0.005
Partner's permission needed*		
No	1	
Yes	0.4 [0.2–0.8]	0.014
Counseling received**		
No	1	
Yes	2.7 [1.2–6.3]	0.019

Notes: *Partner's permission needed before adopting a contraceptive method; **Counseling during pregnancy/in the delivery room/in postpartum period; Hosmer Lemeshow test: $p=0.374$.

Abbreviation: OR, odds ratio.

month,¹² respectively, of the postpartum period, whereas our study looked at women in the first 12 months after delivery. In addition, these studies were conducted in hospital settings, while the current study was conducted with the general population. Abera and al. reported that 48.4% of the general population of Gondar (in the north-west of Ethiopia) used contraceptives in 2013.¹³ This lower prevalence as compared to our study could be due to the fact that Gondar is a more urbanized area where the population has increased access to family planning, as opposed to Parakou. In 2015, in Uganda, Sileo and al. reported the prevalence of postpartum contraceptive use at 25.0%.¹⁴ This rate is high as compared to our results and was obtained among women seen for antenatal care (ANC), postnatal visits, and infant immunizations. Such results suggest the potentially positive impact of integrated postnatal care on increasing contraceptive uptake.

Factors Associated With Postpartum Contraceptive Use

Our study revealed that women who had used contraceptives prior to their most recent pregnancy were more likely to use contraceptives during the postpartum period. This is most likely due to the fact that such women were already familiar

with and had a better understanding of contraception. This same finding was reported by Achyut and al in India in 2012,¹⁵ and by Sileo and al.¹⁴ Women who had resumed sexual activities post-partum were more likely to use contraceptives than others. This could indicate a perception among women that contraceptives are not useful until they resume their sexual relationships. Findings from Abraha and al¹⁶ and Bwazi and al¹² are also consistent with our results. Women who had discussions around family planning with their partner were the most likely to use contraceptives in the postpartum period. Sileo and al¹⁴ revealed that partner communication about the topic of FP was significantly associated with postpartum period family planning. Bwazi and al found a relationship between the extent to which husbands assist women in accessing family planning services during the post-partum period and the use of contraceptive methods.¹² These findings could be explained by the value placed on the husband's input into major decisions in the family, including contraceptive use. Furthermore, in this study, women who did not need their husband's permission were more likely to use contraceptives in the postpartum period. Our study found that socio-cultural factors play a key role in decisions around adopting a contraception method. Paradoxically, we found that a majority of women who did not discuss contraceptives with their partner reported that they still needed permission prior to using contraceptives. These women may fear their husbands' reaction, due to inaccurate perceptions around contraceptive use. In a study by Eliason and al,¹⁷ 82% of women reported needing their husband's permission before beginning a contraceptive method. These findings highlight the role that husbands could play in encouraging the use of health services in general and family planning services in particular. It is therefore essential to involve men in interventions to promote contraceptive use. This study found that women who reported having received counselling during their pregnancy/in the delivery room/at postpartum visits were more likely to use contraceptives method in the postpartum period. This is consistent with other findings in the literature.^{13,15,18–22} At antenatal and postnatal visits, women can receive counselling around family planning and have the opportunity to start using contraception. Women who receive antenatal and postnatal care adequately are more likely to be informed about contraceptive use and its benefits.

Strengths And Limitations

The purpose of this study was to identify the factors related to postpartum contraceptive use in Parakou in 2018. From a

methodological standpoint, the sampling technique allowed for a representative sample of the population of postpartum women in Parakou. A potential limitation of the study is selection bias, due to the probability of non-response or refusal to participate in the survey. To prevent such refusals and to ensure a minimum sample size, a non-response rate of 10% was included when calculating the sample size. It should be noted, however, that no causal conclusions can be drawn from the results of this study because of the instantaneous nature of cross-sectional studies. Because of the various ethical provisions put in place to ensure free and informed consent as well as the confidentiality of the data, the study remains in compliance with the rules of study in the general population. We therefore believe that the results are valid, reliable and can be generalized in Parakou.

Conclusion

Birth control is a common issue after childbirth. Based on the study results, it appears that the use of contraception in the postpartum period in the city of Parakou is low. It is therefore necessary to strengthen existing actions and improve strategies focusing on the factors identified that are associated with postpartum contraceptive use in order to increase the prevalence in Parakou.

Acknowledgments

The authors wish to thank Miss Raïssa Chaffa and Megan O'Brien for their contributions to the translation of this manuscript.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Alkema L, Chou D, Hogan D, et al. Global, regional, and national levels and trends in maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a systematic analysis by the UN Maternal Mortality Estimation Inter-Agency Group. *Lancet*. 2016;387(10010):462–474. doi:10.1016/S0140-6736(15)00838-7
2. United States Agency International Development. Déclaration pour l'action collective pour la planification familiale du postpartum. Available from: <https://www.mchip.net/sites/default/files/PPFP%20Statement%20for%20Action%20French.pdf>. Accessed February 12, 2018.
3. Cleland J, Bernstein S, Ezeh A, Faundes A, Glasier A, Innis J. Family planning: the unfinished agenda. *Lancet*. 2006;368(9549):1810–1827. doi:10.1016/S0140-6736(06)69480-4
4. Ross JA, Winfrey WL. Contraceptive use, intention to use and unmet need during the extended postpartum period. *Int Fam Plan Perspect*. 2001;27(1):20–27. doi:10.2307/2673801

5. Institut National de la Statistique et de l'Analyse Economique (Bénin). *Enquête Démographique et de Santé et à Indicateurs Multiples du Bénin EDS-MICS-IV 2011–2012*. Cotonou: INSAE; 2012.
6. Organisation Mondiale de la Santé. *Relever le défi de la santé de la femme en Afrique: Rapport de la Commission de la Santé de la femme dans la Région africaine*. Brazzaville: OMS; 2012.
7. Cleland J, Conde-Agudelo A, Peterson H, Ross J, Tsui A. Contraception and health. *Lancet*. 2012;380(9837):149–156. doi:10.1016/S0140-6736(12)60609-6
8. Conde-Agudelo A, Rosas-Bermúdez A, Kafury-Goeta AC. Effects of birth spacing on maternal health: a systematic review. *Am J Obstet Gynecol*. 2007;196(4):297–308. doi:10.1016/j.ajog.2006.05.055
9. Institut National de la Statistique et de l'Analyse Economique. *Cahier des villages et quartiers de ville du département du Borgou (RGPH-4, 2013)*. Cotonou: INSAE; 2016.
10. Filmer D, Pritchett LH. Estimating wealth effects without expenditure data—or tears: an application to educational enrollments in states of India. *Demography*. 2001;38(1):115–132. doi:10.1353/dem.2001.0003
11. Jalang'o R, Thuita F, Barasa SO, Njoroge P. Determinants of contraceptive use among postpartum women in a county hospital in rural Kenya. *BMC Public Health*. 2017;17(604):1–8. doi:10.1186/s12889-017-4510-6
12. Bwazi C, Maluwa A, Chimwaza A, Pindani M. Utilization of postpartum family planning services between six and twelve months of delivery at Ntchisi District Hospital, Malawi. *Health (N Y)*. 2014;6(14):1724–1737. doi:10.4236/health.2014.614205
13. Abera Y, Mengesha ZB, Tessema GA. Postpartum contraceptive use in Gondar town, Northwest Ethiopia: a community based cross-sectional study. *BMC Womens Health*. 2015;15(19):1–8. doi:10.1186/s12905-015-0178-1
14. Sileo KM, Wanyenze RK, Lule H, Kiene SM. Determinants of family planning service uptake and use of contraceptives among postpartum women in rural Uganda. *Int J Public Health*. 2015;60(8):987–997. doi:10.1007/s00038-015-0683-x
15. Achyut P, Mishra A, Montana L, Sengupta R, Calhoun LM, Nanda P. Integration of family planning with maternal health services: an opportunity to increase postpartum modern contraceptive use in urban Uttar Pradesh, India. *J Fam Plann Reprod Health Care*. 2016;42(2):107–115. doi:10.1136/jfprhc-2015-101271
16. Abraha TH, Teferra AS, Gelagay AA. Postpartum modern contraceptive use in northern Ethiopia: prevalence and associated factors. *Epidemiol Health*. 2017;39(12):1–9. doi:10.4178/epih.e2017012
17. Eliason S, Baiden F, Quansah-Asare G, et al. Factors influencing the intention of women in rural Ghana to adopt postpartum family planning. *Reprod Health*. 2013;10(34):1–8. doi:10.1186/1742-4755-10-1
18. Tawfik Y, Rahimzai M, Ahmadzai M, Clark PA, Kamgange E. Integrating family planning into postpartum care through modern quality improvement: experience from Afghanistan. *Glob Health Sci Pract*. 2014;2(2):226–233. doi:10.9745/GHSP-D-13-00166
19. Speizer IS, Fotso JC, Okigbo C, Faye CM, Seck C. Influence of integrated services on postpartum family planning use: a cross-sectional survey from urban Senegal. *BMC Public Health*. 2013;13(752):1–9. doi:10.1186/1471-2458-13-1
20. Zapata LB, Murtaza S, Whiteman MK, et al. Contraceptive counseling and postpartum contraceptive use. *Am J Obstet Gynecol*. 2015;212(171):e1–e8. doi:10.1016/j.ajog.2014.07.059
21. Do M, Hotchkiss D. Relationships between antenatal and postnatal care and post-partum modern contraceptive use: evidence from population surveys in Kenya and Zambia. *BMC Health Serv Res*. 2013;13(6):1–14. doi:10.1186/1472-6963-13-1
22. Wilson EK, Fowler CI, Koo HP. Postpartum contraceptive use among adolescent mother in seven state. *J Adolesc Health*. 2013;52(3):278–283. doi:10.1016/j.jadohealth.2012.05.004

Open Access Journal of Contraception**Dovepress****Publish your work in this journal**

Open Access Journal of Contraception is an international, peer-reviewed, open access, online journal, publishing original research, reports, reviews and commentaries on all areas of contraception. In addition to clinical research, demographics and health-related aspects, the journal welcomes new findings in animal and preclinical

studies relating to understanding the biological mechanisms and practical development of new contraceptive agents. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/open-access-journal-of-contraception-journal>