Potential role of the antimicrobial peptide *Tachyplesin III* against multidrug-resistant *P. aeruginosa* and *A. baumannii* coinfection in an animal model

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Jialong Qi¹
Ruiyu Gao¹
Cunbao Liu¹
Bin Shan²
Fulan Gao¹
Jinrong He¹
Mingcui Yuan¹
Hanghang Xie¹
Shumei Jin³
Yanbing Ma¹

¹Department of Molecular Immunology, Institute of Medical Biology, Chinese Academy of Medical Sciences and Peking Union Medical College, Kunming, People's Republic of China; ²Department of Clinical Lab, The First Affiliated Hospital of Kunming Medical University, Kunming, People's Republic of China; ³Yunnan Institute of Materia Medica, Kunming, People's Republic of China

Background: *Tachyplesin III*, an antimicrobial peptide (AMP), provides protection against multidrug-resistant (MDR) bacterial infections and shows cytotoxicity to mammalian cells. Mixed bacterial infections, of which *P. aeruginosa* plus *A. baumannii* is the most common and dangerous combination, are critical contributors to the morbidity and mortality of long-term in-hospital respiratory medicine patients. Therefore, the development of effective therapeutic approaches to mixed bacterial infections is urgently needed.

Methods and results: In this study, we demonstrated that compared with individual infections, mixed infections with MDR bacteria *P. aeruginosa* and *A. baumannii* cause more serious diseases, with increased pro-inflammatory cytokines (IL-1 β , IL-6, TNF- α) and chemokines (MCP-1/MIP-2) and reduced mouse survival. In vitro treatment with *Tachyplesin III* enhanced phagocytosis in a mouse alveolar macrophage cell line (MH-S). Strikingly, in vivo, *Tachyplesin III* demonstrated a potential role against mixed-MDR bacterial coinfection. The bacterial burden in bronchoalveolar lavage fluid (BALF) was significantly reduced in the *Tachyplesin III*-treated group. In addition, a systemic reduction in pro-inflammatory cytokines and decreased lung injury occurred with *Tachyplesin III* therapy.

Conclusion: Therefore, our study demonstrated that *Tachyplesin III* represents a potential therapeutic treatment against mixed-MDR bacterial infection in vivo, which sheds light on the development of therapeutic strategies against mixed-MDR bacterial infections.

Keywords: *Tachyplesin III*, antimicrobial peptides, multidrug-resistant bacterial, coinfection, phagocytosis

Introduction

The use of antibiotics, first discovered approximately 80 years ago,¹ is a double-edged sword.² Antibiotics play an extremely important role in the human against various bacterial and viral infections by killing pathogens directly, reducing tissue damage and inhibiting the production of pro-inflammatory cytokines. However, long-term antibiotic misuse induces the emergence of multidrug-resistant (MDR) bacteria.³ Decreased antibiotic uptake, like reducing the entry of antibiotics into the bacteria and activating efflux system, and inactivation antibiotics targets are the most common resistance mechanism for bacterial. Frequently, ventilator-associated pneumonia (VAP) and intensive care unit (ICU) patients suffer lung bacterial pneumonia and are infected with multiple bacterial pathogens.⁴ Among these, Gram-negative *P. aeruginosa* and *A. baumannii* are the worst MDR bacteria that

Correspondence: Yanbing Ma
Institute of Medical Biology, Chinese Academy of Medical Sciences and Peking Union Medical College, NO935#, Jiaoling Road, Kunming, People's Republic of China
Tel +86 6 833 9287
Email yanbingma@126.com

generally affect individuals with bacterial pneumonia-induced morbidity and mortality, particularly opportunistic and cross-infected bacteria within long-term in-hospital respiratory medicine patients.^{5,6} *Staphylococcus aureus* alpha toxin (AT) has recently been identified as a risk factor for host mucosal defense and macrophage function. Coinfection with *S. aureus* expressing AT improves the dissemination and propagation of gram-negative bacteria, such as *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, and *Acinetobacter baumannii*.⁷ No drugs can effectively eliminate and clear MDR bacteria. Therefore, therapeutic strategies for MDR bacterial coinfection remain to be elucidated.

Antimicrobial peptides (AMPs), which are widely present in a range of organisms, exhibit a variety of biological activities, including antibacterial,^{4,8,9} antifungal,¹⁰ antiparasitic,¹¹ antiviral,¹² and anticancer¹³ activities. Generally, AMPs are distinguished into four structural classes: α -helical peptides, β -sheet peptides, extended peptides, and loop peptides,^{14–17} of which α -helical peptides represent the largest proportion of the AMPs that have been found. β -Sheet peptides are also widely studied, because of their intramolecular disulfide bond-mediated rigidity in aqueous solution. *Tachyplesin III*, a β -sheet AMP,¹⁸ not only shows potential activity against MDR bacterial infection but also exhibits low cytotoxicity to mammalian cells.¹⁹ The cationic β -hairpin structure of *Tachyplesin III* provides strong stability under serum conditions and provides the foundation for bacteria killing by directly interacting with lipid membranes²⁰ or binding to bacterial DNA.²¹ Recently, our group reported a novel mechanism by which *Tachyplesin III* may target FabG in bacteria killing.⁹

Except the capability of AMPs killing bacterial directly, AMPs also show potential role in innate immune cell regulation.¹⁶ *Cathelicidin-BF*, an α -helix structure AMP, shown a broad and strong ability in anti-MDR bacterial infection in vitro and in vivo. Pretreatment with *Cathelicidin-BF* significantly recruit neutrophils and macrophage to the lung and induce neutrophil extracellular traps formation which reduced bacterial burden and improve mice survival.⁸ LL-37 regulates the differentiation and phagocytosis of DC cells which is dependent on overexpression of phagocytosis receptor and costimulatory molecule CD86/CD80/CD40.²² Whether *Tachyplesin III* mediate innate immune cells especially monocytes remains unclear, we doubt that *Tachyplesin III* could remarkably increase mice against MDR bacterial infection

by the directly killing mechanism collaboration with regulate innate immune cells infiltration to the infection site. Thus, it is essential to evaluate the ability of *Tachyplesin III* to achieve mixed bacterial clearance in vivo. Herein, we identified a potential role for the AMP *Tachyplesin III* in host defense against *P. aeruginosa* and *A. baumannii* coinfection.

Materials and methods

Ethics statement

The animal experiments were approved by the Ethics Committee of Animal Care and Welfare of the Institute of Medical Biology, Chinese Academy of Medical Sciences (CAMS) and Peking Union Medical College (PUMC) (Permit Number: SYXK (dian) 2010-0007) in accordance with the animal ethics guidelines of the Chinese National Health and Medical Research Council (NHMRC) and the Office of Laboratory Animal Management of Yunnan Province, China. All efforts were made to minimize animal.

Cell culture and treatment

MH-S (mouse alveolar macrophage), which was purchased from Chinese Academy of Sciences cell bank, was cultured in RPMI 1640 medium containing 10% FBS, 1% penicillin-streptomycin, 1% nonessential amino acids, and 1% L-glutamine (Solarbio® Life Science). Cells were seeded onto 12-well plates at a million cells per well and incubated overnight. One hour before *Tachyplesin III* (Southeast Asian horseshoe crab, KWCFRVCYRGICYRKCR-NH2, Sangon® Biotech, China) treatment, the cells were washed three times with PBS, incubated with fresh RPMI 1640 medium without antibiotics, and then treated with *Tachyplesin III* at concentrations of 0 μ g/mL, 10 μ g/mL, and 100 μ g/mL for 3 hrs. *Tachyplesin III* was resuspended in fresh RPMI 1640 medium.

Neutral red phagocytosis

Briefly, after pretreatment with *Tachyplesin III*, MH-S cells were washed three times with PBS before neutral red staining. Next, 500 μ L fresh RPMI 1640 medium containing 0.05% neutral red medium that was filtered with a 0.22 μ m filter was added to the 12-well plates and incubated for 4 mins. After removing the medium and washing the cells three times with PBS, multiple red bubbles could be observed inside the treated cells under a microscope (Nikon, Japan). Acid-ethanol solution consisting of 50% ethanol, 1% acetic

acid, and 49% ddH₂O was subsequently used for neutral red extraction. Finally, the results were determined by absorbance measurements with a NanoDrop 2000 at 540 nm, which is the absorption peak for neutral red solution. The peptide Cramp (GLLRKGGEKIGEKLLKIGQKIKNFFQK LVPQPEQ) was employed as control peptide.

Animals

Six- to eight-week-old female *BALB/c* mice were purchased from Beijing Vital River Laboratory Animal Technology Co. Ltd. and maintained in the Central Animal Care Services of our institute under specific pathogen-free (SPF) conditions.

Bacterial strains and infection of mice

Two MDR clinical isolates that were inventory from our own laboratory, *P. aeruginosa* 1409 and *A. baumannii* 1408, were identified with a Vitek 32 system (bioMerieux, France) and further verified by 16S rDNA sequencing with the universal primers 27f and 1492R as previously described.⁴ Both *P. aeruginosa* 1409 and *A. baumannii* 1408 were grown overnight with shaking in Luria-Bertani (LB) medium at 37°C and then subcultured with fresh LB medium for 4 hrs. When the bacterial concentration corresponding to an OD₆₀₀ value reached 0.5, the culture was centrifuged at 5000× g for 10 mins, and the bacteria were washed twice with PBS (8 g/L NaCl, 0.2 g/L KCl, 1.44 g/L Na₂HPO₄, 0.24 g/L KH₂PO₄, pH 7.4) and then resuspended in PBS.

Bacterial infection of mice and survival

To establish the murine bacteremia model, mice were intranasally injected with a 50 µL total volume bacterial solution. For *P. aeruginosa* mono-infection group, 2.5×10⁶ CFU bacterial were used for mice challenge. For *A. baumannii* mono-infection group, 1.5×10⁸ CFU bacterial were used for mice challenge. For coinfection group 2.5×10⁶ CFU *P. aeruginosa* plus 1.5×10⁸ CFU *A. baumannii* were used for mice challenge. To generate *P. aeruginosa* plus *S. aureus* co-infection model, mice were also intranasally injected with a 50 µL total volume bacterial solution. A lethal dose of *P. aeruginosa* which contains 5×10⁶ CFU bacterial were used for mice challenge. For sub-lethal *P. aeruginosa* mono-infection group, 2.5×10⁶ CFU bacterial were used for mice challenge. A total of 1.5×10⁹ CFU and 7.5×10⁸ CFU *S. aureus* were used for mono-infection group, respectively. For coinfection group, 2.5×10⁶ CFU *P. aeruginosa* plus 7.5×10⁸ CFU *S. aureus* were used for mice challenge. After bacterial challenge, mice survival rate was determined by continuous monitoring for 7 days,

which was considered the experiment endpoint, and then all living animals were euthanized.

BALF bacterial burden

To evaluate the BALF bacterial burden, mice were sacrificed 24 hrs after treatment. One milliliter fresh PBS was injected to the lung via minimally invasive tracheostomy, and then repeated lavage of lung tissues for three times. BALF bacterial burden was measured with a 10-fold dilution on LB plates, which were incubated at 37°C overnight. The next day, the CFU was counted, and 30–300 spots/plate were selected for statistical analysis and expressed as CFU/mL BALF.

Cytokine and chemokine measurements

The BALF extraction supernatants were analyzed for IL-6, TNF-α, MCP-1 (eBioscience), IL-1β, and MIP-2 (Abcam) concentration by an enzyme-linked immune sorbent assay (ELISA) kit following the manufacturer's instructions.¹⁶

Lung histological analysis

The lung tissue histology analysis was performed as previously described.³⁵ Briefly, the left lung tissues were harvested from mice the day after bacterial infection and fixed in 4% paraformaldehyde at room temperature for 24 hrs. Subsequently, the tissues were embedded in paraffin and stained with a hematoxylin and eosin (H & E) kit purchased from Solarbio® Life Science.

Statistical analysis

All data were analyzed using GraphPad Prism 7.0 and presented as the mean±standard deviation (SD). The non-parametric log-rank test was used for survival rate comparisons, and the other comparisons were performed using Student's *t*-test or one-way ANOVA between two or three groups, respectively. *indicates *P*-value<0.05, **indicates *P*-value<0.01, ***indicates *P*-value<0.001.

Results

Mixed infection with *P. aeruginosa* and other bacteria reduces mouse survival

To identify the damage of *P. aeruginosa* and *A. baumannii* mixed infection in vivo, we mimicked the pathogenic pattern of these two MDR bacteria. Therefore, three infected BALB/c mouse groups, representing *P. aeruginosa* and *A. baumannii* single and mixed infections, containing eight mice per group, were employed. The mixed-infection mouse group were

intranasally infected with a 50 μ L total volume of mixed bacteria consisting of 2.5×10^6 CFU of live *P. aeruginosa* and 1.5×10^8 CFU of live *A. baumannii*; the same doses were used in the single infection groups. Although the mice were coinfecting with sub-lethal doses of bacteria, fewer than 40% of the mice survived to the 7-day study endpoint in the coinfection group, whereas only one *P. aeruginosa*-infected mouse died ($P < 0.05$), and none of the *A. baumannii*-infected mice died ($P < 0.01$) (Figure 1). To investigate whether *P. aeruginosa* could potentiate Gram-positive bacterial infections, mice were intranasally infected with *P. aeruginosa* in combination with *S. aureus*. As expected, most animals died in the coinfection group (Figure S1), which is consistent with recent studies.^{7,23} Our data indicate that compared with single infection, mixed bacterial infection with *P. aeruginosa* and Gram-negative *A. baumannii* or gram-positive *S. aureus* significantly reduced the survival rate.

Coinfection of *P. aeruginosa* and *A. baumannii* causes serious disease

Gram-negative bacterial pathogen infection can trigger a robust systemic response of pro-inflammatory cytokines and chemokines.²⁴ To investigate the pathogenic inflammatory reaction caused by bacterial infection, we first detected the levels of pro-inflammatory cytokines and chemokines in BALF collected at 24 hrs after bacterial infection. During *P. aeruginosa* and/or *A. baumannii* infection, the expression of IL-1 β , IL-6, TNF- α , MIP-2, and MCP-1 was significantly induced in the infection groups compared with the control (Ctrl) groups. The production of cytokine IL-1 β and chemokine MCP-1 in the mixed-infection group was significantly higher than that in the mono-infection groups. Interestingly,

Figure 1 Coinfection of *P. aeruginosa* and *A. baumannii* reduces mouse survival. Survival curve of BALB/c mice ($n=8$) infected with *P. aeruginosa* (2.5×10^6 CFU) (Pa) alone, *A. baumannii* (1.5×10^8 CFU) (Aba) alone, or *P. aeruginosa* + *A. baumannii*. Survival rates were monitored daily for a week, and survival rate significance was determined by log-rank test. * $P < 0.05$ compared to mono-*P. aeruginosa* infection, ** $P < 0.01$ compared to *A. baumannii* mono-infection, and data represent at least three trials. The animals were euthanized the day after observation endpoint.

there were no differences in IL-6, TNF- α , and MIP-2 production between the *P. aeruginosa* plus *A. baumannii* mixed-infection group and the *P. aeruginosa* mono-infection group, but the expression of IL-6 and TNF- α in the mixed-infection group was higher than that in the *A. baumannii* mono-infection group (Figure 2A). Histological analysis show that, in contrast to WT mice, bacterial infection induced histological lung injury and the coinfection group presented the most serious injury. Increased vascular leakage and bleeding, inflammatory cell infiltration and alveolar disruption occurred in the single-bacterial infection group, and this condition was worse in the mixed-infection group (Figure 2B). In line with these results, the H & E score also consist with pathological section results (Figure 2C). In summary, our results demonstrate that pro-inflammatory cytokine IL-1 β and chemokine MCP-1 were significantly increased and lung injury was significantly increased in the coinfection group.

Pretreatment with *Tachyplesin III* protects mice against mixed bacterial infection

Tachyplesin III is cytotoxic to mammalian cells and shows MDR bacterial-killing activity.¹⁹ To test the toxicity of *Tachyplesin III* in vivo, the mice were intravenously (i.v.) injected with a 5, 10, and 20 mg/kg gradient peptide dose. Only one mouse in the 20 mg/kg group died, and none of the mice in the 5 and 10 mg/kg dose groups died (Figure 3A). The 10 mg/kg dose was chosen for the infection study. First, we try a therapeutic plan to i.v. inject with 10 mg/kg *Tachyplesin III* and control peptide *Cathelicidin-BF* 2 hrs after lethal dose of *P. aeruginosa* infection, *Cathelicidin-BF* significantly increased the survival rate as we described earlier, but *Tachyplesin III* did not show any potential activity in vivo (Figure S2). Then, BALB/c mice were treated (i.v.) with a single dose of 10 mg/kg *Tachyplesin III*, and *Meropenem* was used as a control. Both *P. aeruginosa* and *A. baumannii* were intranasally injected 24 hrs after peptide inoculation. As expected, no deaths occurred within 24 hrs after peptide and bacterial inoculation. Strikingly, in the preventive treatment condition, the mortality rate was significantly reduced within a week in the *Tachyplesin III* peptide-treated group but not in the *Meropenem*-treated group (Figure 3B). To further investigate the pivotal role of *Tachyplesin III* in the coinfection model, the bacterial burden in BALF was measured at 24 hrs after infection. As expected, the bacterial loads in the *Tachyplesin III* peptide-treated group were significantly lower than those in the model and control groups (Figure 3C). Taken together, our findings strongly suggest a potential role for *Tachyplesin III* in triggering an

Figure 2 Bacterial coinfection induced more cytokines production in BALF and worse lung injury in vivo than mono-infection. **(A)** Mice were intranasally infected with co- and mono-bacterial solutions, and BALF was collected in 1 mL cold phosphate buffer saline (PBS) at 24 hrs after injection. BALF was collected after inoculation, and the levels of pro-inflammatory cytokines IL-1 β , IL-6, and TNF- α and chemokines MIP-2 and MCP-1 were determined by ELISA. Coinfection enhanced cytokine and chemokine production. Bars represent the mean $\pm$ SEM for six mice. **(B)** Hematoxylin-eosin (H & E) staining of the lung tissues from mice at 24 hrs after infection. Representative sections of control mice and mono- and co-infected mice are shown. Obvious bleeding and inflammatory cell infiltration were observed, indicating that bacterial coinfection enhances worse lung injury in vivo. **(C)** H & E pathological score results are shown as histograms. The data are shown as the mean $\pm$ SEM. Comparisons among two groups were calculated using paired t-test and one-way ANOVA for three or more groups (* P <0.05, ** P <0.01, *** P <0.001, **** P <0.0001).

Abbreviation: ns, no significance.

antimicrobial response against mixed bacterial infection, reducing the bacterial burden in lung tissues and improving survival.

Tachyplesin III reduces pro-inflammatory cytokines and chemokines production and ameliorates tissue damage in vivo

To determine whether *Tachyplesin III* reduces the systemic response of pro-inflammatory cytokines induced by the

mixed bacterial infection with *P. aeruginosa* plus *A. baumannii*, the level of serum cytokines and chemokines was measured. As expected, *Tachyplesin III* therapy reduced the production of pro-inflammatory cytokines as IL-1 β , IL-6, and TNF- α , but there was no difference in the production of chemokine MCP-1 and MIP-2 (Figure 4A). Histological examination showed no difference between the *Tachyplesin III* group and the wild-type group, which indicated that injection (i.v.) with 10 mg/kg *Tachyplesin III* did not obviously cause lung injury (Figure 4B). Treatment with

Figure 3 Protective efficacy of *Tachyplesin III* therapy in a mixed bacterial infection murine model of pneumonia. **(A)** Survival curve of BALB/c mice (n=6) intravenously infected with *Tachyplesin III* at different doses of 5, 10, and 20 mg/kg. Only one mouse died in the 20 mg/kg group, and no mice died in the 5, 10 mg/kg group, indicating that *Tachyplesin III* was safe through intravenous injection at 10 mg/kg. **(B)** The therapeutic potential of *Tachyplesin III* in bacterial coinfection pneumonia. Pretreatment with 10 mg/kg *Tachyplesin III* significantly increased the mouse survival rate compared with that of the model group, while meropenem-treated mice had no effect on MDR bacterial infection. **(C)** BALF bacterial burden in the bacterial coinfection group. Viable bacteria in the BALF that were collected at 24 hrs after bacterial infection were decreased in the pretreatment with *Tachyplesin III* group compared with the non-treatment (Model) and Meropenem-treated (Ctrl) groups. Bars represent the mean±SEM, survival rates were monitored daily for a week, and survival rate significance was determined by log-rank test, * $P < 0.05$; BALF comparisons among two groups were calculated using paired t-test and one-way ANOVA for three or more groups (* $P < 0.05$, ** $P < 0.01$). The animals were euthanized the day after observation endpoint.

Tachyplesin III resulted in decreased inflammatory cell infiltration, vascular leakage, and alveolar disruption compared with those in the coinfecting groups or in the Meropenem-treated group (Figure 4B). In line with these results, the H & E score also consist with pathological section results (Figure 4C). Collectively, these data indicated that *Tachyplesin III* plays a crucial role in the prophylaxis of MDR bacterial mixed infection-induced lung injury and inflammatory cytokine production.

Tachyplesin III enhances the phagocytic function of mouse alveolar macrophages

Macrophage phagocytosis, which is a secondary line of defense line in host innate immunity against bacterial infection, plays an important role in bacterial clearance.²⁵ To evaluate the effect of *Tachyplesin III* on macrophage phagocytosis, we treated the mouse alveolar macrophage cell line MH-S with different *Tachyplesin III* concentrations in

vitro. After MH-S cells were exposed to *Tachyplesin III* for 3 hrs, a significant dose-dependent phagocytosis of neutral red was observed (Figure 5A). In addition, after extracting the neutral red from MH-S cells, the concentration of neutral red was significantly increased in *Tachyplesin III*-treated cells than that in control peptide- (Cramp) treated cells (Figure 5B). Macrophage phagocytosis was increased in a dose-dependent manner (Figure 5C). Altogether, these results indicated that *Tachyplesin III* induced obvious alveolar macrophage phagocytosis.

Discussion

For nearly 40 years, no novel antibiotics have been discovered or developed, while multiple natural substances, such as AMPs, have recently been described.¹ However, the entire process from identifying to refining antimicrobial products, which are urgently needed for contemporary clinical therapy, is slow and time-consuming, although several

Figure 4 Pulmonary pathological changes of *Tachyplesin III*-treated mice. **(A)** Cytokine and chemokine production was reduced in *Tachyplesin III*-pretreated mice. Serum was collected 24 hrs after bacterial infection, and IL-1 β , IL-6, TNF- α , MIP-2, and MCP-1 were identified by ELISA. **(B)** Lung tissue was stained with H & E, no significant tissue damage occurred in *Tachyplesin III* (10 mg/kg), and less bleeding and inflammatory immune cell infiltration occurred in *Tachyplesin III*-treated mice than in the model and control groups. **(C)** H & E pathology was scored blindly in mouse lungs after infection with *P. aeruginosa* and *A. baumannii*. The data are shown as the mean $\pm$ SEM. Serum cytokine and lung pathology score comparisons among the two groups were calculated using paired t-test and one-way ANOVA for three or more groups (* P <0.05, ** P <0.01, *** P <0.001).

representative AMPs, such P-113, MSI-78, MBI-226, IB-367, and rBPI21, have been applied in clinical trials.^{26,27,28} According to a report by the World Health Organization (WHO), the use of antibiotics extends the human life span at least 20 years. Nevertheless, this protective effect is becoming increasingly weaker with bacterial evolution, particularly the emergence and spread of MDR bacteria, which may contribute to 30 million deaths by 2050.²⁹

Clinically, the Gram-positive bacterium *S. aureus* is often observed in coinfections with gram-negative bacteria, including *P. aeruginosa*, *K. pneumoniae*, and *A. baumannii*, which depend on the influence of toxins on mucosal host defense.⁷ Carbapenem-resistant (CSE) *A. baumannii* and *P. aeruginosa*, two Gram-negative bacteria, are the most critical global priority opportunistic pathogens. In the clinic, *P. aeruginosa* plus *A. baumannii* is the most common and dangerous combination³⁰ and a critical contributor to the morbidity

and mortality of long-term in-hospital respiratory medicine patients, particularly VAP patients.^{5,6} Our study revealed that compared with monoinfection, coinfection with *P. aeruginosa* plus *A. baumannii* significantly reduced mouse survival, and only one mouse died with *P. aeruginosa* infection alone (Figure 1A). Interestingly, none of the mice intranasally infected with 1.5×10^8 CFU live *A. baumannii* died, whereas the addition of less than 3×10^6 CFU *P. aeruginosa* significantly reduced the survival rate, suggesting that the synergistic interactions of these bacteria increased disease severity, which coincides with the clinical phenomenon.³¹ Therefore, this lethal pneumonia model was further analyzed. Gram-negative bacterial pathogen infections can trigger a robust systemic response of pro-inflammatory cytokines and chemokines.³² In the current study, we indicated that, in the case of mixed bacterial infection, the pro-inflammatory cytokines IL-1 β , IL-6, and TNF- α , known as a “cytokine storm”,

Figure 5 *Tachyplesin III* enhanced the phagocytosis of mouse alveolar macrophages in vitro. **(A)** Bright-field image showing neutral red phagocytosis of MH-S cells in a dose-dependent manner with *Tachyplesin III* treatment. **(B)** Quantification of phagocytic capacity via the OD₅₄₀ nm measurement value of neutral red extract. **(C)** MH-S cells were incubated with *Tachyplesin III* and control peptides. The levels of neutral red phagocytosis were measured by OD₅₄₀ nm detection. The data are shown as the mean±SD, and comparisons among two groups were calculated using paired t-tests and one-way ANOVA for three or more groups (* $P < 0.05$, ** $P < 0.01$).

were significantly increased compared with those in the single-bacterial infection group, which may be the main cause of multiple organ failure or death. Furthermore, MDR bacterial coinfection induced worse lung injury as well as multiple inflammatory cell infiltration, alveolar disruption, and vascular leakage (Figure 2).

Our previous work demonstrated that *Tachyplesin III* provides an antibacterial effect against MDR bacteria in vitro³³ which may present strong anti-MDR bacteria activity by interacting with FabG to trigger bacterial death⁹ and this treatment is hypothesized to provide bacterial clearance in vivo. Therefore, to further examine the influence of *Tachyplesin III* on mixed bacterial infection, we proposed a murine pneumonia model that enabled us to evaluate *Tachyplesin III* activity in vivo. Herein, we found that

although *Tachyplesin III* showed cytotoxicity to mammalian cells in vivo, no animal death or lung injury occurred in mice injected (i.v.) with a 10 mg/kg dose (Figures 3A and 4A–E). Strangely, we found that *Tachyplesin III* i.v. injection had non-therapeutic effect in the lethal dose of *P. aeruginosa* challenge (Figure S2) which reminds us that future research should focus on the development of detoxified form of *Tachyplesin III*. In a murine pneumonia model, a low lethal dose of *P. aeruginosa* significantly reduced the survival rate of *A. baumannii*-infected and *S. aureus*-infected mice (Figure S1), which was prevented by pretreatment with *Tachyplesin III* (Figure 3B). Consistent with the reduced mortality, pretreatment with *Tachyplesin III* significantly reduced the BALF bacterial burden, decreased the cytokine storm, and reduced lung injury, which may be the main

causes of animal death (Figures 3B and 4A–C). Compared with mono-infections, mixed infections with *P. aeruginosa* and *A. baumannii* trigger even more serious disease in vivo.

Alveolar macrophages and neutrophils are the first immune cells in the host innate response to bacterial pathogen infection.³⁴ Our recent work indicated that *Cathelicidin-BF*, another AMP subtype, enhanced neutrophil extracellular traps, which are the main facilitators of the capture and clearance of bacteria through the activation of innate immunity.³⁵ Other studies have indicated that macrophages also contribute to the clearance of bacterial pathogen infection.³⁶ In this study, our results demonstrated that *Tachyplesin III* improved alveolar macrophage phagocytosis in a dose-dependent manner in vitro (Figure 5), which may directly reduce the BALF burden (Figure 3C). Thus, our study reports a novel therapeutic strategy for gram-negative MDR bacterial coinfection using the AMP *Tachyplesin III*.

Taken together, the present work illustrated that coinfection with *P. aeruginosa* plus *A. baumannii* causes more serious disease than individual bacterial infections, especially by the release of pro-inflammatory cytokine IL-1 β and a decrease in immune cell infiltration, thereby triggering lung injury. Dramatically, the prophylactic capability of *Tachyplesin III* against clinical MDR bacterial isotype infection was identified. This phenomenon could be partially explained by significantly enhanced survival, reduced BALF bacterial burden, decreased pro-inflammatory cytokine and chemokine levels, and reduced lung injury. Simultaneously, the phagocytic function of macrophages was significantly increased under *Tachyplesin III* in vitro.

Conclusion

In conclusion, our findings indicate that the AMP *Tachyplesin III* plays a potential role against MDR bacterial coinfection in vivo. Pretreatment with *Tachyplesin III* ameliorates bacterial coinfection in mice by enhancing macrophage phagocytosis and reducing the production of pro-inflammatory cytokines. Our results highlight the importance of external AMPs against MDR bacterial mixed infection and present a possible therapeutic strategy for controlling multiple bacterial coinfections that should be confirmed in clinical studies.

Acknowledgments

This work was financially supported by the National Natural Science Foundation of China (grant numbers 81503117 and 81460322), the Innovation Fund of

IMBCAMS grant number 2018018001, the Fundamental Research Funds for the Central Universities of China (grant numbers 2018018001), the CAMS Initiative for Innovative Medicine (grant number 2017-I2M-3-022 and 2016-I2M-019), and the Opening Foundation of the Key Laboratory of Bioactive Peptides of Yunnan Province (grant number AMHD-2017-1). The co-first authors of this paper are Jialong Qi, Ruiyu Gao, and Cunbao Liu.

Disclosure

The authors declare no conflicts of interest in this work.

References

- Kang S-J, Park SJ, Mishig-Ochir T, Lee B-J. Antimicrobial peptides: therapeutic potentials. *Expert Rev Anti Infect Ther*. 2014;12(12):1477–1486. doi:10.1586/14787210.2014.976613
- Francis AM. The wages of sin: how the discovery of penicillin reshaped modern sexuality. *Arch Sex Behav*. 2013;42(1):5–13. doi:10.1007/s10508-012-0018-4
- Laxminarayan R, Duse A, Wattal C, et al. Antibiotic resistance—the need for global solutions. *Lancet Infect Dis*. 2013;13(12):1057–1098. doi:10.1016/S1473-3099(13)70318-9
- Liu C, Shan B, Qi J, Ma Y. Systemic responses of multidrug-resistant *Pseudomonas aeruginosa* and *Acinetobacter baumannii* following exposure to the antimicrobial peptide cathelicidin-BF imply multiple intracellular targets. *Front Cell Infect Microbiol*. 2017;7. doi:10.3389/fcimb.2017.00517
- Lacy MK, Stryjewski ME, Wang W, et al. Telavancin hospital-acquired pneumonia trials: impact of Gram-negative infections and inadequate Gram-negative coverage on clinical efficacy and all-cause mortality. *Clin Infect Dis*. 2015;61(Suppl 2):S87–S93. doi:10.1093/cid/civ536
- Karlowsky JA, Draghi DC, Jones ME, Thornsberry C, Friedland IR, Sahm DF. Surveillance for antimicrobial susceptibility among clinical isolates of *Pseudomonas aeruginosa* and *Acinetobacter baumannii* from hospitalized patients in the United States, 1998 to 2001. *Antimicrob Agents Chemother*. 2003;47(5):1681–1688. doi:10.1128/aac.47.5.1681-1688.2003
- Cohen TS, Hilliard JJ, Jones-Nelson O, et al. *Staphylococcus aureus* α toxin potentiates opportunistic bacterial lung infections. *Sci Transl Med*. 2016;8(329):329ra31–329ra31. doi:10.1126/scitranslmed.aad9922
- Cai S, Qiao X, Feng L, et al. Python cathelicidin CATHPb1 protects against multidrug-resistant staphylococcal infections by antimicrobial-immunomodulatory duality. *J Med Chem*. 2018;61(5):2075–2086. doi:10.1021/acs.jmedchem.8b00036
- Liu C, Qi J, Shan B, Ma Y. Tachyplesin causes membrane instability that kills multidrug-resistant bacteria by inhibiting the 3-Ketoacyl carrier protein reductase FabG. *Front Microbiol*. 2018;9:825.
- Wu H, Ong ZY, Liu S, et al. Synthetic beta-sheet forming peptide amphiphiles for treatment of fungal keratitis. *Biomaterials*. 2015;43:44–49. doi:10.1016/j.biomaterials.2014.11.052
- Lacerda AF, Pelegrini PB, de Oliveira DM, Vasconcelos EA, Grossi-de-Sa MF. Anti-parasitic peptides from arthropods and their application in drug therapy. *Front Microbiol*. 2016;7:91. doi:10.3389/fmicb.2016.00091
- Luo X, Guo Y, Bao J, et al. Characterization and antiviral activities of a novel bovine IFN- ω 24. *Mol Immunol*. 2015;66(2):357–363. doi:10.1016/j.molimm.2015.04.009
- Pan CY, Lin CN, Chiou MT, Yu CY, Chen JY, Chien CH. The antimicrobial peptide pardaxin exerts potent anti-tumor activity against canine perianal gland adenoma. *Oncotarget*. 2015;6(4):2290–2301. doi:10.18632/oncotarget.2959

14. Karal MA, Alam JM, Takahashi T, Levadny V, Yamazaki M. Stretch-activated pore of the antimicrobial peptide, magainin 2. *Langmuir*. 2015;31(11):3391–3401. doi:10.1021/la503318z
15. Cirioni O, Simonetti O, Pierpaoli E, et al. Enhanced efficacy of combinations of pexiganan with colistin versus acinetobacter baumannii in experimental sepsis. *Shock*. 2016;46(2):219–225.
16. Zhao L, Lu W. Defensins in innate immunity. *Curr Opin Hematol*. 2014;21(1):37–42. doi:10.1097/MOH.0000000000000005
17. Kuzmin DV, Emelianova AA, Kalashnikova MB, et al. Comparative in vitro study on cytotoxicity of recombinant beta-hairpin peptides. *Chem Biol Drug Des*. 2018;91(1):294–303. doi:10.1111/cbdd.13081
18. Miyata T, Tokunaga F, Yoneya T, et al. Antimicrobial peptides, isolated from horseshoe crab hemocytes, tachyplesin II, and polyphemusins I and II: chemical structures and biological activity. *J Biochem*. 1989;106(4):663–668. doi:10.1093/oxfordjournals.jbchem.a122913
19. Nakamura T, Furunaka H, Miyata T, et al. Tachyplesin, a class of antimicrobial peptide from the hemocytes of the horseshoe crab (*Tachyplesus tridentatus*). Isolation and chemical structure. *J Biol Chem*. 1988;263(32):16709–16713.
20. Katsu T, Nakao S, Iwanaga S. Mode of action of an antimicrobial peptide, tachyplesin I, on biomembranes. *Biol Pharm Bull*. 1993;16(2):178–181. doi:10.1248/bpb.16.178
21. Yonezawa A, Kuwahara F, Fujii N, Sugiura Y. Binding of tachyplesin I to DNA revealed by footprinting analysis: significant contribution of secondary structure to DNA binding and implication for biological action. *Biochemistry*. 1992;31(11):2998–3004. doi:10.1021/bi00126a022
22. Davidson DJ, Currie AJ, Reid GS, et al. The cationic antimicrobial peptide LL-37 modulates dendritic cell differentiation and dendritic cell-induced T cell polarization. *J Immunol*. 2004;172(2):1146–1156. doi:10.4049/jimmunol.172.2.1146
23. Cohen TS, Hilliard JJ, Jones-Nelson O, et al. Staphylococcus aureus alpha toxin potentiates opportunistic bacterial lung infections. *Sci Transl Med*. 2016;8(329):329ra31. doi:10.1126/scitranslmed.aaf0746
24. Shang D, Meng X, Zhang D, Kou Z. Antibacterial activity of chensinin-1b, a peptide with a random coil conformation, against multiple-drug-resistant *Pseudomonas aeruginosa*. *Biochem Pharmacol*. 2017;143:65–78. doi:10.1016/j.bcp.2017.07.017
25. Camberlein E, Cohen JM, Jose R, et al. Importance of bacterial replication and alveolar macrophage-independent clearance mechanisms during early lung infection with *Streptococcus pneumoniae*. *Infect Immun*. 2015;83(3):1181–1189. doi:10.1128/IAI.02788-14
26. Aloush V, Navon-Venezia S, Seigman-Igra Y, Cabili S, Carmeli Y. Multidrug-resistant *Pseudomonas aeruginosa*: risk factors and clinical impact. *Antimicrob Agents Chemother*. 2006;50(1):43–48. doi:10.1128/AAC.50.1.43-48.2006
27. Giles FJ, Rodriguez R, Weisdorf D, et al. A phase III, randomized, double-blind, placebo-controlled, study of isegagan for the reduction of stomatitis in patients receiving stomatotoxic chemotherapy. *Leuk Res*. 2004;28(6):559–565. doi:10.1016/j.leukres.2003.10.021
28. Bommarius B, Jenssen H, Elliott M, et al. Cost-effective expression and purification of antimicrobial and host defense peptides in *Escherichia coli*. *Peptides*. 2010;31(11):1957–1965. doi:10.1016/j.peptides.2010.08.008
29. Tucker AT, Leonard SP, DuBois CD, et al. Discovery of next-generation antimicrobials through bacterial self-screening of surface-displayed peptide libraries. *Cell*. 2018;172(3):618–628.e13. doi:10.1016/j.cell.2017.12.009
30. Lu Q, Luo R, Bodin L, et al. Efficacy of high-dose nebulized colistin in ventilator-associated pneumonia caused by multidrug-resistant *Pseudomonas aeruginosa* and *Acinetobacter baumannii*. *Anesthesiology*. 2012;117(6):1335–1347. doi:10.1097/ALN.0b013e31827515de
31. Korgaonkar A, Trivedi U, Rumbaugh KP, Whiteley M. Community surveillance enhances *Pseudomonas aeruginosa* virulence during polymicrobial infection. *Proc Natl Acad Sci U S A*. 2013;110(3):1059–1064. doi:10.1073/pnas.1214550110
32. Li S-A, Xiang Y, Wang Y-J, Liu J, Lee W-H, Zhang Y. Naturally occurring antimicrobial peptide OH-CATH30 selectively regulates the innate immune response to protect against sepsis. *J Med Chem*. 2013;56(22):9136–9145. doi:10.1021/jm401134n
33. Liu CB, Shan B, Bai HM, Tang J, Yan LZ, Ma YB. Hydrophilic/hydrophobic characters of antimicrobial peptides derived from animals and their effects on multidrug resistant clinical isolates. *Zool Res*. 2015;36(1):41–47. doi:10.13918/j.issn.2095-8137.2015.1.41
34. Jones MCY, Romela IR, Hebroni F, Seema BP, Caiyun X, Lloyd SM. Neutrophil-derived IL-1b is sufficient for abscess formation in immunity against *Staphylococcus aureus* in mice. *Plos Pathogen*. 2012;8(11):e1003047. doi:10.1371/journal.ppat.1003047
35. Liu CB, Qi JL, Shan B, et al. Pretreatment with cathelicidin-BF ameliorates *Pseudomonas aeruginosa* pneumonia in mice by enhancing NETosis and the autophagy of recruited neutrophils and macrophages. *Int Immunopharmacol*. 2018;65:382–391. doi:10.1016/j.intimp.2018.10.030
36. Wan M, van der Does AM, Tang X, Lindbom L, Agerberth B, Haeggström JZ. Antimicrobial peptide LL-37 promotes bacterial phagocytosis by human macrophages. *J Leukoc Biol*. 2014;95(6):971–981. doi:10.1189/jlb.0513304

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

Dovepress