Vitamin D Deficiency Is Associated With Impaired Muscle Strength And Physical Performance In Community-Dwelling Older Adults: Findings From The English Longitudinal Study Of Ageing

This article was published in the following Dove Press journal:
Clinical Interventions in Aging

Niamh Aspell¹
Eamon Laird¹
Martin Healy²
Brain Lawlor³
Maria O'Sullivan¹

¹Department of Clinical Medicine, School of Medicine, Trinity College Dublin, Trinity Centre for Health Sciences, Dublin 8, Ireland; ²Department of Biochemistry, St James's Healthcare Campus, Dublin 8, Ireland; ³Department of Medical Gerontology and Institute of Neurosciences, Trinity College, Dublin, Ireland

Purpose: Maintaining skeletal muscle function throughout life is a crucial component of successful ageing. Adequate vitamin D status may be important in preserving muscle function. We aimed to determine the association between impaired muscle function and serum vitamin D status in community-dwelling older adults. Falls were explored as a secondary aim.

Methods: Data were analyzed from adults aged ≥ 60 years, from Wave 6 of the English Longitudinal Study of Aging (ELSA). Handgrip strength (HGS) and the short physical performance battery (SPPB) were employed as measures of muscle strength and physical function, respectively. Serum 25-hydroxyvitamin D [25(OH)D] was assessed with concentration <30 nmol/L classed as vitamin D deficient.

Results: The study comprised 4157 community-dwelling adults with a mean age of 69.8 (SD 6.9). Overall, 30.6% had low HGS and 12.7% had low SPPB (≤ 6). Participants with the lowest serum 25(OH)D (<30 nmol/L) had the highest prevalence of impaired muscle strength and performance (40.4% and 25.2%) compared with participants with levels ≥ 50 nmol/L (21.6% and 7.9%). Consistent with this, vitamin D deficiency (<30 nmol/L) was a significant determinant of low HGS (OR 1.44 [1.22, 1.71], $p < 0.001$) and poor physical performance (OR 1.65 [1.31, 2.09], $p < 0.001$) in the logistic regression models. Older adults partaking in regular moderate physical activity had significantly lower odds of impaired muscle strength (OR 0.65 [0.58, 0.79]) and physical function (OR 0.30 [0.24, 0.38]), $p < 0.001$, respectively. Single or multiple falls (15.8% and 10.5% in past year) were not associated with vitamin D status.

Conclusion: Vitamin D deficiency was associated with impaired muscle strength and performance in a large study of community-dwelling older people. It is generally accepted that vitamin D deficiency at the <30 nmol/L cut-off should be reversed to prevent bone disease, a strategy that may also protect skeletal muscle function in ageing.

Keywords: vitamin D, 25-hydroxyvitamin D, ageing, muscle function, short physical performance battery

Introduction

Maintaining muscle mass and function throughout life is a crucial component of successful ageing. The progressive loss of skeletal muscle mass accompanied by a loss of muscle strength or performance, in ageing, are features of primary sarcopenia.^{1,2} This functional decline negatively impacts on mobility, falls risk, independence, quality of life and healthcare costs.²⁻⁵ The condition is a major

Correspondence: Maria O'Sullivan
Human Nutrition, School of Medicine,
Trinity College Dublin, Trinity Centre for
Health Sciences, St James's Healthcare
Campus, Dublin 8, Ireland
Tel +353 1 896 4039
Email maria.osullivan@tcd.ie

public health issue, estimated to effect between 6% and 22% of older adult populations⁶ and up to half of adults over 80 years. Resistance exercise is well evidenced in maintaining muscle function⁶ while nutritional factors such as adequate dietary protein,^{6–9} and potentially adequate vitamin D status,^{8–10} have shown benefit. Globally, vitamin D deficiency is widespread¹¹ particularly in older populations^{12,13} and optimizing vitamin D status may be a practical and cost-effective approach that supports skeletal muscle function.

Findings from animal models and laboratory studies show that vitamin D can improve muscle quality and strength and reduce intramuscular fat and inflammation.¹⁴ Epidemiological^{15–17} and intervention studies^{8,10} further support associations between circulating 25-hydroxyvitamin D [25(OH)D] or vitamin D₃ supplementation and muscle function in older adults, though more recent findings suggest no benefit.^{18–20} For falls outcomes, vitamin D supplementation appears beneficial for older people living in care homes²¹ or with low vitamin D status,²² but the findings are inconsistent for the wider population of community-dwelling older adults.²² Some studies, conversely, have found an increase in falls rate with large bolus vitamin D supplementation.²³

Several methodological differences are apparent across the studies that have investigated vitamin D and musculoskeletal outcomes in community-dwelling older adults; these include the population age, health and lifestyle characteristics, vitamin D status and the outcome measures employed for skeletal muscle. A key aspect is the degree to which study samples capture participants with established vitamin D deficiency or at-risk of deficiency. Such risk factors include advanced age, lower socioeconomic status, poorer health status, non-vitamin D supplement users, non-white ethnicity^{12,24} or mobility impairment.¹⁰ An important consideration is if older adults with vitamin D deficiency have impaired muscle function and, if so, would raising 25(OH)D status or reversing deficiency improve or preserve function. Skeletal muscle measures employed in studies are typically brief assessment of muscle strength or function including handgrip strength (HGS), gait speed or sit to stand tests.⁹ More comprehensive assessments, such as the short physical performance battery (SPPB)²⁵ or imaging techniques, that assess lower limb muscle function or muscle quality^{2,10,16} are important to add value to vitamin D research in this field.

The present study aimed to estimate the prevalence of impaired muscle function (HGS and SPPB) and determine

the association between these variables and serum vitamin D status in community-dwelling older adults from the English Longitudinal Study of Aging (ELSA). As a secondary aim, we explored the association between vitamin D status and falls.

Materials And Methods

Study Design And Population

The study was based on data from community-living older people who participated in ELSA²⁶ Wave 6 (2012–2013) as this represented the first analysis of 25(OH)D. The ELSA study was conducted in line with the Declaration of Helsinki and was approved by the London Multicentre Research Ethics Committee (MREC 01/2/91). Written informed consent was obtained from all participants (<https://www.elsa-project.ac.uk/>).

For the present study, the inclusion criteria were as follows: measured serum 25(OH)D, adults aged ≥ 60 years and participation in assessments of skeletal muscle function, namely HGS and components of the SPPB. Lower limb measures, such as gait speed, were assessed only in adults aged 60 years or over in ELSA. Overall, 7731 participants completed the nurse health visits, allowing the collection of biological data such as blood samples. The most common reasons for missing or unavailable blood samples were for participants who did not wish to consent to, or were considered ineligible for, blood sampling (e.g. bleeding disorders or anticoagulant use). After applying the inclusion criteria, the final analytical study sample comprised of 4157 subjects (Figure 1).

Serum 25(OH)D Measurement And Deficiency

Serum 25(OH)D was assessed from fasting blood sample and analyzed using Diasorin Liaison 25-hydroxyvitamin D immunoassay. All assays were performed in duplicate, the CV values ranged from 8.7% to 9.4%. The laboratory responsible for analysis of vitamin D for ELSA (Royal Victoria Infirmary, Newcastle upon Tyne, UK) participated in the Vitamin D External Quality Assessment Schemes (DEQAS). All blood sampling occurred from January 2012 to July 2013. Vitamin D status was classified according to the Institute of Medicine (IOM) guidelines,²⁷ with deficiency compatible with bone health risk defined as 25(OH)D < 30 nmol/L, and insufficiency as < 50 nmol/L.²⁷ Season was categorized using the extended vitamin D calendar for summer (April–Sept) and winter (Oct–March).²⁸

Figure 1 Participant enrolment in the study and reasons for exclusion.

Measures Of Muscle Strength And Physical Function

Handgrip strength was recorded as the mean grip strength of three readings using the participant's dominant hand. Gender and BMI-specific cut-offs were applied to categorize muscle weakness.¹ Physical performance was measured using the short physical performance battery (SPPB). The SPPB²⁵ is a composite battery of three physical tasks: an 8 foot (2.44 m) gait speed test, a repeated chair rise task and a static balance test and is supported by EWGSOP¹ as a measure of physical performance for research. Individual tests scores were combined to give a composite SPPB score, where 0–6 represented poor physical performance, 7–9 “at-risk” and >10 represented high performers.²⁵ We applied the cut-off of 0–6 to indicate poor or impaired lower extremity performance. Falls rate was reported as one fall or multiple falls in past 12 months; medical attention sought in relation to the fall was also noted.

Potential Confounders And Related Variables

Background socio-demographic, health status, lifestyle vitamin D and muscle function related data were recorded. This included: gender, educational attainment [minimum of O-level (secondary school) education achieved] and employment status (employed/retired). Health-related factors; self-reported health (compared to others of the same age); excellent, very good, good, fair and poor. Self-reported long-standing illness (Yes/No), whether the self-reported longstanding illness was limiting (Yes/No) and medication use (polypharmacy defined as taking 5 or more medications daily). BMI and waist

circumference were recorded. Lifestyle factors included physical activity categorized as mild, moderate and vigorous activity more than once per week. Alcohol intakes, high alcohol frequency (drinks alcohol most days), and current smoking status (Yes/No). Health behaviors which may influence serum 25(OH)D status in the ELSA dataset were season of blood sampling, sun holiday within the past 12 months (Yes/No) and vitamin D-containing supplement use, available as prescribed vitamin D and calcium for the treatment of osteoporosis. Vitamin D intake from non-prescribed vitamin D supplements or from diet was not available.

Data Analysis And Statistics

Descriptive statistics are presented as means and SD or counts and proportions for demographic and lifestyle characteristics. Normality was assessed visually using histogram matrix plots and ladder tables to identify appropriate transformations,²⁹ and Monte Carlo simulations to identify outliers.³⁰ Multiple logistic regression analysis was used to explore the determinants of poor physical performance and poor muscle strength and presented as odds ratios (OR) and 95% confidence intervals (CI). We applied multivariate analysis of variance (MANOVA) to all explanatory variables to test the assumptions of the model, before including them in a stepwise regression. Only variables that reached statistical significance in the initial analysis were considered for the models and after testing for collinearity by scatter plots and Variance Inflation Factors Tests for linear relationships. Tests were performed by the inclusion of continuous predictor variables to the model and were reported as OR for a single unit change with 95% CI. In adjusted models, covariates were selected based on previous research findings and following stepwise regression to test the contribution of each predictor (excluded *Pr.* >0.25). The contribution of covariates to muscle function was examined by including each set of covariates in the following order; model 1; poor physical performance defined as a score of 0–6 in the SPPB, socio-demographic factors, health-related behaviors, physical factors and health conditions. In model 2, low HGS strength defined by gender and BMI-specific cut-offs. Corresponding unadjusted results are also reported. All analyses were performed in STATA 14.0 (Stata Corp LP).

Results

Characteristics Of Study Population

The present study comprised 4157 community-dwelling adults with a mean age of 69.8 (SD 6.9) years. Demographic,

lifestyle and health characteristics are presented in Table 1. The prevalence of polypharmacy was 29.1% and longstanding limiting disease was 54.3%, though most participants (76.3%) rated their health as “good” or above. Overweight and obesity were common (73%), while 41.8% of participants reported partaking regularly in vigorous physical activity.

Mean serum 25(OH)D concentration was 49.7 (SD 23.5) nmol/L (Table 1). Over half of participants (53.7%) had serum 25(OH)D concentrations <50 nmol/L (insufficient) of which 21.8% were vitamin D deficient (<30 nmol/L). Vitamin D supplement use was low at 5.1%, though only prescribed vitamin D for the treatment of osteoporosis was available. Almost half (48.6%) of the study group reported sun holiday travel within the previous 12 months.

Prevalence Of Impaired Muscle Strength, Poor Physical Performance And Falls

In the present study, 30.6% of older adults had low HGS, as classified by gender and BMI-specific cut-offs (Table 2). Poor physical performance (SPPB ≤6) was identified in 12.7%, with a further 19.3% classed as “at-risk” (SPPB 7–9); most participants (67.9%) were considered high performers with scores of 10–12. A single fall or multiple falls occurred in 15.8% and 10.5%, respectively, in the past 12 months. The prevalence of impaired muscle strength, reduced performance and falls was consistently higher in women than men (Table 2).

Muscle Strength And Physical Performance According To Serum 25 (OH)D Concentration

Low HGS and SPPB according to 25(OH)D quintiles are presented in Figures 2 and 3 and Table 3. The lowest quintile (Q1, ≤28 nmol/L) was compatible with vitamin D deficiency, while the highest quintiles, Q4 and Q5, were compatible with sufficiency at 25(OH)D >50 nmol/L. In Q1, 40.4% of older adults had impaired HGS compared with 25.2% and 26.8% in Q4 and Q5, respectively. Consistent with this, the prevalence of poor physical performance significantly decreased in line with increasing 25(OH)D concentration (Figures 2 and 3). Poor performance (SPPB) was almost three-fold higher (21.6%) in Q1 (≤28 nmol/L) compared with levels ≥50 nmol/L (7.9–9.0%). In contrast, the rate of recurrently falls did not significantly differ between vitamin D deficiency (Q1) and sufficiency (Q5) at 11.6% and 9.9%, respectively (Table 3).

Factors Associated With Low SPPB In Community-Dwelling Older Adults

After adjusting for pre-defined covariates, multivariate logistic regression analysis (Table 4) supported the findings that poor physical performance (SPPB score ≤6) was significantly associated with vitamin D deficiency (<30 nmol/L) [OR 1.65 (1.31, 2.09), $p<0.001$]. Regular moderate physical activity [OR 0.30 (0.24, 0.38), $p<0.001$] was the only variable significantly associated with lower odds of poor physical performance. BMI, however, was not a statistically significant correlate of poor physical performance, unadjusted OR for obese BMI was 1.87 (1.46, 2.39, $p<0.001$), but this was no longer statistically significant [OR 1.03 (0.71, 1.25), $p=0.84$] on adjusted analysis. Other determinants of impaired SPPB included advanced age (80+ years), female gender, being retired, poor self-reported health, falls (>1) and polypharmacy.

Factors Associated With Poor Muscle Strength In Community-Dwelling Older Adults

After adjusting for pre-defined covariates, multivariate logistic regression analysis (Table 5) showed that vitamin D deficiency (<30 nmol/L) was a significant determinant of impaired HGS [OR 1.44 (1.22, 1.71) $p<0.001$]. Consistent with the SPPB findings, engaging in moderate physical activity was the only positive correlate of muscle strength identified [OR 0.65 (0.58, 0.79)]. The model showed that aged 70+, female gender, retired, fair self-reported health, falls >1, polypharmacy and overweight or obesity were significantly associated with low muscle strength.

Discussion

Vitamin D deficiency [25(OH)D <30 nmol/L] was significantly associated with both reduced muscle strength and physical performance in a large sample ($n = 4157$) of community-dwelling adults aged 60 years and over, as determined by handgrip strength and the Short Physical Performance Battery, respectively. Given that vitamin D deficiency is prevalent in older populations^{12,13} focusing on correcting deficiency at the 30 nmol/L cut-off¹¹ may be important for protecting muscle function in ageing, in addition to the established effects on preventing bone disease.²⁷ We identified impaired muscle strength and physical performance in 30.6% and 12.7%, respectively, of community-dwelling adults aged 60 and over. This is

Table I Characteristics Of The Study Population Overall And By Gender (n=4157)

	Overall (n=4157)		Males (n=1874)		Females (n=2283)	
	N (%)	SD	N (%)	SD	N (%)	SD
Socio-demographic factors						
Age , mean, SD, years	69.8	6.9	69.7	7.0	69.8	6.9
60–69	2268 (54.6)		1033 (55.1)		1235 (54.1)	
70–79	1450 (34.9)		641 (34.2)		809 (35.4)	
80+	439 (10.6)		200 (10.7)		239 (10.5)	
^a Education ≥ O level	3099 (74.9)		1483 (79.5)		1616 (71.1)	
Employment – retired	3090 (74.3)		1362 (72.7)		1728 (75.7)***	
Ethnicity (non-white)	87 (2.1)		46 (2.5)		41 (1.8)	
Health and lifestyle factors						
BMI kg/m ² , mean, SD	28.0	4.8	27.9	4.1	28.1	5.3
^b Under (<18.5)	37 (0.9)		7 (0.4)		30 (1.3)***	
Normal (≥18.5 to <25)	1085 (26.1)		432 (23.1)		653 (28.6)***	
Overweight (≥25 to <30)	1823 (43.9)		940 (50.2)		883 (38.7)***	
Obese (≥30)	1212 (29.1)		495 (26.4)		717 (31.4)***	
Physical activity						
Vigorous (>1/wk)	1739 (41.8)		923 (49.3)		816 (35.7)***	
Moderate (>1/wk)	2703 (65.0)		1283 (68.5)		1420 (62.2)***	
Mild (>1/wk)	3463 (83.3)		1450 (77.4)		2013 (88.2)***	
Current smoker	391 (9.4)		174 (9.3)		217 (9.5)	
Alcohol – 5–6 days/week	254 (6.1)		148 (7.9)		106 (4.6)***	
SR health						
Excellent	485 (11.7)		212 (11.3)		273 (11.9)	
Very good	1323 (31.8)		607 (32.4)		716 (31.4)	
Good	1365 (32.8)		617 (32.9)		748 (32.8)	
Fair	770 (18.5)		346 (18.5)		424 (18.6)	
Poor	213 (5.1)		92 (4.9)		121 (5.3)	
Long standing illness	2258 (54.3)		1007 (53.7)		1251 (54.8)	
^c Limiting	1328 (58.8)		557 (55.3)		771 (61.7)**	
Polypharmacy (≥5 meds)	1209 (29.1)		548 (29.2)		661 (28.9)	
Poor QoL (CASP <36)	772 (18.6)		378 (20.2)		394 (17.3)*	
Vitamin D related factors						
25(OH)D nmol/L, mean, SD	49.7	23.5	50.5	23.1	49.1*	23.8
Low 25(OH)D status						
<30nmol/L	908 (21.8)		359 (19.2)		549 (24.1)***	
<50nmol/L	2234 (53.7)		977 (52.1)		1257 (55.1)	
^d VitD supp use, n (%)	212 (5.1)		24 (1.3)		188 (8.2)***	
^e Sun holiday travel, n (%)	1904 (48.6)		858 (48.9)		1046 (48.4)	

Notes: Results are presented as means and standard deviation (SD) or counts and proportions, n (%). ^aO-level, approximately equivalent to secondary school. ^bBMI categories. ^cIf the long-standing illness is limiting. ^dVitamin D supplement use, with or without calcium, prescribed for osteoporosis. ^eTravel abroad within the previous 12 months, yes or no. *, **, ***Denote statistically significant differences between groups: *P<0.05, ** P<0.01, ***P<0.001.

Abbreviations: kg/m², kilograms per meter squared; SR, self-reported; CASP-19, Control, Autonomy, Self-realization and Pleasure, a four domain older-adult quality of life measure.

Table 2 HGS, SPPB And Falls In Community-Dwelling Older Adults (n = 4157)

	Overall (n=4157)	Males (n= 1874)	Females (n= 2283)
	N (%)	N (%)	N (%)
^a Low HGS %	1271 (30.6)	505 (26.9)	766 (33.6)***
SPPB poor (≤ 6)	527 (12.7)	192 (10.3)	335 (14.7)***
At-risk (7–9)	804 (19.3)	340 (18.1)	464 (20.3)***
Best (10–12)	2826 (67.9)	1342 (71.6)	1484 (65.0)
^b Falls one fall	658 (15.8)	267 (14.3)	391 (17.1)**
Multiple falls (>1)	438 (10.5)	173 (9.2)	265 (11.6)**
^c Medical attention	813 (74.2)	354 (80.5)	459 (69.9)***

Notes: ^aClassified by gender and BMI-specific cut-offs. ^bFall in the past 12 months. ^cMedical attention was sought for fall. ** Denote statistically significant differences between groups: ** $P < 0.01$, *** $P < 0.001$.

Abbreviations: HGS, handgrip strength; SPPB, short physical performance battery.

broadly in line with the prevalence of sarcopenia, estimated at 1–29%, in community-dwelling populations aged 50 years and over.³¹ The prevalence and incidence of sarcopenia is reported to increase further in hospitalized and long-term care for older adult populations.^{31,32} It should be noted, however, that the present study focused on impaired muscle function only and not specifically on sarcopenia (since muscle mass was not available in the dataset). Muscle function, however, is

considered a principal determinant of sarcopenia in recently updated definitions.²

Based on the analysis by vitamin D quintiles, older adults with the lowest serum 25(OH)D, had the highest prevalence of impaired muscle strength and performance (40.4% and 25.2%) compared with people with adequate levels (21.6% and 7.9%). This finding was further supported by the regression analysis, which showed that vitamin D deficiency (<30 nmol/L) was significantly associated with both poor physical performance and low handgrip strength. Of note, engagement in moderate physical activity was associated with lower odds of impaired muscle strength and low SPPB, which is in line with the evidence supporting physical activity as a modifiable factor in preserving muscle mass, strength and potentiating muscle protein synthesis.^{6,9} Other factors associated with low HGS and low SPPB included advanced age, female gender and variables indicative of poorer health status such as lower self-rated health, polypharmacy and recurrent falls reflecting the complex and multifaceted nature of age-related muscle impairment. Our findings identify vitamin D deficiency as a potentially modifiable lifestyle factors associated with poor muscle function and further supports the established inverse association with physical activity.^{6,10}

Falls were not significantly associated with 25(OH)D concentrations in the present study, though this was a

Figure 2 Prevalence of poor physical performance (SPPB ≤ 6) across serum 25(OH)D quintiles, n = 527. There was a significantly higher proportion of older adults with low SPPB in Q1 [25(OH)D ≤28nmol/L, vitamin D deficient] relative to Q 4 and 5 [vitamin sufficiency 25(OH)D >50nmol/L].

Figure 3 Prevalence of poor muscle strength (low HGS) across serum 25(OH)D quintiles, n= 1271. There was a significantly higher prevalence of impaired muscle strength in Q1 [25(OH)D ≤28nmol/L, vitamin D deficient] relative to quintiles 4 and 5 [vitamin D sufficiency 25(OH)D >50nmol/L].

secondary aim. The occurrence of single or multiple falls was 15.8% and 10.5%, respectively, in the past year but did not significantly differ by 25(OH)D status. A Cochrane review,²² and similarly a recent large RCT,³³ showed no effect for vitamin D supplementation on falls in the community, however, supplementation in older adult subgroups with vitamin D deficiency might protect against falls.²² Findings from a recent Cochrane review suggested that vitamin D supplementation may reduce falls rate in residential care facilities,²¹ which represent a more dependent and vulnerable group compared with community-living populations in the present study. While vitamin D²¹ and sarcopenia³ have been implicated in falls risk, it is acknowledged that falls in ageing have a complex and

multifactorial etiology involving health, sensory, medication and living environmental risk factors.^{21,22}

To date, intervention studies into the effects of vitamin D supplementation on skeletal muscle in ageing have yielded mixed findings^{8,10,18-20} though a trend towards improved lower extremity muscle performance is emerging. Lower limb function is commonly assessed by practical and clinical measures such as gait speed or the Timed-Up-and-Go test (TUG). The SPPB, applied in the present study, is a measure of physical performance of the lower extremities, acting as a comprehensive indicator of balance, gait, muscle strength and endurance.¹ We applied a restrictive score of <6 for the SPPB for muscle impairment and vitamin D deficiency of <30 nmol/L. A recent intervention study among older adults

Table 3 Low Handgrip Strength And Poor Muscle Performance According To Serum 25(OH)D Quintiles In Community-Dwelling Older Adults

25(OH)D Quintiles nmol/L	Q1 ≤28	Q2 29-41	Q3 42-54	Q4 55-69	Q5 ≥70
n, %	856 (20.6)	836 (20.1)	845 (20.3)	821 (19.8)	799 (19.2)
Low HGS %	346 (40.4)***	268 (32.1)	236 (27.9)	207 (25.2)***	214 (26.8)**
SPPB (0-12)					
Poor (≤6)	185 (21.6)***	115(13.8)	90(11.9)	65 (7.9)***	72 (9.0)***
At risk (7-9)	200 (23.4)***	171 (20.5)	149 (17.6)	146 (17.8)	138 (17.3)
High (≥10)	471 (55)***	550 (65.8)	606 (71.7)	610 (74.3)***	589 (73.7)***
^a Falls (>1)	99 (11.6)	86 (10.3)	84 (9.9)	90 (10.9)	79 (9.9)

Notes: ^aFalls, reported to have had multiple (>1) falls in past 12 months. *** **Denote statistically significant differences between groups: ** P<0.01, ***P<0.001.

Abbreviations: Q, quintiles; low HGS, low handgrip strength established by BMI and gender-specific cut-offs; SPPB, short physical performance battery.

Table 4 Logistic Regression Analysis Showing Unadjusted And Adjusted Odds Ratios (95% CI) For Poor Physical Performance (SPPB ≤ 6) In Community-Dwelling Older Adults (n=4157)

	Unadjusted Poor SPPB (≤ 6)			Adjusted Poor SPPB (≤ 6)		
	OR	(95% CI)	P	OR	(95% CI)	P
Female	1.50	(1.25, 1.82)	<0.001	1.39	(1.11, 1.74)	0.004
Age 60–69 (reference)	–	–	1	–	–	1
70–79	1.86	(1.49, 2.31)	<0.001	1.29	(0.99, 1.69)	0.050
80–89	7.38	(5.76, 9.47)	<0.001	3.78	(2.78, 5.15)	<0.001
Employment-retired	4.78	(2.94, 7.69)	<0.001	1.98	(1.18, 3.34)	0.010
Vitamin D deficiency (<30nmol/L)	2.43	(1.99, 2.95)	<0.001	1.65	(1.31, 2.09)	<0.001
^a Vitamin D supplement use	2.10	(1.56, 2.88)	<0.001	1.37	(0.91, 2.05)	0.134
Self-reported health – poor	3.25	(2.67, 3.96)	<0.001	1.55	(1.22, 1.97)	<0.001
Physical activity-moderate	0.16	(0.13, 0.19)	<0.001	0.30	(0.24, 0.38)	<0.001
^b Falls (>1)	3.78	(3.00, 4.75)	<0.001	2.55	(1.94, 3.36)	<0.001
Polypharmacy (≥ 5 medications/d)	4.90	(4.1, 5.9)	<0.001	2.42	(1.93, 3.02)	<0.001
Low HGS	5.10	(4.2, 6.1)	<0.001	2.17	(1.73, 2.72)	<0.001
Poor QoL (CASP-19 <36)	1.95	(1.58, 2.39)	<0.001	1.25	(0.97, 1.61)	0.080
BMI – obese	1.87	(1.46, 2.39)	<0.001	1.03	(0.71, 1.25)	0.84

Notes: ^aUse as prescribed treatment of osteoporosis. ^bFalls, reported as multiple (>1) falls in past 12 months. Reference category: male, age 60–69; employed, 25(OH)D >30 nmol/L, no vitamin D supplement use, excellent self-reported health; low physical activity, no falls, no polypharmacy, i.e. medications <4/d, HGS in normal range by gender and BMI classification, CASP >36, BMI in the underweight category. Adjusted analysis adjusted for age, gender, and physical activity. All models included the same variables and categorizations/groupings, for visual presentation only significant variable categories are presented.

Abbreviations: OR, odds ratio; CI, confidence interval; HGS, handgrip strength; QoL, quality of life; CASP-19, Control, Autonomy, Self-realisation, Pleasure, which is a 4 domain QoL measure for older adults.

with impaired muscle function, defined as SPPB ≤ 9 and vitamin D deficiency of 25(OH)D less than 50 nmol/L, showed that a vitamin D and protein supplement reduced

intramuscular fat and improved muscle density compared with placebo.¹⁰ Future intervention studies into the effects of reversing vitamin D deficiency at 30 nmol/L level on

Table 5 Logistic Regression Analysis Showing Unadjusted And Adjusted Odds Ratios (95% CI) For Low Handgrip Strength In Community-Dwelling Older Adults (n=4157)

	Unadjusted Low HGS			Adjusted Low HGS		
	OR	(95% CI)	P	OR	(95% CI)	P
Female	1.37	(1.19, 1.57)	<0.001	1.25	(1.08, 1.46)	0.003
70–79	2.32	(2.00, 2.69)	<0.001	1.92	(1.62, 2.26)	<0.001
80–89	7.49	(6.00, 9.35)	<0.001	4.71	(3.69, 6.02)	<0.001
Employment-retired	2.81	(2.19, 3.62)	<0.001	1.49	(1.15, 1.96)	0.003
Vitamin D deficiency (<30)	1.79	(1.54, 2.09)	<0.001	1.44	(1.21, 1.71)	<0.001
Self-reported health – poor	2.20	(1.87, 2.59)	<0.001	1.39	(1.15, 1.67)	0.001
Physical Activity – Moderate	0.41	(0.36, 0.48)	<0.001	0.68	(0.58, 0.79)	<0.001
^a Falls (>1)	2.08	(1.69, 2.54)	<0.001	1.47	(1.17, 1.86)	0.001
Polypharmacy (≥ 5 medications/d)	2.63	(2.29, 3.03)	<0.001	1.51	(1.28, 1.78)	<0.001
Poor SPPB (≤ 6 points)	5.07	(4.18, 6.15)	<0.001	2.22	(1.77, 2.77)	<0.001
Poor QoL (CASP-19, <36)	1.36	(1.16, 1.61)	<0.001	0.97	(0.81, 1.17)	0.785
BMI overweight	1.29	(1.09, 1.55)	0.003	1.31	(1.08, 1.59)	0.006
Obese	2.42	(2.02, 2.91)	<0.001	2.26	(1.84, 2.78)	<0.001

Notes: ^aFalls, reported to have had multiple (>1) falls in past 12 months. Reference category: male, age 60–69; employed, 25(OH)D >30 nmol/L, no vitamin D supplement use, excellent self-reported health; low physical activity, no falls, no polypharmacy, i.e. medications <4/d, SPPB >6, CASP >36, BMI in the underweight category. Adjusted analysis adjusted for age, gender, and physical activity. All models included the same variables and categorizations/groupings, for visual presentation only significant variable categories are presented.

Abbreviations: OR, odds ratio; CI, confidence interval; SPPB, short physical performance battery; QoL, quality of life; CASP-19, Control, Autonomy, Self-realization, Pleasure, a four domain QoL measure for older adults.

comprehensive muscle outcomes among vulnerable older adults in the community would be of interest.

In the present study, we identified that older people in the lowest quintile of vitamin D (compatible with deficiency), had the poorest muscle strength and physical performance and represent an important group to target for interventions to reverse deficiency. The recent EWGSOP updated consensus paper on sarcopenia emphasizes the public health and clinical importance of maintaining muscle function in ageing.² The threshold of <30 nmol/L for vitamin D deficiency is established for adverse effects on bone health²⁷ and may also accelerate age-related muscle decline. The recent Vitamin D Assessment (ViDA) study showed beneficial effects of vitamin D supplementation in aged 50+ adults for intermediate outcomes such as bone mineral density, but not for falls, and only in people with 25(OH)D <30 nmol/L.³³ The low usage of vitamin D supplements at 5.1%, which was prescribed for bone health in the present study, is compatible with other community-dwelling cohorts¹² and was not significantly associated with either impaired physical performance or muscle strength. This suggests that eliminating vitamin D deficiency in older populations may require other strategies such as food fortification approaches.^{34,35} Focusing interventions on older people most at risk of vitamin D deficiency (<30 nmol/L)^{12,13} seems sensible, even though such sub-groups may be “hard to reach” in society and to engage in clinical trials. These sub-groups may include, for example, the oldest old, those with poorer health status, of non-white ethnicity, living alone or living with socioeconomic disadvantage. The latter may be particularly important, given that the evidence is currently deemed inadequate to make practical recommendations for vitamin D supplementation for the prevention of age-related loss of muscle mass and function.⁶

The study had several strengths, for example, the findings are derived from a large (n = 4157) well-defined population of community-dwelling older people aged 60 years and over with comprehensive assessments and measures. Moreover, handgrip strength and a composite measure of muscle performance (SPPB) were used, both supported by the EWGSOP.¹ A number of limitations need to be mentioned, namely serum 25(OH)D in the ELSA cohort was assessed by immunoassay (DiaSorin), while the gold standard is considered LC-MS/MS. Analysis, however, was conducted in a laboratory which is a member of DEQAS, reported CVs and QCs were in the expected range and consistent with data from ELSA previously published.^{10,36,37} Furthermore, in line with the study design, our findings cannot demonstrate

cause and effect, and future intervention studies conducted in vitamin D deficient older populations would be required.

In conclusion, impaired muscle strength and physical performance were identified in 30.6% and 12.7%, respectively, of older adults, rising to 40.4% and 25.2% amongst those with the lowest vitamin D concentrations. Consistent with this, vitamin D deficiency [25(OH)D <30 nmol/L] was associated with both impaired muscle strength and impaired physical performance based on regression analysis models. It is accepted, that at a minimum, we should seek to eliminate vitamin D deficiency of ≤ 30 nmol/L, the cut-off for the prevention of bone disease, this strategy that may also protect skeletal muscle function in ageing populations.

Abbreviations

ELSA, English Longitudinal Study of Aging; HGS, hand-grip strength; SPPB, short physical performance battery.

Author Contributions

All authors contributed to data analysis, drafting and revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Funding

This work was supported by an Irish Research Council post-graduate scholarship for NA.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al. Sarcopenia: European consensus on definition and diagnosis: report of the European Working Group on Sarcopenia in Older People. *Age Ageing*. 2010;39(4):412–423. doi:10.1093/ageing/afq034
2. Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*. 2019;48(1):16–31. doi:10.1093/ageing/afy169
3. Landi F, Liperoti R, Russo A, et al. Sarcopenia as a risk factor for falls in elderly individuals: results from the iSIRENTE study. *Clin Nutr*. 2012;31(5):652–658. doi:10.1016/j.clnu.2012.02.007
4. Rizzoli R, Reginster JY, Arnal JF, et al. Quality of life in sarcopenia and frailty. *Calcif Tissue Int*. 2013;93(2):101–120. doi:10.1007/s00223-013-9758-y
5. Benjumea AM, Curcio CL, Duque G, Gomez F. Dynapenia and sarcopenia as a risk factor for disability in a falls and fractures clinic in older persons. *Open Access Maced J Med Sci*. 2018;6(2):344–349. doi:10.3889/oamjms.2018.087
6. Dent E, Morley JE, Cruz-Jentoft AJ, et al. International clinical practice guidelines for sarcopenia (ICFSR): screening, diagnosis and management. *J Nutr Health Aging*. 2018;22(10):1148–1161. doi:10.1007/s12603-018-1139-9

7. Paddon-Jones D, Leidy H. Dietary protein and muscle in older persons. *Curr Opin Clin Nutr Metab Care*. 2014;17(1):5–11. doi:10.1097/MCO.000000000000011
8. Antoniak AE, Greig CA. The effect of combined resistance exercise training and vitamin D3 supplementation on musculoskeletal health and function in older adults: a systematic review and meta-analysis. *BMJ Open*. 2017;7(7):e014619. doi:10.1136/bmjopen-2016-014619
9. Bosaeus I, Rothenberg E. Nutrition and physical activity for the prevention and treatment of age-related sarcopenia. *Proc Nutr Soc*. 2016;75(2):174–180. doi:10.1017/S002966511500422X
10. Englund DA, Kirn DR, Koochek A, et al. Nutritional supplementation with physical activity improves muscle composition in mobility-limited older adults, the VIVE2 study: a randomized, double-blind, placebo-controlled trial. *J Gerontol A Biol Sci Med Sci*. 2017;73(1):95–101. doi:10.1093/gerona/glx141
11. Roth DE, Abrams SA, Aloia J, et al. Global prevalence and disease burden of vitamin D deficiency: a roadmap for action in low- and middle-income countries. *Ann N Y Acad Sci*. 2018;1430(1):44–79. doi:10.1111/nyas.13968
12. Laird E, O'Halloran AM, Carey D, et al. The prevalence of vitamin D deficiency and the determinants of 25(OH)D concentration in older Irish adults: data from The Irish Longitudinal Study on Ageing (TILDA). *J Gerontol A Biol Sci Med Sci*. 2018;73(4):519–525. doi:10.1093/gerona/glx168
13. Aspell N, Laird E, Healy M, Shannon T, Lawlor B, O'Sullivan M. The prevalence and determinants of vitamin D status in community-dwelling older adults: results from the English Longitudinal Study of Ageing (ELSA). *Nutrients*. 2019;11(6):1253. doi:10.3390/nu11061253
14. Ceglia L. Vitamin D and skeletal muscle tissue and function. *Mol Aspects Med*. 2008;29(6):407–414. doi:10.1016/j.mam.2008.07.002
15. Dewansingh P, Melse-Boonstra A, Krijnen WP, van der Schans CP, Jager-Wittenaar H, van Den Heuvel E. Supplemental protein from dairy products increases body weight and vitamin D improves physical performance in older adults: a systematic review and meta-analysis. *Nutr Res*. 2018;49:1–22. doi:10.1016/j.nutres.2017.08.004
16. Shardell M, Semba RD, Kalyani RR, Hicks GE, Bandinelli S, Ferrucci L. Serum 25-hydroxyvitamin D, plasma klotho, and lower-extremity physical performance among older adults: findings from the InCHIANTI study. *J Gerontol A Biol Sci Med Sci*. 2015;70(9):1156–1162. doi:10.1093/gerona/glv017
17. Visser M, Deeg DJ, Lips P. Longitudinal aging study A. Low vitamin D and high parathyroid hormone levels as determinants of loss of muscle strength and muscle mass (sarcopenia): the longitudinal aging study amsterdam. *J Clin Endocrinol Metab*. 2003;88(12):5766–5772. doi:10.1210/jc.2003-030604
18. Bolland MJ, Grey A, Avenell A. Effects of vitamin D supplementation on musculoskeletal health: a systematic review, meta-analysis, and trial sequential analysis. *Lancet Diabetes Endocrinol*. 2018;6(11):847–858. doi:10.1016/S2213-8587(18)30265-1
19. Ranathunga R, Hill TR, Mathers JC, et al. No effect of monthly supplementation with 12000 IU, 24000 IU or 48000 IU vitamin D3 for one year on muscle function: the vitamin D in older people study. *J Steroid Biochem Mol Biol*. 2018.
20. Stockton KA, Mengersen K, Paratz JD, Kandiah D, Bennell KL. Effect of vitamin D supplementation on muscle strength: a systematic review and meta-analysis. *Osteoporos Int*. 2011;22(3):859–871. doi:10.1007/s00198-010-1407-y
21. Cameron ID, Dyer SM, Panagoda CE, et al. Interventions for preventing falls in older people in care facilities and hospitals. *Cochrane Database Syst Rev*. 2018;9:CD005465.
22. Gillespie LD, Robertson MC, Gillespie WJ, et al. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev*. 2012;9:CD007146.
23. Sanders KM, Stuart AL, Williamson EJ, et al. Annual high-dose oral vitamin D and falls and fractures in older women: a randomized controlled trial. *JAMA*. 2010;303(18):1815–1822. doi:10.1001/jama.2010.594
24. Sohl E, Heymans MW, de Jongh RT, et al. Prediction of vitamin D deficiency by simple patient characteristics. *Am J Clin Nutr*. 2014;99(5):1089–1095. doi:10.3945/ajcn.113.076430
25. Guralnik JM, Simonsick EM, Ferrucci L, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol*. 1994;49(2):M85–M94. doi:10.1093/geronj/49.2.m85
26. Steptoe A, Breeze E, Banks J, Nazroo J. Cohort profile: the English longitudinal study of ageing. *Int J Epidemiol*. 2013;42(6):1640–1648. doi:10.1093/ije/dys168
27. Institute of Medicine. *Dietary Reference Intakes for Calcium and Vitamin D*. Washington DC: The National Academies Press; 2011.
28. Holick MF. Sunlight, UV-radiation, vitamin D and skin cancer: how much sunlight do we need? *Adv Exp Med Biol*. 2008;624:1–15. doi:10.1007/978-0-387-77574-6_1
29. Tukey JW. *Exploratory Data Analysis*. Reading, MA: Addison-Wesley; 1977.
30. Hoaglin DC, Iglewicz B, Tukey JW. Performance of some resistant rules for outlier labeling. *J Alzheimers Dis*. 1986;81:991–999.
31. Cruz-Jentoft AJ, Landi F, Schneider SM, et al. Prevalence of and interventions for sarcopenia in ageing adults: a systematic review. Report of the International Sarcopenia Initiative (EWGSOP and IWGS). *Age Ageing*. 2014;43(6):748–759. doi:10.1093/ageing/afu115
32. Martone AM, Bianchi L, Abete P, et al. The incidence of sarcopenia among hospitalized older patients: results from the Glisten study. *J Cachexia Sarcopenia Muscle*. 2017;8(6):907–914. doi:10.1002/jcsm.12224
33. Scragg RKR. Overview of results from the Vitamin D Assessment (ViDA) study. *J Endocrinol Invest*. 2019. doi:10.1007/s40618-019-01056-z
34. Hayes A, Cashman KD. Food-based solutions for vitamin D deficiency: putting policy into practice and the key role for research. *Proc Nutr Soc*. 2017;76(1):54–63. doi:10.1017/S0029665116000756
35. Jaaskelainen T, Itkonen ST, Lundqvist A, et al. The positive impact of general vitamin D food fortification policy on vitamin D status in a representative adult Finnish population: evidence from an 11-y follow-up based on standardized 25-hydroxyvitamin D data. *Am J Clin Nutr*. 2017;105(6):1512–1520. doi:10.3945/ajcn.116.151415
36. de Oliveira C, Hirani V, Biddulph JP. Associations between vitamin D levels and depressive symptoms in later life: evidence from the English Longitudinal Study of Ageing (ELSA). *J Gerontol A Biol Sci Med Sci*. 2018;73(10):1377–1382. doi:10.1093/gerona/glx130
37. de Oliveira C, Biddulph JP, Hirani V, Schneider IJC. Vitamin D and inflammatory markers: cross-sectional analyses using data from the English Longitudinal Study of Ageing (ELSA). *J Nutr Sci*. 2017;6:e1. doi:10.1017/jns.2016.37

Clinical Interventions in Aging

Dovepress

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier

Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>