Open Access Full Text Article

CASE REPORT

Eosinophilic Pneumonia Associated With Natalizumab In A Patient With Multiple Sclerosis: A Case Report And Literature Review

This article was published in the following Dove Press journal: Therapeutics and Clinical Risk Management

Yuichiro Yasuda¹ Tatsuya Nagano ¹ Motoko Tachihara¹ Norio Chihara² Kanoko Umezawa¹ Naoko Katsurada ¹ Masatsugu Yamamoto ¹ Kenji Sekiguchi² Kazuyuki Kobayashi¹ Yoshihiro Nishimura¹

¹Division of Respiratory Medicine, Department of Internal Medicine, Kobe University Graduate School of Medicine, Kobe, Hyogo 650-0017, Japan; ²Division of Neurology, Department of Internal Medicine, Kobe University Graduate School of Medicine, Kobe, Hyogo 650-0017, Japan

Correspondence: Tatsuya Nagano Division of Respiratory Medicine, Department of Internal Medicine, Kobe University Graduate School of Medicine, 7-5-2 Kusunoki-cho, Kobe, Hyogo 650-0017, Japan Tel +81-783-82-5660 Fax +81-783-82-5661 Email tnagano@med.kobe-u.ac.jp



Abstract: We herein report the case of a 39-year-old Japanese female with eosinophilic pneumonia associated with natalizumab. The patient with bronchial asthma had multiple sclerosis and was treated using natalizumab. The patient was referred to our department because of a persistent cough. A chest computed tomography (CT) scan revealed bilateral patchy consolidation surrounded by ground-glass opacity. A bronchoalveolar lavage (BAL) was performed. Eosinophil levels in the BAL fluid were increased and the patient was consequently diagnosed as eosinophilic pneumonia associated with natalizumab. Therefore, natalizumab treatment was discontinued. Subsequent chest CT findings showed a remarkable improvement without any treatment.

Keywords: eosinophilic pneumonia, natalizumab, multiple sclerosis

Introduction

Multiple sclerosis (MS) is an inflammatory demyelinating disease of the central nervous system (CNS). MS is considered to involve genetic and environmental factors.¹ Recently, the number of MS patients in Japan has been increasing.² The clinical symptoms of MS include loss of vision, cognitive impairment, and motor and sensory disturbance. Moreover, relapse and remission of the disease frequently occur.³ Therefore, preventing recurrence is important. In Japan, five types of disease-modifying drugs such as IFNβ-1b, IFNβ-1a, fingolimod, natalizumab, and glatiramer acetate are currently available. Revealing the unknown adverse effects of these drugs is important. Therefore, we report a case of eosinophilic pneumonia (EP) associated with natalizumab, as a third-line treatment for MS and review the overall respiratory adverse events associated with natalizumab in Introduction to avoid confusion. Natalizumab is a humanized immunoglobulin G4 monoclonal antibody, which binds to the α 4 subunit of α 4 β 1 and α 4 β 7 integrins and blocks their binding to endothelial receptors, thereby inhibiting the recruitment of immune cells to inflammatory tissues. The common adverse effects of natalizumab are fatigue and allergic reactions. Moreover, there is progressive multifocal leukoencephalopathy as a serious adverse effect.^{4,5}

Case Report

A 39-year-old Japanese female underwent treatment using natalizumab for relapsing-remitting multiple sclerosis (RRMS) in the division of neurology in our hospital. The patient was referred to division of respiratory medicine with a 3month history of cough.



© 2019 Yasuda et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms. work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission for Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please esp aragraphs 4.2 and 5 of our Terms (http://www.dovepress.com/terms.php).

The patient was diagnosed with MS 9 years ago. She was treated in the neurology division since diagnosis. An initial treatment of interferon beta (IFN-B)-1b was discontinued because she was diagnosed as EP by bronchoscopy and needed steroid treatment. A second-line treatment using fingolimod was initiated 7 years ago. Treatment using natalizumab was initiated 5 months before the first visit to our department, because of poor control of MS. The patient was a never smoker, drank alcoholic beverages occasionally, and had a history of bronchial asthma. She did not have a history of keeping pets or taking new drugs including Chinese herbal medicines or supplements. There was no exposure that could trigger asthma exacerbations. Her vital signs were within a healthy range and no abnormalities were found in a physical examination. Her symptoms such as fever, difficulty breathing, night sweats, wheezing and weight loss were not observed. The results of an arterial blood gas test were also within a healthy range. A chest X-ray revealed patchy consolidation in the bilateral upper and left lower lung field (Figure 1). A chest computed tomography (CT) scan showed bilateral patchy consolidation surrounded by ground-glass opacity (Figure 2). Furthermore, laboratory tests demonstrated increased levels of white blood cells (13,500/µL), eosinophils (5220/µL), serum IgE (600.9 IU/mL), and surfactant protein D (112 ng/ mL). No elevation in tumor markers (carcinoembryonic antigen (1.0 ng/mL), soluble cytokeratin-19 fragments (0.5 ng/



Figure I A chest X-ray revealed patchy consolidation in the bilateral upper and left lower lung field (white arrow).

mL), neuron-specific enolase (15.7 ng/mL)), serum β -D-glucan, anti-neutrophil cytoplasmic antibody (ANCA), and sialylated carbohydrate antigen KL-6 (130 U/mL) was noted (Table 1). The drug-induced lymphocyte stimulation tests were not performed in this case. The results of a pulmonary functional test were normal including carbon monoxide diffusing capacity, and fractional exhaled nitric oxide was 23 ppb. The normal results of the arterial blood gas test and pulmonary functional test might suggest inflammation of the lung was relatively mild. Bronchoalveolar lavage (BAL) from the right upper lobe was performed without transbronchial lung biopsy. No bacteria, fungi, and acid-fast bacillus were detected in the BAL fluid culture. Eosinophilia was revealed in the cell fraction of the BAL fluid (61% macrophages, 35% eosinophils, 4% lymphocytes) (Figure 3). As a result, chest CT showed EP pattern. Moreover, other underlying diseases causing interstitial lung disease (ILD), such as cancer, fungal or bacterial infection, and ANCA-associated vasculitis, were excluded. Furthermore, there were no obvious abnormalities in the chest X-rays before natalizumab administration and no increase in peripheral blood eosinophils before natalizumab treatment. Therefore, natalizumab treatment was discontinued due to the possibility of EP associated with it. Peripheral blood eosinophilia and pulmonary infiltrates improved after the 6week cessation of natalizumab treatment (Figures 4 and 5). Therefore, the patient was diagnosed as EP probably associated with natalizumab. Her symptoms of multiple sclerosis remained stable for a while after the cessation of natalizumab treatment. However, methylprednisolone pulse therapy was started because multiple sclerosis was exacerbated on MRI after EP subsided. After three cycles of methylprednisolone pulse therapy, fingolimod was started.

Discussion

Recently, MS phenotypes have been classified into five distinct subtypes: RRMS, clinically isolated syndrome, radiologically isolated syndrome, primary-progressive MS, and secondary progressive MS. The majority of patients with MS follow the course of RRMS.⁶ Early treatment is effective in the relapsing-remitting phase, and it has been reported that IFN- β improves prognosis compared with placebo.⁷ However, when IFN- β cannot be used because of side effects or if it is poorly effective, a second-line treatment should be initiated. Natalizumab and fingolimod are reported to be effective as a drug after second-line treatments.^{8,9} Natalizumab is a humanized immunoglobulin G4 monoclonal antibody, which binds to the α 4 subunit of α 4 β 1 and α 4 β 7 integrins and blocks their binding to endothelial receptors. As a result, natalizumab can



Figure 2 CT scans showed bilateral patchy consolidation surrounded by ground-glass opacity (white arrow).

inhibit the influx of lymphocytes to the CNS and attenuate inflammation. It has been reported that the number of circulating lymphocytes, monocytes, eosinophils, and basophils increase during natalizumab treatment.4,5 A few cases of hypereosinophilia (≥1500/µL) have been reported as an adverse event of natalizumab treatment.¹⁰ On the other hand, it has also been reported that natalizumab could induce adverse events such as nontuberculous mycobacteria, mycobacterium tuberculosis and paracoccidioidomycosis, sarcoidosis and Crohn's disease.^{11–15} However, there has been only one case of suspected EP.16 In that case, the proportion of eosinophils in BAL was 12%, and it did not meet the criteria of EP. Hence, this was the first case with concurrent hypereosinophilia and EP as adverse event of natalizumab. The summary of cases with respiratory adverse events associated with natalizumab is shown in Table 2. However, the biologic agents are reported to have something to do with EP.^{17,18}

A diagnosis of drug-induced ILD requires correct identification of the drug, singularity of the drug, temporal eligibility, characteristic clinical symptoms, imaging results, BAL results and pathologic patterns of the reaction to the specific drug and the exclusion of other causes for the ILD.¹⁹ A diagnosis of EP requires both characteristic clinical imaging features and a demonstration of alveolar eosinophilia consisting of at least 25% of the eosinophils in the BAL.²⁰ In the current case, a chest CT scan showed lung infiltrates consistent with an EP pattern, and eosinophils in the BAL were increased. Furthermore, the patient had no other causes for the ILD such as cancer, fungal or bacterial infection, and ANCAassociated vasculitis. It is noteworthy that hypereosinophilia and lung infiltrates were improved only by the discontinuation of natalizumab, which seemed to be the causative agent of the EP. Therefore, the patient was diagnosed with EP associated with natalizumab. Although the details of the pathophysiology of EP were unknown, it was thought to be that there was an involvement of T helper 2 (Th2) cell-mediated allergic reaction. Asthma is a chronic inflammatory airway disease associated with Th2 cell cytokines, such as interleukin-4 (IL-4), IL-5, and IL-13, which promote airway eosinophilia.²¹ Moreover, Marchand E et al reported that the prevalence of asthma in patients diagnosed with idiopathic chronic EP was 64%²² It should be noted that she had a past history of IFN- β treatment-induced eosinophilia, although there is little structural similarity between IFN-B and natalizumab. Drug

Blood count			Biochemistry					
WBC	13,500	/μl	ТР	7	g/dl	PR3-ANCA	<0.1	U/mL
Neut	36.2	%	Alb	4.5	mg/dl	MPO-ANCA	<0.1	U/mL
Lymph	18.8	%	T-Bil	0.5	mg/dl	β-D-glucan	4	pg/mL
Mono	5.2	%	AST	17	IU/I	serum IgE	600.9	IU/mL
Eosino	38.7	%	ALT	22	IU/I			
RBC	471×10 ⁴	/μl	LDH	189	IU/I	Arterial blood gas test (room air)		
НЬ	13.5	g/dl	СК	77	IU/I	рH	7.424	
Ht	41.4	%	BUN	9.3	mg/dl	PaO ₂	115	mmHg
MCV	88	fl	Cre	0.48	mg/dl	PaCO ₂	36	mmHg
МСНС	32.6	%	Glu	94	mg/dl	HCO₃ [−]	23.1	mmol/l
Plt	25×10 ⁴	/μl	Na	140	mEq/l			
			к	4	mEq/l	Tumor marker		
			CI	107	mEq/l	CEA	1	ng/mL
			CRP	0.02	mg/dl	CYFRA	<0.5	ng/mL
			KL-6	130	U/mL	NSE	15.7	ng/mL
			SP-D	112	ng/mL			

Table I Laboratory Findings On The Initial Visit

hypersensitivity in the past appears to have an increased tendency to develop sensitivity to new drugs.²³ Therefore, it is possible that the patient could be susceptible to an allergic reaction to natalizumab. As for possible molecular-biological mechanism of natalizumab-induced EP, it binds to $\alpha 4\beta 1$ and

 $\alpha 4\beta 7$ integrins and could induce endothelial injury. As a result, IL-33 and vascular endothelial growth factor (VEGF) are released from injured endothelial cells, recruit eosinophil to the lung, and activate macrophage and dendritic cells, leading to T-helper cell type 2 (Th2) immune responses.

А





В

Figure 3 BAL fluid showed increased numbers of eosinophils with Papanicolaou staining (A, ×400). Under high magnification, the eosinophils were stained light green with nuclear localization (B, ×800).



Figure 4 A chest X-ray and chest CT images showed an improvement in lung infiltrates after the discontinuation of natalizumab.



Figure 5 The clinical course of the patient.

Reference	Respiratory Adverse Event	Age	Sex	Underlying Disease	Peripheral Blood Eosinophil	Eosinophil
						In BALF
11	NTM	28	F	RRMS	N/A	N/A
12	NTM	N/A	N/A	RRMS	N/A	N/A
13	ТЬ	45	F	RRMS	N/A	N/A
13	ТЬ	47	F	RRMS	N/A	N/A
14	PCM	46	м	RRMS	N/A	N/A
15	Sarcoidosis	38	м	CD	N/A	N/A
15	Sarcoidosis	37	м	CD	N/A	N/A
16	Suspected EP	41	F	RRMS	490/µL	12%
Our case	EP	39	F	RRMS	5520/µL	35%

 Table 2 Summary Of Cases With Respiratory Adverse Events Associated With Natalizumab

Abbreviations: NTM, nontuberculous mycobacteria; RRMS, relapsing-remitting multiple sclerosis; N/A, not available; Tb: mycobacterium tuberculosis; PCM, paracoccidioidomycosis; CD, crohn's disease.

T-lymphocytes further promote recruitment and activation of eosinophil in the lung via production of IL-5. Indeed, Abbas M et al reported 3 cases of hypereosinophilia in patients with multiple sclerosis treated with natalizumab and they show a Th2 cell-driven condition, suggesting that natalizumab could activate Th2 lymphocytes in genetically predisposed individuals.¹⁰

Conclusion

In conclusion, clinicians should consider the possibility of EP after the initiation of natalizumab treatment. It is important to highlight new adverse events when they are encountered.

Ethics

The Ethics Committee at Kobe University/Kobe University Graduate School of Medicine confirms that ethics approval for case report or case series are waived.

Consent For Publication

Written informed consent for publication of the case details included accompanying images was obtained from the patient.

Disclosure

All of the authors report that they have no conflicts of interest to disclose in this work.

References

- 1. Sawcer S, Franklin RJ, Ban M. Multiple sclerosis genetics. *Lancet Neurol*. 2014;13:700–709. doi:10.1016/S1474-4422(14)70041-9
- Osoegawa M, Kira J, Fukazawa T, et al. Temporal changes and geographical differences in multiple sclerosis phenotypes in Japanese: nationwide survey results over 30 years. *Mult Scler*. 2009;15:159–173. doi:10.1177/1352458508098372

- Compston A, Coles A. Multiple sclerosis. Lancet. 2008;372:1502– 1517. doi:10.1016/S0140-6736(08)61620-7
- Polman CH, O'Connor PW, Havrdova E, et al. A randomized, placebo-controlled trial of natalizumab for relapsing multiple sclerosis. *N Engl J Med.* 2006;354:899–910. doi:10.1056/NEJMoa044397
- 5. Ransohoff RM. Natalizumab for multiple sclerosis. *N Engl J Med.* 2007;356:2622–2629. doi:10.1056/NEJMct071462
- Katz Sand I. Classification, diagnosis, and differential diagnosis of multiple sclerosis. *Curr Opin Neurol*. 2015;28:193–205. doi:10.1097/ WCO.000000000000206
- 7. Goodin DS, Reder AT, Ebers GC, et al. Survival in MS: a randomized cohort study 21 years after the start of the pivotal IFNβ-1b trial. *Neurology*. 2012;78:1315–1322. doi:10.1212/ WNL.0b013e3182535cf6
- Spelman T, Kalincik T, Zhang A, et al. Comparative efficacy of switching to natalizumab in active multiple sclerosis. *Ann Clin Transl Neurol.* 2015;2:373–387. doi:10.1002/acn3.180
- Bergvall N, Makin C, Lahoz R, et al. Relapse rates in patients with multiple sclerosis switching from interferon to fingolimod or glatiramer acetate: a US claims database study. *PLoS One*. 2014;9:e88472. doi:10.1371/journal.pone.0088472
- Abbas M, Lalive PH, Chofflon M, Simon HU, Chizzolini C, Ribi C. Hypereosinophilia in patients with multiple sclerosis treated with natalizumab. *Neurology*. 2011;77:1561–1564. doi:10.1212/ WNL.0b013e318233b391
- 11. de Masson A, Maillart E, Veziris N, Meyssonnier V, Papeix C, Caumes E. Cavitary pulmonary disease in a patient treated with natalizumab. *Presse Med.* 2014;43:1009–1012. doi:10.1016/j. lpm.2013.12.012
- Hradilek P, Zeman D, Tudik I, Zapletalova O, Ulmann V. Asymptomatic lung disease caused by Mycobacterium kansasii as an opportunistic infection in a patient treated with natalizumab for relapsing-remitting multiple sclerosis. *Mult Scler.* 2014;20:639–640. doi:10.1177/1352458513501572
- Dahdaleh D, Altmann DM, Malik O, Nicholas RS. Breathlessness, night sweats, and weight loss on natalizumab. *Lancet*. 2012;380:726– 727. doi:10.1016/S0140-6736(12)61401-9
- Almeida KJ, Barreto-Soares RV, Campos-Sousa RN, Campos-Sousa MG, Bor-Seng-Shu E. Pulmonary paracoccidioidomycosis associated with the use of natalizumab in multiple sclerosis. *Mult Scler*. 2018;24:1002–1004. doi:10.1177/1352458518763091
- Parisinos CA, Lees CW, Wallace WA, Satsangi J. Sarcoidosis complicating treatment with natalizumab for Crohn's disease. *Thorax*. 2011;66:1109–1110. doi:10.1136/thx.2010.155762
- Curto E, Munteis-Olivas E, Balcells E, Domínguez-Álvarez MM. Pulmonary eosinophilia associated to treatment with natalizumab. *Ann Thorac Med.* 2016;11:224–226. doi:10.4103/1817-1737.185762

- Bartal C, Sagy I, Barski L. Drug-induced eosinophilic pneumonia: A review of 196 case reports. *Medicine (Baltimore)*. 2018;97:e9688. doi:10.1097/MD.00000000009688
- Jin F, Wang ST. Chronic eosinophilic pneumonia after trastuzumab and radiation therapy for breast cancer: a case report. *Medicine (Baltimore)*. 2019;98:e14017. doi:10.1097/ MD.000000000014017
- Camus P, Fanton A, Bonniaud P, Camus C, Foucher P. Interstitial lung disease induced by drugs and radiation. *Respiration*. 2004;71:301–326. doi:10.1159/000079633
- Cottin V. Eosinophilic lung diseases. Clin Chest Med. 2016;37:535– 556. doi:10.1016/j.ccm.2016.04.015
- Lambrecht BN, Hammad H, Fahy JV. The cytokines of asthma. Immunity. 2019;50:975–991. doi:10.1016/j.immuni.2019.03.018
- 22. Marchand E, Etienne-Mastroianni B, Chanez P, Lauque D, Leclerc P, Cordier JF. Idiopathic chronic eosinophilic pneumonia and asthma: how do they influence each other? *Eur Respir J.* 2003;22:8–13. doi:10.1183/09031936.03.00085603
- Leslie C, Paul A. Patterson's Allergic Diseases. 7th ed. Wolters Kluwer Health;2009: 239–314.

Therapeutics and Clinical Risk Management

Dovepress

Publish your work in this journal

Therapeutics and Clinical Risk Management is an international, peerreviewed journal of clinical therapeutics and risk management, focusing on concise rapid reporting of clinical studies in all therapeutic areas, outcomes, safety, and programs for the effective, safe, and sustained use of medicines. This journal is indexed on PubMed Central, CAS, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/therapeutics-and-clinical-risk-management-journal