

Association of *IFNL3* rs12979860 polymorphism with HCV-related hepatocellular carcinoma susceptibility in a Chinese population

This article was published in the following Dove Press journal:
Clinical and Experimental Gastroenterology

Wei Hou^{1,2}
Kunyan Qiao¹
Zhixiao Huo¹
Yanan Du³
Cindy Wang²
Wing-Kin Syn^{2,4}

¹Tianjin Second People's Hospital and Tianjin Institute of Hepatology, Tianjin, People's Republic of China; ²Division of Gastroenterology and Hepatology, Department of Medicine, Medical University of South Carolina, Charleston, SC, USA; ³Department of Biomedical Engineering, School of Medicine, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, Tsinghua University, Beijing, People's Republic of China; ⁴Section of Gastroenterology, Ralph H Johnson Veterans Affairs Medical Center, Charleston, SC, USA

Background: The association between interferon lambda-3 (*IFNL3*, also known as interleukin 28B, *IL28B*) rs12979860 polymorphism and the development of hepatocellular carcinoma (HCC) has been investigated in recent studies with inconclusive and inconsistent results. *IFNL3* rs12979860 polymorphism has been shown a marked differential distribution with regional and ethnic variation. Whether this single nucleotide polymorphism influences susceptibility to hepatitis C virus (HCV)-related HCC remains elusive.

Methods: In this case-control study, a total of 157 Chinese Han patients with chronic HCV infection were enrolled, including 62 HCV-related HCC patients and 95 chronic hepatitis C (CHC) patients without HCC, and the genetic polymorphism of *IFNL3* rs12979860 was genotyped via a DNA microarray-based assay. The logistic regression analysis was employed to determine the correlation between the genetic polymorphism and risk of HCV-related HCC.

Results: A higher proportion of CT/TT genotype and T allele was observed in HCC patients compared to the CHC group. Under the genetic model of allele frequency, the T allele was associated with elevated risk of HCV-related HCC in the Chinese population compared to C allele after an adjustment for age, gender, body mass index, HCV infection duration, and HCV genotypes ($P=0.046$). In the subgroup analysis stratified by HCV genotype, subjects with CHC genotype 1b infection carrying rs12979860 T allele and CT+TT genotype had higher susceptibility to HCC than those with C allele and CC genotype ($P=0.020$, $P=0.037$, respectively).

Conclusion: *IFNL3* rs12979860 polymorphism with T allele could be a factor that increases the risk of HCV-related HCC in the Chinese population, especially those subjects with CHC genotype 1b infection.

Keywords: *IFNL3*, rs12979860, polymorphism, HCV, HCC

Introduction

Hepatitis C virus (HCV) infection is a major public health problem throughout the world. Chronic HCV infection may eventually progress to liver cirrhosis (LC) or even develop to hepatocellular carcinoma (HCC).^{1,2} The risk of development from chronic hepatitis C (CHC) to HCC involves a complex interplay between the viral and host genetic factors.³ Our previous study⁴ together with others' have revealed that host gene polymorphisms may play important roles in the development and progression of HCV-related HCC.

IFNL3 encodes IFN lambda-3 (IFN- λ_3), also known as interleukin 28B, which belongs to the type III IFN- λ family consisting of IL29/IFNL1, IL28A/IFNL2, IL28B/IFNL3, and newly discovered IFNL4.⁵⁻⁷ It has been investigated that

Correspondence: Wei Hou
Division of Gastroenterology and
Hepatology, Department of Medicine,
Medical University of South Carolina,
Strom Thurmond Building, 30 Courtenay
Drive, Charleston, SC 29425, USA
Email houweicn@163.com

Wing-Kin Syn
Section of Gastroenterology, Ralph H
Johnson Veterans Affairs Medical Center,
109 Bee Street, Charleston, SC 29401, USA
Email synw@muscc.edu

rs12979860 (C/T), located near the *IFNL3* gene, involves in a number of aspects of HCV infection and disease progress,^{8–19} including response to therapy, natural elimination of the virus, viral clearance rate, changes in gene expression and lipid metabolism, hepatocyte death rate, inflammatory activity, fibrosis risk, cirrhosis and hepatocarcinogenesis.

Several studies have been recently carried out to explore the association between *IFNL3* rs12979860 polymorphism and HCC risk in different geographic regions and ethnic populations, but with inconclusive and inconsistent results (Table 1).^{20–36} Some reports demonstrated that the carriage of T allele was a risk factor for HCC development.^{20–29} In contrast, one study showed opposite results,³⁰ while other studies did not find any significant association between *IFNL3* rs12979860 polymorphism and HCC risk.^{31–36} Therefore, we sought to conduct a case-control study to investigate the role of genetic polymorphism of *IFNL3* rs12979860 in the susceptibility to HCV-related HCC in a Chinese population.

Materials and methods

Patients

A total of 157 patients with chronic HCV infection, including 62 HCV-related HCC patients, and 95 HCV-infected patients without HCC as the control subjects, were recruited from Tianjin Second People's Hospital and Tianjin institute of Hepatology from July 2015 to

February 2017. The diagnostic criteria for CHC with or without HCC were based on the combination of clinical history, physical examination, imaging, and laboratory data, and/or histology. Seropositive patients for other hepatitis viruses such as hepatitis B virus or human immunodeficiency virus were excluded from this study. The written informed consent and the ethical approval for this study were obtained from all subjects and the Faculty of Health Science Ethics Committee of Tianjin Second People's Hospital. The following data were collected from all HCC patients and control subjects including gender, age, body mass index (BMI), the known duration of HCV infection, and infected HCV genotype.

Genotyping

Genotyping for rs12979860 polymorphism was performed via a DNA microarray-based assay, as described in our previous study.³⁷

Statistical analysis

All analyses were performed using the SPSS 17.0 software (SPSS Inc., Chicago, IL, USA). Chi-square test and Student *t*-test were used where appropriate. The logistic regression analysis was employed to analyze the correlation between the genetic polymorphism and risk of HCC. A *P*-value of less than 0.05 was considered to indicate a significant difference.

Table 1 Characteristics of previous studies on the association of *IFNL3* rs12979860 polymorphism and HCC risk

Study	Year	HCV genotype	Risk factor associated with HCC	Reference
Buivydiene et al.	2018	GT1 GT2 GT3	rs12979860 CC genotype	30
Suo et al.	2013	ND	rs12979860 TT genotype	20
Attallah et al.	2018	GT4	rs12979860 TT genotype	21
Chang et al.	2015	GT1 non-GT1	rs12979860 CT+TT genotype	22
Lee et al.	2015	GT1 non-GT1	rs12979860 CT+TT genotype	23
Ibrahim et al.	2016	ND	rs12979860 CT+TT genotype	24
Chang et al.	2018	GT1 non-GT1	rs12979860 CT+TT genotype	25
Fabris et al.	2011	ND	rs12979860 T allele	26
El-Awady et al.	2012	GT4a	rs12979860 T allele	27
Eurich et al.	2012	GT1b non-GT1b	rs12979860 T allele	28
Zhang et al.	2016	ND	rs12979860 T allele	29
Agúndez et al.	2012	GT1 non-GT1	No association	31
Bochud et al.	2012	GT1 GT2 GT3 GT4	No association	32
Joshita et al.	2012	GT1 GT2	No association	33
Miura et al.	2012	GT1b	No association	34
Akkiz et al.	2014	ND	No association	35
Zekri et al.	2014	ND	No association	36

Abbreviations: *IFNL3*, interferon lambda-3; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; GT, genotype; ND, not determined.

Results

Demographic and clinical characteristics of the study population

The demographic and clinical characteristics of HCV-related HCC patients and control subjects are reported in Table 2. No significant differences were found between the two groups in terms of gender, age, BMI, duration of HCV infection, and HCV genotype distribution ($P>0.05$, all).

Genotype and allele frequencies of *IFNL3* rs12979860

The genotype and allele frequencies of *IFNL3* rs12979860 polymorphism are shown in Figure 1. Three genotypes were detected in rs12979860, CC, CT, and TT. The frequencies of these three genotypes were 91.94%, 4.84%, and 3.22% in HCC group (Figure 1A), 97.90%, 1.05%, and 1.05% in control group (Figure 1A). The C and T allele frequencies of the two groups were 94.35% and 5.65%, 98.42% and 1.58%, respectively (Figure 1C). Though the distribution of *IFNL3* rs12979860 genotypes and alleles was not statistically different between HCC patients and control subjects ($P>0.05$), a higher proportion of CT/TT genotype (Figure 1B) and T allele (Figure 1D) was observed in HCC patients.

Table 2 Demographic and clinical characteristics of the study population

Characteristics	HCC patients N=62 (%)	CHC patients N=95 (%)	P-value
Gender			0.467
Male	35(56.45%)	48(50.53%)	
Female	27(43.55%)	47(49.47%)	
Age (years)			0.118
≤60	30(48.39%)	58(61.05%)	
>60	32(52.61%)	37(38.95%)	
BMI			0.329
<24	33(53.23%)	43(45.26%)	
≥24	29(46.77%)	52(54.74%)	
Duration of HCV infection (years)			0.825
≤20	25(40.32%)	40(42.11%)	
>20	37(59.68%)	55(57.89%)	
HCV genotype			0.416
1b	51(82.26%)	73(76.84%)	
Non-1b	11(17.74%)	22(23.16%)	

Abbreviations: HCC, hepatocellular carcinoma; CHC, chronic hepatitis C; BMI, body mass index; HCV, hepatitis C virus.

Association analysis of HCV-related HCC risk

We further analyzed the correlation between *IFNL3* rs12979860 polymorphism and susceptibility to HCV-related HCC using logistic regression analysis according to four models (i.e., allele frequency, general genotype, dominant, and recessive models). The OR and P -values of all the genetic models were adjusted on age, gender, BMI, HCV infection duration, and HCV genotypes. Under the genetic model of allele frequency (Figure 2A), the T allele of rs12979860 was associated with elevated risk of HCC compared to the C allele (OR=4.166, 95% CI: 1.024–16.959; $P=0.046$). Under the dominant model (Figure 2C), the CT+TT genotype increased the risk of developing HCC with a tendency toward statistical significance (OR=4.643, 95% CI: 0.826–26.098; $P=0.081$). No statistical significances were found under genetic model of general genotype (Figure 2B) as well as recessive model (Figure 2D).

Furthermore, to explore whether HCV genotype would play a role in the association, we separated all the subjects of this study into HCV genotype 1b and non-1b groups (Figure 3). We found that rs12979860 T allele and CT+TT genotype were associated with elevated risk of HCC compared to C allele and CC genotype in HCV 1b genotype group ($P=0.020$; $P=0.037$, respectively).

Discussion

In this case-control study, we attempted to investigate the association between *IFNL3* rs12979860 polymorphism and the HCV-related HCC in a Chinese population. As a result, we found a higher proportion of CT/TT genotype and T allele in HCC patients compared to the CHC group. The T allele was associated with elevated risk of HCV-related HCC compared to C allele under the genetic model of allele frequency. In the subgroup analysis stratified by HCV genotype, subjects with CHC genotype 1b infection carrying rs12979860 T allele and CT+TT genotype had higher susceptibility to HCC than those with C allele and CC genotype.

Our data add new insights into the association between genetic polymorphism of *IFNL3* rs12979860 and the susceptibility to HCV-related HCC, and provide further evidence in a Chinese population. Our findings are consistent with the previous observations,^{20–29} whereas different from some other reports.^{30–36} The following factors might account for the inconsistent results: 1) geographic and racial differences, 2) diversity of viral genotypes.




Figure 1 Genotype and allele frequencies of *IFNL3* rs12979860 in CHC and HCC groups.

Notes: (A) Genotype frequencies of *IFNL3* rs12979860 in CHC and HCC groups; (B) comparison of genotype proportions between CHC group and HCC group; (C) allele frequencies of *IFNL3* rs12979860 in CHC and HCC groups; (D) comparison of allele proportions between CHC group and HCC group.

Abbreviations: CHC, chronic hepatitis C; HCC, hepatocellular carcinoma; *IFNL3*, interferon lambda-3.

IFNL3 encodes IFN- λ 3, one of the type III IFN- λ family member, inducing signaling through binding to the specific IFN- λ receptor chain 1 and the shared IL-10 receptor chain 2. Accumulating evidence strongly suggests

that IFN- λ plays a major role in the control of viral infection and antitumor activity,³⁸⁻⁴⁰ including activation immune cells and suppression of HCC cell proliferation and growth in HCC models.^{41,42} The allelic variants of the




Figure 2 Association between *IFNL3* rs12979860 polymorphism and HCC risk under different genetic models.

Notes: (A) Genetic model of allele frequency; (B) genetic model of general genotype; (C) dominant model; (D) recessive model.

Abbreviations: HCC, hepatocellular carcinoma; CHC, chronic hepatitis C; *IFNL3*, interferon lambda-3.




Figure 3 Association between *IFNL3* rs12979860 polymorphism and HCC risk stratified by genotype.

Notes: (A) HCV genotype 1b group; (B) HCV genotype non-1b group.

Abbreviations: HCC, hepatocellular carcinoma; CHC, chronic hepatitis C; *IFNL3*, interferon lambda-3; HCV, hepatitis C virus.

IFNL3 polymorphism may affect the efficiency of the immunomodulatory process, which could lead to HCC. The exact molecular mechanisms by which *IFNL3* rs12979860 polymorphism influence susceptibility to HCV-related HCC warrant further elucidation.

There are also some potential limitations in the study, principally limited to a relatively small sample size in a Chinese population at a single center. Further studies involving large-scale samples from different centers should be performed.

Conclusion

In summary, our results indicate that *IFNL3* rs12979860 T allele could be a factor that increases the risk of HCV-related HCC, especially those subjects with CHC genotype 1b infection in the Chinese population. Screening genetic polymorphism of *IFNL3* rs12979860 might be helpful in designing effective and efficient HCC surveillance programs for chronic HCV-infected patients.

Abbreviations

IFNL3, interferon lambda-3; IL28B, interleukin 28B; HCC, hepatocellular carcinoma; SNP, single nucleotide polymorphism; CHC, chronic hepatitis C; HCV, hepatitis C

virus; LC, liver cirrhosis; *IFN-λ3*, *IFN* lambda-3; SNP, single nucleotide polymorphism; HBV, hepatitis B virus; HIV, human immunodeficiency virus; BMI, body mass index.

Availability of data and materials

All data generated or analyzed during this study are included in this article.

Ethics approval and consent to participate

Written informed consent was obtained from all participants included in the present study. Ethical approval to carry out the study was obtained from the Ethics Committee of Tianjin Second People's Hospital. This study was conducted in accordance with the Declaration of Helsinki.

Acknowledgments

We would like to thank all participants for their participation in this study. This work was presented in the 27th Annual Conference of APASL, March 14–18, 2018, New Delhi, India and was published in abstract form in *Hepatol Int* (2018) 12 (Suppl 2): S388, HCC-24. This work was supported by the National Natural Science Foundation of

China (grant nos. 30800974 and 81271845) and Tianjin Municipal Health Bureau of Science and Technology Fund (grant nos. 2012KR02 and 12KG118). Resources were provided by the Ralph H Johnson VAMC, Charleston, South Carolina.

Author contributions

WH and WKS conceived and designed the study. All authors contributed to data analysis, drafting or revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- Pradat P, Virlogeux V, Trépo E. Epidemiology and elimination of HCV-related liver disease. *Viruses*. 2018;10(10):pii: E545. doi:10.3390/v10100545
- Axley P, Ahmed Z, Ravi S, Singal AK. Hepatitis C virus and hepatocellular carcinoma: a narrative review. *J Clin Transl Hepatol*. 2018;6(1):79–84. doi:10.14218/JCTH.2017.00067
- Matsuura K, Tanaka Y. Host genetic variations associated with disease progression in chronic hepatitis C virus infection. *Hepatol Res*. 2018;48(2):127–133. doi:10.1111/hepr.13042
- Zhang S, Qiao K, Trieu C, et al. Genetic polymorphism of epidermal growth factor rs4444903 influences susceptibility to HCV-related liver cirrhosis and hepatocellular carcinoma in a Chinese Han population. *Clin Lab*. 2017;63(4):845–850. doi:10.7754/Clin.Lab.2016.161203
- Prokunina-Olsson L, Muchmore B, Tang W, et al. A variant upstream of IFNL3 (IL28B) creating a new interferon gene IFNL4 is associated with impaired clearance of hepatitis C virus. *Nat Genet*. 2013;45(2):164–171. doi:10.1038/ng.2521
- Bibert S, Roger T, Calandra T, et al.; Swiss Hepatitis C Cohort Study. IL28B expression depends on a novel TT/G polymorphism which improves HCV clearance prediction. *J Exp Med*. 2013;210(6):1109–1116. doi:10.1084/jem.20130012
- Terczyńska-Dyla E, Bibert S, Duong FH, et al.; Swiss Hepatitis C Cohort Study Group. Reduced IFN λ 4 activity is associated with improved HCV clearance and reduced expression of interferon-stimulated genes. *Nat Commun*. 2014;5:5699. doi:10.1038/ncomms5972
- Ge D, Fellay J, Thompson AJ, et al. Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature*. 2009;461(7262):399–401. doi:10.1038/nature08309
- Thomas DL, Thio CL, Martin MP, et al. Genetic variation in IL28B and spontaneous clearance of hepatitis C virus. *Nature*. 2009;461(7265):798–801. doi:10.1038/nature08463
- Tanaka Y, Nishida N, Sugiyama M, et al. Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nat Genet*. 2009;41(10):1105–1109. doi:10.1038/ng.449
- Suppiah V, Moldovan M, Ahlenstiel G, et al. IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nat Genet*. 2009;41(10):1100–1104. doi:10.1038/ng.447
- Rauch A, Kutalik Z, Descombes P, et al.; Swiss Hepatitis C Cohort Study; Swiss HIV Cohort Study. Genetic variation in IL28B is associated with chronic hepatitis C and treatment failure: a genome-wide association study. *Gastroenterology*. 2010;138(4):1338–45, 1345.e1–7. doi:10.1053/j.gastro.2009.12.056
- Balogopal A, Thomas DL, Thio CL. IL28B and the control of hepatitis C virus infection. *Gastroenterology*. 2010;139(6):1865–1876. doi:10.1053/j.gastro.2010.10.004
- Hofmann WP, Zeuzem S. A new standard of care for the treatment of chronic HCV infection. *Nat Rev Gastroenterol Hepatol*. 2011;8(5):257–264. doi:10.1038/nrgastro.2011.49
- Hayes CN, Imamura M, Aikata H, Chayama K. Genetics of IL28B and HCV-response to infection and treatment. *Nat Rev Gastroenterol Hepatol*. 2012;9(7):406–417. doi:10.1038/nrgastro.2012.101
- Eslam M, George J. Genome-wide association studies and hepatitis C: harvesting the benefits of the genomic revolution. *Semin Liver Dis*. 2015;35(4):402–420. doi:10.1055/s-0035-1567830
- Eslam M, McLeod D, Kelaeng KS, et al.; International Liver Disease Genetics Consortium (ILDGC). IFN- λ 3, not IFN- λ 4, likely mediates IFNL3-IFNL4 haplotype-dependent hepatic inflammation and fibrosis. *Nat Genet*. 2017;49(5):795–800. doi:10.1038/ng.3836
- Chinnaswamy S. Gene-disease association with human IFNL locus polymorphisms extends beyond hepatitis C virus infections. *Genes Immun*. 2016;17(5):265–275. doi:10.1038/gene.2016.24
- Heim MH, Bochud PY, George J. Host - hepatitis C viral interactions: the role of genetics. *J Hepatol*. 2016;65(1 Suppl):S22–S32. doi:10.1016/j.jhep.2016.07.037
- Suo GJ, Zhao ZX. Association of the interleukin-28B gene polymorphism with development of hepatitis virus-related hepatocellular carcinoma and liver cirrhosis: a meta-analysis. *Genet Mol Res*. 2013;12(3):3708–3717. doi:10.4238/2013.September.19.1
- Attallah AM, Omran D, Marie MS, et al. IL-28B rs12979860 polymorphism affect the course of chronic hepatitis and the development of HCC in Egyptian patients with hepatitis C type 4. *Br J Biomed Sci*. 2018;75(4):157–162. doi:10.1080/09674845.2018.1489599
- Chang KC, Tseng PL, Wu YY, et al. A polymorphism in interferon L3 is an independent risk factor for development of hepatocellular carcinoma after treatment of hepatitis C virus infection. *Clin Gastroenterol Hepatol*. 2015;13(5):1017–1024. doi:10.1016/j.cgh.2014.10.035
- Lee MH, Yang HI, Lu SN, et al. Polymorphisms near the IFNL3 gene associated with HCV RNA spontaneous clearance and hepatocellular carcinoma risk. *Sci Rep*. 2015;5:17030. doi:10.1038/srep17030
- Ibrahim MK, Salama H, Abd El Rahman M, et al. Three gene signature for predicting the development of hepatocellular carcinoma in chronically infected hepatitis C virus patients. *J Interferon Cytokine Res*. 2016;36(12):698–705. doi:10.1089/jir.2016.0042
- Chang KC, Ye YH, Wu CK, et al. Risk factors for development of hepatocellular carcinoma in patients with chronic hepatitis C without sustained response to combination therapy. *J Formos Med Assoc*. 2018;117(11):1011–1018. doi:10.1016/j.jfma.2017.11.008
- Fabris C, Falletti E, Cussigh A, et al. IL-28B rs12979860 C/T allele distribution in patients with liver cirrhosis: role in the course of chronic viral hepatitis and the development of HCC. *J Hepatol*. 2011;54(4):716–722. doi:10.1016/j.jhep.2010.07.019
- El-Awady MK, Mostafa L, Tabll AA, et al. Association of IL28B SNP with progression of Egyptian HCV genotype 4 patients to end stage liver disease. *Hepat Mon*. 2012;12(4):271–277. doi:10.5812/hepatmon.835
- Eurich D, Boas-Knoop S, Bahra M, et al. Role of IL28B polymorphism in the development of hepatitis C virus-induced hepatocellular carcinoma, graft fibrosis, and posttransplant antiviral therapy. *Transplantation*. 2012;93(6):644–649. doi:10.1097/TP.0b013e318244f774
- Zhang Y, Zhu SL, Chen J, Li LQ. Meta-analysis of associations of interleukin-28B polymorphisms rs8099917 and rs12979860 with development of hepatitis virus-related hepatocellular carcinoma. *Onco Targets Ther*. 2016;9:3249–3257. doi:10.2147/OTT.S104904
- Buiydyiene A, Liakina V, Kashuba E, et al. Impact of the Uridine-Cytidine kinase like-1 protein and IL28B rs12979860 and rs8099917 SNPs on the development of hepatocellular carcinoma in cirrhotic chronic hepatitis C patients – a pilot study. *Medicina (Kaunas)*. 2018;54(5):pii: E67. doi:10.3390/medicina54050067

31. Agúndez JA, García-Martin E, Maestro ML, et al. Relation of IL28B gene polymorphism with biochemical and histological features in hepatitis C virus-induced liver disease. *PLoS One*. 2012;7(5):e37998. doi:10.1371/journal.pone.0037998
32. Bochud PY, Bibert S, Kotalik Z, et al.; Swiss Hepatitis C Cohort Study Group; ANRS HC EP 26 Genoscan Study Group. IL28B alleles associated with poor hepatitis C virus (HCV) clearance protect against inflammation and fibrosis in patients infected with non-1 HCV genotypes. *Hepatology*. 2012;55(2):384–394. doi:10.1002/hep.24678
33. Joshita S, Umemura T, Katsuyama Y, et al. Association of IL28B gene polymorphism with development of hepatocellular carcinoma in Japanese patients with chronic hepatitis C virus infection. *Hum Immunol*. 2012;73(3):298–300. doi:10.1016/j.humimm.2011.12.021
34. Miura M, Maekawa S, Kadokura M, et al. Analysis of viral amino acids sequences and the IL28B SNP influencing the development of hepatocellular carcinoma in chronic hepatitis C. *Hepatol Int*. 2012;6(1):386–396. doi:10.1007/s12072-011-9307-6
35. Akkiz H, Kuran S, Akgöllü E, Uskudar O, Bekar A, Bayram S. The role of interleukin 28B gene polymorphism in Turkish patients with hepatocellular carcinoma. *Ann Hepatol*. 2014;13(6):788–795.
36. Zekri AR, Salama H, Medhat E, et al. IL28B rs12979860 gene polymorphism in Egyptian patients with chronic liver disease infected with HCV. *Asian Pac J Cancer Prev*. 2014;15(17):7213–7218. doi:10.7314/apjcp.2014.15.17.7213
37. Qiao K, Trieu C, Huo Z, Du Y, Hou W. Distribution of IL28B rs12979860 and rs8099917 genotypes in patients with chronic hepatitis C virus infection in Tianjin, China. *Clin Lab*. 2018;64(4):543–550. doi:10.7754/Clin.Lab.2017.171036
38. Wack A, Terczyńska-Dyla E, Hartmann R. Guarding the frontiers: the biology of type III interferons. *Nat Immunol*. 2015;16(8):802–809. doi:10.1038/ni.3212
39. Lasfar A, Zloza A, de la Torre A, Cohen-Solal KA. IFN-λ: a new inducer of local immunity against cancer and infections. *Front Immunol*. 2016;7:598. eCollection 2016. doi:10.3389/fimmu.2016.00598
40. Lasfar A, Zloza A, Silk AW, Lee LY, Cohen-Solal KA. Interferon lambda: toward a dual role in cancer. *J Interferon Cytokine Res*. 2019;39(1):22–29. doi:10.1089/jir.2018.0046
41. Abushahba W, Balan M, Castaneda I, et al. Antitumor activity of type I and type III interferons in BNL hepatoma model. *Cancer Immunol Immunother*. 2010;59(7):1059–1071. doi:10.1007/s00262-010-0831-3
42. Yan Y, Wang L, He J, et al. Synergy with interferon-lambda 3 and sorafenib suppresses hepatocellular carcinoma proliferation. *Biomed Pharmacother*. 2017;88:395–402. doi:10.1016/j.biopha.2017.01.077

Clinical and Experimental Gastroenterology

Dovepress

Publish your work in this journal

Clinical and Experimental Gastroenterology is an international, peer-reviewed, open access, online journal publishing original research, reports, editorials, reviews and commentaries on all aspects of gastroenterology in the clinic and laboratory. This journal is indexed on American Chemical Society's Chemical Abstracts Service (CAS).

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-and-experimental-gastroenterology-journal>