Prevalence And Associated Factors Of Visual Impairment Among School-Age Children In Bahir Dar City, Northwest Ethiopia

This article was published in the following Dove Press journal:
Clinical Optometry

Yosef Antehun Merrie¹
Mebratu Mulusew Tegegne²
Minychil Bantihun Munaw²
Haile Woretaw Alemu²

¹Felege Hiwot Referral Hospital, Bahir Dar, Ethiopia; ²Department of Optometry, School of Medicine, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia

Introduction: Visual impairment (VI) is one of the major public health problems in the world. It is highly prevalent among children in sub-Saharan countries, including Ethiopia. Worldwide, the magnitude of VI among school-age children is 1%–10%. However, there was limited information regarding the prevalence and associated factors of VI among school-age children in the study area, which is essential to plan and implement appropriate interventions.

Objective: The aim of this study was to determine the prevalence and associated factors of VI among school-age children living in Bahir Dar city, northwest Ethiopia.

Methods: A community-based cross-sectional study was done on a sample of 632 school-age children selected by multistage sampling in Bahir Dar from April 30 to May 15, 2018. Data were collected through interviews and physical examinations. Face-to-face interviews were done with a pretested semistructured questionnaire. Physical examinations were done with visual acuity measures and assessment of ocular pathology by optometrists. Data were entered into Epi Info 7 and exported to and analyzed with SPSS 20. Binary logistic regression was fitted, and variables with $P < 0.05$ in the multivariate model were considered statistically significant.

Results: A total of 601 study subjects were included in this study, giving a response rate of 95.2%. The median age was 13 (IQR 11–16) years, and 303 (50.3%) were male. Prevalence of VI was 52 (8.7%, 95% CI 6.2%–10.7%). In multivariate analysis, prematurity [AOR 2.8 (95% CI 1.19–6.83)], admission to a neonatal intensive-care unit (AOR 5.5, 95% CI 2.01–15.15), having a parent with VI (AOR 1.8, 95% CI 0.13–0.97), watching television from < 2 m (AOR 8.7, 95% CI 1.49–18.24), and mobile-phone exposure > 4 hours per day (AOR 1.6, 95% CI 1.32–4.45) were factors significantly associated with VI.

Conclusion: The prevalence of VI among school-age children in Bahir Dar was significant. Premature birth, admission to a neonatal intensive-care unit, having a parent with VI, watching television from < 2 m, and mobile exposure > 4 hours per day were significantly associated.

Keywords: school-age children, northwest Ethiopia, visual impairment

Introduction

Visual impairment (VI) is a significant loss of vision, clinically defined as presenting distance visual acuity $< 6/18$ in the better eye. It can be classified as mild, moderate, severe, and blindness 1–5.^{1,2} Of 285 million visually impaired people worldwide,³ 18 million are younger than 15 years and a child goes blind every minute.⁴ The prevalence of VI is higher in developing countries and has been estimated to be 5.3% in Ethiopia.^{5,6} Worldwide, the magnitude of VI among school-age children is

Correspondence: Mebratu Mulusew Tegegne
Department of Optometry, School of Medicine, College of Medicine and Health Sciences, University of Gondar, Gondar, Ethiopia PO Box 196
Email mebratieophta@gmail.com

1%–10%.^{2,7–15} In reality, 75%–90% of all learning activity is done with vision in the classroom, and thus VI drastically affects the academic performance and social activity of children.⁷ The effects of VI in children further extend to the well-being of individuals and families and social welfare. Loss of productivity and payment for treatment and visual aids has human and socioeconomic consequences, which in turn perpetuate ill health, leading to death.¹⁰ If early diagnosis and treatment are made, the majority of VI can be corrected easily.^{11,12} Low birth weight,¹³ white ethnicity, premature birth, and admission to a neonatal intensive-care unit (NICU),¹⁴ large head circumference, presence of congenital abnormalities and cesarean-section delivery,^{15–17} maternal alcohol consumption, families with lower parental education, and positive family history of VI,^{18,19} being female, age range of 10–13 years, learning in public schools, watching television from <2 m,^{8,9} duration of television exposure, distance of television exposure, mobile-phone exposure, medical visit history,^{2,5,7} and lower family monthly income²⁰ have been found to be positively associated with VI.

Although VI and its associated factors have well studied in developed countries, there are limited data for Ethiopia, and none for northwest Ethiopia. Identifying risk factors and estimating the prevalence of VI among school-age children are important for the establishment and implementation of prevention strategies and programs. More action needs to be undertaken in preventing VI, because ocular problems may cause permanent blindness and child mortality.^{8,15–17,21} The prevalence of VI among school-age children in Ethiopia is not well established, and there have been limited surveys done at the community level to identify factors associated with VI in school-age children. This study determines the prevalence of VI and identifies factors associated with VI among school-age children in Bahir Dar city, which could be used as baseline data to facilitate health-care planning and implementation.

Methods

A community based cross sectional study was conducted in Bahir Dar city, northwest Ethiopia from April 30 to May 15, 2018. Bahir Dar is located 578 km from the capital city, Addis Ababa, with a population of about 243,300.²² It has six subcities and 17 *kebele*, hosting approximately 53,725 households and 100,984 children <19 years old.²³ There are three governmental hospitals, three private general hospitals, and two private eye clinics that provide eye-care services.²²

All children aged 6–18 years old who had been living in Bahir Dar > 6 months, except those with recent ocular trauma or surgery, were included in the study.

Sample size was determined with a single population-proportion formula: $n = \frac{(Z_{\alpha/2})^2 P(1-P)}{d^2}$

where Z had a 95% CI of 1.96, P the proportion of VI (7.24%) from a similar study in Adama, Ethiopia of school-age children (0.0724),⁹ and d maximum tolerable error (marginal error) of 3% = 0.03). By adding a 10% nonresponse rate and using a design effect of 2, the final sample size was 632. Multistage sampling using two sampling stages process was used. First, four of 17 *kebele* were selected using simple random sampling from a census list obtained from the Bahir Dar City Statistical Agency. In four selected *kebele*, there was a total population of 44,438 and total households of 12,015.²³ Then, systematic random sampling was used to select participant households proportionally with a sampling fraction of 19 to get one child. If there were more than one child in the selected house, one child was selected randomly, and if the selected child was not present, the selected house was revisited. If the selected house had no school-age children, the next household was visited for school-age children.

Presenting distance visual acuity was defined as distance visual acuity without any correction in each eye. VI was considered when presenting distance visual acuity in the better eye was <6/18.^{25,26} This was categorized in to three categories: moderate VI — presenting distance VA <6/18 and ≥6/60 in the better eye (category 1),²⁷ severe VI — presenting distance VA <6/60 and ≥3/60 in the better eye (category 2),²⁵ and blindness — presenting distance VA <3/60 in the better eye (category 3).²⁶ Positive family ocular history was defined as history of ocular problems in any first-degree relative.²⁸ School-age children were defined as age 6–18 years.²⁹ Drinking alcohol was categorized as nondrinkers (people who explicitly recorded zero for current consumption of any alcoholic beverage and zero or blank for previous drinking) and drinkers (those who reported drinking any alcoholic beverage at least three times per week and above).³⁰

An English version of the questionnaire was prepared and translated into Amharic (local language). The Amharic version of the questionnaire, Snellen charts, and torchlight were used to collect data. Interviews were conducted with children aged 12 years and above, while for those <12 years of age, the parent or guardian was contacted. Five trained MSc optometrists participated in data

collection. Pretesting was conducted in 5% of the sample in an area with a similar setting outside the study area (Gondar city). Training was given to data collectors and supervisors. There was close supervision by supervisors. The collected data were checked for completeness, accuracy, and clarity by the principal investigator and supervisors on a daily basis. Data cleanup and cross-checking was done before analysis.

Collected data were entered into Epi Info 7 and exported to and analyzed with SPSS 20. Summary statistics, frequencies, and cross-tabulations were performed. Bivariate and multivariate logistic regression models were employed to identify associations between independent variables and a dichotomous outcome variable. Variables with $p < 20\%$ in the bivariate analysis were taken for further multivariate analysis with backward logistic regression. The Hosmer–Lemeshow model was used to check model fitness. Multicollinearity between independent variables were checked by variance inflation factor. AORs with 95% CIs were used to identify significant predictors of the outcome variable. $P < 0.05$ was considered statistically significant

Before the study, ethical clearance was obtained from the University of Gondar College of Medicine and Health Sciences and comprehensive specialized hospital and School Of Medicine Ethical Review Committee, and a support letter was obtained from Bahir Dar kebele administrators. Verbal assent was obtained from all study participants and consent was obtained from parents/guardians after a detailed explanation of the study. The University of Gondar College of Medicine and Health Sciences and comprehensive specialized hospital and School Of Medicine Ethical Review Committee had approved the taking of verbal consent from parents and guardians. Those children with VI were managed at Felege Hiwot Referral Hospital after creation of a referral system by communication with the hospital administrators.

Results

Sociodemographic Characteristics Of The Study Participants

A total of 601 study subjects were included for a response rate of 95.1%. The mean age of participants was 13.14 ± 2.82 year. Half the study participants (50.3%) were males. The majority (84.9%) of study participants were Orthodox Christians and Amhara (93.8%) in ethnicity. About three-quarters of children (75%) and less than a

Table 1 Sociodemographic Characteristics Of Study Participants (n=601)

	Frequency	Percentage
Age, years		
6–9	45	7.5
10–13	258	42.9
14–18	298	49.6
Sex		50.3
Male	302	
Female	299	49.7
Educational status of household head		
Illiterate	21	3.4
Can read and write	104	17.3
Primary school	174	29
Secondary school	158	26.3
College and above	144	24
Educational level of the child		
Kindergarten	9	1.5
Primary school	451	75
Secondary school and above	141	23.5
Religion		
Orthodox	510	84.9
Muslim	67	11.1
Protestant	23	3.8
Others*	1	0.2
Ethnicity		
Amhara	564	93.8
Tigre	15	2.5
Agew	14	2.3
Oromo	6	1.00
Others**	2	0.30

Notes: *Jehovah's witness; **Dire Dawa and Benishangul-Gumuz.

third of household heads (29%) had finished primary school (Table 1).

About 27.3% of children had been admitted to an NICU, and 22.1% were born prematurely (Table 2).

With regard to maternal history during pregnancy, nearly 10% of study participants' mothers had had diabetes mellitus, hypertension, pneumonia, malaria, or syphilis (Table 3).

Only eight children had had no exposure to television, while 194 watched television from ≤ 2 m (Table 4).

Prevalence Of Visual Impairment

VI was diagnosed in 52 children (8.7%, 95% CI 6.2%–10.7%) 27 (52%) of whom were male. Of those children

Table 2 Children's Birth History (n=601)

	Frequency	Percentage
Gestational age		
<37 weeks	133	22.1
≥37 weeks	468	77.9
Mode of delivery		
SVD	542	90.2
Cesarean section	59	9.8
Admission to NICU		
Yes	164	27.3
No	437	72.7
Exclusive breast-feeding		
Yes	492	81.9
No	109	18.1

Abbreviations: SVD, spontaneous vaginal delivery; NICU, neonatal intensive-care unit.

Table 3 Maternal History During Pregnancy (n=601)

	Frequency	Percentage
Maternal alcohol consumption during pregnancy		
Yes	17	2.8
No	584	97.2
Maternal cigarette smoking during pregnancy		
Yes	0	
No	601	100
Systemic illness during pregnancy		
Yes*	62	10.3
No	590	89.7

Note: *Hypertension, diabetes, malaria, pneumonia, syphilis.

who had VI, 43 (7.2%) had moderate VI, six (1%) severe VI, and four (0.5%) were blind (Table 5).

The majority of study participants were born maturely (85.2%), 82.5% had been exclusively breast-fed >6 months, 4.9% had been admitted to an NICU, and 32.3% were watching television from 2 m or closer, of which 12.9% had VI. However, 168 (28%) study participants watched television from >4 m, of which only 3.7% had VI (Figure 1).

Factors Associated With Visual Impairment

In the bivariate analysis, history of medical visits, siblings with VI, educational status of household head,

Table 4 Socioeconomic And Behavioral Factors (n=601)

	Frequency	Percentage
Family monthly income, ETB		
<2,000	140	23.3
2,001–5,000	283	47.1
5,001–10,000	129	21.5
10,001–15,000	22	4.5
>15,000	27	3.7
Family size		
<4	272	45.3
4–6	240	39.9
>6	89	14.9
School type		
Public	344	57.2
Private	257	42.8
Medical visits (for children)		
Yes	282	46.9
No	319	53.1
Frequency of medical visits (for children)		
Yearly	92	15.3
Symptoms seen and when traumatized	190	31.6
History of television exposure		
Yes	593	98.6
No	8	1.40
Duration of television exposure		
<2 hours/day	142	23.6
2–4 hours/day	253	42.1
>4 hours/day	198	32.9
Television-exposure distance		
<2 m	194	32.3
2–4 m	231	38.3
>4 m	168	28.0
Duration of mobile/computer exposure		
<2 hours/day	260	43.3
2–4 hours/day	201	33.4
>4 hours/day	97	16.1

Abbreviation: ETB, Ethiopian birr.

exclusive breast-feeding, admission to an NICU, mode of delivery, maternal alcohol consumption, frequency of medical visits, history of systemic illness, having chronic disease (DM/Htn), parents with VI, mobile-exposure duration, gestational age, duration of television exposure, television-exposure distance, and age of the child were significantly associated with VI. However, in

Table 5 Prevalence Of VI In Terms Of Age And Sex (n=601)

Age, Years	Visual Impairment		No VI	
	Male	Female	Male	Female
6–9	0	3	20	22
10–13	18	10	119	111
14–18	9	12	136	141

the multivariable logistic analysis, only five variables — admission to an NICU, prematurity (gestational age <37 weeks), children from parents with VI, television-exposure distance <2 m and mobile/computer/video-game-exposure >4 hours/day — were independently significantly associated with VI (Table 6).

Discussion

VI is a common ophthalmic problem in young children.³ An understanding of the prevalence of VI and associated factors adds new knowledge about the risk factors of VI, leading to better understanding and management of the condition. The overall prevalence of VI in this study was 8.7% (95% CI 6.5%–11.1%), which is in line with other studies reported across the world: (Gombak, Malaysia [7.1%]³¹ and Sanaa, Yemen [9.9%]),²⁶ but higher than previous studies in Sekoru, southwest Ethiopia (0.062%),³² the whole of Ethiopia (5.3%),³³ Arada, Addis Ababa, Ethiopia (5.8%),² Puducherry, India (6.37%),⁷ Ogun, Nigeria (2.09%),³⁴ Ashanti, Ghana (3.7%),³⁵ and Sydney, Australia (2.7%).²⁹ This discrepancy might be due to the differences in study design, study setting, and inclusion criteria. In this community-based cross-sectional study, school-age children within the community were included, while in the Addis Ababa and Sekoru studies, a school-based cross-sectional study was conducted, which missed those children having VI who stay at home due to its significant effect on learning ability. In addition to these differences, there are also economic

and ethnic differences between the population of this study and other reported studies, which greatly influences VI prevalence, as reported in many studies.^{2,15} According to a study conducted in Nigeria, 2.09% had VI.³⁴ This difference could be due to the fact that in this study, there was no adequate provision of eye-care services, which might have increased prevalence.

The definition of VI influences reported prevalence rates. The definition used in some studies (best-corrected visual acuity <6/18 in the better eye)^{2,12} differs from this study. This study used the new ICD10 definition of VI, ie, presenting distance visual acuity ≤6/18. Therefore, taking the visual acuity of the participant as present definitely increases prevalence. The prevalence found in this study was lower than in Selangor, Malaysia (25%).²⁸ This may be due to variation in study populations. Most Asian nations are myopic because of complex genetic traits responsible for myopia, and in turn myopic cases outnumber VI cases in this population.⁹

This study revealed that those study participants who watched television from a distance <2 m had nearly nine times the odds of developing VI than those were watching television from >4 m, which is in agreement with a study done in Adama, Ethiopia.⁹ This might be related to the fact that watching television from close range creates a visual strain on the eyes of the children and increases the chance of refractive error. However, it is impossible to speculate on the temporal relationship between distance from the television and the development of VI, as myopic children move more closely to the television to see the picture more clearly. Study participants who had a history of admission to an NICU had nearly six times the odds of developing VI than those who had had no admissions to an NICU. This might be due to phototherapy for prematurity and development of pathological jaundice in neonates, which exposes their eyes to radiation and in turn may cause VI.³⁵

Children who had parents with VI had nearly double the odds of being visually impaired compared to those with parents who had no VI, which is in agreement with a study in southwest Nigeria.¹⁸ This is due to the inheritance nature of VI. A family history of VI is often present, but well-defined genetic patterns are unusual.⁶ Prematurity was associated with VI in this study. Children who were born prematurely had nearly triple the odds of being visually impaired compared to those who were born maturely. Studies from Denmark^{14–21} and Australia¹⁷ support this result. This could be due to risk of refractive error in premature infants and prematurely developed sensory and motor systems, which could be associated with VI.

Figure 1 Television exposure distance characteristics of study participants in Bahir Dar, northwest Ethiopia, June 2018 (n=601).

Table 6 Bivariate And Multivariate Logistic Regression Output Of Factors Associated With VI (n=601)

	Visual Impairment		COR (95% CI)	AOR (95% CI)
	Yes (n=52)	No (n=549)		
Age, years				
6–9	7	42	1.00	1.00
10–13	24	230	1.59 (0.3–3.371)	0.21 (0.05–0.89)
14–18	21	277	2.19 (0.34–1.13)	0.50 (0.20–1.04)
Sex				
Male	27	275	0.92 (0.60–1.90)	
Female	25	274	1.00	
Educational level of household head				
Illiterate	7	18	6.61 (0.68–11.66)	
Can read and write	8	95	1.71 (0.44–6.54)	
Primary school	15	156	1.44 (0.38–5.38)	
Secondary school	14	144	1.75 (0.43–7.14)	
College and above	8	136	1.00	
Maternal illness during pregnancy				
Yes	8	107	1.33 (1.82–22.8)	
No	44	442	1.00	
Type of illness during pregnancy				
Chronic disease (DM/Htn)	22	24	2.40 (0.02–0.08)	
Others	11	5	1.00	
Drank alcohol during pregnancy				
Yes	12	122	0.95 (2.25–18.04)	
No	40	427	1.00	
Gestational age				
<37 weeks	22	249	1.13 (1.61–5.21)	2.81 (1.19–6.83)***
≥37 weeks	30	300	1.00	1.00
Mode of delivery				
SVD	39	503	1.00	
Cesarean section	13	46	0.27 (1.81–7.32)	
Admission to NICU				
Yes	21	296	1.72 (3.56–11.64)	5.52 (2.01–15.15)**
No	31	253	1.00	1.00
Sibling with VI				
Yes	10	41	2.95 (1.58–6.83)	
No	42	508	1.00	
Parents with VI				
Yes	16	247	1.84 (1.97–6.78)	1.8 (0.13–0.97)*
No	36	302	1.00	1.00
Family history of spectacle use				
Yes	19	54	5.29 (2.80–9.90)	
No	33	495	1.00	
History of medical visits				
Yes	10	272	1.00	
No	42	277	0.3 (2.03–8.39)	

(Continued)

Table 6 (Continued).

	Visual Impairment		COR (95% CI)	AOR (95% CI)
	Yes (n=52)	No (n=549)		
Frequency of medical visits				
Yearly	10	82	1.00	
Symptoms seen	19	128	0.12 (0.04–0.33)	
Duration of watching TV				
<2 hours/day	20	98	1.00	
2–4 hours/day	12	198	3.36 (0.63–2.75)	
>4 hours/day	6	155	5.27 (0.59–3.53)	
TV-exposure distance				
<2 m	26	223	1.47 (0.69–12.81)	8.71 (1.49–18.24)*
2–4 m	17	293	2.97 (0.86–4.87)	2.32 (2.42–25.26)
>4 m	5	29	1.00	1.00
Duration of mobile exposure				
<2 hours/day	20	245	1.00	1.00
2–4 hours/day	12	154	1.04 (2.44–17.11)	2.98 (1.30–6.78)
>4 hours/day	6	99	1.95 (1.10–7.93)	1.6 (1.32–4.45)*

Notes: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Abbreviations: TV, television; VI, visual impairment; DM, diabetes mellitus; Htn, hypertension; SVD, spontaneous vaginal delivery; NICU, neonatal intensive-care unit.

The results we found were more extreme than other studies, and this is significant for public health. This implies that the prevention techniques applied were at a low level. This might be related to poor utilization of eye-care services and the limitation of technologically advanced health-care services. According to the Vision 2020 plan, the major priority area to act on is elimination of avoidable VI.³⁶ To achieve the goal, a regular school screening program and community service is required.

Limitations

Assessment of VI was done with visual acuity only, which did not include visual field testing. The majority of questions in the data-collection instrument asked about subject history, and would have been exposed to recall bias. As this study was cross-sectional study, it was impossible to identify which comes first: the associated factors or the outcome. For example, according to this study, watching television from <2 m may cause a visual strain and development of myopia, and logically myopic children move more closely to the television, so it is impossible to determine which comes first. Causality and such temporal relationships between dependent and other independent variables were not clearly identified either, due to the design of the study. In order to identify

such temporal relationships clearly, we recommend other studies be conducted with better design.

Conclusion

The prevalence of VI among school-age children in Bahir Dar was 8.7%, which is higher than other studies. Moderate VI was most common among visually impaired children. Premature birth, admission to NICU, parents with VI, television exposure <4 m, and mobile-exposure duration >2 and 4 hours per day were independently significantly variables associated with VI.

Availability Of Data

The hard copy of the collected primary data used in this research is securely locked so that is accessible to the authors only. For the sake of privacy of the participants, the data are not totally open to all readers; however, data can be made available from one of the authors upon reasonable request.

Acknowledgments

We are deeply indebted to the University of Gondar, which gave ethical clearance to conduct this research. We would also like to acknowledge the study participants for their cooperation and willingness to participate in this study.

Disclosure

The authors report no conflicts of interest with respect to the research, authorship, or publication of this article.

References

1. Resnikoff S, Pascolini D, Etya'ale D, et al. Global data on visual impairment in the year 2002. *Bull World Health Organ.* 2004;82(11):844–851.
2. Darge HF, Shibru G, Mulugeta A, et al. The prevalence of visual acuity impairment among school children at Arada Subcity primary schools in Addis Ababa, Ethiopia. *J Ophthalmol.* 2017;2017:9326108.
3. Pascolini D, Mariotti SP. Global estimates of visual impairment: 2010. *Br J Ophthalmol.* 2012;96(5):614–618. doi:10.1136/bjophthalmol-2011-300539
4. Resnikoff S, Pascolini D, Mariotti SP, et al. Global magnitude of visual impairment caused by uncorrected refractive errors in 2004. *Bull World Health Organ.* 2008;86(1):63–70. doi:10.2471/BLT.00.000000
5. Demissie BS, Solomon AW. Magnitude and causes of childhood blindness and severe visual impairment in Sekoru District, Southwest Ethiopia: a survey using the key informant method. *Trans R Soc Trop Med Hyg.* 2011;105(9):507–511. doi:10.1016/j.trstmh.2011.04.007
6. Weingeist T, Liesegang T, Grand M. American academy of ophthalmology, basic and clinical science course 2000–2001 lens and cataract. *Biochemistry.* 10–17.
7. Vishnuprasad R, Bazroy J, Madhanraj K, et al. Visual impairment among 10–14-year school children in Puducherry. *J Family Med Prim Care.* 2017;6(1):58. doi:10.4103/2249-4863.214983
8. Aniza I, Nawi AM, Jamsiah M, et al. Prevalence of visual acuity impairment and its associated factors among secondary school students in Beranang, Selangor. *Malaysian J Publ Health Med.* 2012;12(1):39–44.
9. Bezabih L, Abebe TW, Fite RO. Prevalence and factors associated with childhood visual impairment in Ethiopia. *Clin Ophthalmol.* 2017;11:1941. doi:10.2147/OPTH.S135011
10. Livingston P, McCarty C, Taylor H. Visual impairment and socioeconomic factors. *Br J Ophthalmol.* 1997;81(7):574–577. doi:10.1136/bjo.81.7.574
11. Faal H, Minassian D, Sowa S, et al. National survey of blindness and low vision in The Gambia: results. *Br J Ophthalmol.* 1989;73(2):82–87. doi:10.1136/bjo.73.2.82
12. Whitfield R, Schwab L, Ross-Degnan D, et al. Blindness and eye disease in Kenya: ocular status survey results from the Kenya Rural Blindness Prevention Project. *Br J Ophthalmol.* 1990;74(6):333–340. doi:10.1136/bjo.74.6.333
13. Pai AS-I, Wang JJ, Samarawickrama C, et al. Prevalence and risk factors for visual impairment in preschool children: the Sydney Paediatric Eye Disease Study. *Ophthalmology.* 2011;118(8):1495–1500. doi:10.1016/j.ophtha.2011.01.027
14. Robaei D, Rose KA, Kifley A, Cosstick M, Ip JM, Mitchell P. Factors associated with childhood visual impairment: findings from a population-based study. *Ophthalmology.* 2006;113(7):1146–1153. doi:10.1016/j.ophtha.2006.02.019
15. Blencowe H, Lawn JE, Vazquez T, et al. Preterm-associated visual impairment and estimates of retinopathy of prematurity at regional and global levels for 2010. *Pediatr Res.* 2013;74(S1):35. doi:10.1038/pr.2013.205
16. Torp-Pedersen T, Boyd HA, Poulsen G, et al. Perinatal risk factors for visual impairment. *Int J Epidemiol.* 2010;39(5):1229–1239. doi:10.1093/ije/dyq092
17. Azonobi I, et al. Risk factors for visual impairment in Southwestern Nigeria. *Pak J Ophthalmol.* 2009;25:3.
18. Chia A, Lin X, Dirani M, et al. Risk factors for strabismus and amblyopia in young Singapore Chinese children. *Ophthalmic Epidemiol.* 2013;20(3):138–147. doi:10.3109/09286586.2013.767354
19. Gilbert CE, Shah SP, Jadoon MZ, et al. Poverty and blindness in Pakistan: results from the Pakistan national blindness and visual impairment survey. *Bmj.* 2008;336(7634):29–32. doi:10.1136/bmj.39395.500046.AE
20. Foster A, Gilbert C. Epidemiology of childhood blindness. *Eye.* 1992;6(2):173. doi:10.1038/eye.1992.34
21. Powlis A, Botting N, Cooke RWI, et al. Visual impairment in very low birthweight children. *Arch Dis Child Educ Pract Ed.* 1997;76(2):F82–F87. doi:10.1136/fn.76.2.F82
22. Goujon A, Barakat B, Goujon A, et al. Projection of populations by level of educational attainment, age, and sex for 120 countries for 2005–2050. *Demogr Res.* 2010;22:383–472. doi:10.4054/DemRes.2010.22.15
23. Huizink AC, Mulder EJ. Maternal smoking, drinking or cannabis use during pregnancy and neurobehavioral and cognitive functioning in human offspring. *BMC.* 2005;30/2006:24–41.
24. Macnaughton J. Eye Essentials: low Vision Assessment. *Clin Exp Optom.* 2006;89(2):118–120.
25. WHO. Change the definition of blindness. WHO, 2008.35. Available From: www.who.int/blindness:ChangeTheDefinitionofBlindness. Accessed October 14, 2019
26. Bamashmus M, Al-Akily S. Profile of childhood blindness and low vision in Yemen: a hospital-based study/Profil de la cecite et de la basse vision chez l'enfant au Yemen: une etude hospitaliere. *East Mediterr Health J.* 2010;16(4):425. doi:10.26719/2010.16.4.425
27. Tielsch JM, Katz J, Sommer A, et al. Family history and risk of primary open angle glaucoma: the Baltimore Eye Survey. *Arch Ophthalmol.* 1994;112(1):69–73. doi:10.1001/archophth.1994.01090130079022
28. Edussuriya K, Sennanayake S, Senaratne T, et al. The prevalence and causes of visual impairment in central Sri Lanka the Kandy Eye study. *Ophthalmology.* 2009;116(1):52–56. doi:10.1016/j.ophtha.2008.08.034
29. Kaminski M, Rumeau C, Schwartz D. Alcohol consumption in pregnant women and the outcome of pregnancy. *Alcohol Clin Exp Res.* 1978;2(2):155–163. doi:10.1111/j.1530-0277.1978.tb04716.x
30. Goh PP, Abqariyah Y, Pokharell GP, Ellwein LB. Refractive error and visual impairment in school-age children in Gombak District, Malaysia. *Am Acad Ophthalmol.* 2004;56(0161–6420/05).
31. Demissie BS, Solomon AW. Solomon, magnitude and causes of childhood blindness and severe visual impairment in Sekoru District, Southwest Ethiopia. *Royal Soc Trop Med Hyg.* 2010;105:507–511
32. Berhane Y, Worku A, Bejiga A, et al. *Ethiop J Health Dev.* 2007;21(3):204–210.
33. Fasina F, Ajaiyeoba A. The prevalence and causes of blindness and low vision in Ogun State, Nigeria. *Afr J Biomed Res.* 2003;6(2).
34. Kumah BD, Ebri A, Abdul-Kabir M, et al. Refractive error and visual impairment in private school children in Ghana. *Am Acad Optometry.* 2013;90(12):1456–1461. doi:10.1097/OPX.0000000000000099
35. Wu PY, Lim RC, Hodgman JE, et al. Effect of phototherapy in preterm infants on growth in the neonatal period. *J Pediatr.* 1974;85(4):563–566. doi:10.1016/S0022-3476(74)80471-3
36. WHO. Global initiative for the elimination of avoidable blindness: action plan 2006–2011. 2007. Available from: <https://apps.who.int/iris/handle/10665/43754>. Accessed October 11, 2019.

Clinical Optometry

Dovepress

Publish your work in this journal

Clinical Optometry is an international, peer-reviewed, open access journal publishing original research, basic science, clinical and epidemiological studies, reviews and evaluations on clinical optometry. All aspects of patient care are addressed within the journal as well as the practice of optometry including economic and business analyses. Basic and clinical research papers are published that cover

all aspects of optics, refraction and its application to the theory and practice of optometry. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-optometry-journal>