

# Downregulated Circular RNA hsa\_circ\_0000291 Suppresses Migration And Proliferation Of Gastric Cancer Via Targeting The miR-183/ITGB1 Axis

This article was published in the following Dove Press journal:  
*Cancer Management and Research*

Chuanwu Cao\*

Shilong Han\*


Yifeng Yuan

Yongfa Wu

Weishuai Lian

Xiaojun Zhang

Long Pan

Maoquan Li 

Department of Interventional and Vascular Surgery, Tenth People's Hospital of Tongji University, Shanghai 200072, People's Republic of China

\*These authors contributed equally to this work

**Background:** Circular RNAs are implicated in a variety of cancers. This investigation found that hsa\_circ\_0000291 expression was upregulated in gastric cancer (GC) cell lines, yet its role in GC has not yet been reported.

**Objective:** To explore the effects of hsa\_circ\_0000291 on GC cell proliferation and invasion.

**Materials and methods:** In the current research, we used the gastric cancer cell lines MGC803 and MKN-28 to study hsa\_circ\_0000291 function. The relationship between hsa\_circ\_0000291, miR-183 and ITGB1 was analyzed by firefly luciferase analysis and Western blots, and qRT-PCR approaches were used for protein and gene expression analysis, respectively. Tumor growth and metastasis were determined in nude mice xenografts using MKN-28 cells, with or without hsa\_circ\_0000291 downregulation.

**Results:** Our data showed that hsa\_circ\_0000291 was upregulated in GC cell lines, whereas hsa\_circ\_0000291 silencing suppressed cell metastasis and proliferation in in vivo and in vitro studies. Our results showed that the downregulation of hsa\_circ\_0000291 suppressed integrin beta 1 (ITGB1) expression via miR-183 "sponging," which was validated by rescue experiments using the luciferase reporter assay. Our observations suggested that hsa\_circ\_0000291 silencing suppressed the aggressive, metastatic GC phenotype.

**Conclusion:** Taken together, hsa\_circ\_0000291 knockdown inhibited GC cell metastasis and growth by regulating the miR-183/ITGB1 axis. Importantly, this approach could provide a therapy target and potential biomarker for the diagnosis and treatment of GC.

**Keywords:** hsa\_circ\_0000291, miR-183, integrin beta 1, proliferation and migration, gastric cancer

## Introduction

Gastric cancer (GC) is the main causal factor of mortality among Asian populations.<sup>1</sup> While there have been improvements in the diagnostics and treatment of GC, a large proportion of patients have poor prognoses.<sup>2</sup> Radical gastric surgery is often associated with high recurrence rates that result in a 5-year overall survival rate of 20%~40%.<sup>3,4</sup> Therefore, it is imperative to seek appropriate markers for therapy and the early diagnosis of GC.

Increasing evidence has found that endogenous noncoding RNAs such as circular RNAs (circRNAs), long noncoding RNAs (lncRNAs), and microRNAs (miRNAs), function in the pathological regulation of different diseases, including cancer.<sup>5,6</sup> Various from linear RNAs, circRNAs are circular structures, having no 5'

Correspondence: Maoquan Li  
Department of Interventional and Vascular Surgery, Tenth People's Hospital of Tongji University, 301 Middle Yan Chang Road, Shanghai 200072, People's Republic of China  
Tel +86-021-66313506  
Email cjr.limaoquan@vip.163.com

caps and 3' tails. Recent reports have illustrated that circRNAs function in the development and progression of GC.<sup>7-9</sup> Several circRNAs appear to regulate gene function in tumorigenesis via sponge miRNA mechanisms.<sup>10,11</sup> Nevertheless, circRNA regulatory functions in GC are unclear.

In this research, hsa\_circ\_0000291 had abnormal expression. Similarly, miR-183 negatively correlated with hsa\_circ\_0000291 levels. Therefore, our aim was to examine the function and biological roles of hsa\_circ\_0000291 during GC. The data suggested that hsa\_circ\_0000291 was upregulated in GC cell lines, and that downregulated hsa\_circ\_0000291 suppressed GC cell migration and proliferation by targeting the miR-183/ITGB1 axis. We believe these data will provide new therapeutic targets for GC treatment going forward.

## Materials And Methods

### Animal And Ethics Statement

BALB/c nude mice aged 4 weeks, weighing 15~20 g, were purchased from Shanghai SLAC Laboratory Animal Co., Ltd (Shanghai, China). All procedures were approved by and conducted following the guidelines of the Ethics Committee of the Shanghai Tenth People's Hospital of China. All surgical procedures were performed under anesthesia, and every effort was made to minimize suffering. Mice were anesthetized by intraperitoneal injection of 30 mg/kg sodium pentobarbital.

### Cell Culture And Transfection

The human GC cell lines MGC-803, BGC823, SGC-7901, MKN-28, and gastric epithelial cell GES-1 were bought from the Type Culture Collection of the Chinese Academy of Sciences, Shanghai, China. Cells were cultivated in DMEM medium (Gibco, Gaithersburg, USA), which contained 10% fetal bovine serum (FBS, Gibco) at 37°C, in 5% CO<sub>2</sub>. Small interfering RNAs for hsa\_circ\_0000291 (sicircRNA) (5'-GGGAAGUUUAAUUGUAUGUUU-3'), miR-183 mimics, miR-183 inhibitors (5'-AGUGAAUUCUACCAGUCCCAUA-3'), negative controls (NC) and the integrin beta 1 (ITGB1) overexpression vector were transfected into cultured MGC-803 or MKN-28 cells using Lipofectamine 2000 (Invitrogen, Carlsbad, USA). To further identify hsa\_circ\_0000291 effects on tumor growth in vivo, we constructed a hsa\_circ\_0000291 knockdown model in lentiviral stabilized MKN-28 cells.

## Fluorescence In Situ Hybridization (FISH) Analysis

We used probes to the hsa\_circ\_0000291 sequence for FISH analysis, as reported previously.<sup>12</sup> Specific FITC-labelled hsa\_circ\_0000291 probes (GTGCTTGTAAGAGGGCCCTGG) were used to counterstain cells following standard protocols (Genepharma, Shanghai, China).

## Bioinformatics Analyses

We used the bioinformatics analysis website Circular RNA Interactome for circRNA and miRNA interaction predictions ([https://circinteractome.nia.nih.gov/miRNA\\_Target\\_Sites/mirna\\_target\\_sites.html](https://circinteractome.nia.nih.gov/miRNA_Target_Sites/mirna_target_sites.html)). The target site between miR-183 and the 3'-UTR of ITGB1 was predicted using the TargetScan web-based tool ([http://www.targetscan.org/vert\\_71/](http://www.targetscan.org/vert_71/)). Tools provided at <http://gepia.cancer-pku.cn/> were used to predict hsa\_circ\_0000291 expression in the prognosis of gastric cancer.

## Cell Proliferation Assay

Following standard protocols provided by Invitrogen (Carlsbad, CA, USA) we used the cell counting Kit-8 (CCK-8) to assess both MGC803 and MKN-28 cell proliferation. Briefly, 2×10<sup>3</sup> cells in 100 µL DMEM were seeded into 96-well plates. We measured cell viability at 0, 24, 48 and 72 h after seeding with 10 µL CCK-8.

For colony formation assays, we seeded MGC803 and MKN-28 cells at a density of 2×10<sup>3</sup> cells/well into 6 well plates. After culturing in DMEM medium, plus 10% FBS for 10 days, we washed cells in sterile PBS and fixed them with 4% paraformaldehyde for 30 mins before staining with Crystal Violet and counting.

## Western Blots

We extracted protein from tumor cells or tissues using RIPA lysis buffer (Sigma-Aldrich, St. Louis, USA). We standardized protein concentrations using the BCA Protein Assay kit (Vigorous Biotechnology Beijing, China) before resolving proteins on SDS-PAGE and transferring them to nitrocellulose membranes (Millipore, Madison, USA). After blocking in nonfat milk (5%) and incubating with primary antibodies, membranes were incubated with horseradish peroxidase-coupled secondary antibodies ITGB1 (1:500, Santa Cruz Biotechnology, Dallas, TX, USA), glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (1:1,000,

Santa Cruz Biotechnology, Dallas, TX, USA). GAPDH was used as an internal control.

## RNA Extraction And qRT-PCR

We performed RNA extractions using TRIzol reagent (Invitrogen), using a previously published protocol.<sup>13</sup> We used pTRUEscript the First Strand cDNA Synthesis Kit (Aidlab, Beijing, China) for cDNA synthesis. We used 2 × SYBR Green qPCR Mix. Aidlab was used for qRT-PCR detection which was conducted using the ABI 7900HT sequence detection system, (Thermo Fisher Scientific). We used the  $2^{-\Delta\Delta CT}$  method to compute results. The hsa\_circ\_0000291 primers were; forward, 5'-GCCCTCGCTTTCATGTGC-3' and reverse, 5'-CTAGAGACTATAGGGAAACC-3'. The miR-183 primers were forward, 5'-CGGTATGGCACTGGTAGAATTCACT-3' and reverse, 5'-GCTTTCCAATGCACTGACCATT-3'. U6; forward 5'-CTCGCTTCGGCAGCAC-3' and reverse, 5'-AACGCTTCACGAATTTGCGT-3'. GAPDH forward, 5'-GGAGCGAGATCCCTCCAAAAT-3' and reverse, 5'-GGCTGTTGTCATACTTCTCATGG-3'. GAPDH and U6 were used as internal references.

## Migration Assay

For cell migration analysis, MGC803 and MKN-28 cells were placed into a transwell upper chamber at a density of  $1 \times 10^5$  cells (8 µm pore membrane, BD Biosciences, Franklin Lakes, USA) in 200 µL serum-free medium. Complete medium (500 µL) was added to the bottom chamber. After 24 h culture, cell numbers in the bottom chamber were calculated after fixing with 4% paraformaldehyde and staining with 0.1% crystal violet.

## Tumor Xenograft Formation And Metastasis Assays

A total of  $2 \times 10^7$  MKN-28 cells with or without hsa\_circ\_0000291 silencing was injected into the right flanks of nude mice, as reported previously.<sup>14</sup> Tumor sizes (volume =  $0.5 \times \text{length} \times \text{width}^2$ ) were measured by a vernier calipers every five days for 30 day before mice were euthanized.

For metastasis analyses,  $2 \times 10^5$  MKN-28 cells transfected with luciferase expression vectors, with or without hsa\_circ\_0000291 silencing, were intravenously injected into mice tails. MKN-28 cell metastasis was analyzed using bioluminescence imaging after intravenously injected for 30 day by luciferin intravenous injection (150 mg luciferin/kg body weight).

## Dual Luciferase Reporter Assay

Reporter plasmids were constructed by inserting wild-type/mutated-type circRNA, or the ITGB1 3'-UTR sequence into a pGL3 vector (Promega, Madison, USA). Then lipofectamine 2000 was used and miR-183 mimics combined with reporter plasmids co-transfected into 239T cells. Firefly and Renilla luciferase activities were captured by a Dual Luciferase Reporter Assay System (Promega, Sunnyvale, USA) after culturing for 48 h.


## Statistics Analyses

We utilized GraphPad Prism (GraphPad, La Jolla, USA) for data analyses. The data were calculated as the mean ± SEM.  $p$  value ≤ 0.05 reflected significant differences.


## Results

### Hsa\_circ\_0000291 Downregulation Suppresses Tumor Progression In Vivo

We observed that hsa\_circ\_0000291 expression was increased in gastric cancer tissues when compared with adjacent normal tissues (Figure 1A). The RT-qPCR detection method also found that hsa\_circ\_0000291 expression in GC cell lines increased when compared to GES1 cells (Figure 1B). Hsa\_circ\_0000291 was derived from a *CD44* gene exon. A fluorescence in situ hybridization assay showed that hsa\_circ\_0000291 localized to the cytoplasm (Figure 1C). To identify if hsa\_circ\_0000291 participated in the progress of GC, lentiviral stable strains of hsa\_circ\_0000291 knockdown (sh-circRNA) in MKN-28 cells were constructed. Our data showed that hsa\_circ\_0000291 expression in sh-circRNA MKN-28 cells was significantly downregulated, when compared to control or negative control (NC) cells (Figure 2A). The lentiviral-stabilized circRNA silenced MKN-28 cells or NCs were used for subcutaneous tumorigenesis analysis. These data indicated that hsa\_circ\_0000291 knockdown suppressed tumor growth (weight and volume) when compared to the NC group (Figure 2B–D). Bioluminescence imaging showed that hsa\_circ\_0000291 silencing suppressed MKN-28 cell metastasis (majority in lung tissue) in mice (Figure 2E). Using qRT-PCR, we found that miR-183 expression was upregulated following hsa\_circ\_0000291 silencing in mouse tumor tissues (Figure 2F). Western blot detection revealed that ITGB1 expression was downregulated after hsa\_circ\_0000291 knockdown (Figure 2G and H). These results suggested that hsa\_circ\_0000291 silencing suppressed tumor metastasis and growth in vivo. The results also showed that both miR-183 and ITGB1 participated in GC progression.



**Figure 1** The expression of hsa\_circ\_0000291 and sub-cellular localization. **(A)** The qRT-PCR assay shows the expression of hsa\_circ\_0000291 in gastric cancer tissues and adjacent normal tissues. Data are denoted by the mean  $\pm$  SD. \*\*\*P < 0.001 versus normal group. **(B)** The qRT-PCR assay shows the expression of hsa\_circ\_0000291 in GC cell lines (MGC803, MKN-28, SGC7901 and BGC823) and normal human gastric epithelial cell GES1. Data are denoted by the mean  $\pm$  SD. \*\*\*P < 0.001 versus GES1 group. **(C)** Fluorescence in situ hybridization was performed to capture the subcellular localization of hsa\_circ\_0000291. DAPI = nuclear staining (bottom, left); hsa\_circ\_0000291 = green fluorescent-tagged hsa\_circ\_0000291 (top, left). Merged images are plotted at right.



**Figure 2** Downregulation hsa\_circ\_0000291 suppressed tumor metastasis and growth in nude mice xenografts. **(A)** The quantitative reverse transcription-polymerase chain reaction assay illustrates the hsa\_circ\_0000291 expression in adenovirus-transfected cells (sh-circRNA) or negative control (NC) transfected MKN-28 cells. Data are denoted by the mean  $\pm$  SD. \*\*\*P < 0.001 versus NC. **(B)** Representative photographs of MKN-28 tumor formation in xenografts of nude mice. **(C)** Tumor volume summary in mice that measured weekly. Data are denoted by the mean  $\pm$  SD. \*\*P < 0.01, \*\*\*P < 0.001 versus NC. **(D)** Tumor weight was captured 30 days from injection. Data are denoted by the mean  $\pm$  SD. \*\*\*P < 0.001 versus NC. **(E)** Live imaging demonstrates the hsa\_circ\_0000291 effects on metastasis of MKN-28 cells 30 day after intravenous tail injection. scale bars, 1 cm. **(F)** qRT-PCR assay showing the miR-183 expression. Data are denoted by the mean  $\pm$  SD. \*\*\*P < 0.001 versus control. **(G and H)** Western blot analysis of the integrin beta 1 (ITGB1) expression in tumor tissues. Data are denoted by the mean  $\pm$  SD. \*\*\*P < 0.001 versus NC.



## Knockdown Of hsa\_circ\_0000291 Inhibits Cell Migration And Proliferation By Regulating The miR-183/ITGB1 Axis

To further explore regulatory mechanisms, MGC803 and MKN-28 cells were transfected with a hsa\_circ\_0000291 silencing vector (sicircRNA), combined with an ITGB1 overexpression vector, or treatment with an miR-183 inhibitor. Data showed that hsa\_circ\_0000291 expression was downregulated after sicircRNA administration, but downregulating miR-183 or overexpression of ITGB1 could not rescue hsa\_circ\_0000291 expression in these cells (Figure 3A and B). Our qRT-PCR data illustrated that downregulated hsa\_circ\_0000291 promoted miR-183 expression. MiR-183 treatment also suppressed miR-183 expression (Figure 3A and B). Overexpressed ITGB1 had no effects on miR-183 expression after hsa\_circ\_0000291 silencing in both MGC803 and MKN-28 cell lines (Figure 3C and D). Western blot analysis demonstrated that downregulated hsa\_circ\_0000291 blocked ITGB1 expression. Inhibited miR-183 rescued the expression of ITGB1, and ITGB1 expression levels were increased after transfection with the ITGB1 overexpression vector (Figure 3E and F). These observations suggested that ITGB1 was downstream of miR-183.

The result from the CCK8 assay and cloning formation revealed that hsa\_circ\_0000291 silencing suppressed MGC803 and MKN-28 cell proliferation, but that miR-183 inhibitor treatment rescued the proliferation of both MGC803 and MKN-28 cells. ITGB1 overexpression further promoted the proliferation of both MGC803 and MKN-28 cell lines (Figure 3G–K). Transwell migration assay data suggested that hsa\_circ\_0000291 silencing-induced migration inhibition was rescued after treatment with the miR-183 inhibitor. The upregulation of ITGB1 promoted the migration of both MGC803 and MKN-28 cell lines (Figure 3L–N). In summary, these data suggest that hsa\_circ\_0000291 silencing suppressed tumor cell metastasis and proliferation by enhancing miR-183 expression, while suppressing ITGB1 expression.

## The Inhibitory Effects Of miR-183 On Cell Proliferation And Migration Were Reversed After ITGB1 Up-Regulation In Vitro

To identify the relationship between miR-183 and ITGB1, we transfected both MGC803 and MKN-28 cell lines with miR-183 mimics, with/without the ITGB1 overexpression vector. Our qRT-PCR data revealed that miR-183 expression was upregulated in both MGC803 and MKN-28 cell lines after

transfection with the miR-183 mimic, and that ITGB1 overexpression did not affect miR-183 expression, (Figure 4A and B). Western blot analysis revealed that ITGB1 expression was downregulated after miR-183 overexpression but was rescued after transfection with the ITGB1 overexpression vector (Figure 4C and D). The results from the CCK8 (Figure 4E and F) and cloning formation assay revealed that cell proliferation was suppressed after miR-183 overexpression, but ITGB1 overexpression rescued and promoted the proliferation of both MGC803 and MKN-28 cell lines (Figure 4G–I). Transwell migration assays revealed that miR-183 expression inhibited cell migration, whereas upregulation of ITGB1 promoted the migration of both MGC803 and MKN-28 cell lines (Figure 4J–L). These results suggested that the overexpression of miR-183 inhibited tumor cell migration and proliferation by targeting ITGB1.


## The Relationship Between hsa\_circ\_0000291, miR-183 And ITGB1

To predict a relationship between hsa\_circ\_0000291, miR-183 and ITGB1, our bioinformatics analyses showed that hsa\_circ\_0000291 appeared to target miR-183. To physically validate this observation, a luciferin reporter vector was constructed. The results of this luciferase reporter assay suggested that miR-183 inhibited luciferase activity after transfection with the wild luciferase reporter vector, but that it did not affect luciferase activity after transfection with a mutated luciferase reporter vector, suggesting that miR-183 was the target of hsa\_circ\_0000291 (Figure 5A and B).

Bioinformatics analysis found that miR-183 directly interacted with the ITGB1 3'-UTR and suppressed post-translational ITGB1 expression (Figure 5C). A luciferase reporter assay showed that miR-183 inhibited luciferase activity, after transfection with a wild-type luciferase reporter vector, but luciferase activity was not affected after transfection with a mutated luciferase reporter vector, suggesting that ITGB1 was the target of miR-183 (Figure 5D). These data indicated that hsa\_circ\_0000291 knockdown inhibited metastasis and the growth of gastric cancer through targeting the miR-183/ITGB1 axis.

## Discussion


Studies have identified that GC is increasingly diagnosed at advanced stages. The prognosis is poor, with a five-year survival rate of 5%~20%,<sup>15</sup> with surgery considered a viable standard option. However, postoperative recurrence and metastasis rates are unsatisfactory.<sup>16</sup> circRNAs are a new



**Figure 3** Knockdown hsa\_circ\_0000291 inhibits gastric cancer cell migration and proliferation by regulation of the miR-193/ITGB1 axis in in vitro experiment. **(A and B)** qRT-PCR assay shows hsa\_circ\_0000291 expression in both MGC803 **(A)** and MKN-28 **(B)** cells after transfection with siRNA against hsa\_circ\_0000291 combined with or not miR-183 inhibitor or the ITGB1 overexpression vector. Data are denoted by the mean  $\pm$  SD. \*\*\* $P < 0.001$  versus NC. #### $P < 0.001$  versus siRNA. **(C and D)** qRT-PCR assay shows miR-183 expression in both MGC803 **(C)** and MKN-28 **(D)** cells. Data are denoted by the mean  $\pm$  SD. \*\*\* $P < 0.001$  versus NC. #### $P < 0.001$  versus siRNA. **(E and F)** Western blots show ITGB1 expression in both MGC803 **(E)** and MKN-28 **(F)** cells. Relative protein levels were analyzed, and data are denoted by the mean  $\pm$  SD. \*\*\* $P < 0.001$  versus NC. #### $P < 0.001$  versus siRNA. **(G and H)** CCK8 assays were performed to assess cell proliferation in both MGC803 **(G)** and MKN-28 **(H)** cells. Data are denoted by the mean  $\pm$  SD. \* $P < 0.05$ , \*\*\* $P < 0.001$  versus NC. #### $P < 0.001$  versus siRNA. **(I–K)** Cloning formation assay showing the MGC803 and MKN-28 cell proliferations. Data are denoted by the mean  $\pm$  SD. \*\*\* $P < 0.001$  versus NC. #### $P < 0.001$  versus siRNA. scale bars, 200  $\mu$ m. **(L–N)** Cell migration was determined in MGC803 and MKN-28 cells by Transwell® assays. Data are denoted by the mean  $\pm$  SD. \*\*\* $P < 0.001$  versus NC. #### $P < 0.001$  versus siRNA. scale bars, 50  $\mu$ m.

family of ncRNAs where biological mechanisms are largely unknown.<sup>17,18</sup> In this study, our data showed that hsa\_circ\_0000291 expression increased in GC cell lines, in comparison to wild-type cells, suggesting that hsa\_circ\_0000291

participates in GC progression. We also found that hsa\_circ\_0000291 silencing suppressed GC metastasis and proliferation in in vitro and in vivo models, suggesting that hsa\_circ\_0000291 may function as an oncogene.




**Figure 4** ITGB1 overexpression reversed miR-183-induced cell migration and growth inhibition in vitro. MGC803 and MKN-28 cells were transfected with miR-183 mimics combined with or without the ITGB1 overexpression vector. (A and B) qRT-PCR assay showing the expression of miR-183 in both MGC803 (A) and MKN-28 (B) cells. Data are denoted by the mean  $\pm$  SD. \*\*\* $P$  < 0.001 versus NC. (C and D) Western blot showing ITGB1 expression in both MGC803 (C) and MKN-28 (D) cells. Relative protein levels were analyzed, and data are denoted by the mean  $\pm$  SD. ### $P$  < 0.001 versus NC. \*\*\* $P$  < 0.001 versus mimic. (E and F) CCK8 assays were performed to assess cell proliferation in both MGC803 (E) and MKN-28 (F) cells. Data are denoted by the mean  $\pm$  SD. \*\*\* $P$  < 0.001 versus NC. ### $P$  < 0.001 versus mimic. (G-I) Cloning formation assay showing the cell proliferation of MGC803 and MKN-28 cells. Data are denoted by the mean  $\pm$  SD. \*\*\* $P$  < 0.001 versus NC. ### $P$  < 0.001 versus mimic. (J-L) Cell invasion and migration were determined in MGC803 and MKN-28 cells by Transwell® assays. Data are denoted by the mean  $\pm$  SD. \*\*\* $P$  < 0.001 versus NC. ### $P$  < 0.001 versus mimic. scale bars, 50  $\mu$ m.

The silencing of hsa\_circ\_0000291 promoted miR-183 expression and suppressed ITGB1 expression. Previous studies have suggested that miR-183 upregulation promotes apoptosis and inhibits human endometrial cancer cell invasion, proliferation, migration and epithelial-mesenchymal transition by targeting ezrin.<sup>19</sup> MiR-183 also promotes canonical Wnt signaling activities and regulates bladder cancer apoptosis and growth by the downregulation of AXIN2.<sup>20</sup> In this study, the hsa\_circ\_0000291 silencing effects on GC proliferation and metastasis were abolished by an miR-183 inhibitor, suggesting that hsa\_circ\_0000291 silencing suppressed GC proliferation and metastasis by absorbing miR-183. This observation appeared to be consistent with a previous report suggesting that circRNAs may function as sponges for microRNAs.<sup>18</sup> CircRNAs-miRNAs

participate in several physiological and pathophysiological processes, including cell differentiation, proliferation and metastasis.<sup>21-23</sup> Bifluorescein studies have also shown that miR-183 is the hsa\_circ\_0000291 target.

We showed that miR-183 overexpression suppressed ITGB1 expression. Previously, it has been shown that ITGB1 plays vital roles in cell growth and motility in non-small cell lung cancer (NSCLC), breast cancer, GC and liver cancer.<sup>24</sup> In NSCLC, miR-134 inhibits tumor cell mobility and invasions by modulating ITGB1.<sup>24</sup> In another study, it was found that ITGB1 is the candidate miR-493-5p target and that ITGB1 expression levels were associated with non-small-cell carcinoma prognoses.<sup>25</sup> In this study, we noted that miR-183 inhibited ITGB1 expression at the post-transcriptional level by interacting with the





**Figure 5** The interaction relationships among miR-183, hsa\_circ\_0000291, and ITGB1. **(A)** The predicted miR-183 binding sites in the hsa\_circ\_0000291. The the hsa\_circ\_0000291 mutated (Mut) version is also given. **(B)** The relative luciferase activity was determined 48 h after transfection with miR-183 mimic/normal control (NC) or with the hsa\_circ\_0000291 wild-type/Mut in HEK293T cells. Data are denoted by the mean  $\pm$  SD. \*\*\* $P < 0.001$ . **(C)** The predicted miR-183 binding sites with the ITGB1 3'-UTR. The 3'-UTR-ITGB1 mutated version is also provided. **(D)** Relative luciferase activity was determined 48 h after transfection with miR-183 mimic/normal control or with the 3'-UTR-ITGB1 wild-type/Mut in HEK293T cells. Data are denoted by the mean  $\pm$  SD. \*\*\* $P < 0.001$ .

ITGB1 3'-UTR. Bifluorescein reports have also confirmed that ITGB1 is the target of miR-183.

## Conclusions

Our study revealed that hsa\_circ\_0000291 expression in GC cell lines was increased, and that hsa\_circ\_0000291 silencing suppressed GC cell metastasis and proliferation both in vivo and in vitro. Our results showed that hsa\_circ\_0000291 downregulation suppressed cell migration and proliferation by targeting the miR-183/ITGB1 axis. In conclusion, we identified hsa\_circ\_0000291 as a promising GC therapeutic target, therefore further investigation of the hsa\_circ\_0000291/miR-183/ITGB1 axis is required and may provide a foundation for developing novel potential therapeutic strategies towards GC treatment.

## Ethics Approval

The Animal Research Committee of Shanghai Tenth People's Hospital, Tongji University approved all experimental protocols and surgical procedures.

## Abbreviations

circRNAs, circular RNAs; GC, gastric cancer; ITGB1, integrin beta 1; miRNA, microRNA; CCK-8, Cell Counting Kit-8.

## Availability Of Data And Materials

All data analyzed and generated in this research report are enclosed in the article.

## Acknowledgment

National Natural Science Foundation of China (No. 81702942) funded this study.

## Author Contributions

CC and ML conceived the research and drafted the manuscript with feedback from all authors. SH, YY and YW carried out the experiments and analyzed the data. WL, LP and XZ conducted experiments and corrected the manuscript. All authors contributed to data analysis, drafting or revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Hartgrink HH, Jansen EP, van Grieken NC, van de Velde CJ. Gastric cancer. *Lancet*. 2009;374(9688):477–490. doi:10.1016/S0140-6736(09)60617-6
- Yagi K, Nozawa Y, Endou S, Nakamura A. Diagnosis of early gastric cancer by magnifying endoscopy with NBI from viewpoint of histological imaging: mucosal patterning in terms of white zone visibility and its relationship to histology. *Diagn Ther Endosc*. 2012;2012:954809. doi:10.1155/2012/954809



3. Patt YZ, Hassan MM, Lozano RD, et al. Phase II trial of cisplatin, interferon alpha-2b, doxorubicin, and 5-fluorouracil for biliary tract cancer. *Clin Cancer Res*. 2001;7(11):3375–3380.
4. Martin R, Jarnagin W. Intrahepatic cholangiocarcinoma. Current management. *Minerva Chir*. 2003;58(4):469–478.
5. Prensner JR, Chinnaiyan AM. The emergence of lncRNAs in cancer biology. *Cancer Discov*. 2011;1(5):391–407. doi:10.1158/2159-8290.CD-11-0209
6. Shi X, Sun M, Liu H, Yao Y, Song Y. Long non-coding RNAs: a new frontier in the study of human diseases. *Cancer Lett*. 2013;339(2):159–166. doi:10.1016/j.canlet.2013.06.013
7. Dang Y, Ouyang X, Zhang F, et al. Circular RNAs expression profiles in human gastric cancer. *Sci Rep*. 2017;7(1):9060. doi:10.1038/s41598-017-09076-6
8. Zhang J, Liu H, Hou L, et al. Circular RNA\_LARP4 inhibits cell proliferation and invasion of gastric cancer by sponging miR-424-5p and regulating LATS1 expression. *Mol Cancer*. 2017;16(1):151. doi:10.1186/s12943-017-0719-3
9. Huang M, He YR, Liang LC, Huang Q, Zhu ZQ. Circular RNA hsa\_circ\_0000745 may serve as a diagnostic marker for gastric cancer. *World J Gastroenterol*. 2017;23(34):6330–6338. doi:10.3748/wjg.v23.i34.6330
10. Qu S, Zhong Y, Shang R, et al. The emerging landscape of circular RNA in life processes. *RNA Biol*. 2017;14(8):992–999. doi:10.1080/15476286.2016.1220473
11. Qu S, Yang X, Li X, et al. Circular RNA: a new star of noncoding RNAs. *Cancer Lett*. 2015;365(2):141–148. doi:10.1016/j.canlet.2015.06.003
12. Wang L, Tong X, Zhou Z, et al. Circular RNA hsa\_circ\_0008305 (circPTK2) inhibits TGF-beta-induced epithelial-mesenchymal transition and metastasis by controlling TIF1gamma in non-small cell lung cancer. *Mol Cancer*. 2018;17(1):140. doi:10.1186/s12943-018-0889-7
13. Wu Z, Huang W, Wang X, et al. Circular RNA CEP128 acts as a sponge of miR-145-5p in promoting the bladder cancer progression via regulating SOX11. *Mol Med*. 2018;24(1):40. doi:10.1186/s10020-018-0039-0
14. Tan S, Sun D, Pu W, et al. Circular RNA F-circEA-2a derived from EML4-ALK fusion gene promotes cell migration and invasion in non-small cell lung cancer. *Mol Cancer*. 2018;17(1):138. doi:10.1186/s12943-018-0887-9
15. Ochendusko S, Puskulluoglu M, Konopka K, et al. Clinical effectiveness and toxicity of second-line irinotecan in advanced gastric and gastroesophageal junction adenocarcinoma: a single-center observational study. *Ther Adv Med Oncol*. 2017;9(4):223–233. doi:10.1177/1758834016689029
16. Benevento I, Bulzonetti N, De Felice F, Musio D, Vergine M, Tombolini V. The role of different adjuvant therapies in locally advanced gastric adenocarcinoma. *Oncotarget*. 2018;9(74):34022–34029. doi:10.18632/oncotarget.26106
17. Liu H, Liu Y, Bian Z, et al. Circular RNA YAP1 inhibits the proliferation and invasion of gastric cancer cells by regulating the miR-367-5p/p27 (Kip1) axis. *Mol Cancer*. 2018;17(1):151. doi:10.1186/s12943-018-0902-1
18. Xie F, Li Y, Wang M, et al. Circular RNA BCRC-3 suppresses bladder cancer proliferation through miR-182-5p/p27 axis. *Mol Cancer*. 2018;17(1):144. doi:10.1186/s12943-018-0892-z
19. Yan H, Sun BM, Zhang YY, et al. Upregulation of miR-183-5p is responsible for the promotion of apoptosis and inhibition of the epithelial-mesenchymal transition, proliferation, invasion and migration of human endometrial cancer cells by downregulating Ezrin. *Int J Mol Med*. 2018;42(5):2469–2480. doi:10.3892/ijmm.2018.3853
20. Chen D, Li SG, Chen JY, Xiao M. MiR-183 maintains canonical Wnt signaling activity and regulates growth and apoptosis in bladder cancer via targeting AXIN2. *Eur Rev Med Pharmacol Sci*. 2018;22(15):4828–4836. doi:10.26355/eurrev\_201808\_15618
21. Qiu L, Chang G, Bi Y, Liu X, Chen G. Circular RNA and mRNA profiling reveal competing endogenous RNA networks during avian leukosis virus, subgroup J-induced tumorigenesis in chickens. *PLoS One*. 2018;13(10):e0204931. doi:10.1371/journal.pone.0204931
22. Luo J, Zhang C, Zhan Q, et al. Profiling circRNA and miRNA of radiation-induced esophageal injury in a rat model. *Sci Rep*. 2018;8(1):14605. doi:10.1038/s41598-018-33038-1
23. Jin C, Bao J, Wang Y, et al. Changes in circRNA expression profiles related to the antagonistic effects of Escherichia coli F17 in lamb spleens. *Sci Rep*. 2018;8(1):14524. doi:10.1038/s41598-018-31719-5
24. Pan B, Guo J, Liao Q, Zhao Y. beta1 and beta3 integrins in breast, prostate and pancreatic cancer: a novel implication. *Oncol Lett*. 2018;15(4):5412–5416. doi:10.3892/ol.2018.8076
25. Qin Q, Wei F, Zhang J, Li B. miR-134 suppresses the migration and invasion of non-small cell lung cancer by targeting ITGB1. *Oncol Rep*. 2017;37(2):823–830. doi:10.3892/or.2017.5350

## Cancer Management and Research

### Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>

Dovepress

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.