Impact Of TP53 Gene Promoter Methylation On Chronic Lymphocytic Leukemia Pathogenesis And Progression

This article was published in the following Dove Press journal:
Journal of Blood Medicine

Waleed Haji Saeed¹
Adil Abozaid Eissa²
Adnan Anwar Al-Doski²

¹Department of Nursing, Akre Technical Institute, Duhok Polytechnic University, Duhok, Iraq; ²Department of Pathology, College of Medicine, University of Duhok, Duhok, Iraq

Background: Chronic lymphocytic leukemia (CLL) is a malignant lymphoid disorder that results from the overgrowth of mature-looking lymphoid cells in the blood and lymphatic tissue. Various clinical presentations have been attributed to the disease as a result of the different underlying genetic and epigenetic alterations. The current study has been initiated to study the role of an epigenetic alteration affecting the promoter of the *TP53* gene on CLL pathogenesis and progression.

Methods: The current study involved 54 newly diagnosed patients presenting with CLL as well as 30 normal individuals as controls. After obtaining verbal consent, data collection was done and the blood collected from all enrolled individuals for hematological investigations as well as for molecular categorization of *TP53* methylation status. Methylation-specific polymerase chain reaction (MS-PCR) technique was used to define the methylation status of the *TP53* gene promoter that encompasses DNA extraction, bisulfite conversion, conventional PCR amplification, running on agarose gel and documentation. Finally, statistical analysis was done to assess any correlation of the *TP53* epigenetic alteration to the disease etiology and the progression.

Results: In the current study, all controls and 42 of 54 patients show unmethylated *TP53* gene promoter; on the other hand, the methylated promoter was detected among 12 patients with a *p*-value of 0.001. *TP53* gene promoter methylation significantly linked to reduced platelet count (*p*-value of 0.047) and advanced stage at presentation (*p*-value of 0.076). No significant differences were seen among both methylated and unmethylated *TP53* promoters in relation to the age of the affected individuals, total white blood cell counts and hemoglobin level of the affected individuals.

Conclusion: The current study revealed a significant correlation of *TP53* gene promoter methylation to chronic lymphocytic leukemia pathogenesis and lower platelet counts.

Keywords: chronic lymphocytic leukemia, epigenetic alteration, *TP53* gene promoter

Background

Chronic lymphocytic leukemia (CLL) is a clonal disorder that results in the accumulation of morphologically mature-looking and immunologically incompetent lymphoid cells in the bone marrow, peripheral blood, and lymphatic tissues.^{1,2} Until recently, it was thought that CLL developed mainly from impaired apoptosis and evasion of cell death; however, it has been acknowledged that CLL-affected tissue comprising the bone marrow contains proliferation centers that support the top-up theory of the CLL cell population.³⁻⁵ CLL is a heterogeneous disease as some patients may progress rapidly, even though they may fail to respond to

Correspondence: Adil Abozaid Eissa
Department of Pathology, College of
Medicine, University of Duhok, 9 Azadi
Hospital Road, Duhok, Iraq
Tel +9647504576051
Email adilkh77@uod.ac

therapy, and others may remain stable for years without any intervention, and this mirrors the genetic configuration and the epigenetic modification of different genes that lead to disease development, stability, progression, and response to different chemotherapeutic agents.^{6,7}

Epigenetic modification may result from different strategies including DNA methylation/hypomethylation, histone alterations, and chromatin conversion. The epigenetic alteration has significant roles in controlling different genes' activities, genomic imprint, reproduction, embryonic development, inactivation of the X chromosome, tissue remodeling, as well as in disease pathogenesis, progression, and chemo-refractoriness.^{8–12} DNA methylation arises as a consequence of the addition of a methyl group to cytosine in the CpG dinucleotides that are characteristically seen as collections in small stretches of DNA termed “CpG islands” in the promoter regions of different housekeeping, the tumor suppressor, and some tissue-specific genes.^{13,14} Hypermethylation in the promoter regions through the action of DNA methyltransferases enzyme results in gene suppression and inactivation, while hypomethylation results in gene activation.^{8,15}

Protein 53 (p53), an end product for the *TP53* gene, plays a dynamic role in sustaining genetic stability and regulating the cellular response to DNA damage and other replicative stresses. Activation of p53 can ultimately lead to either DNA repair or programmed cell death, thus preventing cancer formation, i.e., it functions as a tumor suppressor and *TP53* is classified as a tumor suppressor gene.¹⁶ Inactivation of the *TP53* gene through different mutations like in Li-Fraumeni syndrome or hypo-activation through epigenetic alteration by methylation impairs apoptosis and may lead to many types of cancers, including CLL.¹⁷

The current study was initiated to estimate the effect of methylation of different genes functions and to correlate the *TP53* non-functioning (through methylation) with the pathogenesis and progression of CLL.

Methods

The current study represents a case-control study. A total of 54 newly diagnosed patients with CLL attending Nanakali hospital for blood diseases in Erbil, Iraq from January 1, 2017 to July 30, 2018 were enrolled. These patients were diagnosed as described by WHO criteria, which include absolute lymphocytosis (absolute lymphocytosis $>5 \times 10^9/L$) with typical immunological expression by flowcytometry.⁵ Also, 30 age- and sex-matched individuals with a negative history of previous “hematological and non-hematological malignancies” were

recruited for the purpose of the current study and used as controls. At first, verbal informed consent was obtained from all enrolled patients and the study was approved by the appropriate ethical committee at the University of Duhok College of Medicine and Duhok Directorate of Health, Duhok, Iraq and followed the Declaration of Helsinki ethical values for medical research concerning human beings.

Supportive information was collected from all enrolled individuals and then blood (also bone marrow sample was taken from patients) were collected in K2-EDTA tubes from each enrollee and all had their DNA extracted by a modified salting-out extraction method adopted by Iranpur-Mubarak and his colleague that yield high quantity with high purity DNA.^{18,19}

The methylation-specific polymerase chain reaction (MS-PCR) technique was used in the current study, which requires an initial conversion of DNA structure and then amplification using specific primer sets for methylated and unmethylated DNA targets.²⁰

At first, DNA bisulfite conversion is done with the aid of the EpiTect fast bisulfite conversion kit from Qiagen (Germany) and all steps were done according to the manufacturer's instructions. Then the converted DNA samples were amplified using M-P53-F 5'-TTCGGTAGGCGGATTATTG-3' and M-P53-R 5'-AAATATCCCCGAAACCC AAC-3' primers for the methylated *TP53* gene and U-P53-F 5'-TTGGTAGGTGGATTATTTGTTT-3' and U-P53-R 5'-CC AATCCAAAAAACATATCAC-3' primers for the unmethylated *TP53* gene.²¹ Initially a master mix was prepared using 2X Hot Start Master Mix from GeNet Bio (South Korea) and then amplification was performed using a PCR thermal cycler (2720) from Applied Biosystems, Hitachi (Japan). PCR conditions were as follows: for methylated *TP53* gene (94°C for 5 min, 35 cycles at 94°C for 30 s, 60°C for 40 s, 72°C for 60 s, 72°C for 7 min) and for unmethylated gene (94°C for 5 min, 35 cycles at 94°C for 30 s, 53°C for 40 s, 72°C for 60 s, 72°C for 7 min). Finally, the amplified products were subjected to 2% agarose gel electrophoresis. As shown in Figure 1, the methylated DNA sample produced a 193-bp DNA product with the M-P53 primers set, while the unmethylated DNA samples produced no band. On the other hand, the unmethylated DNA samples produced a 247-bp DNA product with the U-P53 primers set, while the methylated DNA samples produced no band. Appropriate methylated and unmethylated DNA controls were obtained from Qiagen [Epitect®Control DNA (Human), unmethylated and bisulfite converted (100)] and ran in parallel to patients' samples in all runs.

Figure 1 Gel electrophoresis of amplified products with U-P53 primers for determination of P53 methylation status. L: represents the 100-bp ladder; CU: unmethylated commercial P53 control DNA (247 bp); C-M: methylated commercial P53 control DNA; 1, 2, & 4 reveals unmethylated P53 DNA patient samples; and 3: represents methylated P53 DNA patient samples.

After data collection, data were analyzed using SPSS 24. Frequency and frequency percentages were used to describe the study sample. Student's *t*-test was used for continuous variables, and chi-square or Fisher exact test were used for categorical variables. *P*-value was considered to be significant if less than 0.05.

Results

A total of 54 patients with CLL were enrolled in the current study. Their ages ranged from 44 to 80 years with a median age of 62.2 ± 8.2 years. Of these, 37 (68.5%) were male and 17 (31.5%) were female, with a male to female ratio of 2.18.

After amplification with specific primers designed for determination of *TP53* gene methylation status, samples from 42 patients (77.8%) and all 30 controls (100%) show a 247-bp pattern with U-P53 (F&R) primers that is consistent with unmethylation status (Figure 1). On the other hand,

samples from 12 patients (22.2%) revealed no band with U-P53 (F&R) primers, but show a 193-bp band with M-P53 (F&R) primers, which is consistent with methylation status (Figure 2). The methylated status was seen among 17.6% of female patients (3 females) and only among 23.1% of male patients (9 males) with a *p*-value of 0.539.

Methylated P53 had been associated significantly with CLL groups in comparison to controls with a *p*-value of <0.001 . Using a *t*-test as shown in Table 1, the methylated status shows a significant correlation with low platelet count (*p*-value 0.047) as well as advanced stage (*p*-value 0.076); despite this, it does not show a significant correlation with hemoglobin level (*p*-value 0.167), total white blood cell count (*p*-value 0.144), and absolute lymphocytes count (*p*-value 0.425).

Discussion

The exact mechanism underlying CLL is still unclear; nevertheless, various molecular changes (genetic and epigenetic) have been implicated in CLL tumorigenesis and disease progression.^{1,22}

From these, *TP53* gene modification disturbs the process of apoptosis and results in prolonged survival of the malignant B cells in tissue and peripheral blood.²³ Also,

Figure 2 Gel electrophoresis of amplified products with M-P53 for determination of P53 methylation status. L: represents the 100-bp ladder; CM: methylated commercial P53 control DNA (193 bp); CU: unmethylated commercial P53 control DNA; 1 & 2 reveal methylated P53 DNA patient samples; and 3: represents unmethylated P53 DNA patient samples.

Table I Correlation Of *TP53* Gene Methylation Status To Hematological Data And Staging Categories

Parameters	Different Categories	Methylated		Un-Methylated		Total	P value
		No.	%	No.	%		
Hemoglobin level (g/dl)	>10.0	9	16.67	38	70.37	47 (87.04%)	0.167
	Hb < 10.0	3	5.55	4	7.41	7 (12.96%)	
Platelets (*10 ⁹ /L)	>100	7	12.96	37	68.52	44 (81.48)	0.047
	< 100	5	9.26	5	9.26	10 (18.52%)	
WBC count (*10 ⁹ /L)	< 50	10	18.53	40	74.07	50 (92.59%)	0.144
	>50	2	3.70	2	3.70	4 (7.41%)	
Lymphocytes count (*10 ⁹ /L)	5–20	5	9.3	21	38.9	26 (48.15%)	0.425
	> 20	7	1.9	21	3.7	28 (51.85%)	
Stage	A	5	9.3	24	44.4	29 (53.70%)	0.076
	B	2	3.7	13	24	15 (27.78%)	
	C	5	9.3	5	9.3	10 (18.52%)	

TP53 inactivation had been criticized for fludarabine resistance in 30–40% of patients with CLL, the backbone drug for this leukemia. Only a quarter of cases had contributed to *TP53* gene deletion and the remaining three-fourths had contributed to gene inactivation through methylation of the promoter region of the gene.^{24,25} As the methylation results in gene inactivation and in accordance with the previous review written by Wang and Wang, the current study revealed a significant association of *TP53* gene promoter methylation with the CLL development.²⁶ Not only the pathogenesis but also the advanced stage (low platelets) had been associated significantly with *TP53* gene suppression as through promoter methylation and this poor prognosis was in accordance with the previous study conducted by Zenz and his coworkers.^{23,25} On the other hand, methylation of the *TP53* gene promoter was not significantly associated with low hemoglobin level or high total white cells counts, and this may explain the heterogeneous nature of CLL and its correlation with other gene malfunctions. The various effect of *TP53* methylation on platelet count, hemoglobin level, and total white cell count may represent the different effects of *TP53* dysfunction on the micro-environment and interaction of different genes with the hemopoietic proliferation and maturation.^{27,28}

Epigenetic modification of promoters of different genes contributes to gene activation or suppression as a result of hypo- or hypermethylation, respectively, and targeting this mechanism, e.g., with 5-aza-2'-deoxycytidine (5-aza-dC) that upregulates expression of *TP53* mRNA and protein, may be valuable in fighting CLL in the future as such drugs have been used extensively in myelodysplastic syndrome with excess

blasts that show a characteristically high ratio of hypermethylation and different gene inactivation.²⁹

Follow-up of the patients involved in the current study has been initiated to determine the effect of the methylation on the progression of the disease and the response to various chemotherapeutic agents.

Conclusion

The current study revealed that *TP53* methylation contributes significantly to CLL development and progression. Further workups are recommended to study their relation with other genetic changes as malignancies are multifactorial and heterogeneous that arise from the interaction of different genetic changes.

Abbreviations

CLL, chronic lymphocytic leukemia; *TP53*, tumor protein 53; WBC, white blood cells; K2-EDTA, dipotassium-ethylenediaminetetraacetic acid; PCR, polymerase chain reaction; CPG, cytosine–guanine dinucleotide.

Ethics Approval And Consent To Participate

The current study was approved by the ethical committee at the University of Duhok College of Medicine and Duhok Directorate of Health and followed the Declaration of Helsinki ethical values for medical research concerning human beings. As the current study does not include any therapeutic modality and has a minor threat of injury to enrolled subjects, only verbal informed consent was taken

from all enrolled individuals according to the general guidelines of Duhok Directorate of Health for obtaining consent.

Disclosure

The authors declare that they have no competing interests in this work.

References

- Gaidano G, Foà R, Dalla-Favera R. Molecular pathogenesis of chronic lymphocytic leukemia. *J Clin Invest*. 2012;122(10):3432–3438. doi:10.1172/JCI64101
- Ghia P, Ferreri AM, Caligaris-Cappio F. Chronic lymphocytic leukemia. *Crit Rev Oncol Hematol*. 2007;64(3):234–246. PMID: 17544290. doi:10.1016/j.critrevonc.2007.04.008.
- Ponzoni M, Doglioni C, Caligaris-Cappio F. Chronic lymphocytic leukemia: the pathologist's view of lymph node microenvironment. *Semin Diagn Pathol*. 2011;28(2):161–166. doi:10.1146/annurev-pathol-020712-163955
- Damle RN, Calissano C, Chiorazzi N. Chronic lymphocytic leukaemia: a disease of activated monoclonal B cells. *Best Pract Res Clin Haematol*. 2010;23(1):33–45. doi:10.1016/j.beha.2010.02.001
- Campo E, Ghia P, Montserrat E, et al. Chronic lymphocytic leukaemia/small lymphocytic lymphoma. In: Swerdlow SH, Campo E, Harris N, et al., editors. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*. Revised 4th ed. Lyon: International Agency for Research on Cancer; 2017:216–221.
- Zenz T, Mertens D, Kuppers R, Dohner H, Stilgenbauer S. From pathogenesis to treatment of chronic lymphocytic leukaemia. *Nat Rev Cancer*. 2010;10:37–50. doi:10.1038/nrc2764
- Zenz T, Gribben JG, Hallek M, Ohner H, Keating MJ, Stilgenbauer S. Risk categories and refractory CLL in the era of chemoimmunotherapy. *Blood*. 2012;119:4101–4107. doi:10.1182/blood-2011-11-312421
- Lee CJ, Evans J, Kim K, Chae H, Kim S. Determining the effect of DNA methylation on gene expression in cancer cells. *Methods Mol Biol*. 2014;1101:161–178. PMID: 24233782. doi:10.1007/978-1-62703-721-1_9
- Cheedipudi S, Genolet O, Dobrev G. Epigenetic inheritance of cell fates during embryonic development. *Front Genet*. 2014;5:19. doi:10.3389/fgene.2014.00019
- Tang W, Ho S-M. Epigenetic reprogramming and imprinting in origins of disease. *Rev Endocr Metab Disord*. 2007;8(2):173–182. doi:10.1007/s11154-007-9042-4
- Hagood J. Beyond the genome: epigenetic mechanisms in lung remodeling. *Physiology (Bethesda)*. 2014;29(3):177–185. doi:10.1152/physiol.00048.2013
- Ross SA, Milner J. Epigenetic modulation and cancer: effect of metabolic syndrome? *Am J Clin Nutr*. 2007;86(3):872S–877S. doi:10.1093/ajcn/86.3.872S
- Robertson KD, Wolffe AP. DNA methylation in health and disease. *Nat Rev Genet*. 2000;1:11–19. doi:10.1038/35049533
- Hube F, Reverdiau P, Iochmann S, Gruel Y. Improved PCR method for amplification of GC-rich DNA sequences. *Mol Biotechnol*. 2005;31:81–84. doi:10.1385/MB:31:1:081
- Esteller M. CpG island hypermethylation and tumor suppressor genes: a booming present, a brighter future. *Oncogene*. 2002;21:5427–5440. doi:10.1038/sj.onc.1205600
- Wang L, Kwong Y, Wong K, et al. Epigenetic inactivation of mir-34b/c in addition to mir-34a and DAPK1 in chronic lymphocytic leukemia. *J Transl Med*. 2014;12:52. PMID: PMC3941938; PMID: 24559316. doi:10.1186/1479-5876-12-52.
- Hollstein M, Sidransky D, Vogelstein B, Harris C. p53 mutations in human cancers. *Science*. 1991;253(5015):49–53. PMID 1905840. doi:10.1126/science.1905840.
- Iranpur-Mubarak V, Esmailzadeh AK. Rapid extraction of high quality DNA from whole blood stored at 4 °C for long period; 2010. Available from: <http://www.protocol-online.org/prot/Protocols/Rapid-Extraction-of-High-Quality-DNA-from-Whole-Blood-Stored-at-4-C-for-Long-Period-4175.html>. Accessed March 14, 2018.
- Kashmoola M, Eissa A, Al-Takay D, Al-Allawi N. Molecular characterization of G6PD deficient variants in Nineveh province, Northwestern Iraq. *Indian J Hematol Blood Transfus*. 2015;3:133–136. doi:10.1007/s12288-014-0368-2
- Galm O, Herman J. Methylation-specific polymerase chain reaction. *Methods Mol Med*. 2005;113:279–291. PMID:15968111. doi:10.1385/1-59259-916-8:279
- Jarzynski A, Papiernik K, Polz-Dacewicz M. Analysis of mutation and promoter methylation of TP53 gene in tumors of the head and neck. *Curr Issues Pharm Med Sci*. 2016;29(2):53–56. doi:10.1515/cipms-2016-0011
- Fernández V, Jares P, Salaverria I, et al. Gene expression profile and genomic changes in disease progression of early-stage chronic lymphocytic leukemia. *Haematologica*. 2008;93(1):132–136. doi:10.3324/haematol.11694
- Zenz T, Eichhorst B, Busch R, et al. TP53 mutation and survival in chronic lymphocytic leukemia. *J Clin Oncol*. 2010;28(29):4473–4479. doi:10.1200/JCO.2009.27.8762
- Zenz T, Kröber A, Scherer K, et al. Monoallelic TP53 inactivation is associated with poor prognosis in chronic lymphocytic leukemia: results from a detailed genetic characterization with long-term follow-up. *Blood*. 2008;112:3322–3329. doi:10.1182/blood-2008-04-154070
- Stilgenbauer S, Zenz T. Understanding and managing ultra high-risk chronic lymphocytic leukemia. *Hematology Am Soc Hematol Educ Program*. 2010;2010:481–488. doi:10.1182/asheducation-2010.1.481
- Wang C, Wang X. The role of TP53 network in the pathogenesis of chronic lymphocytic leukemia. *Int J Clin Exp Pathol*. 2013;6(7):1223–1229. ISSN:1936-2625/IJCEP1304037.
- Gaiti F, Chaligne R, Gu H, et al. Epigenetic evolution and lineage histories of chronic lymphocytic leukaemia. *Nature*. 2019;569:576–580. doi:10.1038/s41586-019-1198-z
- Hudson J, Shoaibi M, Maestro R, Carnero A, Hannon G, Beach D. A proinflammatory cytokine inhibits p53 tumor suppressor activity. *J Exp Med*. 1999;190(10):1375–1382. doi:10.1084/jem.190.10.1375
- Schroeder T, Rautenberg C, Haas R, Germing U, Kobbe G. Hypomethylating agents for treatment and prevention of relapse after allogeneic blood stem cell transplantation. *Int J Hematol*. 2018;107(2):138–150. doi:10.1007/s12185-017-2364-4

Journal of Blood Medicine**Dovepress****Publish your work in this journal**

The Journal of Blood Medicine is an international, peer-reviewed, open access, online journal publishing laboratory, experimental and clinical aspects of all aspect pertaining to blood based medicine including but not limited to: Transfusion Medicine; Blood collection, Donor issues, Transmittable diseases, and Blood banking logistics; Immunohematology; Artificial and alternative blood based

therapeutics; Hematology; Biotechnology/nanotechnology of blood related medicine; Legal aspects of blood medicine; Historical perspectives. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/journal-of-blood-medicine-journal>