Long Noncoding RNA KCNQ1OT1 Promotes the Progression of Non-Small Cell Lung Cancer via Regulating miR-204-5p/ATG3 Axis

This article was published in the following Dove Press journal:
OncoTargets and Therapy

Yan Kang¹
Yaoli Jia²
Qilong Wang¹
Qianru Zhao³
Meng Song⁴
Ran Ni¹
Jing Wang¹

¹Department Two of Respiratory Medicine, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, Henan, People's Republic of China;

²Department of Respiratory Medicine, Xuchang Central Hospital, Xuchang, Henan, People's Republic of China;

³Department of Geriatric Medicine, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, Henan, People's Republic of China; ⁴Department of Epidemiology and Health Statistics, College of Public Health, Zhengzhou University, Zhengzhou, Henan, People's Republic of China

Purpose: Non-small cell lung cancer (NSCLC) is the first leading cause of cancer-related death globally. Long noncoding RNA KCNQ1 overlapping transcript 1 (KCNQ1OT1) was involved in the progression of multiple cancers by sponging target miRNA. We aimed to explore the pathological mechanism of KCNQ1OT1 in NSCLC progression.

Methods: The expression of KCNQ1OT1, miR-204-5p and autophagy-related gene 3 (ATG3) was measured by quantitative real-time polymerase chain reaction (qRT-PCR). 3-(4, 5-Dimethyl-2-thiazolyl)-2, 5-diphenyl-2-H-tetrazolium bromide (MTT) assay and flow cytometry assay were conducted for the detection of cell proliferation and apoptosis, respectively. Western blot assay was performed to examine the protein levels of B-cell lymphoma-2 (BCL-2), BCL2-Associated X (Bax), cleaved caspase-3, cleaved caspase-9 and LC3 II/LC3 I and P62. The interaction between miR-204-5p and KCNQ1OT1 or ATG3 was validated by dual-luciferase reporter system and RNA immunoprecipitation (RIP) assay. Murine xenograft assay was conducted to explore the function of KCNQ1OT1 in vivo. Immunohistochemistry (IHC) staining assay was used for the analysis of ki67-positive cell percentage.

Results: The expression of KCNQ1OT1 and ATG3 was up-regulated whereas miR-204-5p was down-regulated in NSCLC tumors and cells. MiR-204-5p was inversely correlated with KCNQ1OT1 or ATG3. In addition, KCNQ1OT1 knockdown facilitated apoptosis, inhibited autophagy and proliferation of NSCLC cells in vitro and blocked tumor growth in vivo. However, the miR-204-5p inhibitor reversed the effects. More importantly, ATG3 was a target gene of miR-204-5p and ATG3 overexpression restored the effect of miR-204-5p on NSCLC cell progression.

Conclusion: KCNQ1OT1 promotes cell proliferation and autophagy and inhibits cell apoptosis via regulating miR-204-5p/ATG3 axis, providing a promising target for NSCLC therapy.

Keywords: KCNQ1OT1, miR-204-5p, ATG3, NSCLC

Introduction

Non-small cell lung cancer (NSCLC) which accounts for more than 80% of lung cancer is the first leading cause of cancer-related death globally.^{1,2} Despite advanced treatment strategies, including surgical resection, chemotherapy, radiotherapy, immunotherapy and combined therapy, have palliated the symptoms, the 5-year overall survival rate is still less than 20% in patients with NSCLC.³⁻⁵ In addition, chemo-resistance and high metastasis of NSCLC vitiated the therapy outcomes.⁶ Therefore, it is urgently needed to elucidate the physiological and pathological mechanism of NSCLC.

Correspondence: Ran Ni; Jing Wang
Department Two of Respiratory Medicine, The First Affiliated Hospital of Zhengzhou University, No. 1 Jianshe East Road, Zhengzhou 450052, Henan, People's Republic of China
Tel +86 13633860030; +86 13838352886
Email anxian218911qco@126.com; shijin199410ut@126.com

Long non-coding RNAs (lncRNAs) are fundamental modulators of cell cycle, proliferation, migration, differentiation, invasion, metastasis, apoptosis, inflammation and autophagy in multiple cancers.^{7–9} LncRNA KCNQ1 overlapping transcript 1 (KCNQ1OT1), mapped in human KCNQ1 locus with a length of 91 kb, plays essential roles in tumor proliferation, metabolism, epithelial–mesenchymal transition (EMT) and growth via functioning as competing endogenous RNA (ceRNA).^{10,11} KCNQ1OT1 mainly acts as an oncogene in different cancers. For instance, KCNQ1OT1 facilitated proliferation, migration and invasion of cholangiocarcinoma cells by interacting with miR-140-5p.¹² Besides, KCNQ1OT1 was reported to improve chemoresistance of oxaliplatin against hepatocellular carcinoma by up-regulating ABCC1 via sponging miR-7-5p.¹³ Though a previous study has demonstrated that KCNQ1OT1 could be an oncogene in NSCLC,¹⁴ the molecular mechanism of KCNQ1OT1 in NSCLC remains obscure.

MicroRNAs (miRNAs) are evolutionarily conserved small non-coding RNAs with 19–23 nucleotides in length.^{15,16} Despite without protein coding capacity, miRNAs are widely involved in the regulation of gene expression at the post-transcriptional level by complementarily binding with messenger RNAs (mRNAs).^{15–19} Ectopic expression of miRNA is closely associated with tumorigenesis and development.²⁰ As a tumor suppressor, miR-204-5p is a pivotal regulator in multiple cancers. For example, miR-204-5p suppressed cell invasion and metastasis by targeting forkhead box C1 in laryngeal squamous cell carcinoma.²¹ Similarly, miR-204-5p attenuated proliferation, migration, invasion and metastasis of hepatocellular carcinoma and oral squamous cell carcinoma though repressing CXCR4 and SIRT1 expression, respectively.^{22,23} Autophagy-related gene 3 (ATG3) could be targeted by several miRNAs to participate in the regulation of NSCLC.^{24,25} However, whether miR-204-5p can target ATG3 in NSCLC requires deep investigation.

In the present study, we attempted to investigate the underlying mechanism of the KCNQ1OT1/miR-204-5p/ATG3 axis in NSCLC cell progression. All the results indicated that KCNQ1OT1 promotes proliferation and autophagy while suppresses apoptosis by regulating the miR-204-5p/ATG3 axis. This study might provide promising biomarkers for the diagnosis and treatment of NSCLC.

Materials and Methods

Tissue Samples

Fresh NSCLC tumor tissues and the corresponding normal tissues were obtained from 35 NSCLC patients recruited from The First Affiliated Hospital of Zhengzhou University. The patients had not received preoperative treatment. All NSCLC patients have signed written informed consents and the experiment protocols were approved by the Ethics Committee of The First Affiliated Hospital of Zhengzhou University.

Cell Transfection

NSCLC cell lines HCC827, H1299, A549, H460 and human bronchial epithelial cell BEAS-2B were purchased from the American Type Culture Collection (ATCC, Manassas, VA, USA). All the cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (FBS; Gibco) and 0.05% penicillin/streptomycin (Gibco). Small hairpin RNA (shRNA) targeting KCNQ1OT1 (sh-KCNQ1OT1), shRNA negative control (sh-NC), pcDNA, and pcDNA-ATG3 overexpression vector (ATG3) were synthesized by Genepharma (Shanghai, China). The miRNA mimics (miR-204-5p), miR-204-5p inhibitor (anti-miR-204-5p), miRNA negative control (miR-NC) and miRNA negative control inhibitor (anti-miR-NC) were purchased from RIBOBIO (Guangzhou, China). The synthetic oligonucleotides or plasmids were transfected in A549 and H460 cells using Lipofectamine 2000 (Invitrogen, Carlsbad, CA, USA).

Quantitative Real-Time Polymerase Chain Reaction (qRT-PCR)

NSCLC tissues and cells were incubated with TRIzol reagent (Invitrogen) to extract total RNA. The cDNA for KCNQ1OT1, miR-204-5p and ATG3 was synthesized by All-in-One™ First-Strand cDNA Synthesis Kit (FulGen, Guangzhou, China). Then, qRT-PCR was performed using SYBR green (Applied Biosystems, Foster City, CA, USA) following the standard procedures. The primers for KCNQ1OT1, miR-204-5p, ATG3, GAPDH and U6 were listed as follows: KCNQ1OT1 (Forward, 5'-TTGGT AGGATTTTGTGAGG-3'; Reverse, 5'-CAACCTTCCC CTACTACC-3'); miR-204-5p (Forward, 5'-CGAAGTT CCCTTTGTCATCCT-3'; Reverse, 5'-GTGCAGGGTCCG AGGTATTC-3'); ATG3 (Forward, 5'-GCAAACAAGAA CCTATGACCTG-3'; Reverse, 5'-GTCTTCATACATGTG CTCAACTG-3'); GAPDH, (Forward, 5'-AGGTCGGTGT GAACGGATTTG-3'; Reverse, 5'-GGGGTCGTTGATG

GCAACA-3'); U6, (Forward, 5'-ACCCTGAGAAATACC CTCACAT-3'; Reverse, 5'-GACGACTGAGCCCCCTGA TG-3').

3-(4, 5-Dimethyl-2-Thiazolyl)-2, 5-Diphenyl-2-H-Tetrazolium Bromide (MTT) Assay and Flow Cytometry Assay

Cell viability and apoptosis were evaluated by MTT assay and flow cytometry assay, respectively. For MTT assay, transfected A549 and H460 cells (5000 cells/well) were seeded on 96-well plates for 24 h, 48 h, 72 h and 96 h. Then, the cells were reacted with 10 μ L MTT reagent (Beyotime, Shanghai, China) and kept for an additional 4 h. The OD value at 490 nm was measured by a spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). For flow cytometry assay, transfected A549 and H460 cells were collected, stained using Annexin V-FITC/PI Apoptosis Detection Kit (Vazyme, Nanjing, China) and analyzed by BD FACS Canto II flow cytometry (BD Biosciences, Franklin Lakes, NJ, USA).

Dual-Luciferase Reporter Assay

The sequences of KCNQ10T1 and 3'-UTR of ATG3 including the wild-type or mutant binding sequences of miR-204-5p were cloned and introduced into pmiR-REPORTTM vectors (RiboBio, Guangzhou, China) to generate luciferase reporter vectors WT-KCNQ10T1, MUT-KCNQ10T1, ATG3 3'UTR-WT and ATG3 3'UTR-MUT, respectively. Then, miR-204-5p or miR-NC was transfected into A549 and H460 cells together with indicated luciferase reporter vector using Lipofectamine 2000 transfection reagent (Invitrogen). Dual-Luciferase Reporter Assay Kit (Vazyme) was utilized for the detection of luciferase activity after 48 h.

RNA Immunoprecipitation (RIP) Assay

Magna RNA immunoprecipitation kit (Millipore, Bedford, MA, USA) was employed for RIP assay. In brief, A549 and H460 cells transfected with KCNQ10T1 and negative control (NC) were lysed by RIP buffer. Then, cell lysis was incubated with magnetic beads coated with Anti-Argonaute2 (Anti-Ago2; Abcam, Cambridge, MA, USA) or immunoglobulin G (Anti-IgG; Abcam). The enrichment of miR-204-5p was subjected to qRT-PCR after RNAs in the magnetic beads were purified.

Western Blot Assay

Western blot assay was performed following the standard protocol. In brief, after total protein was isolated and quantified, the proteins were separated by sodium dodecyl sulfonate-polyacrylamide gel (SDS-PAGE; Solarbio, Beijing, China) and then transferred onto polyvinylidene difluoride membranes (PVDF; Pall Corporation, New York, NYC, USA). Next, the membranes were blocked in skim milk for 2 h and then incubated with primary antibodies against ATG3 (ab108251; 1:2000; Abcam, Cambridge, MA, USA), LC3 II/LC3 I (ab128025; 1:200; Abcam), P62 (ab56416; 1:500; Abcam), B-cell lymphoma-2 (BCL-2; ab196495; 1:1000; Abcam), cleaved caspase-3 (ab49822; 1:500; Abcam), cleaved caspase-9 (ab2324; 1:2000; Abcam), BCL2-Associated X (Bax; ab182733; 1:2000; Abcam) or GAPDH (ab181602; 1:5000; Abcam) at 4°C overnight followed by incubation with HRP-conjugated secondary antibody (D110150; 1:5000; Sangon, Shanghai, China) for 2 h. The protein bands were analyzed using an enhanced chemiluminescence reagent (Beyotime).

Murine Xenograft Assay

Male nude mice of 5 weeks old were purchased from Vital River Laboratory Animal Technology (Beijing, China). Our animal experiment protocols were approved by the Ethics Committee of The First Affiliated Hospital of Zhengzhou University and performed according to the NIH guidelines for the care and use of laboratory animals (NIH Publication No. 85-23 Rev.1985). In brief, A549 and H460 cells stably transfected with sh-KCNQ10T1 and sh-NC were subcutaneously injected in mice to establish xenograft mice model. Tumor volume was measured every 4 d from day 8 and calculated using the formula: (length $\times$ width²)/2. Tumor weight was measured after 28 d and tumors were harvested and preserved at -80°C.

Immunohistochemistry (IHC) Staining Assay

The tissue sections were deparaffinized, rehydrated using a graded ethanol series and heated in citrate buffer for antigen retrieval. Then, the sections were washed, blocked and incubated with normal goat serum. Then, the samples were incubated with a primary antibody anti-Ki67 (ab15580; 1:500; Abcam) at 4°C overnight and secondary antibody (ab205718; 1:5000; Abcam). The sections were stained

with diaminobenzidine (DAB) and then counterstained with hematoxylin, dehydrated and mounted. At last, the section was photographed using a digital microscope camera (Nikon, Tokyo, Japan).

Statistical Analysis

All the experiments were conducted at least in triplicate and data were presented as means $\pm$ standard deviation (SD). Statistical analysis was carried out using SPSS 13.0 software (SPSS, Chicago, IL, USA) and GraphPad Prism 7 (GraphPad Inc., San Diego, CA, USA). The correlation between miR-204-5p and KCNQ1OT1 or ATG3 was analyzed by Pearson's correlation coefficient analysis. *P* value less than 0.05 ($P < 0.05$) was considered as statistically significant.

Results

KCNQ1OT1 Depletion Induces Apoptosis and Suppresses Proliferation and Autophagy in NSCLC Cells

The functions of KCNQ1OT1 on NSCLC cell proliferation, apoptosis and autophagy were assessed by MTT assay, flow cytometry analysis and Western blot assay, respectively. As illustrated in Figure 1A and B, KCNQ1OT1 expression was extremely higher in NSCLC tumor tissues than that in the corresponding normal tissues. Likewise, KCNQ1OT1 expression was up-regulated in NSCLC cell lines (HCC827, H1299, A549, H460) compared with human bronchial epithelial cell BEAS-2B (Figure 1C). Furthermore,

Figure 1 KCNQ1OT1 knockdown repressed proliferation and autophagy and induced apoptosis in NSCLC. (A, B) KCNQ1OT1 expression in 35 pairs of NSCLC tumor tissues and normal tissues. (C) KCNQ1OT1 expression in NSCLC cell lines (HCC827, H1299, A549, H460) and human bronchial epithelial cell BEAS-2B. (D–K) A549 and H460 cells were stably transfected with sh-KCNQ1OT1 or sh-NC. (D) KCNQ1OT1 expression in stably transfected A549 and H460 cells. (E, F) Cell viability of transfected A549 (E) and H460 cells (F). (G) Cell apoptosis of transfected A549 and H460 cells. (H, I) The expression of apoptosis-related protein cleaved caspase-3, cleaved caspase-9, Bax and anti-apoptosis protein BCL-2 in transfected A549 (H) and H460 cells (I). (J, K) Protein expression of autophagy markers LC3 and P62 in transfected A549 (J) and H460 cells (K). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

loss-of-function experiments were employed by knocking down KCNQ1OT1 to explore the regulatory effects of KCNQ1OT1 on NSCLC cell progression. An obvious reduction of KCNQ1OT1 expression was noticed in A549 and H460 cells stably transfected with sh-KCNQ1OT1, indicating the transfection efficiency was relatively high (Figure 1D). Moreover, cell growth was inhibited apparently in NSCLC cells after KCNQ1OT1 silencing (Figure 1E and F). On the contrary, the cell apoptosis rate was enhanced in the sh-KCNQ1OT1 transfected group compared with sh-NC group (Figure 1G). Hence, the expression of apoptosis-related proteins was measured. We found that Bax, cleaved caspase-3 and cleaved caspase-9 were dramatically elevated whereas anti-apoptosis protein BCL-2 was decreased in both A549 and H460 cells stably transfected with sh-KCNQ1OT1 (Figure 1H and I). We also analyzed the expression of autophagy markers LC3 and P62 and observed that deficiency of KCNQ1OT1 repressed LC3II/LC3I expression and boosted P62 expression (Figure 1J and K). Collectively, KCNQ1OT1

knockdown induced apoptosis and suppressed proliferation and autophagy in NSCLC cells.

KCNQ1OT1 is a Sponge of miR-204-5p

Growing evidence has validated that lncRNA KCNQ1OT1 exerts its function by sponging the target miRNA. As searched by the online prediction tool StarBase v2.0, miR-204-5p includes the binding sites of KCNQ1OT1 (Figure 2A). To confirm the prediction, wild type (WT-KCNQ1OT1) and mutant type (MUT-KCNQ1OT1) vectors were constructed and co-transfected into A549 and H460 cells with miR-204-5p or miR-NC to establish dual-luciferase reporter system. Luciferase activity was reduced evidently in NSCLC cells co-transfected with WT-KCNQ1OT1 and miR-204-5p compared to WT-KCNQ1OT1 and miR-NC co-transfected cells, while luciferase activity in MUT-KCNQ1OT1 transfection group remained unchanged (Figure 2B and C). Additionally, the RIP assay indicated that the enrichment of miR-204-5p in the Anti-Ago2 RIP group was elevated in NSCLC cells transfected

Figure 2 KCNQ1OT1 directly interacted with miR-204-5p. (A) The putative binding sites between KCNQ1OT1 and miR-204-5p. (B, C) Luciferase activity of A549 (B) and H460 cells (C) co-transfected with WT-KCNQ1OT1 or MUT-KCNQ1OT1 and miR-204-5p or miR-NC. (D) The enrichment of miR-204-5p in Anti-Ago2 immunoprecipitation complex in A549 and H460 cells transfected with KCNQ1OT1 or pcDNA. (E, F) The expression of miR-204-5p in A549 (E) and H460 cells (F) transfected with KCNQ1OT1, sh-KCNQ1OT1, pcDNA or sh-NC. (G, H) The expression of miR-204-5p in NSCLC tumor tissues and normal tissues. (I) The expression of miR-204-5p in NSCLC cell lines (HCC827, H1299, A549, H460) and human bronchial epithelial cell BEAS-2B. (J) The correlation between expression of KCNQ1OT1 and miR-204-5p ($r = -0.7785$, $P < 0.0001$). * $P < 0.05$, *** $P < 0.001$.

with KCNQ1OT1 (Figure 2D). More importantly, overexpression of KCNQ1OT1 repressed miR-204-5p expression whereas depletion of KCNQ1OT1 facilitated miR-204-5p expression (Figure 2E and F). Subsequently, we discovered that the expression of miR-204-5p was down-regulated in NSCLC tumor tissues and cell lines compared with normal tissues (Figure 2G and H) and cells (Figure 2I). Pearson's correlation coefficient analysis exhibited that KCNQ1OT1 was negatively correlated with miR-204-5p ($r=-0.7785$, $P<0.0001$) in NSCLC tissues (Figure 2J). These findings demonstrated that KCNQ1OT1 acted as a sponge of miR-204-5p.

KCNQ1OT1 Regulates Proliferation, Apoptosis and Autophagy by Binding to miR-204-5p in NSCLC Cells

To explore whether KCNQ1OT1 could regulate NSCLC progression by binding to miR-204-5p, A549 and H460

cells were transfected with sh-KCNQ1OT1+anti-miR-204-5p, sh-KCNQ1OT1+anti-miR-NC, sh-KCNQ1OT1 or sh-NC. As exhibited in Figure 3A, all the vectors were transfected into A549 and H460 cells successfully with high transfection efficiency. MTT results revealed that miR-204-5p inhibitor could abrogate KCNQ1OT1 silencing induced inhibition on cell proliferation in NSCLC (Figure 3B and C). Conversely, the KCNQ1OT1 knockdown promoted the apoptosis of A549 and H460 cells and miR-204-5p inhibitor rescued the effect (Figure 3D). Consistently, KCNQ1OT1 knockdown enhanced the expression of apoptosis-related protein Bax, cleaved caspase-3 and cleaved caspase-9, and reduced the expression of anti-apoptosis protein BCL-2. However, the miR-204-5p inhibitor reversed the trends (Figure 3E and F). Furthermore, the level of LC3II/LC3I was decreased whereas P62 was increased after KCNQ1OT1 silencing; however, the miR-204-5p inhibitor transfected group overturned the effects (Figure 3G and H). All the data implicated that KCNQ1OT1 regulated

Figure 3 miR-204-5p inhibitor attenuated the effects of KCNQ1OT1 silencing on cell proliferation, apoptosis and autophagy in NSCLC. A549 and H460 cells were transfected with sh-KCNQ1OT1+anti-miR-204-5p, sh-KCNQ1OT1, sh-KCNQ1OT1+anti-miR-NC or sh-NC. (A) The expression of miR-204-5p in transfected A549 and H460 cells. (B, C) Cell viability of transfected A549 and H460 cells. (D) Cell apoptosis of transfected A549 and H460 cells. (E, F) The expression of apoptosis-related protein cleaved caspase-3, cleaved caspase-9, Bax and anti-apoptosis protein BCL-2 in transfected A549 (E) and H460 cells (F). (G, H) Protein expression of autophagy markers LC3 and P62 in transfected A549 (G) and H460 cells (H). * $P<0.05$.

Figure 4 ATG3 is a target of miR-204-5p. (A) The putative binding sites between ATG3 and miR-204-5p. (B, C) Luciferase activity of A549 (B) and H460 cells (C) co-transfected with ATG3 3'UTR-WT or ATG3 3'UTR-MUT and miR-204-5p or miR-NC. (D, E) Protein expression of ATG3 in A549 (D) and H460 cells (E) transfected with miR-204-5p, anti-miR-204-5p, anti-miR-NC or miR-NC. (F, G) The expression of ATG3 in NSCLC tumor tissues and normal tissues. (H) The correlation between ATG3 and miR-204-5p ($r = -0.6014$, $P < 0.0001$). (I, J) The expression of ATG3 mRNA (I) and protein (J) in NSCLC cell lines (HCC827, H1299, A549, H460) and human bronchial epithelial cells BEAS-2B. * $P < 0.05$, *** $P < 0.001$.

proliferation, apoptosis and autophagy by interacting with miR-204-5p in NSCLC cells.

ATG3 is a Target of miR-204-5p

Based on bioinformatics analysis using StarBase v2.0, miR-204-5p comprised the binding sites of 3'UTR ATG3 (Figure 4A). Dual-luciferase reporter assay exhibited that luciferase activity was relatively lower in NSCLC cells co-transfected with ATG3 3'UTR-WT and miR-204-5p than ATG3 3'UTR-WT and miR-NC co-transfected group, confirming the interaction between ATG3 and miR-204-5p (Figure 4B and C). Besides, ATG3 protein expression was reduced by the introduction of miR-204-5p while enhanced by miR-204-5p inhibitor in A549 and H460 cells (Figure 4D and E). As expected, ATG3 expression was up-regulated in NSCLC tumor tissues compared with normal tissues (Figure 4F and G). Then, we found ATG3 expression was negatively correlated with miR-204-5p expression in NSCLC tissues, as illustrated by Person's correlation coefficient analysis ($r = -0.6014$, $P < 0.0001$) (Figure 4H). Similarly, the mRNA and protein levels of ATG3 were considerably higher in NSCLC cell lines (HCC827, H1299, A549, H460) than those in BEAS-

2B cells (Figure 4I and J). Altogether, ATG3 was a target of miR-204-5p.

ATG3 Overexpression Reverses miR-204-5p Mediated Regulatory Effects on Proliferation, Apoptosis and Autophagy in NSCLC Cells

Following confirmation that ATG3 is a target of miR-204-5p, A549 and H460 cells were transfected with miR-204-5p+ATG3, miR-204-5p+pcDNA, miR-204-5p or miR-NC to illuminate the effects of miR-204-5p/ATG3 axis on NSCLC cell development. As presented in Figure 5A, the expression of ATG3 protein was raised in NSCLC cells transfected with ATG3 compared with the pcDNA group. Moreover, ATG3 restored the suppressive effect of miR-204-5p on the proliferation of A549 and H460 cells (Figure 5B and C). Likewise, ATG3 elevation reversed miR-204-5p induced promotional effect on NSCLC cell apoptosis (Figure 5D). As expected, the expression levels of apoptosis-related protein Bax, cleaved caspase-3 and cleaved caspase-9 were enhanced by miR-204-5p and

Figure 5 ATG3 abrogated miR-204-5p mediated acceleration on cell apoptosis and suppression on cell proliferation and autophagy in NSCLC. A549 and H460 cells were transfected with miR-204-5p+ATG3, miR-204-5p+pcDNA, miR-204-5p or miR-NC. (A) The expression of ATG3 protein in transfected A549 and H460 cells. (B, C) Cell viability of transfected A549 (B) and H460 cells (C). (D) Cell apoptosis of transfected A549 and H460 cells. (E, F) The expression of apoptosis-related protein caspase-3, caspase-9, Bax and anti-apoptosis protein BCL-2 in transfected A549 (E) and H460 cells (F). (G, H) Protein expression of autophagy markers LC3 and P62 in transfected A549 (G) and H460 cells (H). * $P < 0.05$.

repressed by the overexpression of ATG3. However, the anti-apoptosis protein BCL-2 expression showed the opposite results (Figure 5E and F). In addition, the ratio of LC3II/LC3I was reduced while P62 protein expression was increased by miR-204-5p; however, the effects were reversed by ATG3 (Figure 5G and H). The results clarified that miR-204-5p could regulate NSCLC cell proliferation, apoptosis and autophagy through ATG3.

Silencing of KCNQ1OT1 Inhibits Tumor Growth in vivo

Subsequently, xenograft model was established by subcutaneously injecting sh-KCNQ1OT1 and sh-NC stably transfected A549 and H460 cells into the nude mice. We found that tumor growth was significantly repressed in the xenograft model stably transfected with sh-KCNQ1OT1 compared with sh-NC group (Figure 6A and B). Similarly, tumor weight was

lower in sh-KCNQ1OT1 transfected mice than sh-NC group (Figure 6C and D). Besides, the expression of KCNQ1OT1 and ATG3 was down-regulated whereas miR-204-5p was up-regulated in the tissue samples collected from sh-KCNQ1OT1 transfected group (Figure 6E and F). More importantly, the levels of ATG3, LC3II/LC3I, BCL-2 were suppressed whereas P62, Bax, cleaved caspase-3, cleaved caspase-9 were promoted in sh-KCNQ1OT1 group (Figure 6G). IHC staining assay indicated that KCNQ1OT1 knockdown led to a significant decrease in the percentage of ki67-positive cells compared to sh-NC group (Figure 6H). Taken together, KCNQ1OT1 silencing suppressed tumor growth in vivo by accelerating apoptosis and suppressing autophagy in NSCLC.

Discussion

LncRNAs have been demonstrated to be associated with the progression of diverse human cancers, including NSCLC. In this study, we aimed to explore the function of lncRNA

Figure 6 Kcnq1ot1 silencing suppressed tumor growth in vivo. Sh-NC or sh-KCNQ1OT1 transfected A549 cells or H460 cells were injected into the nude mice. **(A, B)** Tumor volume of xenograft mice was measured every 4 d from day 8. **(C, D)** Tumor weight was measured when xenograft mice were sacrificed after 28 days. **(E, F)** The expression of KCNQ1OT1, miR-204-5p, ATG3 in xenograft mice tumor tissues. **(G)** Protein expression of ATG3, LC3, P62, BCL-2, cleaved caspase-3, cleaved caspase-9 and Bax in xenograft mice tumor tissues. **(H)** The percentage of Ki67-positive cells in xenograft mice tumor tissues. * $P < 0.05$.

KCNQ1OT1 in NSCLC development. Previous studies have identified that lncRNA KCNQ1OT1 is an essential prognosis biomarker in a variety of diseases, including diabetic retinopathy, osteolysis and breast cancer,^{26–28} while the function and the molecular mechanism of KCNQ1OT1 in NSCLC progression were not fully validated. KCNQ1OT1 was up-regulated in breast cancer and overexpression of KCNQ1OT1 accelerated cell growth in vitro and repressed tumor growth in vivo by enhancing CCNE2 via sponging miR-145.²⁹ Upregulation of KCNQ1OT1 stimulated NSCLC cell progression by inhibiting miRNA-27b-3p and boosting HSP90AA1 expression.³⁰ Additionally, KCNQ1OT1 promoted cell development as well as cisplatin resistance against tongue cancer by targeting miR-211-5p to alter Ezrin/Fak/Src signaling pathway.³¹

Consistently, KCNQ1OT1 enhanced oxaliplatin resistance by modulating the miR-34a/ATG4B axis in colon cancer.³² Therefore, we guessed that KCNQ1OT1 exerted its functions by sponging the specific miRNA in NSCLC.

MiR-204-5p was identified to be a target of KCNQ1OT1 in this study. Interestingly, miR-204-5p was found to be an autophagy-associated gene in several tumor types, such as rhabdomyosarcoma, endometrial and colorectal cancer.^{33–35} In addition, Luan et al reported that miR-204-5p improved cell sensitivity to chemotherapy and inhibited melanoma growth by targeting MMP9 and B-cell lymphoma-2.³⁶ Similarly, miR-204-5p enhanced chemotherapeutic sensitivity and repressed invasion of colorectal cancer cells by interacting with RAB22A.³⁷ Moreover, miR-204-5p was clarified

to weaken cell proliferation and induce cell apoptosis in papillary thyroid carcinoma as well as expedite apoptosis and repress cell metastasis in osteosarcoma by targeting IGFBP5 and EBF2, respectively.^{38,39} Therefore, we speculated that miR-204-5p functioned as a tumor suppressor to suppress proliferation, migration, invasion and autophagy as well as enhance apoptosis of NSCLC cells.

Wang et al declared that miR-16 could target the 3' UTR of ATG to hamper TGF- β 1-induced epithelial-to-mesenchymal transition (EMT) via the activation of autophagy in NSCLC.²⁴ Hua et al reported that miR-1/ATG3 participated in the regulation of chemoresistance in NSCLC.²⁵ Thus, we speculated that KCNQ1OT1 modulated proliferation, apoptosis and autophagy of NSCLC cells via modulating miR-204-5p/ATG3 axis. Initially, we noticed that the expression of KCNQ1OT1 and ATG3 was up-regulated whereas miR-204-5p was down-regulated in NSCLC tumors and cells compared with the corresponding normal tissues and cells. As expected, KCNQ1OT1 knockdown expedited apoptosis and repressed proliferation and autophagy of NSCLC cells in vitro and in vivo. However, the miR-204-5p inhibitor reversed the effects. The rescue experiments also displayed that ATG3 elevation abolished miR-204-5p mediated regulatory effects on cell proliferation, apoptosis and autophagy in NSCLC.

In conclusion, we clarified that KCNQ1OT1 contributes to cell proliferation and autophagy and suppresses cell apoptosis by altering the miR-204-5p/ATG3 axis in NSCLC. Our study represented potential biomarkers for targeted therapy of NSCLC.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Ai X, Mao F, Shen S, et al. Bexarotene inhibits the viability of non-small cell lung cancer cells via slc10a2/PPARGgamma/PTEN/mTOR signaling pathway. *BMC Cancer*. 2018;18:407. doi:10.1186/s12885-018-4224-x
2. You J, Cheng J, Yu B, et al. Baicalin, a Chinese herbal medicine, inhibits the proliferation and migration of human non-small cell lung carcinoma (NSCLC) cells, A549 and H1299, by activating the SIRT1/AMPK signaling pathway. *Med Sci Monit*. 2018;24:2126–2133. doi:10.12659/MSM.909627
3. Xie Y, Lv Y, Zhang Y, et al. LATS2 promotes apoptosis in non-small cell lung cancer A549 cells via triggering Mff-dependent mitochondrial fission and activating the JNK signaling pathway. *Biomed Pharmacother*. 2019;109:679–689. doi:10.1016/j.biopha.2018.10.097
4. Li Q, Yang Z, Chen M, et al. Downregulation of microRNA-196a enhances the sensitivity of non-small cell lung cancer cells to cisplatin treatment. *Int J Mol Med*. 2016;37:1067–1074. doi:10.3892/ijmm.2016.2513
5. Tang Q, Li M, Chen L, et al. miR-200b/c targets the expression of RhoE and inhibits the proliferation and invasion of non-small cell lung cancer cells. *Int J Oncol*. 2018;53:1732–1742.
6. Moravcikova E, Krepela E, Donnenberg VS, et al. BOK displays cell death-independent tumor suppressor activity in non-small-cell lung carcinoma. *Int J Cancer*. 2017;141:2050–2061. doi:10.1002/ijc.30906
7. Cui M, Chen M, Shen Z, et al. LncRNA-UCA1 modulates progression of colon cancer through regulating the miR-28-5p/HOXB3 axis. *J Cell Biochem*. 2019.
8. Li C, Lv Y, Shao C, et al. Tumor-derived exosomal lncRNA GAS5 as a biomarker for early-stage non-small-cell lung cancer diagnosis. *J Cell Physiol*. 2019.
9. Li N, Zhan X, Zhan X. The lncRNA SNHG3 regulates energy metabolism of ovarian cancer by an analysis of mitochondrial proteomes. *Gynecol Oncol*. 2018;150:343–354. doi:10.1016/j.ygyno.2018.06.013
10. Wang CG, Liao Z, Xiao H, et al. LncRNA KCNQ1OT1 promoted BMP2 expression to regulate osteogenic differentiation by sponging miRNA-214. *Exp Mol Pathol*. 2019;107:77–84. doi:10.1016/j.yexmp.2019.01.012
11. Bian Y, Gao G, Zhang Q, et al. KCNQ1OT1/miR-217/ZEB1 feedback loop facilitates cell migration and epithelial-mesenchymal transition in colorectal cancer. *Cancer Biol Ther*. 2019;20:886–896. doi:10.1080/15384047.2019.1579959
12. Sun H, Li Y, Kong H, et al. Dysregulation of KCNQ1OT1 promotes cholangiocarcinoma progression via miR-140-5p/SOX4 axis. *Arch Biochem Biophys*. 2018;658:7–15. doi:10.1016/j.abb.2018.09.019
13. Hu H, Yang L, Li L, et al. Long non-coding RNA KCNQ1OT1 modulates oxaliplatin resistance in hepatocellular carcinoma through miR-7-5p/ABCC1 axis. *Biochem Biophys Res Commun*. 2018;503:2400–2406. doi:10.1016/j.bbrc.2018.06.168
14. Zheng L, Zhang FX, Wang LL, et al. LncRNA KCNQ1OT1 is over-expressed in non-small cell lung cancer and its expression level is related to clinicopathology. *Eur Rev Med Pharmacol Sci*. 2019;23(16):6944–6950. doi:10.26355/eurrev_201908_18734
15. Chang RM, Yang H, Fang F, et al. MicroRNA-331-3p promotes proliferation and metastasis of hepatocellular carcinoma by targeting PH domain and leucine-rich repeat protein phosphatase. *Hepatology*. 2014;60:1251–1263. doi:10.1002/hep.27221
16. Luan X, Wang Y. LncRNA XLOC_006390 facilitates cervical cancer tumorigenesis and metastasis as a ceRNA against miR-331-3p and miR-338-3p. *J Gynecol Oncol*. 2018;29:e95. doi:10.3802/jgo.2018.29.e95
17. Ji H, Sang M, Liu F, et al. miR-124 regulates EMT based on ZEB2 target to inhibit invasion and metastasis in triple-negative breast cancer. *Pathol Res Pract*. 2019;215:697–704. doi:10.1016/j.prp.2018.12.039
18. Wang YX, Zhu HF, Zhang ZY, et al. MiR-384 inhibits the proliferation of colorectal cancer by targeting AKT3. *Cancer Cell Int*. 2018;18:124. doi:10.1186/s12935-018-0628-6
19. Wang Y, Li J, Dai L, et al. MiR-17-5p may serve as a novel predictor for breast cancer recurrence. *Cancer Biomark*. 2018;22:721–726. doi:10.3233/CBM-181228
20. Liu T, Song Z, Gai Y. Circular RNA circ_0001649 acts as a prognostic biomarker and inhibits NSCLC progression via sponging miR-331-3p and miR-338-5p. *Biochem Biophys Res Commun*. 2018;503:1503–1509. doi:10.1016/j.bbrc.2018.07.070
21. Gao W, Wu Y, He X, et al. MicroRNA-204-5p inhibits invasion and metastasis of laryngeal squamous cell carcinoma by suppressing forkhead box C1. *J Cancer*. 2017;8:2356–2368. doi:10.7150/jca.19470
22. Wang X, Li F, Zhou X. miR-204-5p regulates cell proliferation and metastasis through inhibiting CXCR4 expression in OSCC. *Biomed Pharmacother*. 2016;82:202–207. doi:10.1016/j.biopha.2016.04.060
23. Jiang G, Wen L, Zheng H, et al. miR-204-5p targeting SIRT1 regulates hepatocellular carcinoma progression. *Cell Biochem Funct*. 2016;34:505–510. doi:10.1002/cbf.3223
24. Wang H, Zhang Y, Wu Q, et al. MiR-16 mimics inhibit TGF- β 1-induced epithelial-to-mesenchymal transition via activation of autophagy in non-small cell lung carcinoma cells. *Oncol Rep*. 2018;39:247–254.

25. Hua L, Zhu G, Wei J. MicroRNA-1 overexpression increases chemosensitivity of non-small cell lung cancer cells by inhibiting autophagy related 3-mediated autophagy. *Cell Biol Int*. 2018;42(9):1240–1249. doi:10.1002/cbin.v42.9
26. Sun X, Xin Y, Wang M, et al. Overexpression of long non-coding RNA KCNQ1OT1 is related to good prognosis via inhibiting cell proliferation in non-small cell lung cancer. *Thorac Cancer*. 2018;9:523–531. doi:10.1111/tca.2018.9.issue-5
27. Shao J, Pan X, Yin X, et al. KCNQ1OT1 affects the progression of diabetic retinopathy by regulating miR-1470 and epidermal growth factor receptor. *J Cell Physiol*. 2019;234:17269–17279. doi:10.1002/jcp.28344
28. Gao X, Ge J, Li W, et al. LncRNA KCNQ1OT1 ameliorates particle-induced osteolysis through inducing macrophage polarization by inhibiting miR-21a-5p. *Biol Chem*. 2018;399:375–386. doi:10.1515/hsz-2017-0215
29. Feng W, Wang C, Liang C, et al. The dysregulated expression of KCNQ1OT1 and its interaction with downstream factors miR-145/CCNE2 in breast cancer cells. *Cell Physiol Biochem*. 2018;49:432–446. doi:10.1159/000492978
30. Dong Z, Yang P, Qiu X, et al. KCNQ1OT1 facilitates progression of non-small-cell lung carcinoma via modulating miRNA-27b-3p/HSP90AA1 axis. *J Cell Physiol*. 2019;234:11304–11314. doi:10.1002/jcp.v234.7
31. Zhang S, Ma H, Zhang D, et al. LncRNA KCNQ1OT1 regulates proliferation and cisplatin resistance in tongue cancer via miR-211-5p mediated Ezrin/Fak/Src signaling. *Cell Death Dis*. 2018;9:742. doi:10.1038/s41419-018-0793-5
32. Li Y, Li C, Li D, et al. lncRNA KCNQ1OT1 enhances the chemoresistance of oxaliplatin in colon cancer by targeting the miR-34a/ATG4B pathway. *Onco Targets Ther*. 2019;12:2649–2660. doi:10.2147/OTT.S188054
33. Tk C, Ts L, Th C, et al. Dysregulation of microRNA-204 mediates migration and invasion of endometrial cancer by regulating FOXC1. *Int J Cancer*. 2012;130:1036–1045. doi:10.1002/ijc.26060
34. Imam JS, Plyler JR, Bansal H, et al. Genomic loss of tumor suppressor miRNA-204 promotes cancer cell migration and invasion by activating AKT/mTOR/Rac1 signaling and actin reorganization. *PLoS ONE*. 2012;7:e52397. doi:10.1371/journal.pone.0052397
35. Sumbul AT, Gogebakan B, Ergun S, et al. miR-204-5p expression in colorectal cancer: an autophagy-associated gene. *Tumour Biol*. 2014;35:12713–12719. doi:10.1007/s13277-014-2596-3
36. Luan W, Qian Y, Ni X, et al. miR-204-5p acts as a tumor suppressor by targeting matrix metalloproteinases-9 and B-cell lymphoma-2 in malignant melanoma. *Onco Targets Ther*. 2017;10:1237–1246. doi:10.2147/OTT
37. Yin Y, Zhang B, Wang W, et al. miR-204-5p inhibits proliferation and invasion and enhances chemotherapeutic sensitivity of colorectal cancer cells by downregulating RAB22A. *Clin Cancer Res*. 2014;20:6187–6199. doi:10.1158/1078-0432.CCR-14-1030
38. Liu L, Wang J, Li X, et al. MiR-204-5p suppresses cell proliferation by inhibiting IGFBP5 in papillary thyroid carcinoma. *Biochem Biophys Res Commun*. 2015;457:621–626. doi:10.1016/j.bbrc.2015.01.037
39. Li M, Shen Y, Wang Q, et al. MiR-204-5p promotes apoptosis and inhibits migration of osteosarcoma via targeting EBF2. *Biochimie*. 2019;158:224–232. doi:10.1016/j.biochi.2018.12.003

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>

Dovepress