

The Role of the Reactive Oxygen Species Scavenger Agent, Astaxanthin, in the Protection of Cisplatin-Treated Patients Against Hearing Loss

This article was published in the following Dove Press journal:
Drug Design, Development and Therapy

Benyu Nan^{1,2}
Xi Gu³
Xinsheng Huang²

¹Department of Otorhinolaryngology-Head and Neck Surgery, Wenzhou Medical University, Affiliated Hospital 2, Wenzhou 325000, People's Republic of China; ²Department of Otorhinolaryngology-Head and Neck Surgery, Zhongshan Hospital, Fudan University, Shanghai 200030, People's Republic of China; ³Department of Otolaryngology-Head and Neck Surgery, The First Affiliated Hospital of Fujian Medical University, Fuzhou 350000, People's Republic of China

Abstract: Emerging evidence of significant hearing loss occurring shortly after cisplatin administration in cancer patients has stimulated research into the causes and treatment of this side effect. Although the aetiology of cisplatin-induced hearing loss (CIHL) remains unknown, an increasing body of research suggests that it is associated with excessive generation of intracellular reactive oxygen species (ROS) in the cochlea. Astaxanthin, a xanthophyll carotenoid, has powerful anti-oxidant, anti-inflammatory, and anti-apoptotic properties based on its unique cell membrane function, diverse biological activities, and ability to permeate the blood-brain barrier. In this review, we summarize the role of ROS in CIHL and the effect of astaxanthin on inhibiting ROS production. We focus on investigating the mechanism of action of astaxanthin in suppressing excessive production of ROS.

Keywords: astaxanthin, oxidative stress, cisplatin, hearing loss

Introduction

Cisplatin, an effective antineoplastic agent commonly used in clinical practice, has many serious adverse effects including nephrotoxic, neurotoxic, and ototoxic effects. These life-long disabling adverse effects are strongly associated with the dosage, frequency, and duration of cisplatin treatment. Cisplatin-induced hearing loss (CIHL), which is permanent and mostly bilateral, can negatively affect academic development and social integration, especially in children.¹ To the best of our knowledge, cisplatin ototoxicity has not been studied in detail, and the mechanisms responsible for the degeneration of cochlear structures are not completely understood.

Emerging evidence indicates that excessive production of reactive oxygen species (ROS) contributes to cisplatin ototoxicity. Mechanistically, cisplatin ototoxicity is associated with the absence of glutathione (GSH) and the inhibition of glutathione peroxidase (GSH.Px) and glutathione reductase activities because cisplatin can covalently bind to the sulfhydryl groups of anti-oxidant enzymes, causing enzyme inactivation.² Increased lipid peroxidation in the cochlea inhibits essential cellular enzymes and membrane transporters, thereby disturbing ion channel function. Increased ROS production eventually results in apoptosis and necroptosis, supporting the hypothesis that ROS play a crucial role in cisplatin ototoxicity and suggesting that inhibiting ROS production could be beneficial for protecting the cochlea and reversing hearing loss.

Correspondence: Xinsheng Huang
Department of Otorhinolaryngology-Head and Neck Surgery, Zhongshan Hospital, Fudan University, Fenglin Road 180, Xuhui District, Shanghai 200030, People's Republic of China
Tel +86 13681971739
Fax +86 021-64041990
Email huang.xinsheng@zs-hospital.sh.cn

Astaxanthin is a red carotenoid agent with potent anti-oxidant properties that can scavenge singlet oxygen and free radicals. These properties confer astaxanthin with anti-inflammatory and immunomodulatory activities, protective effects against neuronal damage, anti-aging and anti-cancer activities, and the ability to inhibit cell membrane peroxidation. The anti-oxidant activity of astaxanthin is 10-fold greater than that of zeaxanthin, lutein, canthaxanthin, and β -carotene, and 100-fold greater than that of α -tocopherol.³ Growing evidence suggests that astaxanthin inhibits the development of oxidative stress-associated diseases and mitochondrial dysfunction.⁴ Moreover, powerful permeation of the blood-brain barrier (BBB) allows astaxanthin to act as a potent neuroprotective agent in mammals.

The use of cisplatin is limited by its ototoxicity and nephrotoxicity. Methods to increase diuresis, such as hydration, have the potential to reduce its nephrotoxicity. However, there are currently no effective FDA-approved treatments for ototoxicity. We reviewed the evidence supporting the ability of astaxanthin to inhibit ROS generation and prevent mitochondrial dysfunction and neurodegeneration. Based on this assessment, we hypothesized that astaxanthin may be effective for the prevention and treatment of CIHL. In this review, we focus on the following topics: (1) The mechanisms underlying cisplatin ototoxicity; (2) astaxanthin-based therapies for diseases related to excessive ROS production; (3) astaxanthin biochemistry and bioactivity; and (4) downstream pathways of astaxanthin contributing to the inhibition of ROS generation.

Mechanisms of Cisplatin Ototoxicity

An increasing body of research suggests that cisplatin ototoxicity is related to cellular hypersensitivity, although the precise cellular and molecular mechanisms remain unclear. Our understanding of the role of cisplatin in ototoxicity is limited; however, research suggests that cisplatin uptake plays a crucial role. A recent study detected residual platinum in the cochleae of mice and cancer patients receiving cisplatin chemotherapy months-to-years after the treatment.⁵

Cisplatin Transportation

Cisplatin is a square planar complex of a bivalent platinum cation with two cis chloride ligands and two cis ammonia ligands.⁶ The complex was originally assumed to enter cells by passive diffusion because its uptake is concentration-dependent and non-saturable.⁷ However, subsequent studies showed that copper transporter 1 (CTR1),^{8,9} organic cation

transporter 2 (OCT2),¹⁰ mechanotransduction (MET)¹¹ and copper-extruding P-type ATPases (ATP7A and ATP7B)¹² coordinate the cellular uptake of cisplatin. Although there may be other channels involved in cisplatin transportation, they have yet to be identified.^{13–16}

CTR1, a high-affinity copper transporter, is highly expressed in outer hair cells, inner hair cells, stria vascularis, and spiral ganglion neurons,⁸ and contributes to drug entry and cell apoptosis.¹⁷ CTR1 is a major entry route for cisplatin in hair cells, and it can enhance the cytotoxicity and cellular uptake of cisplatin in cells and in mouse.⁸ Coactivity of both CTR1 and OCT2 may lead to secondary damage in the stria vascularis and spiral ganglion.⁸ Knockout of CTR1 in yeast was reported to increase cisplatin resistance and decrease the intracellular concentration of cisplatin.¹⁸ Although increased expression of CTR1 may affect the intracellular concentration and distribution of cisplatin, it does not affect the ability of cisplatin to target DNA.¹⁹

OCTs belong to the solute carrier (SLC) 22A family,²⁰ and three isoforms (OCT1–3) have been identified, which are mainly expressed in the kidneys and liver.^{21–23} OCT2 plays a key role in cisplatin transportation and is also expressed in the organ of Corti and stria vascularis.^{10,24} In OCT1/2 double-knockout (KO) mice, cisplatin shows no ototoxicity and only mild nephrotoxicity.¹⁰ OCT2 variants were also reported to impede CIHL in children and adult, which highlights its key role in cisplatin transportation in the cochlea.²⁵

The MET channel is a nonselective cation channel that allows calcium and other ions to cross the membrane.¹⁸ In zebrafish, CIHL is associated with functional MET channels, and inhibition of MET channels by quinine or EGTA has protective effects against CIHL.¹¹ Furthermore, zebrafish mutants lacking MET channels are resistant to CIHL.¹¹ These studies suggest that MET channels contribute to the entry of cisplatin into hair cells.

A small increase in the expression of the copper transporter ATP7A resulted in resistance to clinically available platinum drugs in human 2008 ovarian carcinoma cells that may be attributed to ATP7A binding and sequestration of platinum compounds.²⁶ The HSC-4-R, OSC-19-R, and HOC313-R cell lines have acquired resistance to cisplatin that is related to ATP7B overexpression.²⁷ Both ATP7A and ATP7B are expressed in the organ of Corti, stria vascularis and spiral ganglion neurons.²⁸ Inhibition of Na^+ - K^+ -ATPase reduces cisplatin accumulation, suggesting the involvement of other transporters in cisplatin resistance.²⁹

It has been suggested that dysfunction of the LRRC8A and LRRC8D subunits of heteromeric volume-regulated

anion channels (VRAC) can lead to cisplatin resistance in KCP-4 human epidermoid cancer cell line and human lung adenocarcinoma cells.^{30,31} In 2015, Planells-Cases reported that the absence of VRAC is related to cisplatin resistance in haploid KBM7 cells based on genome-wide loss-of-function screening.³² However, the role of VRAC in cisplatin resistance is still unclear. Overexpression of multidrug resistance protein 2 (MRP2) increases the efflux of cisplatin in hepatocellular carcinoma and oesophageal squamous cell carcinoma.^{33,34}

After entry into the cell, cisplatin undergoes aquation to form $[\text{Pt}(\text{NH}_3)_2\text{Cl}(\text{OH}_2)]^+$ and $[\text{Pt}(\text{NH}_3)_2(\text{OH}_2)_2]^{2+}$ (Figure 1). The complex interacts with various reactive groups, blocking DNA replication and transcription and resulting in the inhibition of DNA repair and cell cycle progression.⁷ Platinum-DNA adducts found in the hair cells of the cochlea and the marginal cells of the stria vascularis contribute to the inhibition of cell proliferation.⁷ DNA damage caused by the

formation of DNA adducts induces apoptosis initiated by activation of p53, which activates apoptotic proteins such as Bax or inhibits anti-apoptotic members of the Bcl-2 family³⁵ (Figure 1). Cisplatin-induced DNA damage in the cochlea may interfere with signal transduction pathways, including the v-akt murine thymoma viral oncogene homologue (AKT), v-abl Abelson murine leukaemia viral oncogene homologue 1 (c-ABL), p53, and mitogen-activated protein kinase/c-Jun N-terminal kinase/extracellular signal-regulated kinase (MAPK/JNK/ERK) pathways.^{7,36}

ROS in CIHL

Normally, the maintenance of cochlea function requires high metabolic activity in areas such as the stria vascularis, spiral ligament, and spiral prominence, leading to leakage of electrons from the mitochondrial respiratory chain, which react with oxygen (O_2) to produce superoxide ($\text{O}_2^{\cdot-}$). Environmental stimuli increase oxidative stress in




Figure 1 Schematic of the proposed mechanism of cisplatin transportation and the generation of ROS in CIHL. Cisplatin is transported into cochlear cells by membrane transporters, including copper transporter 1 (CTR1), organic cation transporter 2 (OCT2), and mechano-electrical transduction (MET), and is excluded by copper-extruding P-type ATPases (ATP7A and ATP7B), multidrug resistance protein 2 (MRP2), and volume-regulated anion channels (VRAC). Cisplatin induces ROS production in the inner ear via NADPH oxidase (NOX), xanthine oxidase (XO), cytochrome P450 (CYP450), induced nitric oxide synthase (iNOS), and disturbances in the mitochondrial electron transport chain.

Abbreviations: IL, interleukin; STAT1, signal transducer and activator of transcription 1.

the cochlea in association with increased metabolic activity. The metabolic requirements of the cochlea lead to vulnerability to hypoxic events and ischemia-reperfusion injury.³⁷ Cisplatin promotes the generation of ROS by stimulating enzyme systems or deactivating anti-oxidant systems.^{38,39} This is supported by decreased cochlear GSH and anti-oxidant enzyme activity in rats treated with cisplatin.⁴⁰ This decrease could result from covalent binding of cisplatin to anti-oxidant enzymes (e.g., superoxide dismutase and catalase), loss of metal cofactors, depletion of anti-oxidant enzymes by increased ROS, and depletion of cochlear anti-oxidant enzyme cofactors such as GSH and NADPH, which protect against the toxicity of ROS and allow the regeneration of GSH.⁴¹

NADPH oxidases (NOXs) are membrane-bound, multi-subunit enzyme complexes that face the extracellular space and transfer electrons across the plasma membrane from NADPH to molecular oxygen, generating superoxide in the cell.⁴² Inactive NOX is present in an unassembled form in which p22phox and gp91phox are present in the membrane, whereas p47phox, p67phox, and p40phox exist in the cytosol. When NOX is activated by phosphorylated p47phox, the cytosolic subunits translocate to the membrane and convert the oxidase into an assembled and active form that can transfer electrons from the substrate to O₂, forming O₂^{•-}.⁴³ Some members of the NOX family may be responsible for the bulk of ROS generation in cochlea cells.^{13,44,45} NOX3-dependent ROS generation may be a target of cisplatin, as indicated by evidence showing that knockdown of NOX3 by trans-tympanic delivery of siRNA attenuates cisplatin ototoxicity in rats.^{44,46} NOX3-derived ROS can activate signal transducer and activator of transcription 1 and 6 (STAT1 and STAT6), increasing the production of the pro-inflammatory cytokines tumour necrosis factor alpha (TNF α), interleukin (IL)-1b, and IL-6.^{47,48} These pro-inflammatory mediators exacerbate CIHL by increasing the activity of NOX isoforms including NOX1 and NOX4.⁴⁵ In addition, NOX3 expression can be suppressed by silencing the transient receptor potential vanilloid 1 (TRPV1) channel, which acts as a key factor contributing to ROS generation in cochlear hair cells.¹³ siRNA-mediated silencing of STAT1 abolishes cisplatin-induced p53 activation, consistent with a study showing that STAT1 regulates cisplatin-induced hair cell death.^{48,49} These findings suggest that the TRPV1 and NOX3 signalling pathways may be associated with STAT1, resulting in inflammation and cisplatin-induced hair cell death leading to hearing loss. Increased NOX2 expression in the cochlea increases ROS production in outer hair cells of

the basal turn, which is an important factor associated with the vulnerability of outer hair cells.⁵⁰

The xanthine/xanthine oxidase (XO) system is another active ROS-generating system found in the cochlea that contributes to both superoxide and hydrogen peroxide generation. XO is a xanthine oxidoreductase isoform that catalyses the reduction of O₂ into O₂^{•-} and H₂O₂. Its activity may be enhanced by ROS derived from mitochondria and NOX.⁵¹ Inhibition of XO in a rat model administered ebselen, a GSH.Px mimetic, decreases cisplatin ototoxicity and nephrotoxicity.⁵² Depletion of intracellular ATP transforms adenine nucleotides into hypoxanthine and xanthine, which are substrates of XO.⁵³ This is proposed as a principal mechanism underlying oxidative injury.

Disturbance of the mitochondrial electron transport chain system by cisplatin increases ROS generation concomitant with a decrease in the mitochondrial membrane potential (MMP), an indicator of mitochondrial malfunction.^{54,55} Disruption of Complex I function resulting in elevated ROS production underlies the destruction of sensory hair cells in the cochlea caused by cisplatin.⁵⁶ In addition, ROS-mediated mitochondrial dysfunction can suppress the amplification of intracellular ROS generation, resulting in hair cell apoptosis and necrosis.⁵⁷

Evidence supports an association between the toxic effects of cisplatin and cytochrome P450 (CYP450) enzymes.⁵⁸ However, there is little research on the relation between CYP450 and CHIL. CYP450 is an important source for the generation of ROS and catalytic iron in cisplatin-induced cytotoxicity of the LLC-PK1 cells.⁵⁹ Liu found that CYP450 subfamily members (CYP2e1) play a pivotal role in cisplatin-induced nephrotoxicity and apoptosis related to the generation of ROS.⁶⁰ CYP2e1 null mice showed significantly functional and histologic protection against renal injury and decrease of apoptosis by cisplatin.⁶⁰ Increasing CYP2E1 enhances the toxicity induced by cisplatin in liver injury models both in vitro and in vivo, and the mechanism might associate with accumulating production of ROS and oxidative stress.⁵⁸ E47 HepG2 cells, overexpress human CYP2E1, were sensitive to cisplatin because of an earlier activation of ERK.⁶¹ These results suggest that the generation of ROS by cisplatin in the cochlea is associated with CYP450.

Induced nitric oxide synthase (iNOS) and direct NO generation can be observed in the inner ear (including the spiral ligament, modiolus, spiral limbus, supporting cells, nerve fibres, stria vascularis, hair cells, and spiral ganglion neurons) in response to cisplatin administration.⁶²⁻⁶⁴ ROS/reactive nitrogen species (RNS), as intracellular second

messengers, play a pivotal role in ototoxicity. ROS/RNS can activate downstream signalling pathways, regulate gene expression, and interfere directly with lipids, DNA, RNA, metal cofactors and proteins. Nitrosative stress can react with and suppress ROS generation, overpowering anti-oxidant defence capacities within the inner ear, triggering the pro-apoptotic pathway, and leading to outer hair cell death.^{65,66}

Although our understanding of the mechanism by which ROS exert their effects in the cochlea is still incomplete, it is well established that oxidative stress can cause DNA damage. It is also evident that the DNA damage caused by ROS may be repaired via several pathways, including base excision repair (BER),^{67–69} mismatch repair (MMR),⁷⁰ and nucleotide excision repair (NER).⁷¹ The molecule 8-oxoguanine (8-oxo-G) is well known to cause DNA damage by creating a gap in the DNA, which can be repaired by the enzyme 8-oxoguanine glycosylase (8-OGG1) in mitochondria and nucleus.⁷² Chronic exposure to a low dose of H₂O₂ induces a DNA protective response in C2C12 cells by activating transcription factors that enhance the expression of DNA repair enzymes.^{73,74} An increase in poly-ADP-ribose transferase 1 (PARP1), an enzyme that recruits DNA repair proteins, has been detected in a mouse model of CHIL, and its inhibition is associated with decreased cisplatin-induced cell death.⁷⁵ HEI-OC1 cells exposed to cisplatin can promote and initiate DNA repair, but cannot prevent or reverse DNA damage,⁷⁶ possibly because of the absence of some DNA repair pathways in mitochondria and BER can only cope with minor damage or process single nucleotides in the mitochondria.⁷²

Thus, the cisplatin complex leads to reciprocal activation of ROS generation, NOX, XO, mitochondria, CYP450, MMP, and iNOS as well as pro-inflammatory signalling, suggesting that cisplatin initiates a series of vicious cycles (Figure 1). In turn, ROS enhance lipid peroxidation and DNA damage, finally leading to auditory cell death.

Astaxanthin-Based Therapies for Diseases Related to ROS

Oxidative stress leads to disease conditions that result from increased production of ROS or depletion of the anti-oxidant enzyme system. Mitochondrial disturbance is often involved in the onset of oxidative stress-associated diseases, because the mitochondria are responsible for energy generation and are important sources of ROS. Astaxanthin, which has strong anti-oxidant activity and the ability to maintain metabolic

efficiency, is a potent anti-oxidant with the potential to target several health conditions.⁷⁷

Because nervous system tissues show intense metabolic aerobic activity, rich irrigation with blood vessels, and are abundant in unsaturated fats and iron, they are particularly susceptible to oxidative damage.⁷⁸ Substantial evidence supports the hypothesis that oxidative stress and impaired mitochondrial efficiency can be causative or at least ancillary factors in the pathogenesis of major neurodegenerative diseases, such as Alzheimer's disease (AD), Parkinson's disease (PD), Huntington's, and amyotrophic lateral sclerosis.^{79–82} Anti-oxidants can improve mitochondrial redox potential, which is a key target of ROS and free radical production.⁸³ A number of studies have shown that diets high in anti-oxidants can reduce these associated risks.⁸⁴ Synthesized docosahexaenoic acid-acrylated astaxanthin diesters can relieve oxidative stress and inflammasome activation in patients with AD.⁸⁵ Astaxanthin inhibits the generation of intracellular ROS and protects against 1-methyl-4-phenylpyridinium (MPP⁺)-induced cytotoxicity in a cellular model of PD, protecting SH-SY5Y cells and substantia nigra neurons from apoptosis in a PD model mouse.^{80,86} H₂O₂-stimulated mouse neural progenitor cells pre-treated with astaxanthin show suppression of apoptosis, resulting in cell proliferation.⁸⁷ Recent rat models show that astaxanthin can ameliorate aluminium-induced impaired memory performance.⁸⁸ According to studies of rats fed with natural astaxanthin, astaxanthin can cross the BBB in mammals, and its anti-oxidant potential may extend beyond the BBB, allowing it to act as a potent neuroprotective agent.⁸⁹ In a murine model of ischemic stroke, pre-treatment with astaxanthin decreased ROS production, lipid peroxidation, and cerebral infarction and promoted locomotor function recovery.⁹⁰ Astaxanthin was therefore suggested as a promising candidate neuroprotective agent in mammals.⁹¹

Elevated oxidative stress associated with ROS/RNS and chronic inflammation can aggravate cardiovascular diseases. Emerging *in vitro*- and *in vivo*-based evidence indicates that astaxanthin can reduce ROS, RNS, lipid peroxidation, and inflammatory signalling, and activate anti-oxidant enzymes in the heart.^{92–96} Investigations of the effect of astaxanthin on the myocardium of ischemic mice demonstrated that astaxanthin can significantly prevent ischemic myocardial injury by alleviating mitochondrial impairment in mitochondria with increased ROS expression, mitochondrial depolarization, and swelling.⁹⁷ Astaxanthin promoted recovery of mitochondrial integrity and blocked mitochondria-mediated apoptosis in a homocysteine-induced cardiotoxicity model

resulting from overexpression of ROS, loss of MMP, and fragmentation of mitochondria.⁹⁸ In addition, astaxanthin balanced the expression of Bcl-2 family proteins, thereby suppressing mediators of mitochondrial apoptosis such as PARP1 and caspases.⁹⁸ Studies also suggest that dietary astaxanthin has beneficial effects during chemotherapy in BALB/c mice, not only because it can counteract induced oxidative stress, but also because it has cardioprotective effects.⁹⁴ Epidemiological and clinical data indicate that dietary anti-oxidants might protect against cardiovascular disease, reducing the risk of atherosclerosis by decreasing plasma LDL-cholesterol oxidation.⁹⁹

Astaxanthin may be more effective than vitamin E for preventing and treating non-alcoholic steatohepatitis in mice and humans.¹⁰⁰ It protects the liver against acetaminophen hepatotoxicity by alleviating hepatocyte necrosis, blocking ROS generation, inhibiting oxidative stress, and reducing apoptosis.¹⁰¹ Astaxanthin accumulates in the liver, especially in the microsomal and mitochondrial fractions of liver tissues.¹⁰² These studies indicate the potential of astaxanthin for the treatment of oxidative stress-related liver and heart diseases.

Astaxanthin also has shown biological activity in dermatology clinical trials, promoting skin health and achieving effective skin cancer chemoprevention.¹⁰³ Randomized double-blinded, controlled studies reported that astaxanthin can ameliorate skin wrinkles, elasticity, texture, and viscoelasticity affected by aging and associated with oxidative metabolism and subsequent ROS production.^{104–107} A recent study using a rodent model of deep burns revealed that astaxanthin protects against early progression of a burn wound by reducing ROS-induced inflammation and apoptosis.¹⁰⁸ Moreover, it was reported that astaxanthin powerfully accelerates wound recovery in mice.¹⁰⁹

In addition to its strong antioxidative effect, there is increasing evidence to show that astaxanthin can inhibit the growth of several types of cancer.¹¹⁰ Studies have shown that ROS may be involved in cancer initiation, progression, and proliferation by acting as messengers of the oxidative stress cascade, which can be inhibited by anti-oxidants such as astaxanthin.^{110,111} In the hamster buccal pouch carcinogenesis model, it was shown that astaxanthin prevents the growth of oral cancer by suppressing cell proliferation and inducing apoptosis.¹¹² In 1,2-dimethyl hydrazine-induced rat colon carcinoma and rat hepatocellular carcinoma CBRH-7919 cells, pre-treatment with astaxanthin exerted anti-cancer effects by inducing cellular apoptosis.^{113,114} Astaxanthin also enhanced apoptosis of leukaemia K562 cells and

interfered with cell cycle progression.¹¹⁵ In ovarian cancer SKOV3 cells, astaxanthin combined with human serum albumin could be a candidate treatment via arresting the cell cycle in the G1 phase and inducing apoptosis.¹¹⁶

These positive effects have been attributed to the ability of astaxanthin to scavenge free radicals, quench singlet oxygen, and inhibit lipid peroxidation. The effect of astaxanthin on protecting cellular membranes against oxidation is related to the protection of the inner part and external surface.^{117–119} Anti-oxidants are indispensable for maintaining cellular health by protecting cellular components against oxidative stress and suppressing ROS-induced inflammation. The effects of anti-oxidants are evident in inflammation-related clinical conditions, such as Crohn's disease, asthma, and exercise-induced muscle damage.^{120,121} In 2000, Bennedson reported that dietary astaxanthin helped reduce the symptoms of ulcerative disease and gastric inflammation.¹²² Because of these effects, astaxanthin should be considered for the treatment of various ROS-related diseases.

Astaxanthin Biochemistry and Bioactivity

Astaxanthin is the main carotenoid pigment produced in marine animals, and it is present in many types of seafood, such as salmon, trout, shrimp, and lobster. Astaxanthin has three natural stereoisomers [(3S, 3'S), (3R, 3'R), and (3R, 3'S)], which vary in the configuration of the two hydroxyl groups on the molecule (Figure 2). The presence of the hydroxyl and keto groups on each ionone ring confer unique features, including conversion into an ester, and providing greater anti-oxidant activity and more polar properties than other carotenoids.^{3,123} Astaxanthin can preserve the integrity of cell membranes by penetrating bilayers and protecting the redox state and functional integrity of mitochondria.¹²⁴

Astaxanthin is present in natural sources as an ester of fatty acids or as a conjugate of proteins in foods. Astaxanthin is absorbed into enterocytes through passive diffusion, and is incorporated into chylomicrons to be transferred to the liver. This natural product is then incorporated into low-density lipoprotein and high-density lipoprotein and transported via the circulation. Moreover, the bioavailability of astaxanthin is enhanced when it is taken with dietary lipids.¹²³ Astaxanthin does not turn into vitamin A in the human body; therefore, it is nontoxic if given orally. Even at low concentrations astaxanthin is effective because of its polar features, which optimize the rate and extent of absorption.¹²⁵ The only study on humans confirmed the bioavailability of




Figure 2 Chemical structure of astaxanthin (3,3'-dihydroxy-β, β'-carotene-4,4R'-dione).

astaxanthin administered at a single high dose of 100 mg, and demonstrated that it is transported to the plasma by lipoproteins.¹²⁶ Astaxanthin can be used as a dietary supplement for human, animal, and fishery consumption. The FDA has validated the use of astaxanthin for food colouring (or colour additive) and different applications in animal and fish food.

Downstream Pathways of Astaxanthin Against the Generation of ROS

Although the anti-oxidant properties of astaxanthin play a crucial role in its biological activity, the concentrations achieved in the blood that protect cells against oxidative injury are well below those required to scavenge free radicals directly. To explain this discrepancy, it has been suggested that astaxanthin increases resistance to oxidative stress by activating signalling pathways associated with cell survival (Figure 3).

MAPK Pathway

Astaxanthin may help to prevent neurotoxicity by promoting the activation of the AKT/cyclic AMP-responsive element binding protein and ERKs, and blocking the activation of p38 MAPKs, which play pro-apoptotic roles, whereas ERKs have anti-apoptotic roles.¹²⁷ This is consistent with the observation that astaxanthin decreases ROS production and inflammatory cytokines to inhibit apoptosis and autophagy, which may be related to inactivation of components of the

MAPK family, such as p38 MAPK, JNK and ERK, in a model of hepatic ischemia reperfusion injury.¹²⁸ In addition, inhibition of the TNF- α -mediated JNK signal pathway and phosphorylation of ERK and p38 MAPK are involved in the process of hepatocyte necrosis induced by acetaminophen.¹⁰¹ In a mouse model of smoke-induced impairment of cognitive function, astaxanthin protects the brain against neuroinflammation, synaptic plasticity impairment, and oxidative stress in the cortex and hippocampus by inhibiting p38 MAPK.¹²⁹ In human umbilical vein endothelial cells (HUVECs), however, Western blot analysis revealed that astaxanthin significantly upregulated p-ERK without affecting p38 MAPK. Although expression of p-Akt was also increased, it was not significant.¹³⁰ These results suggest that astaxanthin protects cells against oxidative stress-induced apoptosis.

Nrf2/ARE Pathway

Hydroxypropyl- β -cyclodextrin-astaxanthin (CD-A) can lead to the dissociation of nuclear factor-erythroid 2-related factor 2 (Nrf2) from Keap1, which allows Nrf2 to translocate to the nucleus and bind to anti-oxidant response elements (AREs), thereby inducing an endogenous anti-oxidant response caused by heme oxygenase-1 (HO-1) and NAD(P)H Quinone Dehydrogenase 1 (NQO1).¹³¹ HO-1 is responsible for many oxidative and cytoprotective functions, and NQO1 has anti-inflammatory effects and can prevent the reduction of quinone.^{132,133} Astaxanthin pre-treatment significantly increased the




Figure 3 Proposed mechanism by which astaxanthin inhibits ROS and apoptosis. Astaxanthin inhibits apoptosis and scavenges ROS, and modulates various intracellular pathways, predominantly MAPK, PI3K/Akt and Nrf2/ARE. It can also activate the specificity protein 1 (Sp1)–NMDA receptor subunit 1 (NR1) signalling pathway, which leads to cell death. **Abbreviations:** AST, astaxanthin; Akt, v-akt murine thymoma viral oncogene homologue; ARE, anti-oxidant response element; CREB, CAMP-responsive element binding protein; ERK, extracellular signal–regulated kinases; JNK, c-Jun N-terminal kinase; Keap1, cytosolic inhibitor of Nrf2; Nrf2, nuclear factor-erythroid 2-related factor 2; NF- κ B, nuclear factor-kappa B subunit; NQO1, NADH Quinone Dehydrogenase I; PTEN, phosphatase and tensin homolog.

expression of Nrf2, HO-1, and NQO1 mRNA, exerting a protective effect against brain injuries. This was demonstrated in a cerebral ischemia rat model, rat hepatocytes, the human retinal pigment epithelial cell line ARPE-19, and early brain injury in a prechiasmatic cistern model of subarachnoid haemorrhage.^{131,134,135} In HUVECs, astaxanthin activates the Nrf-2/ARE signalling pathway by developing small amounts of ROS, whereas knockdown of Nrf-2 by siRNA inhibits HO-1 mRNA expression.¹³⁰ However, the direct molecular targets responsible for induction of the Nrf2/HO-1/NQO1 pathway remain undefined, as astaxanthin has an indirect anti-oxidant protective effect against ROS.

PI3K/AKT Pathway

Previous studies indicate that cell survival is affected by intracellular ROS generation through the modulation of the phosphatase and tensin homolog (PTEN)/phosphoinositide

3-kinase (PI3K)/AKT pathway.¹³⁶ Astaxanthin protects against isoflurane-induced neuroapoptosis in a rat model, as indicated by decreased brain damage, inhibition of caspase-3 activity, and upregulation of the PI3K/AKT pathway.¹³⁷ Zuluaga recently reported that the generation of ROS induced by stressors (AAPH and t-BuOOH, which are free radical donors that generate a burst of ROS) upregulates PTEN gene expression, which causes cellular apoptosis by deactivating AKT.¹³⁸ Conversely, astaxanthin treatment significantly suppressed PTEN expression and reduced both eNOS and Bax gene expression in endothelial cells under oxidative stress.¹³⁸ Astaxanthin can activate the PI3K/Akt pathway, protecting against H₂O₂-induced oxidative stress through the Nrf2/ARE pathway in ARPE-19 cells.¹³⁹

Astaxanthin activates the specificity protein 1 (Sp1) and NMDA receptor subunit 1 (NR1) signalling pathway, inhibiting the upregulation and nuclear transfer of Sp1

resulting from MPP⁺-induced production of intracellular ROS and cytotoxicity in PC12 cells.¹⁴⁰

Conclusion

Astaxanthin possesses ROS scavenging and anti-oxidant activities, and thus inhibits oxidative stress-induced mitochondrial dysfunction and ROS production in cells caused by various stimuli. We propose that astaxanthin treatment might be a viable approach for the effective mitigation and prevention of CIHL associated with ROS. Astaxanthin may be an excellent candidate for treating CHIL, as it is a safe nutrient with no toxicity when consumed with food, and it has the ability to pass through the BBB because of its lipid solubility. Future studies should investigate the protective properties and underlying mechanisms of astaxanthin, which may contribute to the use of astaxanthin as an otoprotective agent.

The otoprotective effects of astaxanthin have been examined in a zebrafish model, although research into its otoprotective effects is limited.¹⁴¹ Future studies should focus on the pharmaceutical potential and effects of astaxanthin for the treatment of hearing loss, particularly because astaxanthin can be easily absorbed, and when metabolized, it may have greater biological activity than its free form.³ Future studies should include a therapeutic time window, reliability of drug administration routes, and the optimal dosages of astaxanthin. The design of clinical trials to assess the potential of astaxanthin for the treatment of ototoxicity is warranted based on evidence showing its safety. In addition, the efficacy of astaxanthin for the treatment of neurological diseases should promote clinical trials of this compound for the treatment of other diseases.

Author Contributions

All authors contributed to data analysis, drafting or revising the article, gave final approval for publication, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Knight KRG, Kraemer DF, Neuwelt EA. Ototoxicity in children receiving platinum chemotherapy: underestimating a commonly occurring toxicity that may influence academic and social development. *J Clin Oncol*. 2005;23(34):8588–8596. doi:10.1200/Jco.2004.00.5355
2. Sheth S, Mukherjee D, Rybak LP, Ramkumar V. Mechanisms of cisplatin-induced ototoxicity and otoprotection. *Front Cell Neurosci*. 2017;11:338. doi:10.3389/fncel.2017.00338
3. Ambati RR, Phang SM, Ravi S, Aswathanarayana RG. Astaxanthin: sources, extraction, stability, biological activities and its commercial applications—a review. *Mar Drugs*. 2014;12(1):128–152. doi:10.3390/md12010128
4. Kuroki T, Ikeda S, Okada T, et al. Astaxanthin ameliorates heat stress-induced impairment of blastocyst development in vitro—astaxanthin colocalization with and action on mitochondria. *J Assist Reprod Genet*. 2013;30(5):623–631. doi:10.1007/s10815-013-9987-z
5. Breglio AM, Rusheen AE, Shide ED, et al. Cisplatin is retained in the cochlea indefinitely following chemotherapy. *Nat Commun*. 2017;8(1):1654. doi:10.1038/s41467-017-01837-1
6. Dasari S, Tchounwou PB. Cisplatin in cancer therapy: molecular mechanisms of action. *Eur J Pharmacol*. 2014;740:364–378. doi:10.1016/j.ejphar.2014.07.025
7. Wang D, Lippard SJ. Cellular processing of platinum anticancer drugs. *Nat Rev Drug Discov*. 2005;4(4):307–320. doi:10.1038/nrd1691
8. More SS, Akil O, Ianculescu AG, Geier EG, Lustig LR, Giacomini KM. Role of the copper transporter, CTR1, in platinum-induced ototoxicity. *J Neurosci*. 2010;30(28):9500–9509. doi:10.1523/JNEUROSCI.1544-10.2010
9. Shen DW, Pouliot LM, Hall MD, Gottesman MM. Cisplatin resistance: a cellular self-defense mechanism resulting from multiple epigenetic and genetic changes. *Pharmacol Rev*. 2012;64(3):706–721. doi:10.1124/pr.111.005637
10. Ciarimboli G, Deuster D, Knief A, et al. Organic cation transporter 2 mediates cisplatin-induced oto- and nephrotoxicity and is a target for protective interventions. *Am J Pathol*. 2010;176(3):1169–1180. doi:10.2353/ajpath.2010.090610
11. Thomas AJ, Hailey DW, Stawicki TM, et al. Functional Mechanotransduction Is Required for Cisplatin-Induced Hair Cell Death in the Zebrafish Lateral Line. *J Neurosci*. 2013;33(10):4405–4414. doi:10.1523/Jneurosci.3940-12.2013
12. La Fontaine S, Mercer JF. Trafficking of the copper-ATPases, ATP7A and ATP7B: role in copper homeostasis. *Arch Biochem Biophys*. 2007;463(2):149–167. doi:10.1016/j.abb.2007.04.021
13. Mukherjee D, Jajoo S, Whitworth C, et al. Short interfering RNA against transient receptor potential vanilloid 1 attenuates cisplatin-induced hearing loss in the rat. *J Neurosci*. 2008;28(49):13056–13065. doi:10.1523/JNEUROSCI.1307-08.2008
14. Takumida M, Ishibashi T, Hamamoto T, Hirakawa K, Anniko M. Age-dependent changes in the expression of klotho protein, TRPV5 and TRPV6 in mouse inner ear. *Acta Oto-Laryngol*. 2009;129(12):1340–1350. doi:10.3109/00016480902725254
15. Takumida M, Anniko M. Expression of canonical transient receptor potential channel (TRPC) 1-7 in the mouse inner ear. *Acta Oto-Laryngol*. 2009;129(12):1351–1358. doi:10.3109/00016480902798350
16. Takumida M, Anniko M. Expression of transient receptor potential channel mucolipin (TRPML) and polycystine (TRPP) in the mouse inner ear. *Acta Oto-Laryngol*. 2010;130(2):196–203. doi:10.3109/00016480903013593
17. Ishida S, Lee J, Thiele DJ, Herskowitz I. Uptake of the anticancer drug cisplatin mediated by the copper transporter Ctr1 in yeast and mammals. *P Natl Acad Sci USA*. 2002;99(22):14298–14302. doi:10.1073/pnas.162491399
18. Waissbluth S, Daniel SJ. Cisplatin-induced ototoxicity: transporters playing a role in cisplatin toxicity. *Hear Res*. 2013;299:37–45. doi:10.1016/j.heares.2013.02.002
19. Holzer AK, Samimi G, Katano K, et al. The copper influx transporter human copper transport protein 1 regulates the uptake of cisplatin in human ovarian carcinoma cells. *Mol Pharmacol*. 2004;66(4):817–823. doi:10.1124/mol.104.001198
20. Budiman T, Bamberg E, Koepsell H, Nagel G. Mechanism of electrogenic cation transport by the cloned organic cation transporter 2 from rat. *J Biol Chem*. 2000;275(38):29413–29420. doi:10.1074/jbc.M004645200

21. Pelis RM, Wright SH. SLC22, SLC44, and SLC47 transporters-organic anion and cation transporters: molecular and cellular properties. *Curr Top Membr.* 2014;73:233–261. doi:10.1016/B978-0-12-800223-0.00006-2
22. Schaeffeler E, Hellerbrand C, Nies AT, et al. DNA methylation is associated with downregulation of the organic cation transporter OCT1 (SLC22A1) in human hepatocellular carcinoma. *Genome Med.* 2011;3:82. doi:10.1186/gm298
23. Bleasby K, Castle JC, Roberts CJ, et al. Expression profiles of 50 xenobiotic transporter genes in humans and pre-clinical species: a resource for investigations into drug disposition. *Xenobiotica.* 2006;36(10–11):963–988. doi:10.1080/00498250600861751
24. Hellberg V, Gahm C, Liu W, Ehrsson H, Rask-Andersen H, Laurell G. Immunohistochemical localization of OCT2 in the cochlea of various species. *Laryngoscope.* 2015;125(9):E320–E325. doi:10.1002/lary.25304
25. Lanvers-Kaminsky C, Sprowl JA, Malath I, et al. Human OCT2 variant c.808G > T confers protection effect against cisplatin-induced ototoxicity. *Pharmacogenomics.* 2015;16(4):323–332. doi:10.2217/Pgs.14.182
26. Samimi G, Safaei R, Katano K, et al. Increased expression of the copper efflux transporter ATP7A mediates resistance to cisplatin, carboplatin, and oxaliplatin in ovarian cancer cells. *Clin Cancer Res.* 2004;10(14):4661–4669. doi:10.1158/1078-0432.CCR-04-0137
27. Yoshizawa K, Nozaki S, Kitahara H, Ohara T et al. Copper efflux transporter (ATP7B) contributes to the acquisition of cisplatin-resistance in human oral squamous cell lines. *Oncol Rep.* 2007;18(4):987–991. PMID:17786364
28. Ding D, He J, Allman BL, et al. Cisplatin ototoxicity in rat cochlear organotypic cultures. *Hear Res.* 2011;282(1–2):196–203. doi:10.1016/j.heares.2011.08.002
29. Tadini-Buoninsegni F, Sordi G, Smeazzetto S, Natile G, Arnesano F. Effect of cisplatin on the transport activity of P-II-type ATPases. *Metallomics.* 2017;9(7):960–968. doi:10.1039/c7mt00100b
30. Min XJ, Li H, Hou SC, et al. Dysfunction of volume-sensitive chloride channels contributes to cisplatin resistance in human lung adenocarcinoma cells. *Exp Biol Med (Maywood).* 2011;236(4):483–491. doi:10.1258/ebm.2011.010297
31. Lee EL, Shimizu T, Ise T, Numata T, Kohno K, Okada Y. Impaired activity of volume-sensitive Cl⁻ channel is involved in cisplatin resistance of cancer cells. *J Cell Physiol.* 2007;211(2):513–521. doi:10.1002/jcp.20961
32. Planells-Cases R, Lutter D, Guyader C, et al. Subunit composition of VRAC channels determines substrate specificity and cellular resistance to Pt-based anti-cancer drugs. *EMBO J.* 2015;34(24):2993–3008. doi:10.15252/embj.201592409
33. Yamasaki M, Makino T, Masuzawa T, et al. Role of multidrug resistance protein 2 (MRP2) in chemoresistance and clinical outcome in oesophageal squamous cell carcinoma. *Br J Cancer.* 2011;104(4):707–713. doi:10.1038/sj.bjc.6606071
34. Korita PV, Wakai T, Shirai Y, et al. Multidrug resistance-associated protein 2 determines the efficacy of cisplatin in patients with hepatocellular carcinoma. *Oncol Rep.* 2010;23(4):965–972. doi:10.3892/or_00000721
35. Vaseva AV, Moll UM. The mitochondrial p53 pathway. *Biochim Biophys Acta.* 2009;1787(5):414–420. doi:10.1016/j.bbabi.2008.10.005
36. Rybak LP, Ramkumar V. Ototoxicity. *Kidney Int.* 2007;72(8):931–935. doi:10.1038/sj.ki.5002434
37. Seidman MD, Quirk WS, Nuttall AL, Schweitzer VG. The protective effects of allopurinol and superoxide dismutase-polyethylene glycol on ischemic and reperfusion-induced cochlear damage. *Otolaryngol Head Neck Surg.* 1991;105(3):457–463. doi:10.1177/01945989110500318
38. Kopke RD, Liu W, Gabaizadeh R, et al. Use of organotypic cultures of Corti's organ to study the protective effects of antioxidant molecules on cisplatin-induced damage of auditory hair cells. *Am J Otol.* 1997;18(5):559–571. PMID:9303151
39. Clerici WJ, Yang LH. Direct effects of intraperilymphatic reactive oxygen species generation on cochlear function. *Hearing Res.* 1996;101(1–2):14–22. doi:10.1016/S0378-5955(96)00126-8
40. Rybak LP, Husain K, Morris C, Whitworth C, Somani S. Effect of protective agents against cisplatin ototoxicity. *Am J Otol.* 2000;21(4):513–520. PMID:10912697
41. DeWoskin R, Riviere J. Cisplatin-induced loss of kidney copper and nephrotoxicity is ameliorated by single dose diethyldithiocarbamate, but not mesna. *Toxicol Appl Pharmacol.* 1992;112(2):182–189. doi:10.1016/0041-008x(92)90186-v
42. Gao HM, Zhou H, Hong JS. NADPH oxidases: novel therapeutic targets for neurodegenerative diseases. *Trends Pharmacol Sci.* 2012;33(6):295–303. doi:10.1016/j.tips.2012.03.008
43. Turrens JF, Boveris A. Generation of superoxide anion by the NADH dehydrogenase of bovine heart mitochondria. *Biochem J.* 1980;191(2):421–427. doi:10.1042/bj1910421
44. Mukherjea D, Jajoo S, Kaur T, Sheehan KE, Ramkumar V, Rybak LP. Transtympanic administration of short interfering (si) RNA for the NOX3 isoform of NADPH oxidase protects against cisplatin-induced hearing loss in the rat. *Antioxid Redox Signal.* 2010;13(5):589–598. doi:10.1089/ars.2010.3110
45. Kim HJ, Lee JH, Kim SJ, et al. Roles of NADPH oxidases in cisplatin-induced reactive oxygen species generation and ototoxicity. *J Neurosci.* 2010;30(11):3933–3946. doi:10.1523/JNEUROSCI.6054-09.2010
46. Banfi B, Malgrange B, Knisz J, Steger K, Dubois-Dauphin M, Krause KH. NOX3, a superoxide-generating NADPH oxidase of the inner ear. *J Biol Chem.* 2004;279(44):46065–46072. doi:10.1074/jbc.M403046200
47. Kim HJ, Oh GS, Lee JH, et al. Cisplatin ototoxicity involves cytokines and STAT6 signaling network. *Cell Res.* 2011;21(6):944–956. doi:10.1038/cr.2011.27
48. Kaur T, Mukherjea D, Sheehan K, Jajoo S, Rybak LP, Ramkumar V. Short interfering RNA against STAT1 attenuates cisplatin-induced ototoxicity in the rat by suppressing inflammation. *Cell Death Dis.* 2011;2:e180. doi:10.1038/cddis.2011.63
49. Schmitt NC, Rubel EW, Nathanson NM. Cisplatin-induced hair cell death requires STAT1 and is attenuated by epigallocatechin gallate. *J Neurosci.* 2009;29(12):3843–3851. doi:10.1523/Jneurosci.5842-08.2009
50. Qi MH, Qiu Y, Zhou XY, et al. Regional up-regulation of NOX2 contributes to the differential vulnerability of outer hair cells to neomycin. *Biochem Biophys Res Co.* 2018;500(2):110–116. doi:10.1016/j.bbrc.2018.03.141
51. McNally JS, Saxena A, Cai H, Dikalov S, Harrison DG. Regulation of xanthine oxidoreductase protein expression by hydrogen peroxide and calcium. *Arterioscler Thromb Vas.* 2005;25(8):1623–1628. doi:10.1161/01.ATV.0000170827.16296.6e
52. Lynch ED, Gu R, Pierce C, Kil J. Reduction of acute cisplatin ototoxicity and nephrotoxicity in rats by oral administration of allopurinol and ebselen. *Hear Res.* 2005;201(1–2):81–89. doi:10.1016/j.heares.2004.08.002
53. Kinugasa Y, Ogino K, Furuse Y, et al. Allopurinol improves cardiac dysfunction after ischemia-reperfusion via reduction of oxidative stress in isolated perfused rat hearts. *Circ J.* 2003;67(9):781–787. doi:10.1253/circj.67.781
54. Jones QR, Warford J, Rupasinghe HP, Robertson GS. Target-based selection of flavonoids for neurodegenerative disorders. *Trends Pharmacol Sci.* 2012;33(11):602–610. doi:10.1016/j.tips.2012.08.002

55. Lee JS, Kang SU, Hwang HS, Pyun JH, Choung YH, Kim CH. Epicatechin protects the auditory organ by attenuating cisplatin-induced ototoxicity through inhibition of ERK. *Toxicol Lett.* 2010;199(3):308–316. doi:10.1016/j.toxlet.2010.09.013
56. Tabuchi K, Nishimura B, Nakamagoe M, Hayashi K, Nakayama M, Hara A. Ototoxicity: mechanisms of cochlear impairment and its prevention. *Curr Med Chem.* 2011;18(31):4866–4871. doi:10.2174/092986711797535254
57. Cai JY, Yang J, Jones DP. Mitochondrial control of apoptosis: the role of cytochrome c. *Bba-Bioenergetics.* 1998;1366(1–2):139–149. doi:10.1016/S0005-2728(98)00109-1
58. Lu YK, Cederbaum AI. Cisplatin-induced hepatotoxicity is enhanced by elevated expression of cytochrome P450 2E1. *Toxicol Sci.* 2006;89(2):515–523. doi:10.1093/toxsci/kfj031
59. Liu H, Baliga M, Baliga R. Effect of cytochrome p450 2E1 inhibitors on cisplatin-induced cytotoxicity to renal proximal tubular epithelial cells. *Anticancer Res.* 2002;22(2a):863–868. PMID:12014663
60. Liu H, Baliga R. Cytochrome P450 2E1 null mice provide novel protection against cisplatin-induced nephrotoxicity and apoptosis. *Kidney Int.* 2003;63(5):1687–1696. doi:10.1046/j.1523-1755.2003.00908.x
61. Lu Y, Cederbaum A. The mode of cisplatin-induced cell death in CYP2E1-overexpressing HepG2 cells: modulation by ERK, ROS, glutathione, and thioredoxin. *Free Radical Bio Med.* 2007;43(7):1061–1075. doi:10.1016/j.freeradbiomed.2007.06.021
62. Watanabe K, Hess A, Bloch W, Michel O. Expression of inducible nitric oxide synthase (iNOS/NOS II) in the vestibule of guinea pigs after the application of cisplatin. *Anti-Cancer Drug.* 2000;11(1):29–32. doi:10.1097/00001813-200001000-00005
63. Watanabe K, Inai S, Jinnouchi K, et al. Nuclear-factor kappa B (NF-kappa B)-inducible nitric oxide synthase (iNOS/NOS II) pathway damages the stria vascularis in cisplatin-treated mice. *Anticancer Res.* 2002;22(6C):4081–4085. PMID:12553036
64. Ruan RS. Possible roles of nitric oxide in the physiology and pathophysiology of the mammalian cochlea. *Ann N Y Acad Sci.* 2002;962:260–274. doi:10.1111/j.1749-6632.2002.tb04073.x
65. Rybak LP. Mechanisms of cisplatin ototoxicity and progress in otoprotection. *Curr Opin Otolaryngol Head Neck Surg.* 2007;15(5):364–369. doi:10.1097/MOO.0b013e3282ee452
66. Rybak LP, Whitworth CA, Mukherjea D, Ramkumar V. Mechanisms of cisplatin-induced ototoxicity and prevention. *Hear Res.* 2007;226(1–2):157–167. doi:10.1016/j.heares.2006.09.015
67. Bauer NC, Corbett AH, Doetsch PW. The current state of eukaryotic DNA base damage and repair. *Nucleic Acids Res.* 2015;43(21):10083–10101. doi:10.1093/nar/gkv1136
68. Friedberg EC. A history of the DNA repair and mutagenesis field The discovery of base excision repair. *DNA Repair (Amst).* 2016;37:A35–A39. doi:10.1016/j.dnarep.2015.12.003
69. Miller JH, Goodman MF. Tomas Lindahl: 2015 Nobel Laureate. *DNA Repair (Amst).* 2016;37:A29–A34. doi:10.1016/j.dnarep.2015.12.006
70. Radman M. Mismatch repair earns Nobel Prize in Chemistry 2015 to Paul Modrich for a biochemical tour de force. *DNA Repair (Amst).* 2016;37:A22–A28. doi:10.1016/j.dnarep.2015.12.004
71. Van Houten B. A tale of two cities: a tribute to Aziz Sancar's Nobel Prize in Chemistry for his molecular characterization of NER. *DNA Repair (Amst).* 2016;37:A3–A13. doi:10.1016/j.dnarep.2015.12.002
72. Cadet J, Davies KJA. Oxidative DNA damage & repair: an introduction. *Free Radic Biol Med.* 2017;107:2–12. doi:10.1016/j.freeradbiomed.2017.03.030
73. Santa-Gonzalez GA, Gomez-Molina A, Arcos-Burgos M, Meyer JN, Camargo M. Distinctive adaptive response to repeated exposure to hydrogen peroxide associated with upregulation of DNA repair genes and cell cycle arrest. *Redox Biol.* 2016;9:124–133. doi:10.1016/j.redox.2016.07.004
74. Van Houten B, Santa-Gonzalez GA, Camargo M. DNA repair after oxidative stress: current challenges. *Curr Opin Toxicol.* 2018;7:9–16. doi:10.1016/j.cotox.2017.10.009
75. Kim HJ, Oh GS, Shen A, et al. Augmentation of NAD(+) by NQO1 attenuates cisplatin-mediated hearing impairment. *Cell Death Dis.* 2014;5:e1292.
76. Gentilin E, Simoni E, Candito M, Cazzador D, Astolfi L. Cisplatin-induced ototoxicity: updates on molecular targets. *Trends Mol Med.* 2019;S1471-4914(19):30210–33012. doi:10.1016/j.molmed.2019.08.002
77. Lorenz RT, Cysewski GR. Commercial potential for Haematococcus microalgae as a natural source of astaxanthin. *Trends Biotechnol.* 2000;18(4):160–167. doi:10.1016/S0167-7799(00)01433-5
78. Facchinetti F, Dawson VL, Dawson TM. Free radicals as mediators of neuronal injury. *Cell Mol Neurobiol.* 1998;18(6):667–682. doi:10.1023/A:1020221919154
79. Kim SH, Kim H. Inhibitory effect of astaxanthin on oxidative stress-induced mitochondrial dysfunction—a mini-review. *Nutrients.* 2018;10(9). doi:10.3390/nu10091137
80. Bae JW, Kim MJ, Jang CG, Lee SY. Protective effects of heme oxygenase-1 against MPP+-induced cytotoxicity in PC-12 cells. *Neurol Sci.* 2010;31(3):307–313. doi:10.1007/s10072-010-0216-6
81. Borlongan CV, Kanning K, Poulos SG, Freeman TB, Cahill DW, Sanberg PR. Free radical damage and oxidative stress in Huntington's disease. *J Fla Med Assoc.* 1996;83(5):335–341. PMID: 8666972
82. Ferrante RJ, Browne SE, Shinobu LA, et al. Evidence of increased oxidative damage in both sporadic and familial amyotrophic lateral sclerosis. *J Neurochem.* 1997;69(5):2064–2074.
83. Rebrin I, Zicker S, Wedekind KJ, Paetau-Robinson I, Packer L, Sohal RS. Effect of anti-oxidant-enriched diets on glutathione redox status in tissue homogenates and mitochondria of the senescence-accelerated mouse. *Free Radical Bio Med.* 2005;39(4):549–557. doi:10.1016/j.freeradbiomed.2005.04.008
84. Chang CH, Chen CY, Chiou JY, Peng RY, Peng CH. Astaxanthin secured apoptotic death of PC12 cells induced by beta-amyloid peptide 25-35: its molecular action targets. *J Med Food.* 2010;13(3):548–556. doi:10.1089/jmf.2009.1291
85. Che HX, Li Q, Zhang TT, et al. Effects of astaxanthin and docosahexaenoic-acid-acylated astaxanthin on Alzheimer's disease in APP/PS1 double-transgenic mice. *J Agr Food Chem.* 2018;66(19):4948–4957. doi:10.1021/acs.jafc.8b00988
86. Lee DH, Kim CS, Lee YJ. Astaxanthin protects against MPTP/MPP plus -induced mitochondrial dysfunction and ROS production in vivo and in vitro. *Food Chem Toxicol.* 2011;49(1):271–280. doi:10.1016/j.fct.2010.10.029
87. Kim JH, Choi W, Lee JH, et al. Astaxanthin inhibits H2O2-mediated apoptotic cell death in mouse neural progenitor cells via modulation of P38 and MEK signaling pathways. *J Microbiol Biotechnol.* 2009;19(11):1355–1363. doi:10.4014/jmb.0906.06003
88. Al-Amin MM, Reza HM, Saadi HM, et al. Astaxanthin ameliorates aluminum chloride-induced spatial memory impairment and neuronal oxidative stress in mice. *Eur J Pharmacol.* 2016;777:60–69. doi:10.1016/j.ejphar.2016.02.062
89. Tso Mark OM, Lam Tim-Tak. Method of retarding and ameliorating central nervous system and eye damage. *Google Patents.* 1996;167.

90. Shen H, Kuo CC, Chou J, et al. Astaxanthin reduces ischemic brain injury in adult rats. *FASEB J*. 2009;23(6):1958–1968. doi:10.1096/fj.08-123281
91. Wu W, Wang X, Xiang Q, et al. Astaxanthin alleviates brain aging in rats by attenuating oxidative stress and increasing BDNF levels. *Food Funct*. 2014;5(1):158–166. doi:10.1039/c3fo60400d
92. Gross GJ, Hazen SL, Lockwood SF. Seven day oral supplementation with Cardax (disodium disuccinate astaxanthin) provides significant cardioprotection and reduces oxidative stress in rats. *Mol Cell Biochem*. 2006;283(1–2):23–30. doi:10.1007/s11010-006-2217-6
93. Pashkow FJ, Watumull DG, Campbell CL. Astaxanthin: a novel potential treatment for oxidative stress and inflammation in cardiovascular disease. *Am J Cardiol*. 2008;101(10A):58D–68D. doi:10.1016/j.amjcard.2008.02.010
94. Nakao R, Nelson OL, Park JS, Mathison BD, Thompson PA, Chew BP. Effect of astaxanthin supplementation on inflammation and cardiac function in BALB/c mice. *Anticancer Res*. 2010;30(7):2721–2725. PMID:20683004
95. Abdelzaher LA, Imaizumi T, Suzuki T, Tomita K, Takashina M, Hattori Y. Astaxanthin alleviates oxidative stress insults-related derangements in human vascular endothelial cells exposed to glucose fluctuations. *Life Sci*. 2016;150:24–31. doi:10.1016/j.lfs.2016.02.087
96. Alam MN, Hossain MM, Rahman MM, et al. Astaxanthin prevented oxidative stress in heart and kidneys of isoproterenol-administered aged rats. *J Diet Suppl*. 2018;15(1):42–54. doi:10.1080/19390211.2017.1321078
97. Pongkan W, Takatori O, Ni YH, et al. beta-Cryptoxanthin exerts greater cardioprotective effects on cardiac ischemia-reperfusion injury than astaxanthin by attenuating mitochondrial dysfunction in mice. *Mol Nutr Food Res*. 2017;61(10). doi:10.1002/mnfr.201601077
98. Fan CD, Sun JY, Fu XT, et al. Astaxanthin attenuates homocysteine-induced cardiotoxicity in vitro and in vivo by inhibiting mitochondrial dysfunction and oxidative damage. *Front Physiol*. 2017;8. doi:10.3389/fphys.2017.01041
99. Gammone MA, Riccioni G, D’Orazio N. Carotenoids: potential allies of cardiovascular health? *Food Nutr Res*. 2015;59:26762. doi:10.3402/fnr.v59.26762
100. Ni Y, Nagashimada M, Zhuge F, et al. Astaxanthin prevents and reverses diet-induced insulin resistance and steatohepatitis in mice: a comparison with vitamin E. *Sci Rep*. 2015;5:17192. doi:10.1038/srep17192
101. Zhang J, Zhang S, Bi J, Gu J, Deng Y, Liu C. Astaxanthin pretreatment attenuates acetaminophen-induced liver injury in mice. *Int Immunopharmacol*. 2017;45:26–33. doi:10.1016/j.intimp.2017.01.028
102. Takahashi K, Watanabe M, Takimoto T, Akiba Y. Uptake and distribution of astaxanthin in several tissues and plasma lipoproteins in male broiler chickens fed a yeast (*Phaffia rhodozyma*) with a high concentration of astaxanthin. *Brit Poultry Sci*. 2004;45(1):133–138. doi:10.1080/00071660410001668950
103. Rao AR, Sindhuja HN, Dharmesh SM, Sankar KU, Sarada R, Ravishankar GA. Effective inhibition of skin cancer, tyrosinase, and antioxidative properties by astaxanthin and astaxanthin esters from the green alga *haematococcus pluvialis*. *J Agr Food Chem*. 2013;61(16):3842–3851. doi:10.1021/jf304609j
104. Park JS, Chyun JH, Kim YK, Line LL, Chew BP. Astaxanthin decreased oxidative stress and inflammation and enhanced immune response in humans. *Nutr Metab*. 2010;7:18. doi:10.1186/1743-7075-7-18
105. Tominaga K, Hongo N, Fujishita M, Takahashi Y, Adachi Y. Protective effects of astaxanthin on skin deterioration. *J Clin Biochem Nutr*. 2017;61(1):33–39. doi:10.3164/jcbn.17-35
106. Yoon HS, Cho HH, Cho S, Lee SR, Shin MH, Chung JH. Supplementing with dietary astaxanthin combined with collagen hydrolysate improves facial elasticity and decreases matrix metalloproteinase-1 and -12 expression: a comparative study with placebo. *J Med Food*. 2014;17(7):810–816. doi:10.1089/jmf.2013.3060
107. Tominaga K, Hongo N, Karato M, Yamashita E. Cosmetic benefits of astaxanthin on humans subjects. *Acta Biochim Pol*. 2012;59(1):43–47. doi:10.18388/abp.2012_2168
108. Fang Q, Guo S, Zhou H, Han R, Wu P, Han C. Astaxanthin protects against early burn-wound progression in rats by attenuating oxidative stress-induced inflammation and mitochondria-related apoptosis. *Sci Rep*. 2017;7:41440. doi:10.1038/srep41440
109. Meepansan J, Rungjang A, Yingmema W, Deenonpoe R, Ponnikorn S. Effect of astaxanthin on cutaneous wound healing. *Clin Cosmet Investig Dermatol*. 2017;10:259–265. doi:10.2147/CCID.S142795
110. Zhang L, Wang H. Multiple mechanisms of anti-cancer effects exerted by astaxanthin. *Mar Drugs*. 2015;13(7):4310–4330. doi:10.3390/md13074310
111. Cairns RA, Harris IS, Mak TW. Regulation of cancer cell metabolism. *Nat Rev Cancer*. 2011;11(2):85–95. doi:10.1038/nrc2981
112. Kavitha K, Kowshik J, Kishore TK, Baba AB, Nagini S. Astaxanthin inhibits NF-kappaB and Wnt/beta-catenin signaling pathways via inactivation of Erk/MAPK and PI3K/Akt to induce intrinsic apoptosis in a hamster model of oral cancer. *Biochim Biophys Acta*. 2013;1830(10):4433–4444. doi:10.1016/j.bbagen.2013.05.032
113. Nagendraprabhu P, Sudhandiran G. Astaxanthin inhibits tumor invasion by decreasing extracellular matrix production and induces apoptosis in experimental rat colon carcinogenesis by modulating the expressions of ERK-2, NFkB and COX-2. *Invest New Drugs*. 2011;29(2):207–224. doi:10.1007/s10637-009-9342-5
114. Song XD, Zhang JJ, Wang MR, Liu WB, Gu XB, Lv CJ. Astaxanthin induces mitochondria-mediated apoptosis in rat hepatocellular carcinoma CBRH-7919 cells. *Biol Pharm Bull*. 2011;34(6):839–844. doi:10.1248/bpb.34.839
115. Zhang X, Zhao WE, Hu L, Zhao L, Huang J. Carotenoids inhibit proliferation and regulate expression of peroxisome proliferators-activated receptor gamma (PPARgamma) in K562 cancer cells. *Arch Biochem Biophys*. 2011;512(1):96–106. doi:10.1016/j.abb.2011.05.004
116. Su XZ, Chen R, Wang CB, Ouyang XL, Jiang Y, Zhu MY. Astaxanthin combine with human serum albumin to abrogate cell proliferation, migration, and drug-resistant in human ovarian carcinoma SKOV3 cells. *Anticancer Agents Med Chem*. 2019;19(6):792–801. doi:10.2174/1871520619666190225123003
117. Goto S, Kogure K, Abe K, et al. Efficient radical trapping at the surface and inside the phospholipid membrane is responsible for highly potent antiperoxidative activity of the carotenoid astaxanthin. *Bba-Biomembranes*. 2001;1512(2):251–258. doi:10.1016/S0005-2736(01)00326-1
118. Santa-Maria AR, Walter FR, Valkai S, et al. Lidocaine turns the surface charge of biological membranes more positive and changes the permeability of blood-brain barrier culture models. *Bba-Biomembranes*. 2019;1861(9):1579–1591. doi:10.1016/j.bbamem.2019.07.008
119. Matsushita Y, Suzuki R, Nara E, Yokoyama A, Miyashita K. Anti-oxidant activity of polar carotenoids including astaxanthin-β-glucoside from marine bacterium on PC liposomes. *Fish Sci*. 2000;66(5):980–985. doi:10.1046/j.1444-2906.2000.0155.x
120. Greene LS. Asthma and oxidant stress - nutritional, environmental, and genetic risk-factors. *J Am Coll Nutr*. 1995;14(4):317–324. doi:10.1080/07315724.1995.10718516

121. Dekkers JC, van Doornen LJP, Kemper HCG. The role of anti-oxidant vitamins and enzymes in the prevention of exercise-induced muscle damage. *Sports Med.* 1996;21(3):213–238. doi:10.2165/00007256-199621030-00005
122. Bennedsen M, Wang X, Willen R, Wadstrom T, Andersen LP. Treatment of H-pylori infected mice with anti-oxidant astaxanthin reduces gastric inflammation, bacterial load and modulates cytokine release by splenocytes. *Immunol Lett.* 1999;70(3):185–189. doi:10.1016/S0165-2478(99)00145-5
123. Higuera-Ciapara I, Felix-Valenzuela L, Goycoolea FM. Astaxanthin: a review of its chemistry and applications. *Crit Rev Food Sci.* 2006;46(2):185–196. doi:10.1080/10408690590957188
124. Wolf AM, Asoh S, Hiranuma H, et al. Astaxanthin protects mitochondrial redox state and functional integrity against oxidative stress. *J Nutr Biochem.* 2010;21(5):381–389. doi:10.1016/j.jnutbio.2009.01.011
125. Kidd P. Astaxanthin, cell membrane nutrient with diverse clinical benefits and anti-aging potential. *Altern Med Rev.* 2011;16(4):355–364.
126. Osterlie M, Bjerkgeng B, Liaen-Jensen S. Plasma appearance and distribution of astaxanthin E/Z and R/S isomers in plasma lipoproteins of men after single dose administration of astaxanthin. *J Nutr Biochem.* 2000;11(10):482–490. doi:10.1016/S0955-2863(00)00104-2. PMID:22214255
127. Yan TT, Zhao Y, Zhang X, Lin XT. Astaxanthin inhibits acetaldehyde-induced cytotoxicity in sh-sy5y cells by modulating Akt/CREB and p38MAPK/ERK signaling pathways. *Mar Drugs.* 2016;14(3):56. doi:10.3390/md14030056
128. Li JJ, Wang F, Xia YJ, et al. Astaxanthin pretreatment attenuates hepatic ischemia reperfusion-induced apoptosis and autophagy via the ROS/MAPK pathway in mice. *Mar Drugs.* 2015;13(6):3368–3387. doi:10.3390/md13063368
129. Yang X, Guo AL, Pang YP, et al. Astaxanthin attenuates environmental tobacco smoke-induced cognitive deficits: a critical role of p38 MAPK. *Mar Drugs.* 2019;17(1):24. doi:10.3390/md17010024
130. Niu TT, Xuan RR, Jiang LG, et al. Astaxanthin induces the Nrf2/HO-1 anti-oxidant pathway in human umbilical vein endothelial cells by generating trace amounts of ROS. *J Agr Food Chem.* 2018;66(6):1551–1559. doi:10.1021/acs.jafc.7b05493
131. Wu Q, Zhang XS, Wang HD, et al. Astaxanthin activates nuclear factor erythroid-related factor 2 and the anti-oxidant responsive element (Nrf2-ARE) pathway in the brain after subarachnoid hemorrhage in rats and attenuates early brain injury. *Mar Drugs.* 2014;12(12):6125–6141. doi:10.3390/md12126125
132. Wu ML, Ho YC, Yet SF. A central role of heme oxygenase-1 in cardiovascular protection. *Antioxid Redox Sign.* 2011;15(7):1835–1846. doi:10.1089/ars.2010.3726
133. Dinkova-Kostova AT, Talalay P. NAD(P)H:quinoneacceptor oxidoreductase 1 (NQO1), a multifunctional anti-oxidant enzyme and exceptionally versatile cytoprotector. *Arch Biochem Biophys.* 2010;501(1):116–123. doi:10.1016/j.abb.2010.03.019
134. Pan L, Zhou Y, Li XF, Wan QJ, Yu LH. Preventive treatment of astaxanthin provides neuroprotection through suppression of reactive oxygen species and activation of anti-oxidant defense pathway after stroke in rats. *Brain Res Bull.* 2017;130:211–220. doi:10.1016/j.brainresbull.2017.01.024
135. Tripathi DN, Jena GB. Astaxanthin intervention ameliorates cyclophosphamide-induced oxidative stress, DNA damage and early hepatocarcinogenesis in rat: role of Nrf2, p53, p38 and phase-II enzymes. *Mutat Res-Gen Tox En.* 2010;696(1):69–80. doi:10.1016/j.mrgentox.2009.12.014
136. Nakanishi A, Wada Y, Kitagishi Y, Matsuda S. Link between PI3K/AKT/PTEN pathway and NOX protein in diseases. *Aging Dis.* 2014;5(3):203–211. doi:10.14336/Ad.2014.0500203
137. Wang CM, Cai XL, Wen QP. Astaxanthin reduces isoflurane-induced neuroapoptosis via the PI3K/Akt pathway. *Mol Med Rep.* 2016;13(5):4073–4078. doi:10.3892/mmr.2016.5035
138. Zuluaga N, Barzegari A, Letourneur D, Gueguen V, Pavon-Djavid G. Oxidative stress regulation on endothelial cells by hydrophilic astaxanthin complex: chemical, biological, and molecular anti-oxidant activity evaluation. *Oxid Med Cell Longev.* 2017;2017:1–15. doi:10.1155/2017/8073798
139. Li ZR, Dong X, Liu HL, et al. Astaxanthin protects ARPE-19 cells from oxidative stress via upregulation of Nrf2-regulated Phase II enzymes through activation of PI3K/Akt. *Mol Vis.* 2013;19:1656–1666.
140. Ye QY, Zhang XD, Huang BX, Zhu YG, Chen XC. Astaxanthin suppresses MPP⁺-induced oxidative damage in PC12 cells through a Sp1/NR1 signaling pathway. *Mar Drugs.* 2013;11(4):1019–1034. doi:10.3390/md11041019
141. Takemoto Y, Hirose Y, Sugahara K, Hashimoto M, Hara H, Yamashita H. Protective effect of an astaxanthin nanoemulsion against neomycin-induced hair-cell damage in zebrafish. *Auris Nasus Larynx.* 2018;45(1):20–25. doi:10.1016/j.anl.2017.02.001

Drug Design, Development and Therapy

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>

Dovepress