Upregulated lncRNA THRIL/TNF- α Signals Promote Cell Growth and Predict Poor Clinical Outcomes of Osteosarcoma

This article was published in the following Dove Press journal:
OncoTargets and Therapy

Bo Xu
Xinmeng Jin*
Tieyi Yang
Yan Zhang
Shuyi Liu
Liang Wu
Hui Ying
Zhi Wang

Department of Orthopaedics, Gongli
Hospital of Pudong New Area, Shanghai
200135, People's Republic of China

*These authors contributed equally to
this work

Background: The immunosuppressive facet and tumorigenic role of TNF- α have been revealed in osteosarcoma (OS). Long noncoding RNA THRIL is identified to regulate TNF- α expression and participates in immune response. Thus, investigations on the clinical expression pattern of THRIL/TNF- α signal in OS would provide a potential target premise for OS patients.

Methods: We collected OS (n=83), nontumor tissues (n=37) and serum samples (n=83 for OS and n=40 for healthy control) to determine the expressions and clinical significance of THRIL/TNF- α signal. Knockdown of THRIL in OS cell lines MG63 and Saos2 in vitro and in vivo was performed to confirm its function in the development of OS.

Results: Elevated expression of THRIL was associated with increased TNF- α levels in OS tissues and serum samples. Combination of THRIL and TNF- α in tissues showed a more efficient diagnostic value for OS patients than either of them. Moreover, high-expressed THRIL was associated with larger tumor size, advanced Enneking stage and lung metastasis, whereas high TNF- α expression was found in patients with high histologic grade and patients who simultaneously harbor high THRIL and TNF- α showed the worst overall survival and metastasis-free survival. TNF- α signals increased OS cell vitalities in vitro but knockdown of THRIL inhibited TNF- α expressions, leading to impaired cell vitality, increased apoptosis and also downregulated epithelial to mesenchymal transition (EMT) phenotype and the ability of invasion, but these processes were restored by the treatment of TNF- α . The oncogenic role of THRIL/TNF- α signal was also confirmed in the xenograft model of MG63 cells.

Conclusion: Overexpressed THRIL and TNF- α promoted OS progression with efficient diagnostic and prognostic value. THRIL/TNF- α signal supported cell growth and EMT phenotype of OS cells in vitro and in vivo.

Keywords: lncRNA, THRIL, TNF- α , osteosarcoma, growth

Introduction

Osteosarcoma (OS) often occurs in children and adolescents with an incidence of 4.4 per million people worldwide and is the most common malignant bone tumor. Approximately 50% of all patients at the time of diagnosis develop metastases, which results in a high mortality rate despite improvements in radiotherapy, adjuvant chemotherapy and surgery.¹⁻³ Elucidation of the molecular basis of OS progression is conducive to developing effective therapeutic agents and improving OS patient's prognosis.

Correspondence: Zhi Wang
Department of Orthopaedics, Gongli
Hospital of Pudong New Area, NO. 219
Miaopu Road, Pudong New Area,
Shanghai 200135, People's Republic of
China
Tel +86 188 1782 1626
Email bbswangzhi@163.com

Persistent inflammation is known to promote and exacerbate malignancy. An intrinsic (driven by genetic events that cause neoplasia) and an extrinsic (driven by inflammatory conditions which predispose to cancer) pathway link inflammation and cancer.⁴ Tumor necrosis factor- α (TNF- α) is a proinflammatory cytokine that coordinates tissue homeostasis by regulating cytokine production, cell survival, and cell death.⁵ Accumulating evidence indicated that, in addition to being produced by immune cells within the tumor-associated microenvironment, TNF- α can be produced by a wide variety of malignant cells, including OS cells.⁶ Li et al reported that TNF- α may enhance the adaptive immune resistance mediated by IFN- γ -induced B7-H1 in hepatocellular carcinoma cells.⁷ TNF- α expression in ovarian carcinoma varies by histologic subtype and provides some support for the role of inflammation in ovarian carcinogenesis.⁸ Besides, TNF- α also promoted OS progression by maintaining tumor cells in an undifferentiated state.⁹ Activation of TNF- α signal enhanced CRL4B E3 ligase activity and regulates cell cycle progression in human OS cells.¹⁰ TNF- α regulated IL-34 expression in osteosarcoma cells and contributes to OS growth by increasing the neo-angiogenesis and the recruitment of M2 macrophage.¹¹ Thus, the regulator of TNF- α is essential for controlling its inflammation imbalance during cancer therapy of OS.

Long noncoding RNAs (lncRNAs) are encoded by a vast less explored region of the human genome and its disruption is intrinsically linked to diverse diseases including cell growth, inflammation and tumor metastasis.¹² lncRNA THRIL (TNF- α and hnRNPL related immune-regulatory lncRNA) is essential for the induction of TNF- α expression. Knockdown of THRIL caused dysregulation of TNF- α during innate activation of THP1 macrophages and was correlated with the severity of symptoms in patients with Kawasaki disease, an acute inflammatory disease of childhood.¹³ Upregulated THRIL was demonstrated to aggravate lipopolysaccharide (LPS)-induced osteoarthritis cell injury model via activating JAK1/STAT3 and NF- κ B pathways.¹⁴ Currently, the function of THRIL/TNF- α signal in cancer is unclear. One of the most prominent features of cancer is hypoxia, which was found to increase the THRIL expression.¹⁵ Therefore, considering the tumorigenic role of TNF- α in OS, the function and its clinical significance of THRIL/TNF- α signal needed to be investigated in OS patients.

In this study, we analyzed the expression pattern of THRIL/TNF- α in OS tissues and serum and further

revealed its diagnostic and prognostic value. We found that THRIL and TNF- α were co-upregulated in OS patients and predicted poor clinical outcomes. Knockdown of THRIL was performed in vitro and in vivo to confirm its role in OS growth and metastasis.

Materials and Methods

Ethics and Sample Collection

All methods used for this study were approved by the Ethics Committee of Gongli Hospital of Pudong New Area, and this study was conducted in accordance with the Declaration of Helsinki. The OS tissues and serum samples (n=83), nontumor tissues (n=37) and serum from healthy subjects were obtained from patients who had undergone surgical resection of OS in the Gongli Hospital of Pudong New Area and other hospitals in Acknowledgments. We confirmed that all patients provided written informed consent. The median age was 22 years and the median of tumor size was 8 cm, and 22 in 83 patients were found with lung metastases. The tumor stages were confirmed according to the Enneking staging system (ESS). The tissues or serum samples were isolated from surgical removal or patients' blood and then stored at -80°C until use. All the patients were followed-up until 20 April 2019. The minimum follow-up time is 3 months and the maximum follow-up time is 58 months (median: 26 months). The high or low expressions of THRIL/TNF- α were defined by the median of its expression.

RNA Extraction and RT-Q-PCR

Total RNA was extracted from clinical specimens or cells via the TRIzol reagent (Invitrogen, USA), A260/A280 nm >1.8 was qualified for quantitative analysis. Then, RNAs were transcribed into cDNA using superscriptase II (Invitrogen, USA) according to the manufacturer's instructions. The cDNA was subjected to real time-polymerase chain reaction using Power SYBR[®] Green PCR Master Mix (Invitrogen, USA). The following primers were used: THRIL-F: GAGTGCAGTGGCGTGATCTC, R: AAAATTAGTCAGGCATGGTGGTG; TNF- α -F: CCTCTCTCTAA TCAGCCCTCTG; R: GAGGACCTGGGAGTAGATGA G; Bcl-2-F: GGTGGGGTCATGTGTGTGG, R: CGGTTCAGGTACTCAGTCATCC; E-cadherin-F: CGAGAGCTA CACGTTACCGG, R: GGGTGTGCGAGGGAAAAATAG G; Vimentin-F: GACGCCATCAACACCGAGTT, R: CTTTGTCGTTGGTTAGCTGGT. The expression levels of genes were estimated via $2^{-\Delta\Delta\text{Ct}}$ method and were

normalized to GAPDH for gene expression. Each sample was measured in triplicate.

ELISA

The serum levels of TNF- α were qualified by Human TNF- α ELISA kit (Cat#:EHC103a) according to the manufacturer's instructions. The detection sensitivity was 7.8 pg/mL. All samples were tested by duplication, and absorbency was measured at 620 nm by a microplate reader.

Cell Culture and Reagents

Human OS cell lines MG63 and Saos2 were obtained from the American Type Culture Collection (ATCC) and cultured in DMEM supplement with 10% FBS (GIBCO) and 1% penicillin/streptomycin. All the cells were cultured at 37°C water-saturated 5% CO₂ atmosphere. siRNA-THRIL or negative control was conducted by RiboBio (Guangzhou, china), siRNA-THRIL-1: 5'- GCTCTATGCACAGATAA ATTT -3'; siRNA-THRIL-2: 5'- GCTCAGTCGTGTTTA TTAATT -3'; siRNA-TNFR1: 5'-CAAAGGAACCUACUU GUACUU-3'. Human recombinant TNF- α was purchased from Preoteintech (#HZ-1014, WUHAN, China). The

antibody to E-cadherin and Vimentin were obtained from Abcam (USA).

Estimation of Cell Growth

MG63 and Saos2 cells were transfected of siRNA-THRIL with/without the treatment of TNF- α (50 ng/mL) for 48 h, the cell vitality was determined by CCK-8. Briefly, cells were harvested and washed with PBS, and then cell counting kit-8 (Kumamoto, Japan) mixed with DMEM was used for cell viability assay according to the manufacturer's instructions., and the absorbance was measured at 450 nm by a microplate reader. In addition, apoptosis was determined by flow cytometry. Briefly, cells were stained with 2 μ L of annexin V mixed with 2 μ L of propidium iodide (PI, eBioscience) were used according to the manufacturer's instructions and were analyzed using a flow cytometry.

Transwell Assay

To determine the role of THRIL in invasion of OS cells, 2×10^4 cells after transfection were suspended with 200 μ L of serum-free medium. The Corning Polycarbonate Membrane Insert transwell chambers (Product #3422, Corning Costar Corp, Cambridge, MA, USA) were used

Figure 1 The expressions of THRIL and TNF- α in OS tissues and serum. (A and B) The mRNA levels of THRIL and TNF- α were analyzed in OS tissues (n=83) and nontumor tissues (n=37) by Q-PCR. (C) The serum TNF- α level was confirmed in the serum derived from OS patients (n=83) and independent healthy subjects (n=40) by ELISA. (D and E) The correlation between THRIL and TNF- α in OS tissues and serum was determined. (F) ROC curves were performed to evaluate the diagnostic role of THRIL and TNF- α alone, or THRIL/TNF- α combination. *** $P < 0.001$, data represent the means $\pm$ SD.

to perform invasion assays. The Transwell membranes were pre-coated with Matrigel (BD Biosciences, USA) at 37°C for at least 2 hrs to form a reconstructed basement membrane. After incubation, cells at the top chamber were removed with cotton swabs, and those on the lower surface were fixed with methanol, stained by crystal violet and counted under an inverted microscope.

Immunofluorescence Assays

MG63 and Saos2 cells with THRIL knockdown were cultured in a plastic chamber on microscope glass slides (Millicell EZ Slide, Millipore, Billerica, MA, USA) with TNF- α (50 ng/mL) stimulation for 48 hrs. The cells were then washed in PBS, fixed in 4% paraformaldehyde for 10 mins at room temperature, permeabilized with triton X-100 0.1% for 20 mins. The cells were incubated with primary antibody against E-cadherin and Vimentin for 120 mins. After washings, Alexa Fluor 488/594 secondary antibody

was added for 90 mins at room temperature. Nuclei were stained with DAPI, which was observed under an FV10i confocal microscope (OLYMPUS, Tokyo, Japan).

Tumor Model

To investigate the role of THRIL in the OS development, the xenograft model of human MG63 cells was established, MG63 cells were transfected with lentivirus vector of siRNA-THRIL, or negative control, 2×10^6 conditional MG63 cells were subcutaneously injected in rear flank of nude mice (6 per group), TNF- α (500 ng/mice) was treated i.p. twice a week for 3 weeks. The tumor sizes were measured 3 days apart and the tumor volumes were calculated: $V \text{ (cm}^3\text{)} = \text{width}^2 \text{ (cm}^2\text{)} * \text{length (cm)}/2$. The animal study was approved by institutional animal research committee of Gongli Hospital of Pudong New Area and that animals were cared for following the guidelines for use and care of laboratory animals.

Table I Correlations Between the Expressions of THRIL and TNF- α with Clinicalpathologic Characteristics in OS Patients

Characteristics	n	THRIL		P	χ^2	TNF- α		P	χ^2
		Low (n%)	High (n%)			Low (n%)	High (n%)		
Gender									
Male	39	19(48.72)	20(51.28)	0.907	0.014	21(53.85)	18(46.15)	0.332	0.942
Female	44	22(50)	22(50)			19(43.18)	25(56.81)		
Age									
<22	45	25(55.5)	20(44.4)	0.222	1.491	23(51.11)	22(48.89)	0.563	0.335
≥22	38	16(42.1)	22(57.89)			17(44.74)	21(55.26)		
Tumor size (cm)									
≤8	35	22(62.85)	13(37.14)	0.036*	4.386	18(51.42)	17(48.57)	0.614	0.254
>8	48	19(39.58)	29(60.42)			22(45.83)	26(54.17)		
Anatomic site									
Femur	52	24(46.15)	28(53.85)	0.61	0.988	21(40.38)	31(59.62)	0.128	2.315
Tibia	18	9(50)	9(50)			11(61.11)	7(38.89)		
Other	13	8(61.54)	5(38.46)			8(61.54)	5(38.46)		
Enneking stage									
I-IIA	34	23(67.65)	11(32.35)	0.006*	7.763	20(58.82)	14(41.17)	0.518	0.417
IIB-III	49	18(36.73)	31(63.26)			20(40.81)	29(59.18)		
Histologic grade									
Low	40	21(52.5)	19(47.5)	0.586	0.297	24(60)	16(40)	0.038*	4.311
High	43	20(46.51)	23(53.49)			16(37.20)	27(62.79)		
Lung metastasis									
No	61	35(57.38)	26(42.62)	0.015*	5.862	29(47.54)	32(52.46)	0.843	0.039
Yes	22	6(27.27)	16(72.73)			11(50)	11(50)		
Total	83	41(100)	42(100)			40(100)	43(100)		

Note: *P <0.05, statistically significant by Chi-square test.

Table 2 Clinical Significance of Simultaneously High-Expressed or Low-Expressed THRIL and TNF- α in OS Patients

Characteristics	n	THRIL ^{Low} /TNF- α ^{Low} (n/%)	THRIL ^{High} /TNF- α ^{High} (n/%)	P	χ^2
Gender					
Male	30	12(40)	18(60)	0.18	1.791
Female	24	14(58.33)	10(41.67)		
Age					
<22	26	14(53.85)	12(46.15)	0.491	0.652
≥22	28	12(42.86)	16(57.14)		
Tumor size(cm)					
≤8	25	18(72)	7(28)	0.001*	10.61
>8	29	8(27.59)	21(72.41)		
Anatomic site					
Femur	42	22(52.38)	20(47.62)	0.467	1.523
Tibia	8	3(37.50)	5(62.50)		
Other	4	1(25)	3(75)		
Enneking stage					
I-IIA	22	16(72.73)	6(27.27)	0.003*	8.984
IIB-III	32	10(31.25)	22(68.75)		
Histologic grade					
Low	24	14(58.33)	10(41.67)	0.18	1.795
High	30	12(40)	18(60)		
Lung metastasis					
No	37	22(59.46)	15(40.54)	0.014*	6.023
Yes	17	4(23.53)	13(76.47)		
Total	54	26(100)	28(100)		

Note: *P <0.05, statistically significant by Chi-square test.

Statistical Analyses

The results are analyzed by the Statistical Package for Social Sciences version 16.0 (SPSS 16.0, SPSS Inc., Chicago, IL, USA) and the Prism statistical software package (Version 5.0, Graphpad Software Inc.). Kolmogorov–Smirnov and Shapiro–Wilk tests showed that the expression of THRIL/TNF- α did not follow a normal distribution. The Mann–Whitney *U*-test was used to compare the two groups (e.g., normal versus OS) and the differences between more than two groups were analyzed by the Kruskal–Wallis test. The receiver operating characteristic (ROC) curve was used to analyze its diagnostic value. Correlations between the expression of THRIL/TNF- α were analyzed by Spearman's Rho analysis. Correlations between THRIL/TNF- α and clinicopathological characters were analyzed by Pearson Chi-square. Kaplan–Meier survival curves and the log-rank statistic were used to analyze the prognostic significance of THRIL/TNF- α . Cox proportional hazard regression was used for univariate and multivariate analysis of overall survival according to THRIL/TNF- α

expression. *P* <0.05 was considered statistically significant. All experiments were performed at least 3 times.

Results

THRIL Is Upregulated in OS and Positively Correlates with TNF- α

We first determined the expression of THRIL and TNF- α during the development of OS. THRIL and TNF- α were found to be upregulated in OS tissues (n=83) when compared to which in nontumor tissues (n=37) (Figure 1A and B). The enhanced TNF- α level was confirmed in the serum derived from OS patients (n=83) when compared with independent healthy subjects (n=40) (Figure 1C). Since THRIL could regulate TNF- α expression, we analyzed the correlation between THRIL and TNF- α . The results showed that THRIL is positively associated with TNF- α in OS tissues and serum (Figure 1D and E), suggesting that THRIL/TNF- α signal participated in the OS development.

Figure 2 The prognostic roles of THRIL and TNF- α in OS. The expressions of THRIL/TNF- α were divided into High and Low groups by median. (A and B) The overall survival time and metastasis-free time of OS patients were analyzed according to the THRIL expression. (C and D) The overall survival time and metastasis-free time of OS patients were analyzed according to the TNF- α expression. (E and F) The overall survival time and metastasis-free time of OS patients were analyzed according to the THRIL and TNF- α expression.

The receiver operating characteristic (ROC) curve suggests that THRIL (AUC=0.773) and TNF- α (AUC=0.794) are good diagnostic markers to discriminate OS patients from healthy people, but combined with two factors showed more predominant diagnostic value (AUC=0.852) (Figure 1F).

Elevated Expression of THRIL and TNF- α Occurs in OS Patients with Advanced Tumor Progression

Considering the upregulated THRIL/TNF- α signal in OS samples, we next determined the clinical significance of THRIL/TNF- α in OS. The expression of THRIL/TNF- α was divided into High and Low groups by median. The results showed that patients with larger tumor size, advanced Enneking stage and distant metastasis have higher THRIL levels in OS tissues. However, THRIL showed no relationships with gender, age, anatomic site and histology grade. Elevated TNF- α level was found in patients with high histology grade, which was consistent with the previous findings that TNF- α promoted OS progression by maintaining tumor cells in an undifferentiated state (Table 1).

When the combination of THRIL and TNF- α was used for analysis, we found that the patients whose simultaneously high-expressed THRIL and TNF- α (THRIL^{High}/TNF- α ^{High}) showed more advanced OS progression with a higher incidence of lung metastasis than patients with low-expressed THRIL and TNF- α (THRIL^{Low}/TNF- α ^{Low}) (Table 2).

High THRIL and TNF- α Levels Predict Short Overall Survival and Metastasis-Free Time

The prognostic role of THRIL/TNF- α signal was also analyzed. The results showed that high expression of THRIL predicted shorter overall survival time ($P=0.024$) and metastasis-free survival (MFS) ($P=0.033$) (Figure 2A and B), but TNF- α showed no impacts on overall survival time and metastasis-free time of OS patients (Figure 2C and D).

We further analyzed the survival time of patients with THRIL^{High}/TNF- α ^{High}, the results indicated that, compared with THRIL alone, combination of THRIL and TNF- α was more efficient to predict the decreased survival time of OS patients (Figure 2E and F).

Table 3 Prognostic Factors in the Cox Proportional Hazard Model for Overall Survival

Variables	Overall Survival					
	HR	Univariate 95% CI	Sig	HR	Multivariate 95% CI	Sig.
Gender Male vs female	0.885	0.563–1.427	0.812			
Age <22 vs ≥22	1.223	0.734–1.891	0.438			
Tumor size(cm) <8 vs ≥8	2.863	1.233–3.675	0.013*	2.473	2.303–4.392	0.006*
Anatomic site Femur vs other	0.935	0.942–1.343	0.826			
Enneking stage I-IIA vs. II B-III	2.173	1.694–4.726	0.031*	1.943	0.813–2.046	0.328
Histologic grade Low vs high	0.917	0.459–1.558	0.923			
Lung metastasis - vs +	3.625	1.301–4.862	0.002*	4.115	2.263–6.345	0.001*
THRIL Low vs high	2.075	1.630–2.684	0.025*	2.347	1.349–3.758	0.023*
TNF-α Low vs high	1.234	0.676–1.602	0.463			

Note: *P <0.05, statistically significant.

Abbreviations: HR, hazard ratio; CI, confidence interval.

THRIL Is an Independently Prognostic Factor for OS Patients

We further performed univariate and multivariate analyses, the results found that gender, age, anatomic site, Enneking stage, histologic grade and TNF- α were not independent prognostic indicators for overall survival time. In addition, the tumor size, lung metastasis and the expression of THRIL were independent prognostic factors for the overall survival time of patients with OS (Table 3). However, THRIL was not an independent prognostic factor for metastasis-free survival of patients with OS (Table 4).

Knockdown of THRIL Inhibits OS Growth and Invasion via TNF- α in vitro and in vivo

The function of TNF- α alone was investigated and found that the treatment of TNF- α (50 ng/mL) elevated the cell vitality of OS cells, which was consistent with previous reports¹⁶ (Figure 3A). Then, we performed knockdown of

THRIL in two OS cell lines MG63 and Saos2 to confirm its pro-tumor function. The results indicated that knockdown of THRIL inhibited the expression of TNF- α by two siRNA. SiRNA-1 was selected for further study (Figure 3B and C). To confirm the role of TNF- α signal for THRIL function, we inhibited TNF receptor (TNFR) in OS cells in response to TNF- α treatment and THRIL knockdown. The results demonstrated that the OS cells with THRIL knockdown showed impaired cell vitality, but the treatment of TNF- α (50 ng/mL) to the OS cells with THRIL knockdown could restore its cell vitality, which was dependent on TNFR signals. Inhibition of TNFR by siRNA suppressed the function of TNF- α in the OS cells with THRIL knockdown. Thus, the function of THRIL is dependent on TNF- α /TNFR signals (Figure 3D and E).

Thus, we further investigated the role of THRIL/TNF- α signal in cell apoptosis and metastasis of two OS cell lines. Inhibition of THRIL induced apoptosis in 48 h, which could be restored by the treatment of TNF- α (50 ng/mL), indicating that THRIL promoted OS growth via TNF- α (Figure 3F). The ability of invasion was also

Table 4 Prognostic Factors in the Cox Proportional Hazard Model for Metastasis-Free Survival

Variables	Metastasis-Free Survival					
	HR	Univariate 95% CI	Sig	HR	Multivariate 95% CI	Sig.
Gender Male vs female	0.923	0.623–1.969	0.903			
Age <22 vs ≥22	1.058	0.822–1.358	0.828			
Tumor size(cm) <8 vs ≥8	2.266	1.384–3.294	0.033*	1.789	1.873–2.545	0.128
Anatomic site Femur vs other	0.867	0.683–1.225	0.774			
Enneking stage I-IIA vs. IIB-III	1.984	1.213–2.685	0.053			
Histologic grade Low vs High	0.881	0.739–1.304	0.893			
Lung metastasis - vs +	2.782	1.426–3.678	0.012*	3.302	1.892–5.367	0.008*
THRIL Low vs high	1.976	1.567–3.869	0.038*	1.347	0.835–2.112	0.236
TNF-α Low vs high	0.789	0.492–1.263	0.673			

Note: *P <0.05, statistically significant.

Abbreviations: HR, hazard ratio; CI, confidence interval.

determined by Transwell assay, we found that knockdown of THRIL inhibited OS invasion but the addition of TNF-α abrogated this suppression and upregulated the ability of invasion (Figure 3G). This process might be attributed to changed EMT phenotype of OS cells. Inhibition of THRIL downregulated the expression of vimentin but TNF-α (50 ng/mL) could restore the levels of vimentin in MG63 and Saos2 cells (Figure 3H).

We also investigated the role of THRIL in vivo. The results showed that the mice with THRIL knockdown by lentivirus have smaller tumor volume than that with a negative control. However, treatment of TNF-α (500 ng/mice) could restore the antitumor function of THRIL knockdown (Figure 3I). In line with this, the antiapoptosis gene Bcl-2 and Vimentin were downregulated by THRIL knockdown, but were recovered by the treatment of TNF-α in vivo (Figure 3J).

Discussion

Continuous exposure to inflammatory cytokines (such as TNF-α, IL-6, IL-8 and IL-1β, etc.) is known to cause tumorigenesis.⁴ TNF-α can be produced by leukemia

cells to mediate cell survival and higher TNF-α concentrations in leukemia patients were correlated with heavier fatigue and poorer quality of life.¹⁷ TNF-α also promoted OS progression by maintaining tumor cells in an undifferentiated state.⁹ Many lncRNAs have been reported to regulate and control the balance of inflammation during several diseases. Li et al reported that THRIL bound specifically to heterogenous nuclear ribonucleoprotein L (hnRNPL) and formed a functional THRIL–hnRNPL complex that regulated transcription of the TNF-α gene by binding to its promoter. Transcriptome analysis revealed that THRIL was required for expression of many immune-response genes, including TNF-α.¹³ We here identified the positive regulatory role of THRIL to TNF-α in OS cells. Knockdown of THRIL inhibited TNF-α expression and TNF-α-induced OS cell growth. Similarly, NF-κB-interacting lncRNA (NKILA) was highly induced in response to IL-1β and TNF-α stimulation.¹⁸ lncRNA Mirt2 functioned as a checkpoint to prevent aberrant activation of inflammation and was a potential regulator of macrophage polarization via attenuating Lys63 (K63)-linked ubiquitination of TRAF6,

Figure 3 Knockdown of THRIL inhibits cell growth and invasion of OS cell via TNF- α . (A) TNF- α (50 ng/mL) was treated to two OS cell lines MG63 and Saos2 and the cell vitality was determined. (B and C) Knockdown of THRIL in two OS cell lines MG63 and Saos2 via siRNA, and the expressions of THRIL/TNF- α were determined at 24 hrs by Q-PCR. (D and E) After knockdown of THRIL/TNFR1, the cells were treated with or without TNF- α (50 ng/mL) for 48 h, and cell vitality and apoptosis of two cell lines were analyzed by CCK-8. (F) After knockdown of THRIL with or without TNF- α stimulation, the cell apoptosis was analyzed by flow cytometry. (G) Transwell assay was performed to determine the invasion of OS cell after THRIL knockdown. Five random views were selected to count the cells. (H) Immunofluorescence assay was performed to analyze the expression of E-cadherin (Green) and Vimentin (Red) in two cell lines. (I) 2×10^6 conditional MG63 cells were transfected with a lentivirus vector of siRNA-THRIL or negative control and were subcutaneously injected in the rear flank of nude mice (6 per group). TNF- α (500 ng/mice) was treated i.p. The mean tumor size (mm³) was analyzed. (J) The expressions of THRIL, TNF- α , Bcl-2, E-cadherin and Vimentin in tumor tissues were determined by Q-PCR. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Data represent the means $\pm$ SD.

thus inhibiting activation of NF- κ B and MAPK pathways.¹⁹ lnc-IL7R was capable of diminishing the LPS-induced inflammatory response via repressing LPS-induced E-selectin, VCAM-1, IL-6, and IL-8.²⁰ However, the inflammatory lncRNAs in during the development of OS remains unclear. TGF- β or TNF- α treatment could enhance the expression of oncogenic lncRNA UCA1 for cell growth, migration and invasion of OS cells.²¹ We also found that TNF- α -induced EMT phenotype could be restored by THRIL knockdown via upregulation of E-cadherin and downregulation of Vimentin. Thus, controlling chronic inflammation is crucial to prevent tumor progression.

Currently, the researches on the role of THRIL focused on inflammation-related diseases. Upregulated THRIL was found to predict increased risk of acute respiratory distress syndrome and positively correlates with disease severity, inflammation, and mortality in sepsis patients.²² In this study, we found that lncRNA THRIL upregulated TNF- α in tumor tissue and serum of OS patients, which could be efficient diagnostic and prognostic biomarker. THRIL was also positively associated with CRP, PCT, TNF- α , and IL-1 β levels in sepsis patients.²² In Kawasaki disease, an acute inflammatory disease of childhood, THRIL was correlated with the severity of symptoms of patients.¹³ In hypoxia-induced injuries, THRIL inhibition represents a protective effect against hypoxia-induced injuries in H9C2 cells.²² However, THRIL RNA expression was significantly decreased in autism spectrum disorder (ASD) children, although TNF- α , IL-1 β and IL-17 were significantly increased in ASD children.²³ THRIL was significantly decreased in patients with type 2 diabetes mellitus compared to control subjects, which were positively correlated with poor glycemic control, insulin resistance.²⁴ But the investigation of THRIL in cancer is limited. We found that the expression of THRIL was increased in OS patients. High-expressed THRIL was positively associated with advanced OS progression including tumor size, clinical stage and lung metastasis. Patients with upregulated THRIL have shorter overall survival time and metastasis-free time. Esfandi et al investigated the role of THRIL in lung cancer, but the expression of THRIL was not changed in lung cancer samples and their corresponding adjacent noncancerous tissues.²⁵ Similar results were also reported in bladder cancer.²⁶ Interestingly, Wang et al found a significantly decreased risk of precancerous cervical lesions for the THRIL rs7133268 AG genotype (odds ratio

adjusted=0.63).²⁷ These findings suggested the content-dependent role of THRIL during disease progression.

The function of THRIL was related to its regulatory role of TNF- α . We here confirmed that THRIL was positively associated with upregulated TNF- α in OS tissues and serum. Combination of THRIL and TNF- α showed prominent diagnostic value for OS patients. However, TNF- α exerts diverse functions in cancer, other than the role of promoting tumor death. For example, TNF- α could promote cell growth, migration and invasion of MCF-7 estrogen receptor (ER)-positive and MDA-MB-231 (triple-negative) breast cancer cell lines, partially by inducing the expression of MMPs.²⁸ Consistently, we here found that knockdown of THRIL in vivo could inhibit cell growth and invasion, which was abrogated by the treatment of TNF- α . The clinical analysis also showed that patients with high expression of TNF- α have high histology grade and THRIL^{High}/TNF- α ^{High} predicted worst overall survival and metastasis-free survival of OS patients.

In sum, we here investigated the clinical role and function of upregulated THRIL/TNF- α signal in OS and found that patients with THRIL^{High}/TNF- α ^{High} showed poor clinical outcomes and short metastasis-free survival time, which was involved with THRIL/TNF- α signal-supported cell growth and EMT phenotype for OS progression.

Acknowledgment

We would like to thank Shanghai Changzheng Hospital, Shanghai Ninth People's Hospital and Shanghai Ruijin Hospital for their precious OS samples.

Author Contributions

All authors contributed to data analysis, drafting or revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Funding

This work was supported by Key Discipline Construction Project of Pudong Health Bureau of Shanghai (Grant No. PWZxk2017-18 to TYY) and Discipline Construction Project of Pudong New Area Commission of Health and Family Planning (Grant No. PWYts2018-03 to TYY).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Toki S, Kobayashi E, Yoshida A, et al. A clinical comparison between dedifferentiated low-grade osteosarcoma and conventional osteosarcoma. *Bone Joint J.* **2019**;101-B(6):745–752. doi:10.1302/0301-620X.101B6.BJJ-2018-1207.R1
2. Ahmed G, Zamzam M, Kamel A, et al. Effect of timing of pulmonary metastasis occurrence on the outcome of metastasectomy in osteosarcoma patients. *Journal of Pediatric Surgery.* **2019**;54(4):775–779. doi:10.1016/j.jpedsurg.2018.06.019
3. Kumar R, Kumar M, Malhotra K, Patel S. Primary osteosarcoma in the elderly revisited: current concepts in diagnosis and treatment. *Curr Oncol Rep.* **2018**;20(2):13. doi:10.1007/s11912-018-0658-1
4. Capece D, Verzella D, Tessitore A, Alesse E, Capalbo C, Zazzeroni F. Cancer secretome and inflammation: the bright and the dark sides of NF-kappaB. *Semin Cell Dev Biol.* **2018**;78:51–61. doi:10.1016/j.semcdb.2017.08.004
5. Salomon BL, Leclerc M, Tosello J, Ronin E, Piaggio E, Cohen JL. Tumor necrosis factor alpha and regulatory T cells in oncoimmunology. *Front Immunol.* **2018**;9:444. doi:10.3389/fimmu.2018.00444
6. Ciebia M, Wlodarczyk M, Zgliczynska M, et al. The role of tumor necrosis factor alpha in the biology of uterine fibroids and the related symptoms. *Int J Mol Sci.* **2018**;19:12. doi:10.3390/ijms19123869
7. Li N, Wang J, Zhang N, et al. Cross-talk between TNF-alpha and IFN-gamma signaling in induction of B7-H1 expression in hepatocellular carcinoma cells. *Cancer Immunol Immunother.* **2018**;67(2):271–283. doi:10.1007/s00262-017-2086-8
8. Gupta M, Babic A, Beck AH, Terry K. TNF-alpha expression, risk factors, and inflammatory exposures in ovarian cancer: evidence for an inflammatory pathway of ovarian carcinogenesis?. *Human pathol.* **2016**;54:82–91.
9. Mori T, Sato Y, Miyamoto K, et al. TNFalpha promotes osteosarcoma progression by maintaining tumor cells in an undifferentiated state. *Oncogene.* **2014**;33(33):4236–4241. doi:10.1038/ncr.2013.545
10. Zhang C, Chen B, Jiang K, Lao L, Shen H, Chen Z. Activation of TNF-alpha/NF-kappaB axis enhances CRL4B(DCAF)(11) E3 ligase activity and regulates cell cycle progression in human osteosarcoma cells. *Mol Oncol.* **2018**;12(4):476–494. doi:10.1002/1878-0261.12176
11. Segaliny AI, Mohamadi A, Dizier B, et al. Interleukin-34 promotes tumor progression and metastatic process in osteosarcoma through induction of angiogenesis and macrophage recruitment. *Int J Cancer.* **2015**;137(1):73–85. doi:10.1002/ijc.29376
12. Lin C, Yang L. Long noncoding RNA in cancer: wiring signaling circuitry. *Trends Cell Biol.* **2018**;28(4):287–301. doi:10.1016/j.tcb.2017.11.008
13. Li Z, Chao TC, Chang KY, et al. The long noncoding RNA THRIL regulates TNFalpha expression through its interaction with hnRNPL. *Proc National Acad Sci USA.* **2014**;111(3):1002–1007. doi:10.1073/pnas.1313768111
14. Liu G, Wang Y, Zhang M, Zhang Q. Long non-coding RNA THRIL promotes LPS-induced inflammatory injury by down-regulating microRNA-125b in ATDC5 cells. *Int Immunopharmacol.* **2019**;66:354–361. doi:10.1016/j.intimp.2018.11.038
15. Xia J, Jiang N, Li Y, Wei Y, Zhang X. The long noncoding RNA THRIL knockdown protects hypoxia-induced injuries of H9C2 cells through regulating miR-99a. *Cardiol J.* **2018**;26:564–574.
16. Tan W, Luo X, Li W, et al. TNF-alpha is a potential therapeutic target to overcome sorafenib resistance in hepatocellular carcinoma. *EBio Med.* **2019**;40:446–456. doi:10.1016/j.ebiom.2018.12.047
17. Zhou X, Li Z, Zhou J. Tumor necrosis factor alpha in the onset and progression of leukemia. *Exp Hematol.* **2017**;45:17–26. doi:10.1016/j.exphem.2016.10.005
18. Lu Z, Chen Z, Li Y, et al. TGF-beta-induced NKILA inhibits ESCC cell migration and invasion through NF-kappaB/MMP14 signaling. *J Mol Med (Berl).* **2018**;96(3–4):301–313. doi:10.1007/s00109-018-1621-1
19. Du M, Yuan L, Tan X, et al. The LPS-inducible lncRNA Mirt2 is a negative regulator of inflammation. *Nature Commun.* **2017**;8(1):2049. doi:10.1038/s41467-017-02229-1
20. Cui H, Xie N, Tan Z, et al. The human long noncoding RNA lnc-IL7R regulates the inflammatory response. *Eur J Immunol.* **2014**;44(7):2085–2095. doi:10.1002/eji.201344126
21. Li Q, Xing W, Gong X, Wang Y. Long non-coding RNA urothelial carcinoma associated 1 promotes proliferation, migration and invasion of osteosarcoma cells by regulating microRNA-182. *Cell Physiol Biochem.* **2018**;51(3):1149–1163. doi:10.1159/000495493
22. Wang Y, Fu X, Yu B, Ai F. Long non-coding RNA THRIL predicts increased acute respiratory distress syndrome risk and positively correlates with disease severity, inflammation, and mortality in sepsis patients. *J Clin Lab Anal.* **2019**;e22882.
23. Xie J, Huang L, Li X, et al. Immunological cytokine profiling identifies TNF-alpha as a key molecule dysregulated in autistic children. *Oncotarget.* **2017**;8(47):82390–82398. doi:10.18632/oncotarget.19326
24. Sathishkumar C, Prabu P, Mohan V, Balasubramanyam M. Linking a role of lncRNAs (long non-coding RNAs) with insulin resistance, accelerated senescence, and inflammation in patients with type 2 diabetes. *Human Genomics.* **2018**;12(1):41. doi:10.1186/s40246-018-0173-3
25. Esfandi F, Taheri M, Omrani MD, et al. Expression of long non-coding RNAs (lncRNAs) has been dysregulated in non-small cell lung cancer tissues. *BMC Cancer.* **2019**;19(1):222. doi:10.1186/s12885-019-5435-5
26. Abdolmaleki F, Ghafouri-Fard S, Taheri M, et al. Expression analysis of a panel of long non-coding RNAs (lncRNAs) revealed their potential as diagnostic biomarkers in bladder cancer. *Genomics.* **2019**. doi:10.1016/j.ygeno.2019.04.020
27. Wang Y, Liu Y, Li Z, et al. Association between MALAT1 and THRIL polymorphisms and precancerous cervical lesions. *Genetic Testing Mol Biomarkers.* **2018**;22(9):509–517. doi:10.1089/gtmb.2018.0097
28. Wolczyk D, Zaremba-Czogalla M, Hryniewicz-Jankowska A, et al. TNF-alpha promotes breast cancer cell migration and enhances the concentration of membrane-associated proteases in lipid rafts. *Cell Oncol (Dordr).* **2016**;39(4):353–363. doi:10.1007/s13402-016-0280-x

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>

Dovepress