LncRNA SNHG1 Regulates the Progression of Esophageal Squamous Cell Cancer by the miR-204/HOXC8 Axis

This article was published in the following Dove Press journal:
OncoTargets and Therapy

Hao Miao Li¹
Yong Kui Yu¹
Qi Liu¹
Xiu Feng Wei¹
Jun Zhang¹
Rui Xiang Zhang¹
Hai Bo Sun¹
Zong Fei Wang¹
Wen Qun Xing¹
Yin Li^{1,2}

¹Department of Thoracic Surgery, The Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou, Henan, People's Republic of China; ²Department of Thoracic Surgery, The Cancer Hospital Chinese Academy of Medical Sciences, Beijing, People's Republic of China

Objective: Long noncoding RNA small nucleolar RNA host gene 1 (SNHG1) has been reported to be aberrantly expressed and plays an important role in human cancers, including esophageal squamous cell cancer. However, the regulatory mechanism underlying SNHG1 in the progression of esophageal squamous cell cancer is poorly defined.

Materials and Methods: Fifty-three esophageal squamous cell cancer patients were recruited and overall survival was analyzed. EC9706 and KYSE150 cells were cultured for study in vitro. The expression levels of SNHG1, microRNA (miR)-204 and homeobox c8 (HOXC8) were detected by quantitative real-time polymerase chain reaction and Western blot. Cell cycle distribution, apoptosis, migration and invasion were determined by flow cytometry and transwell assays, respectively. The target interaction among SNHG1, miR-204 and HOXC8 was validated by luciferase reporter assay and RNA immunoprecipitation. Xenograft model was established to investigate the role of SNHG1 in vivo.

Results: High expression of SNHG1 was exhibited in esophageal squamous cell cancer and indicated poor outcomes of patients. SNHG1 silence led to cell cycle arrest at G0-G1 phase, inhibition of migration and invasion and increase of apoptosis. miR-204 was validated to sponge by SNHG1 and target HOXC8 in esophageal squamous cell cancer cells. miR-204 knockdown or HOXC8 restoration reversed the inhibitive role of SNHG1 silence in the progression of esophageal squamous cell cancer cells. Furthermore, inhibiting SNHG1 decreased xenograft tumor growth by regulating miR-204 and HOXC8.

Conclusion: SNHG1 knockdown suppresses migration and invasion but induces apoptosis of esophageal squamous cell cancer cells by increasing miR-204 and decreasing HOXC8.

Keywords: esophageal squamous cell cancer, SNHG1, miR-204, HOXC8

Introduction

Esophageal cancer with the sixth cancer deaths consists of esophageal squamous cell cancer and esophageal adenocarcinoma, and esophageal squamous cell cancer predominates worldwide.¹ Therefore, this study focuses on esophageal squamous cell cancer. Recently, great advances have been gained for the pathogenesis, diagnosis and treatment of esophageal squamous cell cancer.² However, the survival of patients remains poor.³ Hence, much hope is placed in understanding the pathogenesis and exploring a novel strategy for the treatment of esophageal squamous cell cancer.

Noncoding RNAs, including long noncoding RNAs (lncRNAs) with more than 200 nucleotides and microRNAs (miRNAs), have been reported to be aberrantly expressed and associated with cancer progression in esophageal squamous cell

Correspondence: Yin Li
Department of Thoracic Surgery, The Affiliated Cancer Hospital of Zhengzhou University, No. 127 Dongming Road, Zhengzhou, Henan 450000, People's Republic of China
Tel +86-0371-65587225
Email lihaomiao1989@163.com

cancer.⁴ lncRNAs are suggested to be involved in the development and therapeutics of esophageal squamous cell cancer.⁵ Moreover, lncRNAs could act as oncogenes or tumor suppressors in esophageal squamous cell cancer through regulating cell processes, such as proliferation, migration, invasion and apoptosis by functioning as competing endogenous RNAs (ceRNAs). For example, Sun et al⁶ reveal that LINC00657 promotes cell proliferation, migration and radioresistance in esophageal squamous cell cancer by regulating miR-615-3p and JunB. Chu et al⁷ report that lncRNA motor neuron and pancreas homeobox 1-antisense RNA1 (MNX1-AS1) regulates cell proliferation, migration, invasion, cell cycle and apoptosis by miR-34a/Sirtuin 1 (SIRT1) axis in esophageal squamous cell cancer. Furthermore, phosphoglucomutase 5 antisense RNA 1 (PGM5-AS1) as a lncRNA suppresses cell proliferation, migration and invasion by regulating miR-466/phosphatase and tensin homolog deleted on chromosome 10 (PTEN) axis in esophageal squamous cell cancer.⁸ Previous study demonstrates that lncRNA small nucleolar RNA host gene 1 (SNHG1) is highly expressed and associated with poor outcomes of patients in multiple cancers.⁹ What's more, accruing evidences suggest SNHG1 as oncogenic lncRNA to promote cell proliferation, migration and invasion in gastric cancer and pancreatic cancer.^{10,11} More importantly, recent works indicate that abnormally expressed SNHG1 is involved in the regulation of esophageal squamous cell cancer progression.^{12,13} However, the mechanism underlying SNHG1 participating in esophageal squamous cell cancer development remains largely unclear.

Intriguingly, starBase (<http://starbase.sysu.edu.cn/>) predicts that SNHG1 and homeobox c8 (HOXC8) have and share the potential complementary sequences of miR-204, which stimulates us to assume the ceRNA network of SNHG1/miR-204/HOXC8. In the present study, we measured the expression of SNHG1 in esophageal squamous cell cancer tissues and cells and investigated the effect of SNHG1 on progression of esophageal squamous cell cancer by detecting migration, invasion, cell cycle distribution and apoptosis. Moreover, we explored the regulatory network of SNHG1/miR-204/HOXC8.

Materials and Methods

The Cancer Genome Atlas (TCGA) Assay

TCGA assay was conducted via the starBase tool. The expression levels of SNHG1, miR-204 and HOXC8 in

esophageal cancer were analyzed via TCGA. The correlation among SNHG1, miR-204 and HOXC8 in esophageal cancer was also analyzed via TCGA.

Patient Tissues and Cell Culture

We recruited 53 patients with esophageal squamous cell cancer from the Tumor Hospital Affiliated to Zhengzhou University and all patients have signed the informed consent. The esophageal squamous cell cancer tissues and corresponding adjacent normal samples were collected during the surgery and then stored at -80°C . This research was approved by the Ethics Committee of the Tumor Hospital Affiliated to Zhengzhou University.

The human esophageal squamous cell cancer cell lines (EC9706, KYSE450, KYSE150 and Eca109) and normal esophageal epithelium cell Het-1A were purchased from BeNa Culture Collection (Beijing, China) and verified by the company. All cells were cultured in DMEM (Sigma, St. Louis, MO, USA) with 10% fetal bovine serum (FBS) and antibiotics at 37°C with 5% CO_2 .

Cell Transfection

The overexpression vectors of SNHG1 and HOXC8 were generated by inserting their full-length sequences into pcDNA3.1 (Thermo Fisher Scientific, Wilmington, DE, USA), with pcDNA3.1 empty vector (pcDNA) as a corresponding control. siRNA against SNHG1 (si-SNHG1-1, 5'-CUUAAAGUGUUAGCAGACATT-3'; si-SNHG1-2, 5'-AUUCCAUUUUUUUAUACACCUU-3'; si-SNHG1-3, 5'-UGUAUCUAAAAACAAAAGGG-3'), si-HOXC8 (5'-AGGAUUAAGAGAAACUCCUU-3'), siRNA negative control (si-NC) (5'-UUCUCCGAACGUGUCACGUTT-3'), miR-204 mimic (miR-204) (5'-UUCCCUUUGUCAUCUAUGCCU-3'), miRNA negative control (miR-NC) (5'-CGAUCGCAUCAGCAUCGAUUGC-3'), miR-204 inhibitor (anti-miR-204) (5'-AGGCAUAGGAUGACAAAGGAA-3') and inhibitor negative control (anti-miR-NC) (5'-CUAACGCAUGCACAGUCGUACG-3') were generated by GenePharm (Shanghai, China). The transfection of oligonucleotides with 20 nM concentration in EC9706 and KYSE150 cells was received by using Lipofectamine 2000 (Thermo Fisher Scientific). Blank is the non-transfected group. After 24 h, transfected EC9706 and KYSE150 cells were collected and used for further experiments.

Quantitative Real-Time Polymerase Chain Reaction (qRT-PCR)

After the extraction using Trizol reagent (Thermo Fisher Scientific), 1 µg RNA was reversely transcribed to cDNA by using the special RT-PCR Kit (Thermo Fisher Scientific) following the manufacturer's instructions. The qRT-PCR was performed on ABI 7500 Real-time PCR system (Bio-Rad, Hercules, CA, USA) using the diluted cDNA along with SYBR Green mix (Thermo Fisher Scientific) and primers generated by Sangon Biotech (Shanghai, China): SNHG1 (Forward, 5'-TAACCTGCTGGCTCAAAGGG-3'; Reverse, 5'-CAGCCTGGAGTGAACACAGA-3'); HOXC8 (Forward, 5'-CACGTCCAAGACTTCTTCCACACGGC-3'; Reverse, 5'-CACTTCATCCTTCGATTCTGGAACC-3'); miR-204 (Forward, 5'-CTGTCACTCGAGCTGCTGGAATG-3'; Reverse, 5'-ACCGTGTCGTGGAGTCGGCAATT-3'). GAPDH (Forward, 5'-CATGAGAAGTATGACAACAGCCT-3'; Reverse, 5'-AGTCCTTCCACGATAACCAAAGT-3') and U6 (Forward, 5'-CTCGCTTCGGCAGCAC-3'; Reverse, 5'-AACGCTTCACGAATTTGCGT-3') were regarded as internal controls, respectively. The relative expression levels of SNHG1, miR-204 and HOXC8 were analyzed by $2^{-\Delta\Delta C_t}$ method.¹⁴

Flow Cytometry

Flow cytometry was used for analysis of cell cycle distribution and apoptosis. For measurement of cell cycle distribution, EC9706 and KYSE150 cells were cultured for 48 h and then fixed with 75% ethanol (Sigma), followed by incubated with RNase A and PI solution for 30 min at 37°C. For apoptosis analysis, after culture for 48 h, the cells were resuspended in binding buffer and then stained with Annexin V-FITC and PI (Beyotime, Shanghai, China) in the dark. The distribution of cells at different phases and apoptosis were analyzed by a flow cytometer. The apoptotic rate of cells was expressed as the percentage of cells stained with Annexin V-FITC positive and PI negative or positive.

Transwell Assay

The motility of cells was investigated by migration and invasion using transwell assay. For migration assay, EC9706 and KYSE150 cells (1×10^4 /well) were suspended in DMEM without serum and added to the upper chambers of the transwell. For invasion assay, 1×10^4 cells were seeded into Matrigel-coated chambers. The lower chambers were

added with 500 µL DMEM with 10% FBS. After the incubation for 24 h, the cells on lower chambers were stained with 0.1% crystal violet. The number of migrated or invasive cells was counted under a 200× magnification microscope with three random fields.

Western Blot

Equal amounts (20 µg) of protein lysates from cells or tumors were subjected to SDS-PAGE and electro-transfer with PVDF membranes (Millipore, Billerica, MA, USA). The membranes were blocked with 5% non-fat milk for 2 h, incubated with specific primary antibodies overnight and secondary antibody (ab97051, 1:20,000 dilution) for 2 h. The antibody against HOXC8 (ab86236, 1:1000 dilution) was purchased from Abcam (Cambridge, MA, USA), with β-actin (ab227387, 1:10,000 dilution) as a loading control. After the incubation of enhanced chemiluminescence solution (Beyotime), the targeted protein signaling was normalized to β-actin in each sample.

Luciferase Reporter Assay and RNA Immunoprecipitation (RIP)

The sequences of SNHG1 or HOXC8 3'UTR containing miR-204 binding sites were inserted into a pmirGLO luciferase reporter vector (Promega, Madison, WI, USA) to generate the wild-type luciferase reporter vectors (SNHG1-WT and HOXC8-WT). Meanwhile, the corresponding mutants (SNHG1-MUT and HOXC8-MUT) were generated via mutating the binding sites. Co-transfection of the constructed luciferase reporter vectors together with miR-204 or miR-NC was performed into EC9706 and KYSE150 cells. The luciferase reporter system (Promega) was used for luciferase activity analysis at 24 h post-transfection.

EC9706 and KYSE150 cells transfected with miR-204 or miR-NC were lysed for RIP assay with a Magna RNA immunoprecipitation kit (Millipore) following the manufacturer's protocols. The enrichment levels of SNHG1 and HOXC8 in Ago2 or IgG RIP complex were determined by qRT-PCR.

Murine Xenograft Model

The animal experiments were performed in accordance with the guide for care and use of laboratory animals and this study was approved by the Ethics Committee of the Tumor Hospital Affiliated to Zhengzhou University. The BALB/c nude mice (4-week-old, male) purchased from Shanghai Animal Laboratory Center (Shanghai, China) were acclimatized for

1 week and then randomly divided into three groups ($n=3$) for xenograft model. EC9706 cells (2×10^6 cells) stably transfected with recombinant lentivirus expressing shRNA for SNHG1 (sh-SNHG1), control (sh-NC) or empty lentivirus were used for subcutaneously infection of nude mice. The tumor volume was monitored every week and calculated with a formula: volume (mm^3) = length $\times$ width² $\times$ 0.5. At 5 weeks after the injection, mice were killed and tumors were collected for weight and molecular analyses.

Statistical Analysis

The data from three independent experiments were processed by GraphPad Prism 7.0 and presented as mean $\pm$ standard deviation. Overall survival was generated by Kaplan–Meier method and analyzed by the Log rank test. The comparison between two or more groups was conducted by student's *t*-test or ANOVA with Tukey's post hoc test. The difference was statistically significant when $P < 0.05$ (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

Results

The Expression of SNHG1 Is Enhanced in Esophageal Squamous Cell Cancer

The data of TCGA showed a high expression of SNHG1 in esophageal cancer (Supplementary Figure 1A). To detect the abundance of SNHG1 in esophageal squamous cell cancer, 53 patients were recruited. As shown in Figure 1A, the expression of SNHG1 in esophageal squamous cell cancer tissues ($n=53$) was increased fourfold compared with that in corresponding normal samples. Meanwhile, when compared to Het-1A cells, esophageal squamous cell cancer cells, especially EC9706 and KYSE150 cells, exhibited higher

level of SNHG1 (Figure 1B). Moreover, the patients were divided into high ($n=31$) or low ($n=22$) SNHG1 level group according to the mean level of SNHG1 in cancer tissues. The patients with high SNHG1 level displayed poor overall survival compared with those in low expression group ($P=0.027$) (Figure 1C).

SNHG1 Knockdown Inhibits Progression of Esophageal Squamous Cell Cancer Cells

To investigate the role of SNHG1 in esophageal squamous cell cancer development, EC9706 and KYSE150 cells with relatively higher level of SNHG1 (5.6-fold and 3.25-fold, respectively) were chosen for in vitro study. Overexpression of SNHG1 had an oncogenic role in esophageal squamous cell cancer by promoting migration and invasion and inhibiting apoptosis (Supplementary Figure 2). Furthermore, we constructed three siRNAs for SNHG1, and the siRNA with the highest efficacy was chosen for further experiments (Supplementary Figure 3). As shown in Figure 2A and B, the abundance of SNHG1 in EC9706 and KYSE150 cells was effectively decreased 60–65% by transfection of si-SNHG1 in comparison to that in the si-NC group and Blank group. Moreover, knockdown of SNHG1 induced cell cycle arrest at G0-G1 phase, revealed by increasing percentage of cells at G0-G1 phase and reducing distribution at S phase (Figure 2C and D). In addition, the abilities of migration and invasion in EC9706 and KYSE150 cells were significantly decreased by SNHG1 silence (Figure 2E–H). Besides, interference of SNHG1 led to great apoptosis production in EC9706 and KYSE150 cells (Figure 2I and J).

Figure 1 The expression of SNHG1 in esophageal squamous cell cancer. (A) qRT-PCR assay was performed to detect SNHG1 expression in esophageal squamous cell cancer tissues compared with normal samples ($n=53$). (B) qRT-PCR assay was carried out to measure SNHG1 level in esophageal squamous cell cancer cells compared with normal esophageal epithelium cell. (C) The overall survival of patients was assessed according to the mean level of SNHG1 in esophageal squamous cell cancer tissues. * $P < 0.05$, ** $P < 0.001$.

Figure 2 The effect of SNHG1 on cell cycle distribution, migration, invasion and apoptosis in esophageal squamous cell cancer cells. (A and B) The level of SNHG1 in EC9706 and KYSE150 cells was detected by qRT-PCR after transfection of si-SNHG1 or si-NC. Cell cycle distribution (C and D), migration (E and F), invasion (G and H) and apoptosis (I and J) of EC9706 and KYSE150 cells transfected with si-SNHG1 or si-NC were determined by flow cytometry and trans-well assays. Blank was non-transfected group. ** $P < 0.01$, *** $P < 0.001$.

SNHG1 Is a Decoy of miR-204

As shown in Figure 3A, starBase database predicted the complementary sequences of SNHG1 and miR-204 at position: chr11: 62620353-62620374. The wild-type and mutant luciferase reporter vectors (SNHG1-WT and SNHG1-MUT) were constructed and transfected into EC9706 and KYSE150 cells. As revealed in Figure 3B and C, overexpression of miR-204 led to more than 60% reduction of luciferase activity in the SNHG1-WT group, while it did not affect the activity in SNHG1-MUT group. Moreover, the addition of miR-204 led to a markedly increased level of SNHG1 enriched by Ago2 RIP in EC9706 and KYSE150 cells (Figure 3D and E). The data of TCGA displayed that miR-204 expression was decreased in esophageal cancer (Supplementary Figure 1B). Additionally, the expression of miR-204 was notably reduced in esophageal squamous cell cancer cells compared with that in Het-1A cells (Figure 3F). Besides, miR-204

level in EC9706 and KYSE150 cells was evidently decreased by overexpressing SNHG1 and increased via silencing SNHG1 (Figure 3G and H). Furthermore, SNHG1 expression was also negatively regulated by miR-204 (Supplementary Figure 4).

Knockdown of miR-204 Reverses the Effect of SNHG1 Silence on Progression of Esophageal Squamous Cell Cancer Cells

In order to explore whether miR-204 was associated with SNHG1-mediated progression of esophageal squamous cell cancer, EC9706 and KYSE150 cells were transfected with si-NC, si-SNHG1, si-SNHG1 and anti-miR-NC or anti-miR-204. After the transfection, the abundance of miR-204 in EC9706 and KYSE150 cells increased by SNHG1 knockdown was obviously decreased by the transfection of anti-miR-204 (Figure 4A and B). Furthermore,

Figure 3 The target association between SNHG1 and miR-204. **(A)** starBase database predicted the binding sites of SNHG1 and miR-204. **(B and C)** Luciferase activity was determined in EC9706 and KYSE150 cells co-transfected with SNHG1-WT or SNHG1-MUT and miR-204 or miR-NC. **(D and E)** qRT-PCR assay was performed to detect SNHG1 abundance in EC9706 and KYSE150 cells transfected with miR-204 or miR-NC after RIP assay. **(F)** The expression of miR-204 was detected in esophageal squamous cell cancer cells compared with normal esophageal epithelium cell. **(G and H)** The level of miR-204 was measured in EC9706 and KYSE150 cells transfected with pcDNA, SNHG1, si-NC or si-SNHG1. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

inhibition of miR-204 significantly attenuated cell cycle arrest at G0-G1 phase induced by SNHG1 silence (Figure 4C and D). In addition, the suppressive effect of SNHG1 knockdown on migration and invasion of EC9706 and KYSE150 cells was weakened via miR-204 deficiency (Figure 4E–H). What is more, the apoptosis production induced by silencing SNHG1 was specially mitigated by down-regulation of miR-204 in the two cells (Figure 4I and J). Besides, miR-204 overexpression also reversed the promoting role of SNHG1 in esophageal squamous cell cancer development in vitro (Supplementary Figure 2).

HOXC8 Is a Target of miR-204

The target of miR-204 was searched by starBase database, which provided the putative binding sites of miR-204 and HOXC8 at position: chr12: 54406036-54406042 (Figure 5A). To validate this association, luciferase reporter assay and RIP assay were performed in EC9706 and KYSE150 cells. As displayed in Figure 5B and C, the luciferase activity was greatly decreased via transfection of miR-204 in the HOXC8-WT group, whereas it was not changed when the seed sites were mutated in HOXC8-MUT group. Meanwhile, the level of HOXC8 enriched by Ago2 RIP

Figure 4 The rescue effect of miR-204 on SNHG1-mediated progression of esophageal squamous cell cancer cells. (A and B) The level of miR-204 in EC9706 and KYSE150 cells was detected by qRT-PCR after transfection of si-NC, si-SNHG1, si-SNHG1 and anti-miR-NC or anti-miR-204. Cell cycle distribution (C and D), migration (E and F), invasion (G and H) and apoptosis (I and J) of EC9706 and KYSE150 cells transfected with si-NC, si-SNHG1, si-SNHG1 and anti-miR-NC or anti-miR-204 were analyzed by flow cytometry and trans-well assays. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

but not IgG RIP was significantly increased by miR-204 overexpression (Figure 5D and E). TCGA database showed that HOXC8 level was elevated in esophageal cancer (Supplementary Figure 1C). Furthermore, the mRNA level of HOXC8 in esophageal squamous cell cancer cells was notably higher than that in Het-1A cells (Figure 5F). In addition, the abundance of HOXC8 protein in EC9706 and KYSE150 cells was conspicuously reduced via miR-204 addition and elevated by miR-204 exhaustion (Figure 5G and H). Besides, there was a positive correlation between SNHG1 and HOXC8, and a negative correlation between miR-204 and SNHG1 or HOXC8 in esophageal squamous cell cancer, although the correlation between miR-204 and SNHG1 or HOXC8 was insignificant in esophageal cancer by TCGA (Supplementary Figure 1D–I). We hypothesized that the inconsistent results might be explained by demographic difference or cancer subtype difference. Moreover, HOXC8 expression was promoted by SNHG1 overexpression and weakened by miR-204 (Supplementary Figure 2).

HOXC8 Restoration Weakens the Effect of SNHG1 Knockdown on Progression of Esophageal Squamous Cell Cancer Cells

To explore whether SNHG1-regulated progression of esophageal squamous cell cancer was mediated by HOXC8, EC9706

and KYSE150 cells were transfected with si-NC, si-SNHG1, si-SNHG1 and pcDNA or HOXC8. As shown in Figure 6A and B, the protein level of HOXC8 in EC9706 and KYSE150 cells was significantly decreased by silencing SNHG1, which was restored by the introduction of HOXC8 overexpression vector. Moreover, the arrest of cell cycle in EC9706 and KYSE150 cells caused by SNHG1 interference was abrogated by the up-regulation of HOXC8 (Figure 6C and D). Meanwhile, the inhibition of migration and invasion mediated by SNHG1 knockdown was abated by the restoration of HOXC8 in EC9706 and KYSE150 cells (Figure 6E–H). Additionally, the introduction of HOXC8 significantly relieved the apoptosis of EC9706 and KYSE150 cells induced by silencing SNHG1 (Figure 6I and J). Besides, knockdown of HOXC8 also weakened the promoting role of SNHG1 in esophageal squamous cell cancer development in vitro (Supplementary Figure 2).

Silencing SNHG1 Decreases Xenograft Tumor Growth

To further evaluate the role of SNHG1 in esophageal squamous cell cancer in vivo, EC9706 cells stably infected with sh-SNHG1 or sh-NC were injected into nude mice. After 5 weeks, the xenograft tumor volume and tumor weight were significantly reduced in the sh-SNHG1 group compared with those in the sh-NC group (Figure 7A and B). Moreover, the

Figure 5 The target association between miR-204 and HOXC8. (A) The seed sites of miR-204 and HOXC8 were predicted by starBase database. Luciferase reporter assay (B and C) and RIP assay (D and E) were conducted to confirm the bind of miR-204 and HOXC8 in EC9706 and KYSE150 cells. (F) The expression of HOXC8 mRNA was measured in esophageal squamous cell cancer cells compared with normal esophageal epithelium cell. (G and H) The protein level of HOXC8 was examined in EC9706 and KYSE150 cells transfected with miR-NC, miR-204, anti-miR-NC or anti-miR-204. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

analyses of qRT-PCR and Western blot revealed that the expression levels of SNHG1 and HOXC8 protein were remarkably decreased while miR-204 abundance was evidently enhanced in tumor tissues of the sh-SNHG1 group compared with those in the sh-NC group (Figure 7C–E).

Discussion

Esophageal squamous cell cancer is a common cancer wreaking havoc on the health of people worldwide.¹⁵ LncRNAs with ceRNA networks exhibit important roles in regulating esophageal squamous cell cancer malignancy.¹⁶ In the current study, we investigated the function of SNHG1 on

esophageal squamous cell cancer cell processes and elucidated a novel ceRNA mechanism associated with miR-204/HOXC8 axis.

The emerging effort summarized the role of SNHG1 in human cancers, including esophageal squamous cell cancer.¹⁷ By detecting the level of SNHG1 in 53 esophageal squamous cell cancer tissues and cell lines, we found that this lncRNA expression was increased compared with matched controls. This indicated the potential oncogenic role of SNHG1 in esophageal squamous cell cancer, which was confirmed by gain-of-function experiments. Furthermore, SNHG1 inhibition induced by si-SNHG1 led to the impaired abilities of

migration and invasion and increased apoptosis by arresting cell cycle at G0-G1 phase. These results indicated the potential therapeutic effect of target SNHG1 inhibition on esophageal squamous cell cancer, which is also in agreement with the previous study.¹³

Former findings suggested that SNHG1 could serve as a ceRNA to be implicated in the development of human cancers.^{18–20} To figure out whether SNHG1 could play the carcinogenic role in esophageal squamous cell cancer, we searched its targets and validated miR-204 as a target of SNHG1. Intriguingly, there was a negative mutual association between SNHG1 and miR-204. This is also in agreement with the dual-regulation of GAS5/miR-212 and PVT1/miR-143.^{21,22} We hypothesized that it might be induced by competitively binding to the miRNA response elements.²³ As described previously, miR-204 functioned as a tumor suppressor in multiple cancers, including papillary thyroid carcinoma, hepatocellular cancer and lung adenocarcinoma.^{24–26} Meanwhile, miR-204 played the suppressive role in esophageal cancer by decreasing proliferation, invasion and epithelial-mesenchymal transition.^{27,28} In our study, we also found the tumor suppressive role of miR-204, revealed by which its knockdown abated silencing SNHG1-mediated inhibition of esophageal squamous cell cancer progression. This also uncovered that SNHG1 mediated esophageal squamous cell cancer development by sponging miR-204.

HOXC8 has been regarded as an oncogene and predicts poor prognosis in multiple cancers, including hepatocellular carcinoma, ovarian cancer, non-small cell lung cancer and cervical cancer.^{29–32} Moreover, previous studies showed that HOXC8 was highly expressed and associated with poor prognosis and tumor development in esophageal squamous cell cancer.^{33,34} Using starBase database, we found that HOXC8 and SNHG1 have the similar binding site of miR-204. Hence, we hypothesized that HOXC8 might be required for the ceRNA network addressed by SNHG1. Here we first using luciferase reporter assay and RIP confirmed HOXC8 as a target of miR-204 and found that restoration or knockdown of HOXC8 abolished the effect of SNHG1 silence or overexpression on cancer progression, indicating that SNHG1 regulated esophageal squamous cell cancer development by increasing HOXC8 via competitively sponging miR-204. Tumor xenograft model is responsible for studying the tumorigenesis process in esophageal squamous cell cancer.³⁵ Hence, we also established the murine xenograft model using esophageal squamous cell cancer cells and found the anti-cancer role

of SNHG1 knockdown in this cancer in vivo by regulating miR-204/HOXC8 axis.

Conclusion

Our research revealed the therapeutic effect of SNHG1 silence on esophageal squamous cell cancer development by decreasing cell migration and invasion and promoting apoptosis, possibly by regulating HOXC8 via competitively binding miR-204. This study indicates SNHG1 as a potential target for the treatment of esophageal squamous cell cancer and provides a new regulatory mechanism for SNHG1.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Alsop BR, Sharma P. Esophageal cancer. *Gastroenterol Clin North Am*. 2016;45(3):399–412. doi:10.1016/j.gtc.2016.04.001
2. Ohashi S, Miyamoto S, Kikuchi O, Goto T, Amanuma Y, Muto M. Recent advances from basic and clinical studies of esophageal squamous cell carcinoma. *Gastroenterology*. 2015;149(7):1700–1715. doi:10.1053/j.gastro.2015.08.054
3. Pasquali S, Yim G, Vohra RS, et al. Survival after neoadjuvant and adjuvant treatments compared to surgery alone for resectable esophageal carcinoma: a network meta-analysis. *Ann Surg*. 2017;265(3):481–491. doi:10.1097/SLA.0000000000001905
4. Sugihara H, Ishimoto T, Miyake K, et al. Noncoding RNA expression aberration is associated with cancer progression and is a potential biomarker in esophageal squamous cell carcinoma. *Int J Mol Sci*. 2015;16(11):27824–27834. doi:10.3390/ijms161126060
5. Shen WJ, Zhang F, Zhao X, Xu J. LncRNAs and esophageal squamous cell carcinoma - implications for pathogenesis and drug development. *J Cancer*. 2016;7(10):1258–1264. doi:10.7150/jca.14869
6. Sun Y, Wang J, Pan S, et al. LINC00657 played oncogenic roles in esophageal squamous cell carcinoma by targeting miR-615-3p and JunB. *Biomed Pharmacother*. 2018;108:316–324. doi:10.1016/j.biopha.2018.09.003
7. Chu J, Li H, Xing Y, et al. LncRNA MNX1-AS1 promotes progression of esophageal squamous cell carcinoma by regulating miR-34a/SIRT1 axis. *Biomed Pharmacother*. 2019;116:109029. doi:10.1016/j.biopha.2019.109029
8. Zhuhua Z, Weiwei W, Lihua N, Jianying Z, Jiang G. p53-induced long non-coding RNA PGM5-AS1 inhibits the progression of esophageal squamous cell carcinoma through regulating miR-466/PTEN axis. *IUBMB Life*. 2019;71:1492–1502. doi:10.1002/iub.2069
9. Yu Y, Yang J, Yang S, et al. Expression level and clinical significance of SNHG1 in human cancers: a meta-analysis. *Oncotargets Ther*. 2019;12:3119–3127. doi:10.2147/OTT.S184803
10. Guo W, Huang J, Lei P, Guo L, Li X. LncRNA SNHG1 promoted HGC-27 cell growth and migration via the miR-140/ADAM10 axis. *Int J Biol Macromol*. 2019;122:817–823. doi:10.1016/j.ijbiomac.2018.10.214
11. Cui L, Dong Y, Wang X, et al. Downregulation of long noncoding RNA SNHG1 inhibits cell proliferation, metastasis, and invasion by suppressing the Notch-1 signaling pathway in pancreatic cancer. *J Cell Biochem*. 2019;120(4):6106–6112. doi:10.1002/jcb.27897

12. Yan Y, Fan Q, Wang L, Zhou Y, Li J, Zhou K. LncRNA Snhg1, a non-degradable sponge for miR-338, promotes expression of proto-oncogene CST3 in primary esophageal cancer cells. *Oncotarget*. 2017;8(22):35750–35760. doi:10.18632/oncotarget.16189
13. Zhang Y, Jin X, Wang Z, Zhang X, Liu S, Liu G. Downregulation of SNHG1 suppresses cell proliferation and invasion by regulating Notch signaling pathway in esophageal squamous cell cancer. *Cancer Biomark*. 2017;21(1):89–96. doi:10.3233/CBM-170286
14. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods*. 2001;25(4):402–408. doi:10.1006/meth.2001.1262
15. Batra R, Malhotra GK, Singh S, Are C. Managing squamous cell esophageal cancer. *Surg Clin North Am*. 2019;99(3):529–541. doi:10.1016/j.suc.2019.02.006
16. Tian W, Jiang C, Huang Z, Xu D, Zheng S. Comprehensive analysis of dysregulated lncRNAs, miRNAs and mRNAs with associated ceRNA network in esophageal squamous cell carcinoma. *Gene*. 2019;696:206–218. doi:10.1016/j.gene.2019.02.051
17. Thin KZ, Tu JC, Raveendran S. Long non-coding SNHG1 in cancer. *Clin Chim Acta*. 2019;494:38–47. doi:10.1016/j.cca.2019.03.002
18. Lan X, Liu X. LncRNA SNHG1 functions as a ceRNA to antagonize the effect of miR-145a-5p on the down-regulation of NUA1 in nasopharyngeal carcinoma cell. *J Cell Mol Med*. 2019;23(4):2351–2361. doi:10.1111/jcmm.13497
19. Xu M, Chen X, Lin K, et al. The long noncoding RNA SNHG1 regulates colorectal cancer cell growth through interactions with EZH2 and miR-154-5p. *Mol Cancer*. 2018;17(1):141. doi:10.1186/s12943-018-0894-x
20. Deng R, Zhang J, Chen J. LncRNA SNHG1 negatively regulates miRNA1013p to enhance the expression of ROCK1 and promote cell proliferation, migration and invasion in osteosarcoma. *Int J Mol Med*. 2019;43(3):1157–1166. doi:10.3892/ijmm.2018.4039
21. Zhang Z, Zhu Z, Watabe K, et al. Negative regulation of lncRNA GAS5 by miR-21. *Cell Death Differ*. 2013;20(11):1558–1568. doi:10.1038/cdd.2013.110
22. Chen J, Yu Y, Li H, et al. Long non-coding RNA PVT1 promotes tumor progression by regulating the miR-143/HK2 axis in gallbladder cancer. *Mol Cancer*. 2019;18(1):33. doi:10.1186/s12943-019-0947-9
23. Zhao Z, Sun W, Guo Z, et al. Mechanisms of lncRNA/microRNA interactions in angiogenesis. *Life Sci*. 2019;116900. doi:10.1016/j.lfs.2019.116900
24. Liu H, Li R, Guan L, Jiang T. Knockdown of lncRNA UCA1 inhibits proliferation and invasion of papillary thyroid carcinoma through regulating miR-204/IGFBP5 axis. *Onco Targets Ther*. 2018;11:7197–7204. doi:10.2147/OTT.S175467
25. Yu Y, Wang Y, Xiao X, et al. MiR-204 inhibits hepatocellular cancer drug resistance and metastasis through targeting NUA1. *Biochem Cell Biol*. 2019;97(5):563–570. doi:10.1139/bcb-2018-0354
26. Hu WB, Wang L, Huang XR, Li F. MicroRNA-204 targets SOX4 to inhibit metastasis of lung adenocarcinoma. *Eur Rev Med Pharmacol Sci*. 2019;23(4):1553–1562. doi:10.26355/eurev_201902_17114
27. Jiao C, Song Z, Chen J, et al. LncRNA-UCA1 enhances cell proliferation through functioning as a ceRNA of Sox4 in esophageal cancer. *Oncol Rep*. 2016;36(5):2960–2966. doi:10.3892/or.2016.5121
28. Sun Y, Yu X, Bai Q. miR-204 inhibits invasion and epithelial-mesenchymal transition by targeting FOXM1 in esophageal cancer. *Int J Clin Exp Pathol*. 2015;8(10):12775–12783.
29. Xu P, Zhang X, Ni W, et al. Upregulated HOXC8 expression is associated with poor prognosis and oxaliplatin resistance in hepatocellular carcinoma. *Dig Dis Sci*. 2015;60(11):3351–3363. doi:10.1007/s10620-015-3774-x
30. Lu S, Liu R, Su M, et al. Overexpression of HOXC8 is associated with poor prognosis in epithelial ovarian cancer. *Reprod Sci*. 2016;23(7):944–954. doi:10.1177/1933719115625845
31. Liu H, Zhang M, Xu S, et al. HOXC8 promotes proliferation and migration through transcriptional up-regulation of TGFbeta1 in non-small cell lung cancer. *Oncogenesis*. 2018;7(2):1. doi:10.1038/s41389-017-0016-4
32. Huang Y, Chen L, Guo A. Upregulated expression of HOXC8 is associated with poor prognosis of cervical cancer. *Oncol Lett*. 2018;15(5):7291–7296. doi:10.3892/ol.2018.8200
33. Du YB, Dong B, Shen LY, et al. The survival predictive significance of HOXC6 and HOXC8 in esophageal squamous cell carcinoma. *J Surg Res*. 2014;188(2):442–450. doi:10.1016/j.jss.2014.01.017
34. Shen LY, Zhou T, Du YB, Shi Q, Chen KN. Targeting HOX/PBX dimer formation as a potential therapeutic option in esophageal squamous cell carcinoma. *Cancer Sci*. 2019;110(5):1735–1745. doi:10.1111/cas.13993
35. Lee NP, Chan CM, Tung LN, Wang HK, Law S. Tumor xenograft animal models for esophageal squamous cell carcinoma. *J Biomed Sci*. 2018;25(1):66. doi:10.1186/s12929-018-0468-7

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>

Dovepress