Matrine Restrains Cell Growth and Metastasis by Up-Regulating LINC00472 in Bladder Carcinoma

This article was published in the following Dove Press journal:
Cancer Management and Research

Linlin Li^{1,*}
Fei Qi^{1,*}
Kaichen Wang²

¹Department of Operating Room, China-Japan Union Hospital of Jilin University, Changchun 130033, Jilin, People's Republic of China; ²Department of Urinary Surgery, China-Japan Union Hospital of Jilin University, Changchun 130033, Jilin, People's Republic of China

*These authors contributed equally to this work

Purpose: Bladder Carcinoma (BC) is a malignant carcinoma with a high incidence in masculinity. We preliminarily researched the efficacy and mechanism of matrine (MAT) in T24 and 5637 cells.

Patients and Methods: CCK-8, flow cytometry, migration and invasion means were adopted to detect cell viability, apoptosis, migratory and invasive potentials. Moreover, LINC00472 expression was changed via transfection assays and was tested by RT-qPCR. Western blot was used for investigating the levels of CyclinD1, p53, Bcl-2, Bax, pro-Caspase-3, Cleaved-Caspase-3, β -actin, programmed cell death protein 4 (PDCD4) and relate-proteins of cell pathways. Tumor volume and weight were tested via animal experiments.

Results: MAT could not affect the growth of SV-HUC-1 cell but MAT promoted tumor cell apoptosis but restrained viability, invasion and migration. Furthermore, LINC00472 was prominently low expressed in BC tissues. MAT positively regulated LINC00472 and transfection with si-00472 could partly reverse the efficacies of MAT. Moreover, MAT enhanced PDCD4 expression by up-regulating LINC00472. Besides, we discovered MAT elevated PTEN but restrained PI3K/AKT proteins. Finally, tumor volume and weight were declined by MAT in vivo via up-regulating LINC00472.

Conclusion: MAT restrained cell growth and metastasis but promoted PDCD4 expression by up-regulating LINC00472 via restraining PTEN/PI3K/AKT pathway in BC.

Keywords: bladder carcinoma, matrine, LINC00472, PDCD4

Introduction

Bladder carcinoma (BC) is a familiar urinary system carcinoma.¹ Its incidence is higher in masculinity than in femininity.² In recent days, the incidence of BC in China has enhanced year by year.³ BC is an illness that has no distinct symptoms and it is prone to recurrence.⁴ At present, the check means for BC are imaging and urine tests and treatment means are still surgery, radiotherapy and drugs.⁵ In addition, most of the drugs for treating BC lacking specificity and radiotherapy have strong side effects.⁶ Therefore, it is vital to screen safer natural drugs.

Natural products have been recognized as critical latent drugs used for a series of deadly diseases encompassing cancers for a long time. For example, previous studies have pointed out that bitter melon extract (BME) played as a tumor suppressor in breast cancer and neck squamous cell carcinoma (HNSCC), where it inhibited tumor growth by modulation of proliferation, autophagy and immunity.^{7,8} What is more, manifold natural products isolated from Chinese herbal medicine also have been widely utilized for cancer treatment due to their excellent activities, such as anti-proliferation, pro-apoptosis and anti-metastasis, as well as regulate autophagy and immunity.⁹ Matrine

Correspondence: Kaichen Wang
Department of Urinary Surgery, China-Japan Union Hospital of Jilin University, No. 126 Xiantai Street, Changchun 130033, Jilin, People's Republic of China
Tel +86-431-84995999
Fax +86-431-89876666
Email wangkc0119@sina.com

(MAT) is a tetracyclic quinolizidine compound extracted from *sophora flavescens*.¹⁰ Its molecular formula is $C_{12}H_{24}N_2O$ and its molecular weight is 248.37 g/M.¹¹ It has been put forward that MAT has anti-carcinoma and anti-inflammatory efficacies in many diseases.¹² With the deepening of research, it has been put forward that MAT can conspicuously restrain the activities of carcinoma cells. Researches have pointed out MAT can conspicuously restrain prostate carcinoma.¹³ In addition, a research has pointed out that MAT restrains the activity of BC cells.¹⁴

Many types of researches have pointed out that the development of malignant carcinomas usually results from abnormal gene expression.¹⁵ It is widely known that aberrant non-coding RNAs encompassing circular (circ) RNAs, long non-coding RNAs (lncRNAs) as well as microRNAs (miRNAs) may have critical latent functions in regulation of gene expression as well as development of cancers.^{16,17} Here, we are interested in the regulation of lncRNAs in BC progress. lncRNAs are a group of RNA more than 200 nt in length.¹⁸ More and more scientific evidence has proven that lncRNAs are maladjusted in BC, gastric carcinoma and so on,¹⁹ and it is involved in the occurrence and progress of carcinomas by regulating cell growth and metastasis.²⁰ In addition, lncRNAs can regulate several signaling pathways and play a promoting or restraining efficacy in carcinomas.²¹ LINC00472, as a lncRNA, had been put forward to have low expression in colorectal carcinomas²² and played a restraining efficacy in breast carcinomas.²³ However, there was no research about LINC00472 in BC until now.

Currently, we first studied the expression of LINC00472 in BC tissues. T24 and 5637 cells were utilized for the experimental investigation of the efficacy of MAT in vitro. We investigated the efficacies of MAT and LINC00472 on the viability, apoptosis, migration and invasion and the relationship between MAT and programmed cell death protein 4 (PDCD4) was explored. Moreover, we explored the possible mechanism of MAT in BC cells.

Materials and Methods

Clinical Specimens

Selected BC tissues and non-tumor tissues ($n = 20$) were acquired from the patients who had not been treated by any therapies before the cut-off surgery from China-Japan Union Hospital of Jilin University (Changchun, China). Every patient agreed with the performance of this current research and writing an informed consent. This survey was supported by the Medical Ethics Committee of this Hospital.

Cell Culture and Treatment

BC cells-T24/5637 and the normal bladder cell-SV-HUC-1 provided by American Type Culture Collection (Manassas, VA, USA) were separately hatched at Dulbecco's modified eagle medium (DMEM, Gibco, Grand Island, NY, US) or Ham's F-12 medium at 37°C with 5% CO₂. The streptomycin/penicillin of 100 units/mL (Invitrogen, Carlsbad, CA, USA) and 10% fetal bovine serum (FBS, Gibco) were also added in. MAT (Figure 1, Sigma, St. Louis, MO, US) was diluted with ddH₂O to IC₅₀²⁴ and was utilized for cells for 24 h through a filter.

CCK-8

After dosing dispose, the CCK-8 solution (Sigma) was appended to cells in line with the instructions. Cells were hatched in dark 1 h, the optical density was quantitated at 450 nm by a microplate reader (Bio-Rad, Sunnyvale, CA, US).

Flow Cytometry

Transfected cells were harvested and washed with cold 1 × phosphate buffered saline (PBS) after centrifugation. Next, those cells were suspended, followed by 5 µL PI and 5 µL AnnexinV-FITC staining solution (Bioswamp). After that, the resulting cultures were held at 25°C for 10 min in the dark. Results were measured via flow cytometry via FACS can (Beckman Coulter, Fullerton, CA, USA).

Migration and Invasion Assay

A 24-well transwell membrane (8 µm)-contained chamber (Millipore, Bedford, Massachusetts, USA) was utilized in the migration and invasion experiments. The difference between migration and invasion experiments was the membrane coated with Matrigel (Franklin Lakes, NJ, USA). The transfected BC cells were seeded in the top chamber filling in 200 µL serum-free medium, while the lower chamber was filled with complete medium. The intruding chamber was treated for 48 h to specification. Non-invasive cells were scrubbed. Invading cells were fixed by Methanol, stained by crystal violet and counted by microscope (Olympus, Japan). Randomly selected 5 groups of cells were used for counting.

Transfection

siRNA targeting LINC00472 (si-00472) and si-negative controls (NC) synthesized by GenePharma Co. (Shanghai, China) were transfected into cells using Lipofectamine 2000

Figure 1 Matrine (MAT) had no effects on normal SV-HUC-1 cell growth. **(A)** MAT had no effects on viability, and **(B, C)** generation of p53 and CyclinD1, also **(D)** apoptotic cells as well as **(E, F)** expressions of Bcl-2, Bax and Cleaved-Caspase-3; **(G)** MAT is a tetracyclic quinolizidinine compound. Its molecular formula is $C_{12}H_{24}N_2O$ and molecular weight is 248.37 g/mol. Pure MAT is white powder, low content is brown and yellow liquor.

Abbreviation: Ns, no significance.

(Invitrogen). Stably transfected cells were selected by G418 medium (Sigma).

RT-qPCR

Trizol (Invitrogen) was used to extract RNA from BC tissues, non-tumor tissues, or cells. LINC00472 expression was detected by One-Step SYBR[®] PrimeScript[®] PLUS RT-RNA PCR Kit (TaKaRa Biotechnology, Dalian, China). β -Actin was as an internal parameter. The data were calculated as $\log 10 (2^{-\Delta\Delta Ct})$. Primer sequences used in our experiments were displayed as follows:

LINC00472: Forward: 5'- CCCAGAGACAAGAGGA GCAA-3', Reverse: 5'- AGCGTCAAGAGTGGAGGT TT-3'; β -actin: Forward: 5'-AGCGAGCATCCCCAAAGT-3'; Reverse: 5'-GGGCACGA AGGCTCATCATT -3'.

Western Blot

By homogenizing the cells in RIPA buffer (Solarbio), proteins were isolated (Solarbio). Protein assay kit (Bioswamp) was used to quantifying proteins before they were separated by SDS-PAGE and transferred to a PVDF membrane (Millipore). Primary antibodies diluted with 5% blocking buffer (1:1000) were cultured the membrane overnight (4°C). Then, it was washed with PBS and hatched with a secondary antibody goat anti-rabbit IgG (ab6721, Abcam, 1:5000) for 1 h,

25°C. Signals were developed and examined by the Image Lab[™] (Bio-Rad). The β -actin was as internal parameters. Primary antibodies (Abcam) contained antibody CyclinD1 (ab40754), p53 (ab131442), Bcl-2 (ab32124), Bax (ab137321), pro-Caspase-3 (ab32150), Cleaved-Caspase-3 (ab49822), β -actin (ab8227), PDCD4 (ab51495), phosphate and tension homology deleted on chromosome ten (PTEN) (ab31392), phosphatidylinositol 3-kinases (PI3K)(ab86714), phosphorylated (p)-PI3K (ab182651), protein kinase B (AKT) (ab64148), p-AKT (ab8933).

Animal Experiment

We performed the in vivo assay by utilizing male athymic BALB/c nude mice (4–6 weeks of age). In detail, mice were divided into 4 groups: control, MAT, MAT + si-NC and MAT + si-00472, of which 8 were in each group. In the control group, the animals were injected intraperitoneally with 100 μ L of normal saline the next day after 2×10^6 T24 cells were subcutaneously injected into the flank of nude mice. In the MAT group, 50 mg/kg/day MAT (dissolved in physiological saline) instead of the saline for injection. For MAT + si-NC group: the stable si-NC-transfected T24 cells (2×10^6) were injected into mice, and the next day after inoculation, the animals started an intraperitoneal injection of 50 mg/kg/day MAT (soluble in physiological saline). For MAT + si-00472

group: the stably si-00472-transfected T24 cells (2×10^6) were injected before starting the intraperitoneal injection of 50 mg/kg/day MAT (dissolved in physiological saline). The tumor volume and weight were measured once a week, continuously for 4 weeks.

Statistical Analysis

Data were shown as mean $\pm$ Standard Deviation (SD). Data analysis was quantified via SPSS 18.0. The *P*-values were calculated by Student's *t*-test or one-way analysis of variance (ANOVA). *P* < 0.05 was conspicuously significant.

Results

MAT Had No Effects on Growth of Normal SV-HUC-1 Cells

We first detected the roles of MAT in normal human uroepithelial cell line-SV-HUC-1. Outcomes were shown in Figure 1, indicating that there were no significant differences

in cell viability (Figure 1A) and generation of p53 and CyclinD1 (Figure 1B and C). Also, similar outcomes were observed in apoptotic cells (Figure 1D) as well as expressions of Bcl-2, Bax and Cleaved-caspase-3 (Figure 1E and F) between the control group and MAT treatment group. In addition, the chemical structure of MAT was displayed in Figure 1G. Those data indicated that MAT had no adverse impacts on the normal uroepithelial cell line.

MAT Restrained Cell Growth, Migration and Invasion

We explored the functions of MAT in T24 and 5637 cells. MAT meaningfully restrained cell viability (Figure 2A, *P* < 0.01 or *P* < 0.001) and promoted apoptosis (Figure 2E, *P* < 0.001). At the molecular level, when MAT was added, Figure 2B–D displayed that the CyclinD1 expression (*P* < 0.01 or *P* < 0.001) was conspicuously declined and p53 expression (*P* < 0.001) was conspicuously enhanced.

Figure 2 (A) Matrine (MAT) restrained cell growth, invasion and migration of T24 and 5637 cells. MAT restrained cell viability; (B–D) the level of the CyclinD1 was conspicuously declined and p53 was conspicuously enhanced when MAT was added; (E) MAT enhanced cell apoptosis; (F–H) Bcl-2 expression was conspicuously declined while the levels of Bax and Cleaved-Caspase-3 were conspicuously enhanced when MAT was added; (I) MAT meaningfully restrained cell migration; (J) MAT meaningfully restrained cell invasion. ***p* < 0.01; ****p* < 0.001.

Figure 2F–H results pointed out that Bcl-2 expression was conspicuously declined while the levels of Bax and Cleaved-Caspase-3 were conspicuously enhanced ($P < 0.001$). Furthermore, MAT meaningfully restrained cell migration (Figure 2I, $P < 0.01$ or $P < 0.001$) and invasion (Figure 2J, $P < 0.001$). In conclusion, MAT restrained BC cells' growth, invasion and migration.

MAT Up-Regulated LINC00472

LINC00472 was detected in BC tissues, corresponding non-tumor tissues, T24 and 5637 cells. The data revealed LINC00472 was prominently low expressed in BC tissues (Figure 3A, $P < 0.01$). Furthermore, the LINC00472 expression was prominently enhanced when MAT was added (Figure 3B, $P < 0.001$). This implied LINC00472 was positively regulated by MAT.

MAT Restrained Cell Growth, Migration and Invasion by Up-Regulating LINC00472

Firstly, we transfected si-NC and si-00472 into T24 and 5637 cells. We put forward si-00472 statistically declined LINC00472 expression (Figure 4A, $P < 0.001$). Further results revealed that si-00472 disturbed the MAT-induced reduction of viability (Figure 4B, $P < 0.05$) and improvement of apoptotic cells (Figure 4C, $P < 0.05$). At the molecular level, Figure 4D–F displayed that the level of CyclinD1 ($P < 0.01$) was conspicuously enhanced and p53 ($P < 0.01$) was conspicuously declined when transfected si-00472, compared to adding MAT alone. Figure 4G–I pointed out that the levels of Bcl-2 ($P < 0.05$ or $P < 0.01$) were conspicuously enhanced while Bax ($P < 0.01$) and Cleaved-Caspase-3 ($P < 0.001$) were conspicuously declined when transfected si-00472, compared to adding MAT alone. Furthermore, si-00472 meaningfully enhanced MAT-induced cell migration

(Figure 4J, $P < 0.05$ or $P < 0.01$) and invasion (Figure 4K, $P < 0.05$). In short, MAT restrained cell growth, migration and invasion by up-regulating LINC00472 in BC.

MAT Strengthened the Expression of PDCD4 by Up-Regulating LINC00472

Figure 5A and B revealed that MAT could prominently strengthen the expression of PDCD4 ($P < 0.001$), which was declined by si-00472 ($P < 0.01$ or $P < 0.001$). In short, MAT positively regulated PDCD4 expression.

MAT Restrained PTEN/PI3K/AKT Pathways

In Figure 6A, MAT enhanced PTEN expression and decreased the ratio of p/t-PI3K/AKT ($P < 0.001$) while si-00472 declined the MAT-induced rise of ratio of p/t-PI3K/AKT in T24 cells ($P < 0.05$ or $P < 0.01$). A similar phenomenon appeared in 5637 cells (Figure 6B). In short, MAT might exert its function via restraining PTEN/PI3K/AKT pathways in BC.

MAT Inhibited Tumor Growth in vivo

In the animal assay, we noticed that compared with the corresponding group, the tumor volume was significantly decreased by MAT treatment separately at 14 d ($P < 0.05$), 21 d ($P < 0.01$) and 28 d ($P < 0.001$) after injection; however, si-00472 reversed those impacts (Figure 7A). Besides, the tumor weight was declined by MAT treatment ($P < 0.01$), while si-00472 led to an opposite influence ($P < 0.05$) (Figure 7B). This result indicated that MAT inhibited tumor growth in vivo.

Schematic Diagram of Our Findings

We designed a schematic diagram for explaining our findings here as shown in Scheme 1. MAT treatment had no effects on SV-HUC-1. But MAT enhanced PDCD4 expression and

Figure 3 Matrine (MAT) up-regulated long non-coding RNA LINC00472 (LINC00472) in T24 and 5637 cells. (A) LINC00472 expression was prominently lower in BC tissues than in non-tumor tissue; (B) LINC00472 expression was prominently enhanced when MAT was added in T24 and 5637 cells. ** $P < 0.01$; *** $P < 0.001$.

Abbreviations: NT, non-tumor tissue; T, tumor tissue.

Figure 4 Matrine (MAT) restrained cell growth, migration and invasion but induced apoptosis by up-regulating long non-coding RNA LINC00472 (LINC00472) in T24 and 5637 cells. (A) LINC00472 expression was conspicuously declined by si-00472; (B) si-00472 conspicuously recovered the MAT-inhibited viability; (C) si-00472 meaningfully declined MAT-induced apoptosis; (D–F) the level of CyclinD1 was conspicuously enhanced and p53 was conspicuously declined when transfected with si-00472 in cells, compared to adding MAT alone; (G–I) the level of Bcl-2 was conspicuously elevated and Bax and Cleaved-Caspase-3 were conspicuously declined when transfected si-00472, compared to single added MAT alone treatment; (J) si-00472 conspicuously reversed MAT-induced decline in cell migration; (K) si-00472 conspicuously reversed MAT-induced decline in cell invasion. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. **Abbreviations:** NC, negative control; si-00472, small interference RNA of LINC00472.

Figure 5 Matrine (MAT) enhanced programmed cell death protein 4 (PDCD4) expressions by up-regulating long non-coding RNA LINC00472 (LINC00472). (A and B) MAT could prominently strengthened the PDCD4, while si-00472 could prominently decline the MAT-induced rise of PDCD4. ** $P < 0.01$; *** $P < 0.001$.

Abbreviations: MAT, matrine; NC, negative control; si-00472, small interference RNA of LINC00472.

Figure 6 Matrine (MAT) restrained PTEN/PI3K/AKT pathways in T24 and 5637 cells. (A) MAT enhanced PTEN expression and restrained the ratio of p/t-PI3K/AKT in T24 cells; (B) MAT enhanced PTEN expression and restrained the ratio of p/t-PI3K/AKT in 5637 cells. * $P < 0.05$, ** $P < 0.01$; *** $P < 0.001$.

Abbreviations: PTEN, phosphate and tension homology deleted on chromosome ten; PI3K, phosphatidylinoside 3-kinases; AKT, protein kinase B.

Figure 7 Matrine (MAT) inhibited tumor growth in vivo. (A) The tumor volume was decreased by MAT treatment; (B) tumor weight was reduced by MAT treatment. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$, comparing with control; # $p < 0.05$ comparing with si-NC.

Scheme 1 Schematic diagram of our findings. A schematic diagram for understanding our findings was made. It showed the anti-tumor roles of MAT in BC.

Abbreviations: PTEN, phosphate and tension homology deleted on chromosome ten; PI3K, phosphatidylinositol 3-kinases; AKT, protein kinase B; MAT, matrine; PDCD4, programmed cell death protein 4; BC, bladder Carcinoma.

inhibited BC cell proliferation, migration, invasion but promoted apoptosis via up-regulating LINC00472 expression by blocking PTEN/PI3K/AKT pathways in vitro. Moreover, MAT treatment restrained tumor growth by reducing tumor weight and volume in vivo.

Discussion

BC is a familiar malignant carcinoma in the urinary system.²⁵ In this research, we put forward that MAT could restrain the

growth, migration and invasion of T24 and 5637 cells, but did not affect normal cell line-SV-HUC-1. LINC00472 was low expressed in bladder carcinoma. In addition, MAT restrained cell growth, migration and invasion by up-regulating LINC00472. Further, the research put forward that MAT enhanced PDCD4 expression by up-regulating LINC00472. Finally, we put forward that MAT restrained PTEN/PI3K/AKT pathway and si-00472 reversed MAT-induced change in PTEN/PI3K/AKT pathway. Besides, the

in vivo experiments focused on mice also indicated an anti-tumor role of MAT. The tumor volume and tumor weight were declined by MAT and were elevated due to LINC00472 knockdown.

As a type of extract of Chinese traditional medicine *radix sophorae flavescentis*, researches had pointed out that MAT could induce cell apoptosis²⁶ and played a protective role in carcinoma. For instance, a research pointed out that MAT restrained cell proliferation in pancreatic carcinoma²⁷ while the other research pointed out that MAT restrained cell migration in lung carcinoma.²⁸ More interesting, in an animal model of bladder tumor, Gao et al pointed out that MAT could suppress bladder tumor invasion via regulation of the cyclooxygenase-2 (COX-2), and cytosolic recombinant phospholipase A2 (cPLA2).²⁹ MAT treatment induced the inhibition of cell proliferation and invasion of bladder cancer, which was mediated by the PI3K/AKT signaling pathway, in vitro.¹⁴ At the same time, a research displayed that MAT restrained prostate carcinoma, which was another kind of carcinoma of the urinary system.³⁰ Unsurprisingly, we also discovered MAT could restrain cell viability, migratory and invasive potentials but promote apoptosis, and played a protective role in BC cells.

Studies have pointed out that MAT usually plays an anti-carcinoma efficacy by regulating RNA. For instance, Wei et al pointed out MAT declined the viability via the miR-19b-3p.³¹ Li et al pointed out MAT exerted anti-carcinoma efficacy in prostate carcinoma via forkhead box O (FOXO).³² Furthermore, Liao Li et al pointed out MAT restrained BC cells by up-regulating vascular endothelial growth factor (VEGF) and Wang et al pointed out LINC00472 could be a diagnostic marker in renal tumor.³³ So we conjectured LINC0047 might be a potential treatment option in BC.

LINC00472 is a newly discovered lncRNA with anti-carcinoma effects and usually has lower expression in carcinoma.³⁴ For instance, a research had pointed out LINC00472 restrained breast carcinoma cell migration,²³ and the other research had pointed out LINC00472 enhanced apoptosis in lung carcinoma.³⁵ On the other side, a research pointed out that LINC00472 could be as a diagnostic marker in nephropathy.³⁶ Unsurprisingly, we put forward MAT positively regulated LINC00472, and si-00472 changed the impacts of MAT on cell viability, apoptosis, migration and invasion. Ulteriorly, we uncovered a research displaying that LINC00472 played a efficacy in carcinoma by promoting PDCD4,³⁷ so we further studied the impacts of PDCD4 on BC.

PDCD4 is first discovered in 1994, and it situates in 10q24.³⁸ PDCD4 is an anti-carcinoma gene related to cell cycle and apoptosis.³⁹ Many researches had pointed out it could be used as a therapeutic target for various carcinomas.⁴⁰ Studies also pointed out that PDCD4 was low expressed in various carcinoma tissues such as gastric carcinoma and so on.⁴¹ However, the reasons for the decrease of PDCD4 expression levels in carcinomas and their molecular mechanisms were not fully understood. Therefore, we further studied the relationship between MAT, LINC00472 and PDCD4. Gratifyingly, we discovered that PDCD4 was positively regulated by MAT, and MAT enhanced PDCD4 expression by up-regulating LINC00472.

The PTEN/PI3K/AKT pathway plays vital efficacies in carcinoma. Previous researches had reported that PTEN/PI3K/AKT pathway participated in the pathogenesis of BC.⁴² In addition, a research had put forward MAT exerted anti-carcinoma roles via PTEN/PI3K/AKT pathway.⁴³ Other research pointed out LINC00641 served in BC via PTEN/PI3K/AKT pathway.⁴⁴ In this paper, the results were completely consistent with those above results. MAT enhanced the expression of PTEN and restrained the ratio of p/t-PI3K and p/t-AKT, while si-00472 reversed MAT-induced effects. In short, MAT restrained PTEN/PI3K/AKT signaling pathways by up-regulating LINC00472.

Conclusion

In summary, MAT restrained tumor growth in vivo and abated cell growth, invasion and migration by up-regulating LINC00472/PDCD4 via restraining PTEN/PI3K/AKT pathways in BC cells. This article might afford a new thought for the clinical treatment of BC.

Ethical Approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. The research was ratified by the Medical Ethics Committee of the China-Japan Union Hospital of Jilin University.

Data Sharing Statement

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgment

Linlin Li and Fei Qi are co-first authors for this study.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Disclosure

The authors declare that they have no conflicts of interest in this work.

References

- Smith AB, Jaeger B, Pinheiro LC, et al. Impact of bladder cancer on health-related quality of life. *BJU Int*. 2018;121(4):549–557. doi:10.1111/bju.14047
- Martinez Rodriguez RH, Buisan Rueda O, Ibarz L. Bladder cancer: present and future. *Med Clin (Barc)*. 2017;149(10):449–455. doi:10.1016/j.medcli.2017.06.009
- He YT, Li DJ, Liang D, et al. [Incidence and mortality of bladder cancer in China, 2014]. *Zhonghua Zhong Liu Za Zhi*. 2018;40(9):647–652. doi:10.3760/cma.j.issn.0253-3766.2018.09.002
- Grayson M. Bladder cancer. *Nature*. 2017;551(7679):S33. doi:10.1038/551S33a
- Bladder cancer: diagnosis and management of bladder cancer: (c) NICE. (2015) Bladder cancer: diagnosis and management of bladder cancer. *BJU Int*. 2017;120(6):755–765. doi:10.1111/bju.2017.120.issue-6
- Kaufman DS, Shipley WU, Feldman AS. Bladder cancer. *Lancet*. 2009;374(9685):239–249. doi:10.1016/S0140-6736(09)60491-8
- Muhammad N, Steele R, Isbell TS, Philips N, Ray RB. Bitter melon extract inhibits breast cancer growth in preclinical model by inducing autophagic cell death. *Oncotarget*. 2017;8(39):66226–66236. doi:10.18632/oncotarget.19887
- Bhattacharya S, Muhammad N, Steele R, Peng G, Ray RB. Immunomodulatory role of bitter melon extract in inhibition of head and neck squamous cell carcinoma growth. *Oncotarget*. 2016;7(22):33202–33209. doi:10.18632/oncotarget.v7i22
- Wang S, Wu X, Tan M, et al. Fighting fire with fire: poisonous Chinese herbal medicine for cancer therapy. *J Ethnopharmacol*. 2012;140(1):33–45. doi:10.1016/j.jep.2011.12.041
- Michael JP. Indolizidine and quinolizidine alkaloids (July 1999 to June 2000). *Nat Prod Rep*. 2001;18(5):520–542. doi:10.1039/b005384h
- Yong J, Wu X, Lu C. Anticancer advances of matrine and its derivatives. *Curr Pharm Des*. 2015;21(25):3673–3680. doi:10.2174/1381612821666150122123748
- Huang J, Xu H. Matrine: bioactivities and structural modifications. *Curr Top Med Chem*. 2016;16(28):3365–3378. doi:10.2174/1568026616666160506131012
- Huang H, Du T, Xu G, et al. Matrine suppresses invasion of castration-resistant prostate cancer cells by downregulating MMP-2/9 via NF-kappaB signaling pathway. *Int J Oncol*. 2017;50(2):640–648. doi:10.3892/ijo.2016.3805
- Yang Y, Guo J-X, Shao Z-Q, Gao J-P. Matrine inhibits bladder cancer cell growth and invasion in vitro through PI3K/AKT signaling pathway: an experimental study. *Asian Pac J Trop Med*. 2017;10(5):515–519. doi:10.1016/j.apjtm.2017.05.009
- Li G-D. Further thoughts on abnormal chromatin configuration and oncogenesis. *arXiv Preprint arXiv:1901.04432*. 2019. Available from: <https://arxiv.xileosou.top/abs/1901.04432>.
- Li Y, Zheng F, Xiao X, et al. CircHIPK3 sponges miR-558 to suppress heparanase expression in bladder cancer cells. *EMBO Rep*. 2017;18(9):1646–1659. doi:10.15252/embr.201643581
- Muhammad N, Bhattacharya S, Steele R, Ray RB. Anti-miR-203 suppresses ER-positive breast cancer growth and stemness by targeting SOCS3. *Oncotarget*. 2016;7(36):58595–58605. doi:10.18632/oncotarget.v7i36
- Jarroux J, Morillon A, Pinskaya M. History, discovery, and classification of lncRNAs. *Adv Exp Med Biol*. 2017;1008:1–46.
- Peng WX, Koirala P, Mo YY. LncRNA-mediated regulation of cell signaling in cancer. *Oncogene*. 2017;36(41):5661–5667. doi:10.1038/onc.2017.184
- Wang Y, Kong D. Knockdown of lncRNA MEG3 inhibits viability, migration, and invasion and promotes apoptosis by sponging miR-127 in osteosarcoma cell. *J Cell Biochem*. 2018;119(1):669–679. doi:10.1002/jcb.26230
- Hu YC, Wang AM, Lu JK, Cen R, Liu LL. Long noncoding RNA HOXD-AS1 regulates proliferation of cervical cancer cells by activating Ras/ERK signaling pathway. *Eur Rev Med Pharmacol Sci*. 2017;21(22):5049–5055. doi:10.26355/eurrev_201711_13817
- Chen L, Zhang W, Li DY, et al. Regulatory network analysis of LINC00472, a long noncoding RNA downregulated by DNA hypermethylation in colorectal cancer. *Clin Genet*. 2018;93(6):1189–1198.
- Liao Q, Shen Y, Katsaros D, et al. Prognostic and predictive values of long non-coding RNA LINC00472 in breast cancer. *Clin Genet*. 2015;6(11):8579–8592.
- Liao XZ, Tao LT, Liu JH, et al. Matrine combined with cisplatin synergistically inhibited urothelial bladder cancer cells via down-regulating VEGF/PI3K/Akt signaling pathway. *Cancer Cell Int*. 2017;17:124. doi:10.1186/s12935-017-0495-6
- Skeldon SC, Larry Goldenberg S. Bladder cancer: a portal into men's health. *Urol Oncol*. 2015;33(1):40–44. doi:10.1016/j.urolonc.2013.12.005
- Tetik Vardarli A, Duzgun Z, Erdem C, Kaymaz BT, Eroglu Z, Cetintas VB. Matrine induced G0/G1 arrest and apoptosis in human acute T-cell lymphoblastic leukemia (T-ALL) cells. *Bosn J Basic Med Sci*. 2018;18(2):141–149. doi:10.17305/bjbm.2017.2457
- Cho YR, Lee JH, Kim JH, et al. Matrine suppresses KRAS-driven pancreatic cancer growth by inhibiting autophagy-mediated energy metabolism. *Mol Oncol*. 2018;12(7):1203–1215. doi:10.1002/1878-0261.12324
- Xie W, Lu J, Lu Q, et al. Matrine inhibits the proliferation and migration of lung cancer cells through regulation of the protein kinase B/glycogen synthase kinase-3beta signaling pathways. *Exp Ther Med*. 2018;16(2):723–729. doi:10.3892/etm.2018.6266
- Gao H, Guo Y, Deng N, et al. Suppressive effect of matrine on tumor invasion in N-Butyl-N-(4-Hydroxybutyl)Nitrosamine-induced urinary bladder carcinogenesis. *Chemotherapy*. 2014;60(2):119–128. doi:10.1159/000371439
- Chang J, Hu S, Wang W, et al. Matrine inhibits prostate cancer via activation of the unfolded protein response/endoplasmic reticulum stress signaling and reversal of epithelial to mesenchymal transition. *Mol Med Rep*. 2018;18(1):945–957. doi:10.3892/mmr.2018.9061
- Wei YP, Wang XH, Liu G, et al. Matrine exerts inhibitory effects in melanoma through the regulation of miR-19b-3p/PTEN. *Int J Oncol*. 2018;53(2):791–800. doi:10.3892/ijo.2018.4414
- Li Q, Li Q, Huang H, et al. Regulatory effects of antitumor agent matrine on FOXO and PI3K-AKT pathway in castration-resistant prostate cancer cells. *Prostate*. 2018;61(5):550–558.
- Wang J, Zhang C, He W, Gou X. Construction and comprehensive analysis of dysregulated long non-coding RNA-associated competing endogenous RNA network in clear cell renal cell carcinoma. 2018.
- Sui J, Li YH, Zhang YQ, et al. Integrated analysis of long non-coding RNA-associated ceRNA network reveals potential lncRNA biomarkers in human lung adenocarcinoma. *Int J Oncol*. 2016;49(5):2023–2036. doi:10.3892/ijo.2016.3716

35. Su C, Shi K, Cheng X, et al. Long noncoding RNA LINC00472 inhibits proliferation and promotes apoptosis of lung adenocarcinoma cells via regulating miR-24-3p/DEDD. *Technol Cancer Res Treat*. 2018;17:1533033818790490.
36. Wang YZ, Zhu DY, Xie XM, et al. EA15, MIR22, LINC00472 as diagnostic markers for diabetic kidney disease. *J Cell Physiol*. 2019;234(6):8797–8803. doi:10.1002/jcp.27539
37. Ye Y, Yang S, Han Y, et al. Linc00472 suppresses proliferation and promotes apoptosis through elevating PDCD4 expression by sponging miR-196a in colorectal cancer. *Aging*. 2018;10(6):1523–1533. doi:10.18632/aging.v10i6
38. Soejima H, Miyoshi O, Yoshinaga H, Masaki Z. Assignment of the programmed cell death 4 gene (PDCD4) to human chromosome band 10q24 by in situ hybridization. *Cytogenet Genome Res*. 1999;87(1/2):113. doi:10.1159/000015408
39. Matsuhashi S, Manirujjaman M. Control mechanisms of the tumor suppressor PDCD4: expression and functions. *Int J Mol Sci*. 2019;20(9):2304.
40. Lankat-Buttgereit B, Goke R. Programmed cell death protein 4 (pdc4): a novel target for antineoplastic therapy? *Biol Cell*. 2003;95(8):515–519. doi:10.1016/j.biocel.2003.09.003
41. Motoyama K, Inoue H, Mimori K, et al. Clinicopathological and prognostic significance of PDCD4 and microRNA-21 in human gastric cancer. *Int J Oncol*. 2010;36(5):1089–1095. doi:10.3892/ijo.0000590
42. Yang X, Cheng Y, Li P, et al. A lentiviral sponge for miRNA-21 diminishes aerobic glycolysis in bladder cancer T24 cells via the PTEN/PI3K/AKT/mTOR axis. *Tumour Biol*. 2015;36(1):383–391. doi:10.1007/s13277-014-2617-2
43. Wang B, Xu J, Wang H, Chang S, Liu N. Effect and mechanism of sophoridine to suppress hepatocellular carcinoma in vitro and vivo. *Biomed Pharmacother*. 2017;95:324–330. doi:10.1016/j.biopha.2017.08.029
44. Li Z, Hong S, Liu Z. LncRNA LINC00641 predicts prognosis and inhibits bladder cancer progression through miR-197-3p/KLF10/PTEN/PI3K/AKT cascade. *Biochem Biophys Res Commun*. 2018;503(3):1825–1829. doi:10.1016/j.bbrc.2018.07.120

Cancer Management and Research

Dovepress

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>