

Management of hyperuricemia in gout: focus on febuxostat

Mattheus K Reinders¹
Tim L Th A Jansen²

¹Clinical Pharmacy, Atrium Medisch Centrum Parkstad, Heerlen, The Netherlands;

²Rheumatology, Medisch Centrum Leeuwarden, Leeuwarden, The Netherlands

Abstract: Gout is the most common inflammatory arthritis in an elderly population, and can be diagnosed with absolute certainty by polarization microscopy. However, diagnosis may be challenging because atypical presentations are more common in the elderly. Management of hyperuricemia in the elderly with gout requires special consideration because of co-medication, contra-indications, and risk of adverse reactions. Urate-lowering agents include allopurinol and uricosuric agents. These also must be used sensibly in the elderly, especially when renal function impairment is present. However, if used at the lowest dose that maintains the serum urate level below 5.0 to 6.0 mg/dL (0.30 to 0.36 mmol/L), the excess urate in the body will eventually be eliminated, acute flares will no longer occur, and tophi will resolve. Febuxostat, a new xanthine oxidase inhibitor, is welcomed, as few alternatives for allopurinol are available. Its pharmacokinetics and pharmacodynamics are not significantly altered in patients with moderate renal function or hepatic impairment. Its antihyperuricemic efficacy at 80 to 120 mg/day is better than “standard dosage” allopurinol (300 mg/day). Long-term safety data and efficacy data on tophus diminishment and reduction of gout flares have recently become available. Febuxostat may provide an important option in patients unable to use allopurinol, or refractory to allopurinol.

Keywords: aging, febuxostat, hyperuricemia, gout, pharmacotherapy, xanthine oxidase

Introduction

Gout is the collective name for several disorders that are characterized by the formation and deposition of monosodium urate (MSUr) crystals. The condition is associated with recurrent episodes of acute joint pain due to the deposition of MSUr crystals in the synovial fluid. In addition to the effects observed in the joints, skin/subcutaneous tissue and kidneys may also be affected by tophaceous deposits, cellulitis, urate nephropathy, and/or kidney stones, respectively. In most cases, no identifiable underlying cause of gout is present, but evident factors are usually present that may contribute to increases in urate (uric acid) levels, such as reduced renal function, obesity, and the use of diuretics. Hyperuricemia may exist for several years to decades before the first symptoms of gout attacks appear; therefore it is a disease associated and correlated with aging.

Gout is one of the most common inflammatory arthritis affecting the elderly; however in general it appears to be poorly managed.¹⁻³ Partly this is due to an absence of sufficient treatment strategies and guidelines. In 2006, the European League against Rheumatism (EULAR) published the first international recommendations for the diagnosis and treatment of gout.^{4,5} The development of these EULAR and British Society of Rheumatology (BSR) guidelines coincided with improved professional and patient education as well

Correspondence: MK Reinders
Atrium Medisch Centrum Parkstad;
Henri Dunantstraat 5, 6941 PC Heerlen,
The Netherlands
Tel +31 45 5767419
Fax +31 45 5767279
Email m.reinders@atriummc.nl

as the urge for improved professional performance. Until recently, allopurinol was the only antihyperuricemic drug worldwide available. Two uricosurics with barriers regarding availability (benzbromarone and probenecid) have been available for years, but their exact place in treatment strategy is not clear. In 2008 and 2009 respectively, European Medicines Agency (EMA) and Food and Drug Administration (FDA) have approved febuxostat, a novel selective xanthine inhibitor, for the treatment of hyperuricemia in gout patients, respectively. Febuxostat greatly expands the treatment options for refractory or allopurinol-intolerant gout, given the huge prevalence of the disorder and good worldwide availability of only one antihyperuricemic drug.

The incidence and prevalence of gout in the elderly is increasing.^{1,6,7} This appears related to improved lifespan leading to similar increases in age-related diseases (eg, cardiovascular diseases) and their associated adverse effects of treatment (eg, diuretics and low-dose salicylates) which can increase the risk of gout. “Elderly onset gout” differs from “classical” gout found in middle-aged men in several respects: no male predominance but an equal gender distribution, polyarticular presentation with upper-extremity-joint involvement, fewer acute gouty episodes, indolent clinical course, and an increased incidence of tophi.^{1,8} Several reviews have addressed the specific challenges of gout treatment in the elderly.^{1,8-11} This review will focus on febuxostat for the management of gout in the elderly.

Gout pathophysiology

Uric acid is formed from nucleic acid either endogenously from cell breakdown or exogenously from metabolism of food. Cooling and acidification of the microenvironment, which can result in acute formation of urate crystals, reduce the solubility of MSUr. The gut excretes one-third of urate and two-thirds are excreted renally. Renal urate transport is typically explained by a 4-component model: glomerular filtration, a near-complete reabsorption of filtered urate, subsequent secretion, and postsecretory reabsorption in the remaining proximal tubule.¹² Recently, several new urate transporters have been identified as playing key roles in urate homeostasis, including URAT-1, and Glut9.^{12,13} The regulation of serum uric acid levels is under a strong genetic control. A recent meta-analysis of genome-wide association scans shows that common DNA variants at 9 different loci are associated with uric acid concentrations.¹⁴

Excessive consumption of alcohol (particularly beer), sweetened soft drinks, fructose, meat, and seafood can also increase levels of serum urate (sUr).¹²

Inhibition of urate transporters can be achieved by uricosurics, and production of uric acid can be inhibited using xanthine oxidase inhibitors, such as allopurinol. Febuxostat is a new selective inhibitor of xanthine oxidase. Uric acid deposits can also be lysed by the enzyme uricase, the coding gene for which became defective in humans in the Miocene because of an evolutionary mutation. The combined absence of uricase and almost total reabsorption of filtered urate explains that humans (and the greater apes) have 10-fold higher sUr levels than other mammals.

Drug-induced hyperuricemia

Chronic diuretic therapy is associated with reduced excretion of uric acid. Mechanisms are increased uric acid reabsorption in the proximal tubule secondary to volume depletion, and competition between the diuretic and uric acid for the organic acid secretory mechanism in the proximal tubule.¹⁵ Low-dose diuretic therapy in hypertensive patients does not seem to alter serum urate levels significantly.^{15,16} Indeed, the requirement for anti-gout therapy in hypertensive patients is doubled for thiazide doses of ≥ 25 mg/day (in hydrochlorothiazide equivalents); no significant increase in risk is seen for lower doses.^{15,17} Similarly, low-dose therapy with a loop diuretic is not associated with hyperuricemia.^{15,18} However, low-dose diuretic therapy may be effective in hypertension but insufficient in patients with chronic heart failure who often additionally suffer from chronic renal failure.

Salicylates are known to interact with renal urate handling and low doses inhibit urate excretion. In one study it was found that even mini-dose aspirin (75 mg/day) was associated with a 15% decrease of urate excretion in elderly patients.¹⁹ Thus, in elderly people with cardiovascular diseases not only diuretics, but also mini-dose aspirin is of importance.

Characteristics of gout presentation

Gout can be diagnosed with certainty only by identification of urate (MSUr) crystals, present in joints during acute attacks of gout, or in tophi. The clinical practitioner can confirm the presence and type of crystals by polarization microscopy.^{4,20,21}

The characteristic profile of gout is that of severe monoarthritis occurring within several hours. The first metatarsophalangeal joint is affected in 50% of gout attacks, and this is known as podagra. Gout may be localized in other joints, but shoulders, hips, and the vertebral column are rarely affected. The initial gout attack usually involves monoarthritis, but long-term gout over several years may

become polyarticular and could lead to increasing joint damage. Similarly, a positive uric acid balance over a number of years can cause tophaceous deposits, possibly with periodic arthritis.

Urate production

Primary gout tends to involve low urate excretion, which is primarily originated in the proximal tubule. Only a minority of cases involves overproduction of urate. In some treatments of cancer (particularly lymphomas and leukemias), patients can develop tumour lysis syndrome including severe hyperuricemia with risk of urate nephropathy.

Urate nephropathy in gout

Aggressive chemotherapy among patients with chronic leukemia or malignant lymphoma could cause an excessive supply of uric acid resulting in acute urate nephropathy due to the deposition of urate crystals in collection ducts and ureters. Several kinds of urate crystals can be found, including uric acid crystals, amorphous urate, monosodium urate, and ammonium urate crystals.

In chronic hyperuricemia, the risk of developing renal crystals increases as serum urate concentrations rise. The risk is about 10% with serum urate 0.42 to 0.48 mmol/L, but can rise to 50% with serum urate concentrations >0.70 mmol/L. In the absence of stones or other risk factors (such as hypertension), the risk of urate nephropathy has generally been considered low.²²

Radiographic presentations of gout

X-ray examination at the initial onset of gout has revealed no abnormalities except for possible pre-existing arthrosis and soft tissue edema. Cartilage and bone might be affected by chronic and/or recurring arthritis, and subsequently exhibit narrowing of the joint cavity because of the disappearance of cartilage, and erosions or cysts because of contact with juxta-articular bone. These abnormalities and the appearance of the erosions may raise suspicions of gout, but erosions are a secondary manifestation and non-diagnostic characteristic of early gout. In chronic gout, however, some of these characteristic changes can help with diagnosis.

Treatment strategies for gout

Several approaches to the treatment of gout are available depending on the patient's presentation of the disease. Optimal treatment often requires a combination of pharmacological intervention and lifestyle changes. Treatment should be tailored to the patient's prognostic factors (high sUr, previous

attacks, and radiographic signs), the clinical phase of the disease (acute, recurrent, tophaceous) and general risk factors, such as obesity, alcohol consumption, renal impairment, use of diuretics or other risk factors for secondary hyperuricemia. Primary prevention of gout often involves changes in lifestyle, such as a low-purine/weight-reducing diet or restricting alcohol intake; however, many patients are unlikely to undertake such changes until they are diagnosed with the disease, which often occurs when the symptoms are presented in the form of an attack of gout. Acute gout is usually treated by reducing the inflammation of the affected joint with non-steroidal anti-inflammatory drugs (NSAIDs), colchicine, corticosteroids, and cooling.^{5,23–26} Once the acute gout has subsided the objective is to prevent disease recurrence. This might involve lifestyle changes and low doses of NSAIDs or colchicine. In patients with high sUr levels who suffer from frequent attacks of gout, the use of urate-lowering drugs is warranted.

The following is an overview of the different drug classes and their potential use as part of the treatment strategies for gout. Information on these drugs is presented in Table 1 and the current therapeutic strategy is summarized in Table 2.

Some drugs have a documented (modest) urate lowering side effect because of enhanced urate clearance by the kidney. These include losartan, fenofibrate, high dose aspirin and vitamin C.^{27–30} In patients with gout, these drugs might be preferred above other drugs in the same class.

Primary prevention of gout

Primary prevention of gout involves changes in lifestyle, such as changes to diet (low-purine/weight-reducing diet) and restricting alcohol consumption. No randomized studies have been conducted evaluating the effect of lifestyle changes on the incidence of attacks in patients with gout. Nevertheless, experts agree that lifestyle changes may have some effect. Physicians in daily practice also give lifestyle advice, when gout symptoms appear. However, fewer than 20% of patients with gout seeking medical advice are prepared to make long-term changes in lifestyle.³¹ Recently, the negative role of meat, seafood and beer consumption, and the protective role of dairy products in the development of gout were demonstrated in a prospective study over a 12-year period among a population of around 47,000 healthy male subjects.³²

Treatment options for prevention of recurrence of gout

When gout has subsided, it is important to reduce the sUr concentration to prevent recurrence of gouty attacks.

Table 1 Antihyperuricemic drugs in gout

Drug	Action: indication	Daily dose: standard	Distinct clinical pharmacological features
Allopurinol po	XOI: all	100–900 mg: 300 mg	Dosage adjustment to renal function
Benzbromarone po	urate transporter: low excretor, subject to intolerance or allergy to allopurinol	50–200 mg: 100 mg	Poor efficacy in severe renal function impairment
Febuxostat po	XOI: all, particularly in allopurinol intolerance	80–120 mg: 80 mg	No dosage adjustment in renal impairment
Probenecid po	urate transporter: low excretor	500–2000 mg: 1000 mg	Poor efficacy in moderate–severe renal function
Rasburicase iv	UrO: lytic effect on tophi	Compassionate use: eg, 0.2 mg/kg in 60 min infusion on day 1, then 1 × per week; (+ methylprednisolone 100 mg iv)	Biological

Abbreviations: iv, intravenous; po, oral; UrO, urate oxidase; XOI, xanthine oxidase inhibitor.

This involves restriction of alcohol, weight-loss in cases of obesity (not too rapidly as this can trigger gout), and ensuring adequate diuresis. Dietary measures are very important.³² However, a strict low-purine diet can achieve only a limited reduction in sUr levels (≤ 0.10 mmol/L or 1.7 mg/dL), and many patients have difficulty with adhering to long-term dietary changes.³¹ If possible, it is recommended that patients should discontinue diuretics if indicated for hypertension (find alternative antihypertensives), but should continue diuretics if indicated for overhydration/congestive heart failure. However, the effectiveness of these measures is limited and is not supported by controlled studies.

With tophaceous or recurrent gout (≥ 2 attacks per annum) the use of urate-lowering therapy may be warranted.³³ In contrast, asymptomatic hyperuricemia does not require specific treatment. The maintenance dose of urate-lowering drugs is preferably adjusted to the clinical effect: (1) prevention of gout attacks without using colchicine and/or NSAIDs, (2) disappearance of tophi based on the sUr-concentration with tophaceous deposits. Tophi may well disappear when lower sUr target values have been achieved to at least “normal” levels, but preferably low to normal (≤ 0.30 mmol/L).³⁴ In order to prevent further attacks of gout, stable sUr target values of at least ≤ 0.36 mmol/L are required, but in some

Table 2 Suggested experience-based strategy for initiation of antihyperuricemic therapy

1. Confirmation of diagnosis: detect urate crystals by means of polarization microscopy
2. Two or more gout flares per annum or tophi/joint destruction due to gout attacks
3. Therapeutic advice¹: allopurinol 100–300 mg/day²
 - a. allopurinol intolerance: consider febuxostat
4. Laboratory monitoring of effectiveness at 6–8 weeks:³
 - (a) sUr < 0.30 mmol/L, then continue allopurinol
 - (b) sUr > 0.30–0.36 mmol/L, but no further attacks (without colchicine/NSAID/corticosteroids), then continue allopurinol
 - (c) sUr > 0.30 mmol/L plus gout attacks/persistent tophi and uUr > 1.5 mmol/24 hours, go to 5
 - (d) sUr > 0.30 mmol/L plus gout attacks/persistent tophi with uUr < 1.5 mmol/24 hours, go to 6
5. Therapeutic advice: increase allopurinol (eg, +100 mg/day or double the dose); then go to 4.
Allopurinol inefficacy at maximum dosage (corrected for renal function): consider febuxostat
6. Therapeutic advice: add uricosuricum, eg, benzbromarone 100 mg/day, or probenecid 500 twice daily; then go to 4
7. Laboratory monitoring of effectiveness sUr and uUr (possibly pH⁴) after 6 months: see 5
8. NB.: when trying to clear tophi, target value is sUr < 0.30 mmol/L

Notes: ¹Subject to motivation and tolerance by patient. ²Subject to calculated creatinine clearance (cCC) > 50 mL/min, if cCC < 50 mL/min, then only increase allopurinol with 100 mg/day. Serum oxipurinol concentrations might be measured in patients with renal insufficiency. ³Target value for sUr ≤ 0.36 mmol/L might be sufficient when there are no further gout attacks despite withdrawing colchicine/NSAID, otherwise lower target value of 0.30 mmol/L. ⁴If experiencing kidney stones or uUr > 6.0 mmol/24 hours and pH < 7.0 consider alkalinising with sodium bicarbonate.

Abbreviations: sUr, serum urate; uUr, 24-hour excretion of urate in urine; NSAID, non-steroidal anti-inflammatory drug.

patients ≤ 0.30 mmol/L may well be preferable; as has also been recommended by the British guidelines.^{35–38}

There is some but not much evidence for these biochemical targets. But in practice they give clarity also to the patient. Another predefined aim of therapy particularly in elderly gout patients may be adhered to by clinicians treating elderly, ie, a normalization/any lowering of sUr to a level at which the gout patient does not need the chronic use of NSAIDs, glucocorticoids or colchicine; which was not specifically claimed for in aforementioned guidelines. Two main classes of drugs that reduce serum urate concentrations exist: xanthine oxidase inhibitors (XOi), such as allopurinol and febuxostat, and uricosuric drugs, such as benzbromarone and probenecid. XOi work by inhibiting uric acid production, thereby reducing serum urate concentrations. Uricosuric drugs inhibit the reabsorption of uric acid mediated by urate transporters. Rather few comparative randomized studies have been carried out comparing the effectiveness of these drugs.^{39–44}

In the elderly, gout is often accompanied by renal function impairment and urate excretion is often low to very low. It must be kept in mind that when starting therapy with an XOi (uricostatics) it can take upto 8 weeks before a new steady state serum urate concentration is achieved. Secondly, when using allopurinol, the dosage should be tailored to renal function. However, since dosing tables for allopurinol are cautious and poorly investigated in clinically trials, these patients are often far from optimally treated.^{45–47} Possibly, febuxostat (in comparison to allopurinol) will provide a much more convenient and effective treatment option in this group of patients.

Special considerations when treating gout

Antihyperuricemic therapy might provoke arthritis or induce an attack of gout. For safety reasons, antihyperuricemic therapy should only be given after a gout attack has subsided, preferably with protection from colchicine, which should be initiated several days to 2 weeks earlier (0.5 mg twice daily).^{5,48} With normal renal function, administration of allopurinol could be started at a dose of 100 mg once daily, probenecid 250 mg twice daily and benzbromarone 100 mg once daily, and titrated every 2 weeks to standard dosage if tolerated. After 6 to 8 weeks, the efficacy can be evaluated and dose can be increased if necessary. Maintenance doses usually range from allopurinol 200 to 600 mg once daily, benzbromarone 100 to 200 mg once daily and probenecid 500 to 1000 mg twice daily. In case of impaired renal function, allopurinol dosage should be started 100 or 50 mg/day (depending on

renal function), evaluated after 6 to 8 weeks and titrated carefully (with 100 or 50 mg/day).

When frequent attacks of gout without joint damage or tophi are present in patients with intolerance or allergy to allopurinol and uricosurics, prophylaxis with colchicine at low doses can be prescribed, eg, 0.5 mg once or twice daily in patients with good renal function. In exceptional cases, corticosteroids, or a combination of a uricosuricum and allopurinol, may be indicated for maintenance therapy. When a history of urolithiasis is present, adequate diuresis should be ensured and alkalization should be considered, especially if a uricosuric is prescribed.

Compliance is also a special consideration when supervising gout patients, and it is crucial to explain the dosing schedule and any potential side effect to the patient in order to prevent early withdrawal.

Xanthine oxidase inhibitors Position of XOi

Allopurinol, a purine-analogue, is widely prescribed as the daily urate-lowering treatment (ULT) of first choice. Allopurinol has been proven to be effective in all cases of hyperuricemia. Because of the long half-life of the active metabolite oxipurinol (14 to 28 hours), allopurinol can be administered once daily. It can be given as a single dose of commonly 300 mg daily and may be titrated up to a maximum 800 mg daily if needed. In some patients only a daily dose of 100 mg may be adequate. Dosages above 300 mg/day are given 2 to 3 times daily.

By application of a dosage of 300 mg allopurinol the (British) target of sUr < 0.30 mmol/L can be reached in 20% to 26% of patients; but if tolerated, a dosage of 600 mg allopurinol is needed to reach aforementioned target in 78% of patients as demonstrated recently in a prospective randomized controlled trial (RCT).^{43,44} Approximately 20% of patients using allopurinol report adverse events, with 5% discontinuing its use.⁴⁹

Rare but life threatening is the allopurinol hypersensitivity syndrome.⁵⁰ Serious adverse reactions to allopurinol have been related to a decreased creatinine clearance rate and prolonged half-life of oxipurinol, and therefore it was proposed to adjust the allopurinol dose according to the rate of creatinine clearance.⁵¹ The dosage of allopurinol in individual patients should be tailored based on serum urate and calculated creatinine clearance.

Currently, an alternative XOi may be prescribed, ie, febuxostat.^{52,53} Up to now the predominant alternative treatment option in cases of allopurinol intolerance has been a

uricosuric therapy. In one recent RCT benzbromarone 100 to 200 mg daily showed similar success rates when compared with allopurinol 300 to 600 mg daily, and superior to the uricosuric probenecid.^{43,44} The long-term safety of benzbromarone however has not been a topic of research. As there was some doubt on its safety, it was withdrawn from the market in 2003.^{54,55}

An emerging, potentially highly effective treatment is the use of uricase enzymes, such as rasburicase and pegloticase. In some individual cases and one clinical trial it is successfully used to dissolve tophi.^{56,57}

Febuxostat

Febuxostat pharmacology

The active substance is a new chemical entity designated as 2-[3-cyano-4-(2-methylpropoxy) phenyl]-4-methylthiazole-5-carboxylic acid (Figure 1). It is a non-purine, selective xanthine oxidase/xanthine dehydrogenase inhibitor. It is packed in a dosage of 80 or 120 mg febuxostat.

Febuxostat 10 to 120 mg/day dose-dependently reduced mean sUr levels from baseline by 25% to 70% in healthy volunteers; 24-hour urinary uric acid excretion at day 8 was decreased by 46% to 66% relative to placebo.⁵⁸ Both effects seemed to plateau at dosages >120 mg/day.⁵⁸ Age (18 to 40 years versus ≥65 years) and sex had no clinically significant effect on the pharmacokinetic and pharmacodynamic properties of oral febuxostat 80 mg/day in healthy volunteers.⁵⁹

Absorption of febuxostat is rapid, with a time to C_{max} of ≈1 hour. Febuxostat can be administered regardless of food or antacid intake.⁶⁰ Pharmacokinetic values are linear in the range of 10 to 120 mg. Febuxostat is highly (>98%) bound to human plasma proteins, mainly at the diazepam binding site. Elimination half-life of febuxostat is approximately 12 hours. Febuxostat is mainly eliminated by glucuronidation in the liver. Also, some quantifiable active metabolites 67M-1,

67M-2, and 67M-4 were found.^{61–63} No dosage adjustments are recommended in patients with mild to moderate renal impairment. Febuxostat inhibits cytochrome P450 isoenzyme 2D6, but interactions with CYP2D6 are not considered clinically significant. Just as in allopurinol, an important drug–drug interaction can occur with mercaptopurine and azathioprine due to inhibition of xanthine oxidase. Although no data are available yet, the combination of these drugs with febuxostat should be avoided.

Two phase I studies investigated the pharmacokinetics, pharmacodynamics, and safety of febuxostat in male and female subjects with normal or mild, moderate or severe renal function impairment.^{64,65} One study (n = 32) used febuxostat in the standard dosage of 80 mg/day.⁶⁴ Although plasma exposure to febuxostat and its metabolites was generally higher in subjects with increasing degrees of renal impairment, the percentages of decrease in serum uric acid were comparable regardless of the renal function group. A once-daily 80-mg dose of febuxostat appears to be safe and well tolerated in different renal function groups and does not appear to require any dose adjustment based on differences in renal function.⁶⁴

One open label phase I study (n = 27) investigated the pharmacokinetics and pharmacodynamics as well as safety of febuxostat in male and female subjects with normal liver function (n = 11) and subjects with mild (n = 8) and moderate (n = 8) hepatic impairment.⁶⁶ Caveat: patients with severe hepatic impairment, according to Child–Pugh classification, were excluded, and febuxostat 80 mg was given once daily for only 7 days. Overall, mild to moderate hepatic impairment did not significantly affect exposure to febuxostat or its active metabolites during this 1-week period.

Clinical efficacy

In a phase II dose-response study the efficacy of 40 mg, 80 mg, and 120 mg/day febuxostat was evaluated in 153 patients with hyperuricemia (baseline sUr ≥ 0.48 mmol/L) and gout.⁶⁷ Patients were aged 23 up to 80 years. Subjects received febuxostat (40 mg, 80 mg, 120 mg) or placebo once daily for 28 days and colchicine prophylaxis for 14 days prior to and 14 days after randomization. Significantly more patients receiving febuxostat than placebo achieved an sUr level of ≤0.36 mmol/L at each visit ($P < 0.001$ for each comparison). The target sUr level (≤0.36 mmol/L) was achieved at study end in 0% of patients in the placebo group and 56%, 76% and 94% of patients in the 40 mg, 80 mg, and 120 mg febuxostat groups, respectively.




Figure 1 Chemical structure of febuxostat.

Subjects who completed this study were entered into a 5-year open-label extension study (FOCUS) and initially received febuxostat 80 mg daily.⁶⁸ Between weeks 4 and 24, dosing could be adjusted to febuxostat 40 or 120 mg. All subjects received gout flare prophylaxis during the first 4 weeks. Gout flares were recorded and treated throughout the study, and sUr, baseline tophi and safety were monitored. Among 116 subjects initially enrolled, dose adjustments were made for 44 (38%) subjects. As a result, 8 subjects received febuxostat 40 mg, 79 received 80 mg, and 29 received 120 mg daily maintenance dose. At 5 years, 93% (54/58) of the remaining subjects had sUr < 6.0 mg/dL (<0.36 mmol/L). Fifty-eight subjects (50%) discontinued prematurely; 38 did so in the first year. The primary reasons for discontinuation were: personal reasons 22 (19.0%), adverse event 13 (11.2%), gout flare 8 (6.9%), lost to follow-up 5 (4.3%), protocol violation 1 (<1%), other 9 (7.8%). Sustained reduction of sUr was associated with nearly complete elimination of gout flares. In 26 subjects with a tophus at baseline, resolution was achieved in 69% (18/26) by last visit on study drug at any point during the study. There were no deaths reported during the study. Long-term treatment with febuxostat resulted in durable maintenance of sUr < 6.0 mg/dL for most subjects. There was nearly complete abolition of gout flares in patients completing the study. Baseline tophi resolved in a majority of subjects.⁶⁸

In the phase III FACT trial, 762 patients with gout and with serum urate concentrations of at least 8.0 mg/dL

(0.48 mmol/L) were randomly assigned to receive either febuxostat (80 mg or 120 mg) or allopurinol (300 mg) once daily for 52 weeks; 760 patients received the study drug.⁶⁹ Prophylaxis against gout flares with naproxen or colchicine was provided during weeks 1 through 8. The primary endpoint was a target sUr < 6.0 mg/dL (0.36 mmol/L) at the last 3-monthly measurements. The secondary endpoints included reduction in the incidence of gout flares and in tophus area. The primary endpoint was reached in 47% to 59% of patients receiving 80 mg of febuxostat, 44% to 74% of those receiving 120 mg of febuxostat, and 8% to 40% of those receiving 300 mg allopurinol ($P < 0.001$ for comparison of each febuxostat group with the allopurinol group). The rates of discontinuation due to adverse events were higher in both the 80-mg febuxostat group and the 120-mg febuxostat group than in the allopurinol group (Table 3). Categories and frequencies of treatment-related adverse events were not linked to discontinuation, but only published for all reported treatment-related adverse events.

Febuxostat was concluded, at a daily dose of 80 mg or 120 mg, to be more effective than 300 mg allopurinol in lowering sUr.

Febuxostat reduced the median tophus area by 83% and 66% in patients in the 80-mg and 120-mg groups compared with 50% in patients receiving allopurinol. Similar reductions in gout flares occurred in all treatment groups: 64% and 70% in the 80-mg and 120-mg febuxostat groups, and 64% of patients receiving allopurinol. It was considered that an

Table 3 Efficacy and tolerability of febuxostat in randomized controlled trials

Study	Treatment, dosage	Number of patients	Duration	Mean decrease of serum urate	sUr < 6 mg/dl (0.36 mmol/L) final visit	sUr < 5 mg/dL (0.30 mmol/L) final visit	Withdrawal due to ADR
Becker ⁶⁷	Febuxostat 40 mg	37	28 days	37%	56%	21%	–
	Febuxostat 80 mg	40		44%	76%	49%	–
	Febuxostat 120 mg	38		59%	94%	88%	–
	Placebo	38		2%	0%	0%	–
FACT ⁶⁹	Febuxostat 80 mg	257	52 weeks	45%	81%	47%	6.2%
	Febuxostat 120 mg	251		52%	82%	66%	9.2%
	Allopurinol 300 mg	254		33%	39%	13%	3.1%
APEX ⁷²	Febuxostat 80 mg	267	28 weeks	48%	72%	–	6.7%
	Febuxostat 120 mg	269		55%	79%	–	5.9%
	Febuxostat 240 mg	134		68%	92%	–	8.2%
	Allopurinol 300 mg	268		34%	39%	–	6.7%
	Placebo	134		4%	1%	–	3.7%
CONFIRMS ⁷⁴	Febuxostat 40 mg	2269 in total	6 months	–	45%	–	–
	Febuxostat 80 mg			–	67%	–	–
	Allopurinol 200/300 mg			–	42%	–	–

8-week period of prophylaxis against gout flares due to urate mobilization probably was too short, as many gout attacks were noticed in the first weeks afterwards in all groups. This might be one of the reasons that no differences in reduction of gout flares were seen. Another point of discussion is the dose limit of allopurinol of 300 mg/day in this study, which is often considered as the “safe” maximum dosage.^{70,71} It is known that dosages up to allopurinol 600 mg/day are more effective, and the licensed maximum dosage is 800 mg/day (or 900 mg/day in some countries).⁴⁴

In the APEX trial, 1072 patients with gout, including persons with impaired renal function, and with sUr concentrations of at least 0.48 mmol/L were randomly assigned to receive either febuxostat (80 mg or 120 mg or 240 mg) versus allopurinol (300 mg or 100 mg) once daily for 28 weeks versus placebo.⁷² Significantly higher percentages of subjects treated with febuxostat 80 mg (48%), 120 mg (65%), and 240 mg (69%) attained the primary endpoint of at least 3 monthly sUr levels <0.36 mmol/L compared with allopurinol (22%) and placebo (0%). A significantly higher percentage of patients with impaired renal function treated with febuxostat 80 mg (4 out of 9 = 44%), 120 mg (5 out of 11 = 45%), and 240 mg (3 out of 5 = 60%) achieved the primary endpoint compared with those treated with 100 mg allopurinol (0 out of 10 = 0%). Serious adverse events occurred similarly in all groups, although diarrhea and dizziness were more frequent in the 240 mg febuxostat group. Primary reasons for withdrawal were similar across groups except for gout flares, which occurred more frequent with febuxostat than with allopurinol. Schumacher et al concluded that at all doses studied febuxostat more effectively lowered and maintained serum urate levels <0.36 mmol/L than did allopurinol or placebo in subjects with hyperuricemia and gout, including in those with mild to moderate impaired renal function.⁷²

Subjects who completed the FACT or APEX trial were invited to enroll in an open-label extension study and assigned to fixed-dose daily ULT with febuxostat (80 mg or 120 mg) or allopurinol (EXCEL).⁷³ The majority of subjects were male, Caucasian, and in the age range of 45 to 65 years. ULT reassignment was permitted during months 1 to 6 to achieve sUr concentrations of 3.0 to 6.0 mg/dL (0.18 to 0.36 mmol/L). Flares requiring treatment, tophus size, safety, and sUr levels were monitored during up to 40 months of ULT maintenance. After 1 month of initial treatment, >80% of subjects receiving either febuxostat dose, but only 46% of subjects receiving allopurinol, achieved sUr < 6.0 mg/dL (<0.36 mmol/L). After ULT reassignment, >80% of all

remaining subjects maintained the primary efficacy endpoint of sUr < 6.0 mg/dL at each visit. More subjects initially randomized to allopurinol required ULT reassignment to achieve sUr < 6.0 mg/dL compared with subjects receiving febuxostat. Maintenance of sUr < 6.0 mg/dL resulted in progressive reduction to nearly 0 in proportion of subjects requiring gout flare treatment. Baseline tophus resolution was achieved by 46%, 36%, and 29% of subjects maintained on febuxostat 80 mg, febuxostat 120 mg, and allopurinol, respectively. Overall adverse event rates (including cardiovascular adverse event rates), adjusted for 10-fold greater febuxostat than allopurinol exposure, did not differ significantly among treatment groups. Durable maintenance of goal range sUr level with either dose of febuxostat or in smaller numbers of subjects with allopurinol resulted in near elimination of gout flares and improved tophus status over time.

In October 2008 the most recent clinical trial of febuxostat was presented at the Annual Meeting of the American College of Rheumatology.⁷⁴ This 6-month phase III randomized, controlled, multicenter, double-blind trial comparing efficacy and safety of daily febuxostat and allopurinol in subjects with gout (CONFIRMS) trial randomized 2269 patients to receive febuxostat 40 mg/day, febuxostat 80 mg/day, or allopurinol 200 or 300 mg/day depending on renal function (patients with glomerular filtration rate of 30 to 59 mL/min received 200 mg/day of allopurinol). The primary endpoint was the proportion of patients achieving an sUr of less than 6 mg/dL (0.36 mmol/L) at the end of follow up. A secondary endpoint analyzed this same outcome in those patients with mild or moderate renal dysfunction defined as glomerular filtration rates of 60 to 89 and 30 to 59 mL/min, respectively. As in most of the previous febuxostat trials, a majority of the subjects were middle-aged, (mean age 52.8 years) white males. Baseline sUr was 9.6 mg/dL and gout duration on average was 11.6 years; 1483 individuals had mild or moderate renal dysfunction. Urate lowering efficacy was similar with allopurinol 300 mg/day (42% achieved primary endpoint) and febuxostat 40 mg/day (45% achieved primary endpoint) but febuxostat 80 mg/day was statistically superior to the other two groups (67% achieved primary endpoint). Among patients with renal dysfunction, febuxostat 80 mg/day achieved the primary endpoint in a higher proportion of patients (72%) than in those taking febuxostat 40 mg/day (50%) or allopurinol (42%). Rates of adverse events were comparable across groups and at all levels of renal function. Specifically, no difference was reported in the rate of cardiovascular events (6 events were reported: 3 in the allopurinol arm and 3 in the febuxostat 80 mg/day group). Five deaths

occurred during the study, 1 in each febuxostat group and 3 in the allopurinol arm. CONFIRMS also provided additional evidence than that provided by the FACT and APEX trials supporting a stronger urate-lowering potency in the tested doses of febuxostat compared with allopurinol for those individuals with sUr greater than 10 mg/dL or with tophi.^{74,75}

Efficacy and tolerability results of febuxostat in randomized controlled trials and long-term follow-up studies are summarized in Tables 3 and 4.

Febuxostat tolerability and safety

In terms of safety, to date, results from clinical trials have shown that febuxostat is well tolerated with a safety profile comparable to that of placebo and allopurinol (Tables 3 and 4). The most commonly reported adverse drug reactions (investigator assessment) are liver function abnormalities (3.5%), diarrhea (2.7%), headache (1.8%), nausea (1.7%), and rash (1.5%).⁷⁶ In both the double-blind and extension phases, liver function test abnormalities were associated with colchicine administration.⁶² Diarrhea, nausea, and vomiting are more frequent in patients concomitantly treated with colchicine.⁷⁶ Some serious rashes were reported with febuxostat in the APEX study.⁷¹ Since allopurinol can cause life-threatening cutaneous reactions (Stevens-Johnson syndrome) in rare cases, special attention to this serious adverse event is necessary.⁷⁶

Some concerns about the safety of febuxostat have been expressed. Initial cardiovascular safety signals raised in the FACT trial and in the FDA clinical review of July 2006 have not been replicated in recently released phase III trial results. Postmarketing surveillance will be necessary to monitor rare but serious adverse events.

Febuxostat clinical utility in the elderly

Based upon currently available data and expert opinion, treatment with febuxostat should be considered indicated for patients with gout and associated hyperuricemia in several circumstances: allopurinol intolerance due to allergy or

gastrointestinal complaints or biochemical side effects in blood counts or liver function tests. Currently, data are lacking about cross-allergy. Febuxostat may also be considered in mild to moderate renal impairment, ie, creatinine clearance 20 to 60 mL/min, as well as in mild to moderate non-alcoholic hepatic impairment, ie, with ALAT < twice upper limit of normal.

Before prescribing febuxostat, one should check documentation of allopurinol intolerance/ineffectivity/contraindication. Tests should be performed regarding full blood count, liver enzymes (ASAT, ALAT), albumen, thyroid function (TSH), renal function (creatinine), plasma uric acid and in 24-hour urine sample: uric acid and creatinine clearance. Prophylaxis should be given using colchicine in appropriate dosage or alternatively an NSAID at least 3 days before starting febuxostat and prophylaxis should be continued until the target plasma uric acid has been reached (usually 3–6 months). When renal function is impaired (<50 mL/min) febuxostat should be started at a lower dose: 40 mg daily. During therapy, patients should be monitored according to national guidelines, eg, Table 5.

Conclusions

Reviewing the data, febuxostat can be regarded as a useful, novel, non-purine, selective and potent inhibitor of xanthine oxidase. Pharmacokinetics of febuxostat 10 to 120 mg daily are linear. No dose adjustment appears to be necessary in patients with mild-to-moderate renal impairment or mild-to-moderate hepatic impairment. Febuxostat 10 to 120 mg daily rapidly and sustainably reduces serum uric acid by 25% to 70%. Prophylaxis with colchicine or a NSAID is indicated preceding and during 6 months after initiating febuxostat. Febuxostat in clinical trials is well tolerated including those patients experiencing hypersensitivity/intolerance to allopurinol.

Its antihyperuricemic efficacy at 80 to 120 mg/day is better than “standard dosage” allopurinol (300 mg/day). Febuxostat 80 mg daily appears to be equipotent to allopurinol 450 mg/day or benzbromarone 100 mg/day. Long-term safety and efficacy data on tophus diminishment and reduction of

Table 4 Efficacy and tolerability of febuxostat in long-term clinical follow-up

Study	N	Duration	sUr < 6 mg/dL (<0.36 mmol/L)	Incidence of recurrent gout flares (%/year)	Complete resolution of tophi	Withdrawal due to therapeutic failure
FOCUS ⁶⁸	116	5 years	93%	<6%	69%	–
EXCEL ⁷³	949	3 years	>80%	<4%	38%–44%	4.8%

Table 5 Suggestions for monitoring febuxostat therapy

1. After 4 weeks: CBC; serum creatinine; serum urate; ALAT; ASAT; 24-hour urate and creatinine clearance
2. After 12 weeks and every 8 weeks thereafter: CBC; serum creatinine; serum urate; ALAT; ASAT; 24-hour urate and creatinine clearance
3. After 6 months and every 6 months thereafter: TSH
4. Urate target according to BSR: <0.30 mmol/L is preferred, if not attainable then target according to EULAR: <0.36 mmol/L or any target at which the patient is free of gout flares
5. If sUr target free of flares cannot be reached by administering febuxostat monotherapy, combination therapy may be considered with a uricosuric
6. In case of ineffectivity (defined as not reaching the predefined target) of febuxostat, it should be stopped

Abbreviations: ALT, alanine aminotransferase; AST, aspartate aminotransferase; BSR, British Society for Rheumatology; CBC, complete blood count; EULAR, European League Against Rheumatology; sUr, serum urate; TSH, thyroid-stimulating hormone.

gout flares have recently become available. Efficacy and tolerability of febuxostat compared to a uricosuric drug is to be determined in future studies.

Elderly onset gout has distinct clinical features in comparison to gout in middle-aged men, and co-medication and co-morbidity should be carefully evaluated. When preventive treatment is considered in this population, the potential benefits should outweigh the risks.

Antihyperuricemic treatment success strongly depends on strategic use of the available drugs. This includes sufficient prophylaxis of gout flares during start of antihyperuricemic treatment, choice of drug tailored to patient characteristics, and dosage titration to target sUr levels. Febuxostat broadens the therapeutic options for antihyperuricemic in patients with gout. This is of clinical importance, especially for gout patients with currently unmet needs.

Disclosures

The authors have no conflict of interest to declare.

References

1. Singh H, Torralba KD. Therapeutic challenges in the management of gout in the elderly. *Geriatrics*. 2008;63(7):13–18, 20.
2. Saag KG, Choi H. Epidemiology, risk factors, and lifestyle modifications for gout. *Arthritis Res Ther*. 2006;8 Suppl 1:S2.
3. Pascual E, Sivera F. Why is gout so poorly managed? *Ann Rheum Dis*. 2007;66(10):1269–1270.
4. Zhang W, Doherty M, Pascual E, et al; for EULAR Standing Committee for International Clinical Studies Including Therapeutics. EULAR evidence based recommendations for gout – Part I Diagnosis: Report of a task force of the EULAR Standing Committee for International Clinical Studies Including Therapeutics (ESCSIT). *Ann Rheum Dis*. 2006;65(10):1301–1311.

5. Zhang W, Doherty M, Bardin T, et al; for EULAR Standing Committee for International Clinical Studies Including Therapeutics. EULAR evidence based recommendations for gout – Part II Management: Report of a task force of the EULAR Standing Committee for International Clinical Studies Including Therapeutics (ESCSIT). *Ann Rheum Dis*. 2006;65(10):1312–1324.
6. Arrondee E, Michet CJ, Crowson CS, O'Fallon WM, Gabriel SE. Epidemiology of gout: is the incidence rising? *J Rheumatol*. 29(11):2403–2406.
7. Wallace K, Riedel A, Joseph-Ridge N, Wortmann R. Increasing prevalence of gout and hyperuricemia over 10 years among older adults in a managed care population. *J Rheumatol*. 2004;31(8):1582–1587.
8. Fam AG. Gout in the elderly. Clinical presentation and treatment. *Drugs Aging*. 1998;13(3):229–243.
9. Ene-Stroescu D, Gorbien MJ. Gouty arthritis. A primer on late-onset gout. *Geriatrics*. 2005;60(7):24–31.
10. Hoskison KT, Wortmann RL. Management of gout in older adults: barriers to optimal control. *Drugs Aging*. 2007;24(1):21–36.
11. De Leonadis F, Govoni M, Colina M, Bruschi M, Trotta F. Elderly-onset gout: a review. *Rheumatol Int*. 2007;28(1):1–6.
12. Choi HK, Mount DB, Reginato AM. Pathogenesis of gout. *Ann Intern Med*. 2005;143(7):499–516.
13. Enomoto A, Kimura H, Chairoungdua A, Shigeta Y, Jutabha P, Cha SH, et al. Molecular identification of a renal urate anion exchanger that regulates blood urate levels. *Nature*. 2002;417(6887):447–452.
14. Kolz M, Johnson T, Sanna S, et al. Meta-analysis of 28,141 individuals identifies common variants within five new loci that influence uric acid concentrations. *PLoS Genet*. 2009;5(6):e1000504.
15. Spieker LE, Ruschitzka FT, Lüscher TF, Noll G. The management of hyperuricemia and gout in patients with heart failure. *Eur J Heart Fail*. 2002;4(4):403–410.
16. Bagatin J, Sardelic S, Pivac N, et al. Comparison of chlorthalidone, propranolol and bopindolol in six-month treatment of arterial hypertension. *Int J Clin Pharmacol Res*. 1998;18:73–78.
17. Gurwitz JH, Kalish SC, Bohn RL, et al. Thiazide diuretics and the initiation of anti-gout therapy. *J Clin Epidemiol*. 1997;50:953–959.
18. Dunn CJ, Fitton A, Brogden RN. Torasemide. An update of its pharmacological properties and therapeutic efficacy. *Drugs*. 1995;49:121–142.
19. Caspi D, Lubart E, Graff E, Habot B, Yaron M, Segal R. The effect of mini-dose aspirin on renal function and uric acid handling in elderly patients. *Arthritis Rheum*. 2000;43(1):103–108.
20. Dieppe PA. Investigation and management of gout in the young and the elderly. *Ann Rheum Dis*. 1991;50(4):263–266.
21. Pascual E, Tovar J, Ruiz MT. The ordinary light microscope: an appropriate tool for provisional detection and identification of crystals in synovial fluid. *Ann Rheum Dis*. 1989;48(12):983–985.
22. Kanellis J, Kang D-H, Feig DI, et al. Asymptomatic hyperuricemia. In: Wortmann RL, Becker MA, Ryan LM, editors. *Crystal-Induced Arthropathies: Gout, Pseudogout and Apatite-Associated Syndromes*. New York: Taylor and Francis Group; 2006:84–85.
23. Schlesinger N, Detry MA, Holland BK, et al. Local ice therapy during bouts of acute gouty arthritis. *J Rheumatol*. 2004;29(2):331–334.
24. Janssens HJ, Janssen M, van de Lisdonk EH, van Riel PL, van Weel C. Use of oral prednisolone or naproxen for the treatment of gout arthritis: a double-blind, randomised equivalence trial. *Lancet*. 2008;371(9627):1854–1860.
25. Rubin BR, Burton R, Navarra S, Smugar SS, Tershakovec AM. Efficacy and safety profile of treatment with etoricoxib 120 mg once daily compared with indomethacin 50 mg three times daily in acute gout: a randomized controlled trial. *Arthritis Rheum*. 2004;50(7):598–606.
26. Schumacher HR Jr, Boice JA, Daikh DI, et al. Randomised double blind trial of etoricoxib and indometacin in treatment of acute gouty arthritis. *BMJ*. 2002;324(7352):1488–1492.
27. Elisaf M, Tsimichodimos V, Bairaktari E, Siamopoulos KC. Effect of micronized fenofibrate and losartan combination on uric acid metabolism in hypertensive patients with hyperuricemia. *J Cardiovasc Pharmacol*. 1999;34(1):60–63.

28. Ka T, Inokuchi T, Tsutsumi Z, Takahashi S, Moriwaki Y, Yamamoto T. Effects of a fenofibrate/losartan combination on the plasma concentration and urinary excretion of purine bases. *Int J Clin Pharmacol Ther.* 2006;44(1):22–26.
29. Gao X, Curhan G, Forman JP, Ascherio A, Choi HK. Vitamin C intake and serum uric acid concentration in men. *J Rheumatol.* 2008;35(9):1853–1858.
30. Choi HK, Gao X, Curhan G. Vitamin C intake and the risk of gout in men: a prospective study. *Arch Intern Med.* 2009;169(5):502–507.
31. Levinson W, Cohen MS, Brady D, et al. To change or not to change: “Sounds like you have a dilemma”. *Ann Intern Med.* 2001;135(5):386–391.
32. Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G. Purine-rich foods, dairy and protein intake, and the risk of gout in men. *N Engl J Med.* 2004;350(11):1093–1103.
33. Ferraz MB, O’Brien B. A cost effectiveness analysis of urate lowering drugs in nontophaceous recurrent gouty arthritis. *J Rheumatol.* 1995;22(5):908–914.
34. Pérez-Ruiz F, Calabozo M, Pijoan JI, Herrero-Beites AM, Ruibal A. Effect of urate-lowering therapy on the velocity of size reduction of tophi in chronic gout. *Arthritis Rheum.* 2002(4);47:356–360.
35. Sarawate CA, Patel PA, Schumacher HR, et al. Serum urate levels and gout flares: analysis from managed care data. *J Clin Rheumatol.* 2006;12(2):61–65.
36. Shoji A, Yamanaka H, Kamatani N. A retrospective study of the relationship between serum urate level and recurrent attacks of gouty arthritis: evidence for reduction of recurrent gouty arthritis with antihyperuricemic therapy. *Arthritis Rheum.* 2004;51(3):321–325.
37. Yamanaka H, Togashi R, Hakoda M, et al. Optimal range of serum urate concentrations to minimize risk of gouty attacks during anti-hyperuricemic treatment. *Adv Exp Med Biol.* 1998;431:13–18.
38. Jordan KM, Cameron JS, Snaith M, et al; for British Society for Rheumatology and British Health Professionals in Rheumatology Standards, Guidelines and Audit Working Group (SGAWG). British society for Rheumatology and British Health Professionals in Rheumatology guideline for the management of gout. *Rheumatology (Oxford).* 2008;46(8):1372–1374.
39. Sutaria S, Katbamna R, Underwood M. Effectiveness of interventions for the treatment of acute and prevention of recurrent gout—a systematic review. *Rheumatology (Oxford).* 2006;45(11):1422–1431.
40. Pérez-Ruiz F, Alonso-Ruiz A, Calabozo M, Herrero-Beites A, García-Erauskin G, Ruiz-Lucea E. Efficacy of allopurinol and benzbromarone for the control of hyperuricaemia. A pathogenic approach to the treatment of primary chronic gout. *Ann Rheum Dis.* 1998;57(9):545–549.
41. Becker MA, Schumacher HR Jr, Wortmann RL, et al. Febuxostat compared with allopurinol in patients with hyperuricemia and gout. *N Engl J Med.* 2005;353(23):2450–2461.
42. Reinders MK, van Roon EN, Houtman PM, Brouwers JR, Jansen TL. Biochemical effectiveness of allopurinol and allopurinol-probenecid in previously benzbromarone-treated gout patients. *Clin Rheumatol.* 2007;26(5):1459–1465.
43. Reinders MK, Van Roon EN, Jansen TLThA, et al. Efficacy and tolerability of urate lowering drugs in gout: a randomised controlled trial of benzbromarone versus probenecid after failure of allopurinol. *Ann Rheum Dis.* 2009;68(1):51–56.
44. Reinders MK, Haagsma C, Jansen TLTA, et al. A randomized controlled trial with dose escalation on the efficacy and tolerability of allopurinol 300–600 mg/day versus benzbromarone 100–200 mg/day in gout patients. *Ann Rheum Dis.* 2009;68(6):892–897.
45. Dalbeth N, Kumar S, Stamp L, Gow P. Dose adjustment of allopurinol according to creatinine clearance does not provide adequate control of hyperuricemia in patients with gout. *J Rheumatol.* 2006;33(8):1646–1650.
46. Chung Y, Stocker SL, Graham GG, Day RO. Optimizing therapy with allopurinol: factors limiting hypouricemic efficacy. *Am J Med Sci.* 2008;335(3):219–226.
47. Panomvana D, Sripradit S, Angtharak S. Higher therapeutic plasma oxypurinol concentrations might be required for gouty patients with chronic kidney disease. *J Clin Rheumatol.* 2008;14(1):6–11.
48. Borstad GC, Bryant LR, Abel MP, Scroggie DA, Harris MD, Alloway JA. Colchicine for prophylaxis of acute flares when initiating allopurinol for chronic gouty arthritis. *J Rheumatol.* 2004;31(12):2429–2432.
49. Wortmann RL. Recent advances in the management of gout and hyperuricemia. *Current Opin Rheumatol.* 2005;17(3):319–324.
50. Arellano F, Sacristán JA. Allopurinol hypersensitivity syndrome: a review. *Ann Pharmacother.* 1993;27(3):337–343.
51. Hande KR, Noone RM, Stone WJ. Severe allopurinol toxicity. Description and guidelines for prevention in patients with renal insufficiency. *Am J Med.* 1984;76(1):47–56.
52. CHMP 2008. Committee for medicinal products for human use summary of positive opinion for adenuric. 21 February 2008. URL: http://www.emea.europa.eu/pdfs/human/opinion/Adenuric_8075108en.pdf.
53. CDER 2009. Center for Drug Evaluation and Research. URL: <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm149534.htm>.
54. Jansen TL, Reinders MK, Van Roon EN, Brouwers JRBJ. Benzbromarone withdrawn from the European market: another case of “absence of evidence is evidence of absence?” *Clin Exp Rheumatol.* 2004;22(5):651.
55. Lee MH, Graham GG, Williams KM, Day RO. A benefit-risk assessment of benzbromarone in the treatment of gout. Was its withdrawal from the market in the best interest of patients? *Drug Saf.* 2008;31(8):643–665.
56. Moolenburgh JD, Reinders MK, Jansen TL. Rasburicase treatment in severe tophaceous gout: a novel therapeutic option. *Clin Rheumatol.* 2006;25(5):749–752.
57. Vogt B. Urate oxidase (rasburicase) for treatment of severe tophaceous gout. *Nephrol Dial Transplant.* 2005;20(2):431–433.
58. Becker MA, Kisicki J, Khosravan R, et al. Febuxostat (TMX-67), a novel, non-purine, selective inhibitor of xanthine oxidase, is safe and decreases serum urate in healthy volunteers. *Nucleosides Nucleotides Nucleic Acids.* 2004;23(8–9):1111–1116.
59. Khosravan R, Kukulka MJ, Wu JT, Joseph-Ridge N, Vernillet L. The effect of age and gender on pharmacokinetics, pharmacodynamics, and safety of febuxostat, a novel nonpurine selective inhibitor of xanthine oxidase. *J Clin Pharmacol.* 2008;48(9):1014–1024.
60. Khosravan R, Grabowski B, Wu JT, Joseph-Ridge N, Vernillet L. Effect of food or antacid on pharmacokinetics and pharmacodynamics of febuxostat in healthy subjects. *Br J Clin Pharmacol.* 2008;65(3):355–363.
61. Schumacher HR Jr. Febuxostat: a non-purine, selective inhibitor of xanthine oxidase for the management of hyperuricaemia in patients with gout. *Expert Opin Investig Drugs.* 2005;14(7):893–903.
62. Hair PI, McCormack PL, Keating GM. Febuxostat. *Drugs.* 2008;68(13):1865–1874.
63. Bruce SP. Febuxostat: a selective xanthine oxidase inhibitor for the treatment of hyperuricemia and gout. *Ann Pharmacother.* 2006;40(12):2187–2194.
64. Mayer MD, Khosravan R, Vernillet L, Wu JT, Joseph-Ridge N, Mulford DJ. Pharmacokinetics and pharmacodynamics of febuxostat, a new non-purine selective inhibitor of xanthine oxidase in subjects with renal impairment. *Am J Ther.* 2005;12(1):22–34.
65. Hoshida S, Takahashi Y, Ishikawa T, et al. PK/PD and safety of a single dose of TMX-67 (febuxostat) in subjects with mild and moderate renal impairment. *Nucleosides Nucleotides Nucleic Acids.* 2004;23(8–9):1117–1118.
66. Khosravan R, Grabowski BA, Mayer MD, Wu JT, Joseph-Ridge N, Vernillet L. The effect of mild and moderate hepatic impairment on pharmacokinetics, pharmacodynamics, and safety of febuxostat, a novel nonpurine selective inhibitor of xanthine oxidase. *J Clin Pharmacol.* 2006;46(1):88–102.

67. Becker MA, Schumacher HR Jr, Wortmann RL, et al. Febuxostat, a novel nonpurine selective inhibitor of xanthine oxidase: a twenty-eight-day, multicenter, phase II, randomized, double-blind, placebo-controlled, dose-response clinical trial examining safety and efficacy in patients with gout. *Arthritis Rheum.* 2005;52(3):916–923.
68. Schumacher HR, Becker MA, Lloyd E, MacDonald PA, Lademacher C. Febuxostat in the treatment of gout: 5-yr findings of the FOCUS efficacy and safety study. *Rheumatology (Oxford).* 2009;48(2):188–194.
69. Becker MA, Schumacher HR Jr, Wortmann RL, et al. Febuxostat compared with allopurinol in patients with hyperuricemia and gout. *N Engl J Med.* 2005;353(23):2450–2461.
70. Moreland LW. Febuxostat – treatment for hyperuricemia and gout? *N Engl J Med.* 2005;353(23):2505–2507.
71. Rundles RW, Metz EN, Silberman HR. Allopurinol in the treatment of gout. *Ann Intern Med.* 1966;64(2):229–258.
72. Schumacher HR Jr, Becker MA, Wortmann RL, et al. Effects of febuxostat versus allopurinol and placebo in reducing serum urate in subjects with hyperuricemia and gout: a 28-week, phase III, randomized, double-blind, parallel-group trial. *Arthritis Rheum.* 2008;59(11):1540–1548.
73. Becker MA, Schumacher HR, MacDonald PA, Lloyd E, Lademacher C. Clinical efficacy and safety of successful longterm urate lowering with febuxostat or allopurinol in subjects with gout. *J Rheumatol.* 2009;36(6):1273–1282.
74. Becker M, Schumacher HR Jr, Espinoza L, et al. A phase 3 randomized, controlled, multicenter, double-blind trial comparing efficacy and safety of daily febuxostat and allopurinol in subjects with gout. Presented at: the Meeting of the American College of Rheumatology; October 24–29, 2008; San Francisco, California. URL: <http://acr.confex.com/acr/2008/webprogram/Paper3414.html>.
75. Gaffo AL, Saag KG. Febuxostat: the evidence for its use in the treatment of hyperuricemia and gout. *Core Evidence.* 2009;4:25–36.
76. European Medicines Agency (EMA). Summary of product characteristics. 19/05/2009 Adenuric-H-C-777-T-13. URL: <http://www.emea.europa.eu/humandocs/PDFs/EPAR/adenuric/emea-combined-h777en.pdf>.

Clinical Interventions in Aging

Dovepress

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, the American Chemical Society's 'Chemical

Abstracts Service' (CAS), Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>