Association of *XRCC1*, *XRCC2* and *XRCC3* Gene Polymorphism with Esophageal Cancer Risk

This article was published in the following Dove Press journal:
Clinical and Experimental Gastroenterology

Jagjeet Kaur¹
Vasudha Sambyal¹
Kamlesh Guleria¹
Neeti Rajan Singh²
Manjit Singh Uppal²
Mridu Manjari³
Meena Sudan⁴

¹Human Cytogenetics Laboratory, Department of Human Genetics, Guru Nanak Dev University, Amritsar, India; ²Department of Surgery, ³Department of Pathology, ⁴Department of Radiotherapy, Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar, Punjab, India

Aim: The X-ray repair cross-complementing (*XRCC*) gene polymorphisms influence esophageal carcinogenesis by altering the DNA repair capacity. The present study was designed to screen five single nucleotide polymorphisms (SNPs) of *XRCC* genes for their susceptibility to esophageal cancer (EC) risk. There is no previous report on these polymorphisms for EC from India, where EC frequency is high.

Methods: The present study included 497 subjects (213 EC patients and 284 healthy controls). The polymorphisms were screened using the PCR-RFLP method and allele and genotype distribution were compared using chi-square test. Association analysis was done by haplotype analysis and linkage disequilibrium (LD) analysis. Gene-gene interactions were identified using multifactor dimensionality reduction (MDR). The risk was calculated using binary logistic regression.

Results: For *XRCC1* p.Arg399Gln, a decreased risk for EC was associated with the AA genotype [OR (95% CI): 0.53 (0.3–0.95), $p=0.03$] even after adjusting for various covariates [OR (95% CI): 0.49 (0.26–0.9), $p=0.024$] and with the recessive model [OR (95% CI): 0.49 (0.27–0.8), $p=0.016$]. The GA genotype of p.Arg280His was associated with an increased risk for EC [OR (95% CI): 1.7 (1.0–2.82), $p=0.045$] after adjustments. The two *XRCC1* polymorphisms, p.Arg399Gln and p.Arg194Trp were in slight LD among EC patients ($D=0.845$, $r^2=0.042$). *XRCC2* and *XRCC3* polymorphisms were not associated with EC risk.

Conclusion: *XRCC1* p.Arg399Gln plays a protective role in the development of the EC. The study is the first report from India, providing baseline data about genetic polymorphisms in DNA repair genes *XRCC1*, *XRCC2* and *XRCC3* modulating overall EC risk.

Keywords: *XRCC1*, *XRCC2*, *XRCC3*, polymorphisms, esophageal cancer

Introduction

Esophageal cancer (EC) ranks as the sixth leading cause of death from cancer constituting 7% of all gastrointestinal cancers.¹ In India, EC is the fourth most common cause of deaths related to cancer and the second most common cancer among men and fourth among women.² Esophageal carcinogenesis is a complex multistep process with poor prognosis where environmental, geographic and genetic factors appear to play a major role.³ Cancer is a disease characterized by the failure of DNA repair mechanisms. *XRCC1* protein as a part of Base excision repair (BER) pathway plays an efficient role in repairing DNA single-strand breaks.⁴ It is encoded by the *XRCC1* gene located on 19q13.2, comprising of 17 exons and 16 introns.⁵ Two *XRCC1* polymorphisms, p.Arg194Trp (exon 6) and p.Arg280His (exon 9) affect the function of the protein.⁶ *XRCC1* p.Arg399Gln polymorphism in exon 10⁷ has been associated with breast,⁸ lung⁹ and head and neck cancers.¹⁰

Correspondence: Vasudha Sambyal
Department of Human Genetics, Guru Nanak Dev University, Amritsar 143001, Punjab, India
Tel +911832258803-09 ext 3448
Email vasudhasambyal@yahoo.co.in

XRCC2 (chromosome location 7q36.1) plays a major role in Homologous recombination repair (HRR) pathway.¹¹ *XRCC2* p.Arg188His polymorphism located in exon3¹² has been associated with cancers like pancreatic,¹³ ovarian,¹⁴ oral¹⁵ and upper aerodigestive tract cancers.¹⁶

XRCC3 (chromosome location 14q32.3) encodes a 346 amino acid polypeptide¹⁷ that participates in DNA double-strand break repair. Variation in expression of *XRCC3* has been reported in various cancers, like gastric, breast, lung, skin and colorectal.^{18–21} The most common polymorphism in *XRCC3* p.Thr241Met in exon 7 can influence the ability to repair DNA.²¹

Allelic variants of *XRCC1*, *XRCC2*, and *XRCC3* are associated with risk of different types of cancer among different populations all over the world.^{22–24} In the world's second-most populous country India, the large and diverse ethnic subpopulations²⁵ need to be studied to understand the genetic basis of several region-specific complex disorders. To the best of our knowledge, studies related to these DNA-repair genes are very few for esophageal cancer in India (Table 1). The state of Punjab, India, has very high age-adjusted rate (AAR) for EC; it ranks second in males after prostate cancer and fourth after breast, ovarian and cervical cancer in females in this region.^{26,27}

Therefore, the present study was carried out to explore the role of five polymorphisms of *XRCC* genes; *XRCC1* (p.Arg399Gln, p.Arg194Trp, p.Arg280His), *XRCC2* (p.Arg188His) and *XRCC3* (p.Thr241Met) in the risk of EC in the population of Punjab, India.

Materials and Methods

Study Population and Sample Collection

The present case-control study was conducted after due ethical clearance according to the tenets of the Declaration of Helsinki by the institutional ethics committee of Guru Nanak Dev University, Amritsar. The study sample included 497 subjects: 213 (125 females and 88 males) esophageal cancer patients without a history of any other cancer, any treatment before sample collection and pre-operative from Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar. Complete information including age, gender, lifestyle, diet, family history, and environmental factors were recorded on a proforma after written informed consent from all the subjects. Two hundred eighty-four (187 females and 97 males), age and gender-matched unrelated healthy controls from the same geographical region were also included in the study (Table 2).

Collection of Blood samples and genotyping: Intravenous blood sample was collected in EDTA from each subject. Genomic DNA from blood was collected using standard Phenol chloroform method.²⁸ Previously published primer sequences for *XRCC1* p.Arg399Gln,²⁹ *XRCC1* p.Arg194Trp,²⁴ *XRCC1* p.Arg280His,²² *XRCC2* p.Arg188His³⁰ and *XRCC3* p.Thr241Met²³ polymorphisms were used to amplify the target region. The total volume of the PCR reaction mixture was 15µL containing the reagents: 50ng of template DNA, 10pmol of each primer (Sigma, St. Louis, MO, USA), 10X PCR buffer with 1.5 mM MgCl₂, 50 µM of dNTPs mix and 0.6 units of Taq polymerase (Bangalore GeNei).

The PCR products were digested with appropriate restriction endonucleases using manufacturer recommendations (NEB, UK) and analyzed on ethidium bromide stained agarose gel to obtain genotypes based on the size of fragment obtained (Table 3). To reassure of the quality of the result 10% of the samples were re-genotyped.

Statistical Analysis

Statistical analysis was done using SPSS software version 20 (SPSS, Chicago, IL, USA). The analysis of categorical variables was done using the chi-square test and the continuous independent variables were expressed as mean ± standard deviation using the Student's *t*-test. Hardy-Weinberg equilibrium was tested using the chi-square test. Binary logistic regression analysis was done to measure the risk of EC with genotypes of selected polymorphisms and the values were adjusted for the possible confounding effect of age, gender, diet, smoking and alcohol consumption. Odds ratios (ORs), *p*-values and 95% confidence interval (CI) were calculated as the measure of the strength of association of the SNPs and esophageal cancer. Haplotypes and linkage disequilibrium patterns were studied using Haploview software. Furthermore, multifactor dimensionality reduction (MDR) factor analysis version 2 was used to calculate the effect of Gene-gene interactions on esophageal cancer.

Results

The detailed characteristics of 497 subjects (213 EC patients and 284 controls) are given in Table 2 and the allele and genotype frequencies are summarised in Table 4 and Table 5. All the studied subjects were residents of Punjab, India. The endogamous caste groups were Jatt Sikh, Khatri and Scheduled Caste. The subjects were ethnic Punjabi with Scythian and Caucasian racial admixture.³¹ In the present study 5 polymorphisms; *XRCC1* (p.Arg399Gln,

Table I Studies Showing Association of *XRCC1*, *XRCC2* and *XRCC3* (p.Thr241Met) Polymorphism in Different Cancers in Populations from Different Geographical Regions of India

Authors	Population of Region (State)	SNPs	Disease	Number of Subjects	Results
GIT Cancers Present Study	North India (Punjab)	<i>XRCC1</i> p.Arg399Gln, p.Arg194Trp and p.Arg280His <i>XRCC2</i> p.Arg188His <i>XRCC3</i> p.Thr241Met	Esophageal Cancer	497	AA genotype of <i>XRCC1</i> p.Arg399Gln associated with reduced ESCC risk
³⁴ *	North India (Punjab/ Chandigarh)	<i>XRCC1</i> p.Arg399Gln	Esophageal Squamous Cell Carcinoma	280	AA genotype associated with reduced ESCC risk
⁴³ *	North India (Uttar pradesh)	<i>XRCC1</i> p.Arg399Gln p.Arg194His	Gallbladder Cancer	377	GA and AA genotype of p.Arg399Gln associated with decreased risk. No association of p.Arg194Trp polymorphism
⁵⁰	North India (Kashmir)	<i>XRCC1</i> p.Arg399Gln	Colorectal Cancer	260	Association of GA genotype with increased cancer risk
⁴⁴ *	North India (Kashmir)	<i>XRCC1</i> p.Arg399Gln	Colorectal Cancer	266	Association of AA genotype with decreased risk
⁵²	Eastern India (West Bengal)	<i>XRCC1</i> - p.Arg399Gln, p. Arg194Trp and p.Arg280His	Gastric Cancer	152	GA and GG genotype of p.Arg399Gln associated with increased cancer risk No association of p.Arg194Trp and p.Arg280His polymorphism
⁷² *	South India	<i>XRCC1</i> p.Arg194Trp	Gastric Cancer	600	No association
Cancers of Other Sites	Eastern India	<i>XRCC1</i> p.Arg399Gln and p. Arg194Trp <i>XRCC3</i> p.Thr241Met	Oral submucous fibrosis (OSF)	188	No association of <i>XRCC1</i> p.Arg399Gln and p.Arg194trp polymorphism CT genotype of <i>XRCC3</i> p.Thr241Met associated with increased risk of OSF
⁴⁶ *	North India	<i>XRCC1</i> p.Arg194Trp, p. Arg280His and p.Arg399Gln	Hepatocellular Carcinoma	407	TT of p.Arg280His associated with increased risk. AA genotype of p.Arg399Gln associated with decreased risk
⁹⁰	North eastern India	<i>XRCC1</i> p.Arg194Trp <i>XRCC3</i> p.Thr241Met	Breast Cancer	998	Arg/Trp and Trp/Trp genotype of <i>XRCC1</i> p.Arg194Trp associated with an increased risk No association of <i>XRCC3</i> p.Thr241Met
¹⁰¹	Western India (Maharashtra)	<i>XRCC1</i> - p.Arg399Gln, p. Arg194His and p.Arg280His <i>XRCC2</i> - p.Arg188His <i>XRCC3</i> - p.Thr241Met	Breast Cancer	350	No association of <i>XRCC1</i> (p.Arg399Gln, p.Arg194His and p.Arg280His), <i>XRCC2</i> (p.Arg188His) and <i>XRCC3</i> (p.Thr241Met) with breast cancer
⁵⁵	South India	<i>XRCC1</i> p.Arg194Trp, p. Arg280His and p.Arg399Gln	Cervical Cancer	250	No association
⁵¹	North India	<i>XRCC1</i> p.Arg399Gln	Prostate Cancer	450	Association of AA genotype with increased risk

(Continued)

Table 1 (Continued).

Authors	Population of Region (State)	SNPs	Disease	Number of Subjects	Results
⁵⁷	South India	<i>XRCC1</i> p.Arg194Trp, p.Arg280His and p.Arg399Gln	Lymphoblastic Leukemia	234	Association of Arg/Gln and Gln/Gln genotype of p.Arg399Gln with increased risk. No association for p.Arg280His and p.Arg194Trp polymorphism
⁵⁸	South India	<i>XRCC1</i> p.Arg399Gln	Acute Myeloid Leukemia	422	Association of GA- genotype with increased risk of AML
¹⁰²	Middle India (Madhya Pradesh)	<i>XRCC1</i> p.Arg194Trp, p.Arg280His and p.Arg399Gln	Sickle Cell Anemia	500	GA and AA genotype of p.Arg280His associated with an increased risk. No association of <i>XRCC1</i> p.Arg194Trp and p.Arg399Gln polymorphisms
^{45*}	North India	<i>XRCC1</i> p.Arg194Trp, p.Arg280His and p.Arg399Gln	Head and Neck Cancer	556	AA genotype of p.Arg399Gln and TT genotype of p.Arg194Trp associated with reduced risk. No association of p.Arg280His polymorphism.

Note: *Studies with similar results as present study.

p.Arg280His, p.Arg194Trp), *XRCC2* (p.Arg188His) and *XRCC3* (p.Thr241Met) were studied for the association with risk of esophageal cancer (EC). For *XRCC1* p.Arg280His polymorphism, A allele was associated with an increased risk of EC (OR=1.60, 95% CI= 1.02–2.52, p=0.04). According to the Bivariate logistic regression for stratification, genotypes GA and GG+GA (OR=1.8, 95% CI =1.1–2.9, p=0.025) provide 2 fold increased risk of EC after adjusting the confounding effect of age, gender, diet pattern, smoking and alcohol consumption. For *XRCC1* p.Arg399Gln polymorphism, AA genotype (OR= 0.53, 95% CI=0.3–0.95, p=0.03) was associated with 50% decreased the risk of EC. This association persisted and became stronger (OR=0.5, 95% CI=0.3–0.9, p=0.024) after adjustments to age, sex, diet pattern, smoking and alcohol consumption. Smoking and Alcohol intake was reported only by male subjects. Smoking was not reported by many subjects probably because of prevalent socio-religious beliefs in the region prohibit smoking and tobacco intake and alcohol intake. No significant difference in allele (p=0.54) and genotype (p=0.52) frequencies of *XRCC1* p.Arg194Trp polymorphism was observed. No association with EC risk was observed for *XRCC2* p.Arg188His and *XRCC3* p.Thr241Met polymorphism (p=0.53) in the subjects.

Genetic model analysis of *XRCC1* p.Arg399Gln and *XRCC3* p.Thr241Met (Table 6) revealed a decreased risk of EC under the recessive model AA vs GG+GA (OR=0.55,

95% CI=0.32–0.95, p=0.027) for *XRCC1* p.Arg399Gln polymorphism which became more significant after adjustment with binomial logistic regression (OR=0.49, 95% CI=0.27–0.88, p=0.016). For p.Thr241Met polymorphism no genotype combination was associated with EC.

Haplotype and Linkage Disequilibrium Analysis

The haplotype frequencies of three polymorphisms of *XRCC1* were examined for their association with EC risk (Table 7). Based on the measures of linkage disequilibrium (LD), the two polymorphisms of *XRCC1* gene, p.Arg399Gln and p.Arg194Trp, were in slight LD among EC patients (D' =0.845, r^2 =0.042) (Figure 1A). However, control subjects were not in LD for any of the polymorphism (Figure 1B). The haplotype GGT (p.Arg399Gln, p.Arg280His, and p.Arg194Trp) was predominant in EC cases as compared to controls, but the difference was statistically non-significant (p=0.1). The possible genotype combinations of three studied polymorphisms of *XRCC1* were also examined (Table 8). The combinations comprising the AA genotype (p.Arg399Gln) occurred significantly more often in controls than patients, with AA-CC-CC combination associated with significantly decreased risk of EC (OR=0.5, 95% CI=0.29–0.91, p=0.020).

Table 2 Clinical Characteristics of the Esophageal Cancer Patients and the Controls

Characteristics	Cases (n=213)	Controls (n=284)	Odds Ratio (95% CI)	p-value
Age				
Below 55	94	144		0.15
55 or above	119	140		
Age(years)	55.7±12.4	54.4±11.1		0.22
Sex				
Females	125	187		0.07
Males	88	97		
Habitat				
Rural	178	241	1.14	0.70
Urban	35	43	(0.67–1.70)	
Diet				
Vegetarians	108	142	0.97	0.87
Non-vegetarians	105	142	(0.68–1.40)	
Smoking				
No	181	279	9.83	<0.0001
Yes	32	5	(3.7–25.72)	
Drinking				
No	153	233	1.79	0.007
Yes	60	51	(1.17–2.14)	
Family history				
No	192	284		
Yes	21	0		
Type of cancer				
SCC	187	–		
ADC	26	–		
Clinical Stage				
(I)	10	–		
(II)	67	–		
(III)	122	–		
IV	14	–		

Notes: p-value less than 0.05 taken as statistically significant (bold values).

Abbreviation: SCC, squamous cell carcinoma, ADC, adenocarcinoma.

Interaction Analysis (MDR)

Table 9 summarises the MDR analysis evaluating the results of all possible combinations of the studied polymorphisms. According to MDR analysis, the best MDR model included all the five studied polymorphisms *XRCC1* (p.Arg399Gln, p.Arg194Trp, p.Arg280His), *XRCC2* (p.Arg188His) and *XRCC3* (p.Thr241Met). The model had a testing balance accuracy of 0.4648 and a training balance accuracy of 0.5905 with a maximum cross-validation consistency of 10/10. However, the model was not statistically significant (p=0.9). Figure 2 depicts the interaction map of all genes, based on entropy measures between individual genes showing a redundancy effect.

Discussion

In India, the esophagus is a major cancer site (NCRP 2016) with a majority being Squamous Cell Carcinoma (SCC) and a comparatively less frequency of adenocarcinoma patients.^{32,33} Similarly, the predominant type of EC in our study was SCC (88%) with only 12% adenocarcinoma patients.

In India, the inter-individual differences in susceptibility to cancer due to the genetic polymorphisms in *XRCC1* previously found.³⁴ As DNA-repair gene polymorphisms play a very important role in carcinogenesis, we carried out this case-control study to evaluate whether *XRCC1* (p.Arg399Gln, p.Arg194Trp and p.Arg280His), *XRCC2* (p.Arg188His), and *XRCC3* (p.Thr241Met) gene polymorphisms modulate the risk of esophageal cancer. The varied results reported world over for these polymorphisms has been attributed to variations in DNA-repair gene polymorphisms amongst different ethnicities.^{35–39} The population of Punjab, North India are an Indo-Aryan ethnic group with Caucasian, Indoscythian racial admixture. The main caste groups prevalent in Punjab are Baniyas (merchant and money-lender), Brahmins (traditionally priestly but now involved in various occupations), Jatt Sikhs, which constitute the largest part of Sikh community (mainly agriculturist), and Scheduled caste (various occupations).⁴⁰ In the present study, about 60% of the subjects belonged to Jatt Sikh Caste group but the subgroup analysis based on castes was not done due to unequal distribution of the samples.

The *XRCC1*p.Arg399Gln polymorphism is involved in various protein–protein interactions⁴¹ and higher sister chromatid exchanges and DNA adducts.⁴² In the present study, we found that A allele and the AA genotype of *XRCC1*p.Arg399Gln polymorphism was associated with a decreased risk of esophageal cancer. Furthermore, a reduced risk was observed in the recessive model (GG +GAvsAA). When compared with previously reported studies from India, the results of our study were consistent with some studies, especially from North India but contrary to others (Table 1). Very few studies relating the *XRCC1* Arg399Gln polymorphism with esophageal cancer risk are available from which only one is from India. Among previous reports from North India on EC; a study of Chandigarh region³⁴ found association with a decreased risk of *XRCC1* p.Arg399Gln, another study from Uttar Pradesh⁴³ also reported Arg/Gln (p=0.03, OR= 0.62) and

Table 3 Details of Amplification and Digestion Conditions Used for Genotyping of *XRCC1*, *XRCC2* and *XRCC3* Polymorphisms

Polymorphism (Ref SNP)	Location	Nucleotide Change	Annealing Temperature	Amplicon Size (bp)	Restriction Enzyme	Size of Digested Fragments
<i>XRCC1</i> p.Arg194Trp (rs1799782)	Exon 6	C>T	59°C	491	<i>MspI</i>	CC(292,174) CT(313,292,174) TT(313,174)
p.Arg280His (rs25489)	Exon 9	G>A	66°C	861	<i>RsaI</i>	GG(597,201,63) GA(660,597,201,63) AA(660,201)
p.Arg399Gln (rs25487)	Exon 10	G>A	59°C	615	<i>MspI</i>	GG(374,241) GA(615,374,241) AA(615)
<i>XRCC2</i> p.Arg188His (rs3218536)	Exon 3	G>A	59°C	290	<i>HphI</i>	GG(290) GA(290,148,142) AA(148,142)
<i>XRCC3</i> p.Thr241Met (rs1625895)	Exon 7	C>T	62°C	552	<i>NlaIII</i>	CC(313,239) CT(313,239,208,10) TT (239,208,105)

Abbreviation: SNP, single nucleotide polymorphism.

Table 4 Allele and Genotype Frequencies of *XRCC1*, *XRCC2* and *XRCC3* Gene Polymorphisms and Their Association with Esophageal Cancer Risk

Gene	SNP	Allele	Allele Frequency		Odds Ratio (95% CI)	p value	Genotype	Cases n (%)	Controls n(%)	Chi Square (χ^2)	p value*
			Cases (n%)	Controls (n%)							
<i>XRCC1</i>	rs25489 p.Arg280His	G	382(89.7)	530(93.3)	Reference 1.60 (1.02–2.51)	0.04	GG GA AA	171(80.3)	246(86.6)	4.9	0.08
		A	44(10.3)	38(6.7)				40(18.8)	38(13.4)		
	rs25487 p.Arg399Gln	G	283(66.5)	344(60.6)	Reference 0.78 (0.6–1.0)	0.06	GG GA AA	92(43.2)	109(38.4)	5.51	0.06
		A	143(33.5)	224(39.4)				99(46.5)	126(44.4)		
<i>XRCC2</i>	rs3218536 p.Arg188His	G	388(91.1)	515(90.7)	Reference 0.97 (0.62–1.51)	0.89	GG GA AA	175(82)	232(81.7)	0.75	0.68
		A	38(8.9)	52(9.3)				38(18)	51(18)		
	rs1625895 p.Thr241Met	C	334(78.4)	449(79)	Reference 1.03 (0.76–1.43)	0.80	CC CT TT	129(60.6)	178(62.7)	0.59	0.74
		T	92(21.6)	119(21)				76(35.7)	93(32.7)		

Notes: Wild type genotype was taken as reference. *Distribution of overall genotypic frequencies between cases and controls done by χ^2 test (df, 2). P-value less than 0.05 taken as statistically significant (bold values).

Abbreviation: SNP, single nucleotide polymorphism.

Gln/Gln (p=0.003, OR=0.37) genotype to be associated with a decreased risk of gall bladder cancer; a study from Kashmir on colorectal cancer⁴⁴ also reported a protective

role of AA genotype. Similar results were reported in three other studies from North India on head and neck cancer,⁴⁵ hepatocellular carcinoma⁴⁶ and lung cancer.⁴⁷

Table 5 Distributions of *XRCC1*, *XRCC2* and *XRCC3* Polymorphisms Genotypes and Development of Esophageal Cancer

Gene	SNP	Genotype	Cases n(%)	Controls n(%)	Unadjusted OR(CI)(p)	AOR* (CI) (p value)	AOR** (CI) (p value)
<i>XRCC1</i>	rs25489 p.Arg280His	GG	171(80.3)	246(86.6)	Reference	Reference	Reference
		GA	40(18.8)	38(13.4)	1.5(0.9–2.4) p=0.09	1.6(1.0–2.6) p=0.06	1.7 (1.0–2.8) p= 0.04
		AA	2(0.9)	0(0)	NC	NC	NC
		GA+AA			1.6(0.9–2.5) p=0.06	1.6(1.0–2.6) p= 0.04	1.8(1.1–2.9) p= 0.02
	rs25487 p.Arg399Gln	GG	92(43.2)	109(38.4)	Reference	Reference	Reference
		GA	99(46.5)	126(44.4)	0.9(0.6–1.3) p=0.7	0.9(0.6–1.4) p=0.7	1.0(0.7–1.5) p=0.9
		AA	22(10.3)	49(17.2)	0.5(0.3–0.9) p= 0.03	0.5(0.3–0.9) p= 0.03	0.5(0.3–0.9)p= 0.024
		GA+AA			0.8(0.6–1.2) p=0.28	0.8(0.6–1.2) p=0.27	0.9(0.6–1.3) p=0.4
	rs1799782 p.Arg194Trp	CC	181(85)	247(87)	Reference	Reference	Reference
		CT	32(15)	37(13)	1.2(0.7–1.9) p=0.52	1.2(0.7–2.1) p=0.4	1.1(0.7–2.0) p= 0.5
		TT	0(0)	0	NC	NC	NC
<i>XRCC2</i>	rs3218536 p.Arg188His	GG	175(82)	232(81.7)	Reference	Reference	Reference
		GA	38(18)	51(18)	0.9(0.6–1.6) p=0.95	1.0(0.6–1.6) p=0.9	0.9(0.6–1.5) p=0.8
		AA	0(0)	1(0.3)	NC	NC	NC
		GA+AA			0.9(0.6–1.5) p=0.89	0.99(0.6–1.6) p=0.9	0.9(0.5–1.5) p=0.7
<i>XRCC3</i>	rs1625895 p.Thr241Met	CC	129(60.6)	178(62.7)	Reference	Reference	Reference
		CT	76(35.7)	93(32.7)	1.1(0.7–1.6) p=0.53	1.1(0.7–1.6) p=0.6	1.1(0.7–1.6) p=0.6
		TT	8(3.7)	13(4.6)	0.8(0.3–2.1) p=0.72	0.8(0.3–2.2) p=0.8	0.8(0.3–2.3) p=0.8
		CT+TT			1.09(0.7–1.6) p=0.63	1.06(0.7–1.5) p=0.75	1.06(0.7–1.6) p=0.77

Notes: Wild type genotype is taken as reference, p<0.05 taken as statistically significant (bold values). AOR* Adjusted Odds Ratio for age, gender and diet pattern using Binary logistic regression. AOR** Adjusted Odds Ratio for age, gender, diet pattern, cigarette smoking and alcohol consumption using Binary logistic regression analysis.

Table 6 Association Analysis of *XRCC1* p.Arg399Gln and *XRCC3*p.Thr241Met Polymorphisms

SNPs	Comparison	Unadjusted OR (CI) (p value)	AOR* (CI) (p value)	AOR** (CI) (p value)
<i>XRCC1</i> p.Arg399Gln	Dominant model (AA+GAvsGG)	0.82(0.57–1.22) p=0.280	0.82(0.56–1.18) p=0.27	0.92(0.59–1.31) p=0.46
	Codominant model(AAvsGA/GGvsGA)	0.78 (0.60 –1.01) p=0.062	1.16(0.71–1.64) p=0.56	1.23(0.82–1.82) p=0.29
	Recessive model (GG+GAvsAA)	0.55(0.32–0.95) p= 0.027	0.53(0.30–0.95) p= 0.02	0.49(0.27–0.83) p= 0.02
<i>XRCC3</i> p.Thr241Met	Dominant model (TT+CTvsCC)	1.09(0.76–1.58) p=0.63	1.06(0.73–1.5) p=0.75	1.06(0.71–1.63) p=0.77
	Codominant model(TTvsCT/TTvsCC)	1.04 (0.76 –1.42) p=0.80	1.14(0.75–1.62) p=0.62	1.13(0.74–1.65) p=0.61
	Recessive model (CC+CTvsTT)	0.81(0.33–1.92) p=0.65	0.84(0.34–2.08) p=0.69	0.82(0.33–2.05) p=0.64

Notes: Wild type genotype is taken as reference, p<0.05 taken as statistically significant (bold values). AOR* Adjusted Odds Ratio for age, gender and diet pattern using binary logistic regression. AOR** Adjusted Odds Ratio for age, gender, diet pattern, cigarette smoking and alcohol consumption using binary logistic regression.

Contrary to the results of the present study, some previous studies from North India have reported an increased risk with AA genotype of *XRCC1*p.Arg399Gln polymorphism in lung cancer,⁴⁸ head and neck cancer,⁴⁹ colorectal cancer⁵⁰ and prostate cancer.⁵¹ Similarly, some studies from South India reported an association of increased risk with the AA genotype in gastric cancer,⁵² colorectal cancer,²⁹ lung cancer,^{53,54} cervical cancer,⁵⁵ Naso-pharyngeal cancer,⁵⁶ acute lymphoblastic leukemia⁵⁷ and acute myeloid leukemia.⁵⁸ However, few studies from South India reported no association of *XRCC1*p.Arg399Gln polymorphism with any of cancer (Table 1).

Results of the present study on *XRCC1* p.Arg399Gln polymorphism is in agreement with the studies from different parts of the globe like esophageal cancer in Han Chinese,⁵⁹ colorectal adenocarcinoma in Norwegian population,⁶⁰ gall-bladder cancer,⁶¹ and non-melanoma skin cancers.⁶² In contrast, some previous studies have reported the association of Arg399Gln polymorphism with an increased risk of esophageal,⁶³ stomach and oral cancers,⁶⁴ colorectal cancers in Korean, Egyptian and Japanese populations,^{65–67} lung cancer⁶⁸ and breast cancer.²² However, three studies did not find any association between p.Arg399Gln polymorphism

Table 7 Haplotype Frequencies of *XRCC1* Gene Polymorphisms and Their Association with the Risk of Esophageal Cancer

Haplotype <i>XRCC1</i> p.Arg399Gln-p.Arg280His-p.Arg194Trp	Studied Population % (n)(497)	Cases N	Controls	χ^2 (p value)*	Odds Ratio (OR)(95% CI) p value**
G-G-C	0.501(249)	0.508(108)	0.496(141)	0.1 (p=0.7)	Reference
A-G-C	0.347(174)	0.314(67)	0.372(107)	3.6 (p= 0.05)	0.82(0.6–1.2) p=0.3
G-G-T	0.075(37)	0.096(20)	0.058(17)	5.0(p= 0.02)	1.5(0.7–3.1) p= 0.1
G-A-C	0.055(27)	0.060(13)	0.052(14)	0.3(p=0.5)	1.2(0.5–2.7) p= 0.6
A-A-C	0.014(7)	0.015(3)	0.014(4)	0.02(p=0.8)	0.9(0.2–4.4) p=0.9
A-G-T	0.008(4)	0.007(2)	0.008(2)	0.07(p=0.7)	1.3(0.2–9.4) p=0.8
LD Measures	Esophageal Cancer Patients				
	$\hat{D}$	r^2	$\hat{D}$		
	p.Arg399Gln-p.Arg194Trp	0.845	0.042	0.596	
	p.Arg280His-p.Arg194Trp	1.0	0.009	0.969	
p.Arg399Gln-p.Arg280His	0.188	0.001	0.633		

Notes: P* Comparison for esophageal cancer patients and controls for each haplotype. P** Comparison for esophageal cancer patients and controls in comparison to the wild type genotype taken as reference, P<0.05 taken as statistically significant (bold values).

and cancer of the esophagus,⁶⁹ gall bladder⁷⁰ and breast.⁷¹ We did not observe any significant association of *XRCC1* p.Arg194Trp polymorphism with EC risk in the present study. The results were similar to two previous Indian studies.^{52,72} Among international studies, no association of *XRCC1* p.Arg194Trp has been reported with EC risk in the population of North Carolina,⁷³ gastric cancer in Korean population⁷⁴ and breast cancer in Caucasian women.⁷⁵ On the contrary, Trp allele has been reported to be associated with an

increased risk of gastric cancer in the Chinese population.⁷⁶ *XRCC1* is a protein involved in Base excision repair and efficient repair of DNA single-strand breaks. In the present study, we observed a lower risk of esophageal cancer associated with p.Arg399Gln polymorphism of *XRCC1*. A relationship between polymorphism in *XRCC1*Arg399Gln and increased rate of apoptosis has been reported in patients of ulcerative colitis⁷⁷ and in schizophrenia patients.⁷⁸ The alterations to the DNA repair system

Figure 1 LD plot showing the position of the three *XRCC1* polymorphisms and pair-wise D' values observed in the study population with respect to (A) Esophageal Cancer (B) Controls.

Table 8 Association of *XRCC1* Genotype Combinations with Esophageal Cancer

Genotype Combination*	Patients	Controls	Odds Ratio (95% CI)	p value
AA-CC-CC	19	45	0.5(0.3–0.9)	0.02
AA-nn-CC	21	47	0.5(0.3–0.9)	0.03
AA-CC-nn	20	47	0.5(0.3–0.9)	0.02
nn-CT-CC	28	34	1.1(0.6–1.9)	0.69
An-Tn-CC	18	20	1.2(0.6–2.4)	0.56
An-nn-Cn	121	175	0.8(0.6–1.2)	0.28
nn-Tn-Cn	32	37	1.2(0.7–1.9)	0.52
Gn-Tn-Cn	30	35	1.2(0.7–1.9)	0.56

Notes: *Genotypes of *XRCC1* p.Arg399Gln, p.Arg194Trp and p.Arg280His. n: any allele. p value <0.05 is taken as statistically significant (bold values)

could not only suppress it to accomplish the repair process, but it would also become more vulnerable to carry apoptosis and help in wiping out the damaged cells. The increased rates of the apoptosis results into the elimination of potential premalignant cells and hence, the *XRCC1* Gln399 may play a protective role in esophageal cancer risk.

The A allele of *XRCC1*p.Arg280His was associated with an increased risk of esophageal cancer in the present study. Reported studies in different populations and different cancers have contradictory results for this polymorphism. From the North Indian population, p.Arg280His has been associated with an increased risk of hepatocellular carcinoma⁴⁶ but no association has been reported with SCC head and neck.⁴⁵ From a South Indian population, no significant association with breast cancer risk was reported.²² A meta-analysis within the Asian population has reported an association of p.Arg280His polymorphism with bladder cancer risk.⁷⁹ Another meta-analysis reported the heterozygote and the dominant model to be associated with prostate cancer risk.⁸⁰ A allele was associated with an increased risk of adenomas in the Norwegian population.⁶⁰ A study on the Chinese population reported no association between p.Arg280His polymorphism and ESCC.⁸¹

Table 9 Multifactor Dimensionality Reduction (MDR) Analysis

Locus Number	Model	Training Balance Accuracy	Testing Balance Accuracy	Cross Validation Consistency	p value*
1	SNP1	0.5367	0.4965	7/10	0.37
2	SNP1-SNP3	0.5494	0.4538	4/10	0.99
3	SNP1-SNP3-SNP5	0.5621	0.4547	5/10	0.98
4	SNP1-SNP2-SNP3-SNP5	0.5764	0.4289	6/10	0.98
5	SNP1-SNP2-SNP3-SNP4-SNP5	0.5905	0.4648	10/10	0.98

Notes: SNP1: *XRCC1* p.Arg399Gln, SNP2: *XRCC1* p.Arg194Trp, SNP3: *XRCC1* p.Arg280His, SNP4: *XRCC2* Arg188His, SNP5: *XRCC2* p.Thr241Met. *p values were based on 1000 permutations.

Figure 2 Gene–Gene interaction map for esophageal cancer risk: Values inside nodes indicate information gain (IG) of individual attributes or main effects, whereas values between nodes show IG of pairwise combinations of attributes or interaction effects. Positive entropy (red) indicates strong interaction (not found), (yellow) weak interaction while negative entropy (green) indicates redundancy and blue shows no interactions.

Notes: SNP1: *XRCC1* p.Arg399Gln, SNP2: *XRCC1* p.Arg194Trp, SNP3: *XRCC1* p.Arg280His, SNP4: *XRCC2* Arg188His, SNP5: *XRCC2* p.Thr241Met.

XRCC2, an essential protein in the HRR pathway is required for the formation of RAD51 focus, with involvement in tumor progression.¹³ No association of *XRCC2* p.Arg188His with EC has been observed in the present study, which is the first report from India for esophageal cancer risk. In previous reports from India, no association of *XRCC2* p.Arg188His polymorphism was reported with nasopharyngeal carcinoma⁴⁷ whereas another study found an association of GA genotype with increased risk of SCC head and neck.⁴⁹ The contradictory reports from different parts of the world have shown that *XRCC2* p.Arg188His polymorphism was associated with a significantly increased risk of pharyngeal cancer¹⁶ and breast cancer¹² but not with bladder cancer,⁸²

colorectal adenoma,⁸³ skin cancer,⁸⁴ thyroid cancer^{85,86} and breast cancer.⁸⁷ Contrary to this, a protective role of GA and AA genotype in Caucasian breast cancer females of Cyprus has been reported.⁸⁸

XRCC3 gene mainly repairs using the HR pathway and in vitro studies revealed high sensitivity to DNA damaging agents in cells with *XRCC3* gene knockouts.⁸⁹ No association of *XRCC3* p.Thr241Met polymorphism with esophageal cancer has been observed in the present study. A previous study from the north-eastern region of India has reported no association of this polymorphism with breast cancer.⁹⁰ From Indian subcontinent, two populations practicing consanguineous marriages reported an association with increased risk, for colorectal cancer in Kashmiri population⁵⁰ and breast cancer in Pakistani population.⁹¹ For *XRCC3* p.Thr241Met polymorphism, no association was reported with gastric cancer in Italian population⁹² and with colorectal cancer in the West Algerian population⁹³ but an association with an increased risk was reported for lung cancer in an Italian population,²³ oral SCC in Thai population,⁹⁴ and gastric cancer in the Chinese population.⁹⁵ On the other hand, a protective role of CT and TT genotype has been reported in colorectal cancer in Polish population⁹⁶ and Basal cell carcinoma of the skin in population from Hungary, Romania and Slovakia.⁹⁷

The haplotype analysis in the present study shows an association of haplotype GGT of *XRCC1* gene (Arg399-Arg280-194Trp of p.Arg399Gln, p.Arg280His, and p.Arg194Trp polymorphisms) with a risk of EC, however, the association was not statistically significant (Table 7). Among previous reports from India, a study from Andhra Pradesh (South India) also found no association of any haplotype with Chronic Myeloid Leukemia.⁹⁸ However, a study from West Bengal (East India) reported CAG haplotype (Arg194-His280-Arg399) to have reduced risk against gastric cancer.⁵² A study from Uttar Pradesh (North India) reported CGA and CAG haplotype of *XRCC1* (Arg194-His280-Arg399) to be associated with prostate cancer and haplotype CGA to be associated with bladder cancer.³⁸

Post radiotherapy in a cancer patient, radiation injury may develop months to years later after treatment, with numerous individual manifestations. The *XRCC1* codon 399 Arg/Arg genotype has been associated with increased risk of acute radiation dermatitis in nasopharyngeal carcinoma patients treated with intensity-modulated radiation therapy.⁹⁹ Hence a patient with the variant AA genotype of

*XRCC1*p.Arg399Gln polymorphism might be at lower risk for radiation injury. The present study indicates a protective role of the *XRCC1* p.Arg399Gln towards the development of EC. Therefore, *XRCC* gene polymorphisms can also be predicting factor in therapeutic decisions for individualized therapy in EC. Some of the results are at variance from reporting studies in Indian Sub-continent and other Asian countries. These might be due to the ethnic variations in the population residing in Punjab, Northwest India, other states of India and the rest of Asia. Inhabitants of Punjab and Northwestern states of India have an admixture of Indo-Scythian and Caucasian racial elements with more similarity with the central Asian populations as compared to populations in Middle India, South India or eastern parts of India.^{25,100}

Conclusion

The present study being the first report from India, providing baseline data about five genetic polymorphisms in three DNA repair genes *XRCC1*, *XRCC2* and *XRCC3* modulating overall esophageal cancer susceptibility in ethnic Punjabi Indian subjects. The future utility of this data in screening the population for EC risk is the major strength of the study. The results have also been adjusted for various confounding factors. The limitation is a lack of stratification of subjects on the basis of endogamous caste groups of Punjab due to small cross-sectional sample in individual caste group. A possible interaction between *XRCC* polymorphisms and environmental factors on cancer risk was investigated in several studies. It is of great interest to evaluate gene–environment interactions to examine the exact roles of genetic polymorphisms in cancer. However, lacking the original data limited our further evaluation of potential gene–environment interactions. The results suggest a role of *XRCC* gene polymorphisms in esophageal cancer risk and a need to confirm our findings with higher sample size in different ethnic groups inhabiting different geographical areas of India.

Acknowledgments

We are grateful to the participants of the study. The research funding from UGC-UPE and UGC- SAP DRS II to Vasudha Sambyal and Kamlesh Guleria and research fellowship to Jagjeet Kaur is highly acknowledged.

Disclosure

The authors declare that they have no competing interests.

References

1. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2015;136(5):359–386. doi:10.1002/ijc.29210
2. Samarasinghe I. Esophageal cancer in India, current status and future perspectives. *Int J Adv Med Health Res*. 2017;4:5–10. doi:10.4103/IJAMR.IJAMR_19_17
3. Murtaza I, Mushtaq D, Margoob MA, et al. A study on p53 gene alterations in esophageal squamous cell carcinoma and their correlation to common dietary risk factors among population of the Kashmir valley. *World J Gastroenterol*. 2006;12:4033–4037. doi:10.3748/wjg.v12.i25.4033
4. Wood RD, Mitchell M, Sgourous J, Lindahl T. Human DNA repair genes. *Science*. 2001;291:1284–1289. doi:10.1126/science.1056154
5. London RE. The structural basis of XRCC1-mediated DNA repair. *DNA Repair (Amst)*. 2015;30:90–103. doi:10.1016/j.dnarep.2015.02.005
6. Shen MR, Zdzienicka MZ, Mohrenweiser H, Thompson LH, Thelen MP. Mutations in hamster single-strand break repair gene XRCC1 causing defective DNA repair. *Nucleic Acids Res*. 1998;26:1032–1037. doi:10.1093/nar/26.4.1032
7. Pramanik S, Devi S, Chowdhary S, Surendran S, Krishnamurthi K, Chakrabarti T. DNA repair gene polymorphisms at XRCC1, XRCC3, XPD and OGG1 loci in Maharashtrian population of central India. *Chemosphere*. 2011;82:941–946. doi:10.1016/j.chemosphere.2010.10.100
8. Moullan N, Cox DG, Angele S, Romestaing P, Gerard JP, Hall J. Polymorphisms in the DNA repair gene XRCC1, breast cancer risk and response to radiotherapy. *Cancer Epidemiol Biomarkers Prev*. 2003;12:1168–1174.
9. Ratnasinge D, Yao SX, Tangrea JA, et al. Polymorphisms of the DNA repair gene XRCC1 and lung cancer risk. *Cancer Epidemiol Biomarkers Prev*. 2001;10:119–123.
10. Parine NR, Pathan AA, Bobbarala V, Abduljaleel Z, Khan W, Alanazi M. DNA repair gene polymorphisms at XRCC1, XRCC3, XPD, and OGG1 Loci in the hyderabad population of India. *Asian Pac J Cancer Prev*. 2012;13:6469–6474.
11. Braybrooke JP, Spink KG, Thacker J, Hickson ID. The RAD51 family member, RAD51L3, is a DNA-stimulated ATPase that forms a complex with XRCC2. *J Biol Chem*. 2000;275:29100–29106. doi:10.1074/jbc.M002075200
12. Kuschel B, Auranen A, McBride S, et al. Variants in DNA double-strand break repair genes and breast cancer susceptibility. *Hum Mol Genet*. 2002;11:1399–1407. doi:10.1093/hmg/11.12.1399
13. Jiao L, Hassan MM, Bondy ML, et al. XRCC2 and XRCC3 gene polymorphism and risk of pancreatic cancer. *Am J Gastroenterol*. 2008;103:360–367. doi:10.1111/ajg.2008.103.issue-2
14. Auranen A, Song H, Waterfall C, et al. Polymorphisms in DNA repair genes and epithelial ovarian cancer risk. *Int J Cancer*. 2005;117(4):611–618. doi:10.1002/(ISSN)1097-0215
15. Yen CY, Liu SY, Chen CH, et al. Combinational polymorphisms of four DNA repair genes XRCC1, XRCC2, XRCC3 and XRCC4 and their association with oral cancer in Taiwan. *J Oral Pathol Med*. 2008;37:271–277. doi:10.1111/j.1600-0714.2007.00608.x
16. Benhamou S, Tuimala J, Bouchardy C, Dayer P, Sarasin A, Hirvonen A. DNA repair gene XRCC2 and XRCC3 polymorphisms and susceptibility to cancers of the upper aerodigestive tract. *Int J Cancer*. 2004;112(5):901–904. doi:10.1002/ijc.20474
17. Talar-wojnarowska R, Gąsiorowska A, Olakowski A, et al. Analysis of XRCC2 and XRCC3 gene polymorphisms in pancreatic cancer. *Biomed Rep*. 2016;4:236–240. doi:10.3892/br.2015.550
18. Zhao Y, Deng X, Wang Z, Wang Q, Liu Y. Genetic polymorphisms of DNA repair genes XRCC1 and XRCC3 and risk of colorectal cancer in Chinese population. *Asian Pac J Cancer Prev*. 2012;13:665–669. doi:10.7314/APJCP.2012.13.2.665
19. Abdel-fatah T, Arora A, Gorguc I, et al. Are DNA repair factors promising biomarkers for personalized therapy in gastric cancer? *Antioxid Redox Signal*. 2013;18(18):2392–2398. doi:10.1089/ars.2012.4873
20. Mao CF, Qian WY, Wu JZ, Sun DW, Tang JH. Association between the XRCC3 Thr241Met polymorphism and breast cancer risk: an updated meta-analysis of 36 case-control studies. *Asian Pac J Cancer Prev*. 2014;15(16):6613–6618. doi:10.7314/APJCP.2014.15.16.6613
21. Liu HX, Li J, Ye BG. Correlation between gene polymorphisms of CYP1A1, GSTP1, ERCC2, XRCC1, and XRCC3 and susceptibility to lung cancer. *Genet Mol Res*. 2016;15(4). doi:10.4238/gmr15048813
22. Chacko P, Rajan B, Joseph T, Methew BS, Pillai MR. Polymorphisms in DNA repair gene XRCC1 and increased genetic susceptibility to breast cancer. *Breast Cancer Res Treat*. 2005;89:15–21. doi:10.1007/s10549-004-1004-x
23. Improta G, Sgambato A, Bianchino G, et al. Polymorphisms of the DNA repair genes XRCC1 and XRCC3 and risk of lung and colorectal cancer, a case-control study in a Southern Italian population. *Anticancer Res*. 2008;28:291–294.
24. Mandarapu R, Konathala G. Base excision repair pathway gene polymorphisms in South Indian population. *Int J Contemp Med Res*. 2016;3:77–83.
25. Basu A, Mukherjee N, Roy S, et al. Ethnic India: a genomic view, with special reference to peopling and structure. *Genome Res*. 2003;13(10):2277–2290. doi:10.1101/gr.1413403
26. Asthana S, Patil RS, Labani S. Tobacco-related cancers in India: A review of incidence reported from population-based cancer registries. *Indian J Med Paediatr Oncol*. 2016;37(3):152–157. doi:10.4103/0971-5851.190357
27. Bal MS, Bodal VK, Kaur J, Kaur M, Sharma S. Patterns of Cancer, A Study of 500 punjabi patients. *Asian Pac J Cancer Prev*. 2016;16:5107–5110. doi:10.7314/APJCP.2015.16.12.5107
28. Adeli K, Ogbonna G. Rapid purification of human DNA from whole blood for potential application in clinical chemistry laboratories. *Clin Chem*. 1990;36(2):261–264. doi:10.1093/clinchem/36.2.261
29. Wang J, Zhao Y, Jiang J. Polymorphisms in the DNA repair genes XRCC1, XRCC3 and XPD, and colorectal cancer risk, a case-control study in an Indian population. *J Cancer Res Clin Oncol*. 2010;136:1517–1525. doi:10.1007/s00432-010-0809-8
30. Gok I, Baday M, Cetinkunar S, Kilic K, Bilgin BC. Polymorphisms in DNA repair genes XRCC2 and XRCC3 risk of gastric cancer in Turkey. *Bosn J Basic Med Sci*. 2014;14(4):214–218. doi:10.17305/bjbm.2014.4.7
31. Bhasin MK. Morphology to molecular anthropology: castes and tribes of India. *Int J Hum Genet*. 2009;9(3–4):145–230. doi:10.1080/09723757.2009.11886070
32. Alsop BR, Sharma P. Esophageal cancer. *Gastroenterol Clin North Am*. 2016;45(3):399–412. doi:10.1016/j.gtc.2016.04.001
33. Kolkman P, Stack A, McCarthy M, et al. Variations in the histopathologic type of esophageal carcinoma between the United States of America and India. *Indian J Surg Oncol*. 2016;7:52–55. doi:10.1007/s13193-015-0468-y
34. Sobti RC, Singh J, Kaur P, Pachouri SS, Siddiqui EA, Bindra HS. XRCC1 codon 399 and ERCC2 codon 751 polymorphism, smoking, and drinking and risk of esophageal squamous cell carcinoma in a North Indian population. *Cancer Genet Cytogenet*. 2007;175:91–97. doi:10.1016/j.cancergencyto.2007.01.001
35. Duarte MC, Jucimara C, Regina A, Rossit B, Silva AE. Polymorphisms of the DNA repair genes XRCC1 and XRCC3 in a Brazilian population. *Genet Mol*. 2005;28:397–401.

36. Vettrisilvi V, Vijayalakshmi K, Solomon PFD, Venkatachalam P. XRCC1 and XPD gene polymorphisms in a South Indian population. *Asian Pac J Cancer Prev*. 2007;8:283–286.
37. Gangwar R, Manchanda PK, Mittal RD. Implications of XRCC1, XPD and APE1 gene polymorphism in North Indian population, a comparative approach in different ethnic groups worldwide. *Genetica*. 2008;136:163–169. doi:10.1007/s10709-008-9329-8
38. Mittal RD, Mandal RK, Gangwar R. Base excision repair pathway genes polymorphism in prostate and bladder cancer risk in North Indian population. *Mech Ageing Dev*. 2012;133:127–132. doi:10.1016/j.mad.2011.10.002
39. Rao SK, Paul A, Sudarsan A, et al. Allele and genotype distributions of DNA repair gene polymorphisms in South Indian healthy population. *Biomark Cancer*. 2014;6:29–35. doi:10.4137/BIC.S19681
40. Sekhon IS. *Punjabis: The People, Their History, Culture and Enterprise*. New Delhi: Cosmo Publisher; 2000.
41. Shen M, Hung RJ, Brennan P, et al. Polymorphisms of the DNA repair genes XRCC1, XRCC3, XPD, interaction with environmental exposures, and bladder cancer risk in a case-control study in northern Italy. *Cancer Epidemiol Biomarkers Prev*. 2003;12:1234–1240.
42. Duell EJ, Wiencke JK, Cheng TJ, et al. Polymorphisms in the DNA repair genes XRCC1 and ERCC2 and biomarkers of DNA damage in human blood mononuclear cell. *Carcinogenesis*. 2000;21:965–971. doi:10.1093/carcin/21.5.965
43. Srivastava A, Srivastava K, Pandey SN, Choudhuri G, Mittal B. Single-nucleotide polymorphisms of DNA repair genes OGG1 and XRCC1, association with gall bladder cancer in North Indian population. *Ann Surg Oncol*. 2009;16:1695–1703. doi:10.1245/s10434-009-0354-3
44. Khan NP, Pandith AA, Yousuf A, et al. The XRCC1 Arg399Gln gene polymorphism and risk of colorectal cancer, a study in Kashmir. *Asian Pac J Cancer Prev*. 2013;14:6779–6782. doi:10.7314/APJCP.2013.14.11.6779
45. Kumar A, Pant MC, Singh HS, Khandelwal S. Associated risk of XRCC1 and XPD cross talk and life style factors in progression of head and neck cancer in north Indian population. *Mut Res*. 2012;729:24–34. doi:10.1016/j.mrfmmm.2011.09.001
46. Kiran M, Saxena R, Chawla YK, Kaur J. Polymorphism of DNA repair gene XRCC1 and hepatitis-related hepatocellular carcinoma risk in Indian population. *Mol Cell Biochem*. 2009;327:7–13. doi:10.1007/s11010-009-0035-3
47. Singh A, Singh N, Behera D, Sharma S. Association and multiple interaction analysis among five XRCC1 polymorphic variants in modulating lung cancer risk in North Indian population. *DNA Repair (Amst)*. 2016;47:30–41. doi:10.1016/j.dnarep.2016.09.006
48. Uppal V, Mehndiratta M, Mohapatra D, Grover RK, Puri D. XRCC-1 gene polymorphism (Arg399Gln) and susceptibility to development of lung cancer in cohort of North Indian population A pilot study. *J Clin Diagn Res*. 2014;8:17–20.
49. Choudhury JH, Choudhury B, Kundu S, Ghosh SK. Combined effect of tobacco and DNA repair genes polymorphisms of XRCC1 and XRCC2 influence high risk of head and neck squamous cell carcinoma in northeast Indian population. *Med Oncol*. 2014;31(8):67. doi:10.1007/s12032-014-0067-8
50. Nissar S, Lone TA, Banday MZ, et al. Arg399Gln polymorphism of XRCC1 gene and risk of colorectal cancer in Kashmir, A case control study. *Oncol Lett*. 2013;5:959–963. doi:10.3892/ol.2013.1104
51. Berhane N, Sobti RC, Mahdi SA. DNA repair genes polymorphism (XPG and XRCC1) and association of prostate cancer in a north Indian population. *Mol Biol Rep*. 2012;39(3):2471–2479. doi:10.1007/s11033-011-0998-5
52. Ghosh S, Ghosh S, Bankura B, et al. Association of DNA repair and xenobiotic pathway gene polymorphisms with genetic susceptibility to gastric cancer patients in West Bengal India. *Tumour Biol*. 2016;37:9139–9149. doi:10.1007/s13277-015-4780-5
53. Sreeja L, Syamala VS, Syamala V, et al. Prognostic importances of DNA repair gene polymorphisms of XRCC1 Arg399Gln and XPD Lys751Gln in lung cancer patients from India. *J Cancer Res Clin Oncol*. 2008;134:645–652. doi:10.1007/s00432-007-0328-4
54. Natukula K, Jamil K, Pingali UR, Attili VS, Madireddy UR. The codon 399 Arg/Gln XRCC1 polymorphism is associated with lung cancer in Indians. *Asian Pac J Cancer Prev*. 2013;14:5275–5279. doi:10.7314/APJCP.2013.14.9.5275
55. Konathala G, Mandarapu R, Godi S. Data on polymorphism of XRCC1 and cervical cancer risk from South India. *Data Brief*. 2017;10:11–13. doi:10.1016/j.dib.2016.11.052
56. Singh SR, Ghosh SK. Polymorphisms of XRCC1 and XRCC2 DNA repair genes and interaction with environmental factors influence the risk of Nasopharyngeal carcinoma in Northeast India. *Asian Pac J Cancer Prev*. 2016;17:2811–2819.
57. Joseph T, Kusumakumary P, Chacko P, Abraham A, Pillai MR. DNA repair gene XRCC1 polymorphisms in childhood acute lymphoblastic leukemia. *Cancer Lett*. 2005;217:17–24. doi:10.1016/j.canlet.2004.06.055
58. Santhi S, Sangeetha V, Sureshkumar R, et al. Risk effects of XRCC1 Arg399Gln and XPD Lys751Gln gene polymorphisms in de novo acute myeloid leukemia - A study from India. *Indian J Biotechnol*. 2017;16:275–283.
59. Xing D, Qi J, Miao X, Lu W, Tan W, Lin D. Polymorphisms of DNA repair genes XRCC1 and XPD and their associations with risk of esophageal squamous cell carcinoma in a Chinese population. *Int J Cancer*. 2002;100:600–605. doi:10.1002/ijc.10528
60. Skjelbred CF, Saebø M, Wallin H, et al. Polymorphisms of the XRCC1, XRCC3 and XPD genes and risk of colorectal adenoma and carcinoma in a Norwegian cohort, a case-control study. *BMC Cancer*. 2006;6:67–76. doi:10.1186/1471-2407-6-67
61. Stern MC, Umbach DM, van Gils CH, Lunn RM, Taylor JA. DNA repair gene XRCC1 polymorphisms, smoking and bladder cancer risk. *Cancer Epidemiol Biomarkers Prev*. 2001;10:125–132.
62. Nelson HH, Kelsey KT, Mott LA, Karagas MR. The XRCC1 polymorphism, sunburn, and non-melanoma skin cancer, evidence of gene-environment interaction. *Cancer Res*. 2002;62:152–160.
63. Li S, Deng Y, You JP, et al. XRCC1 Arg399Gln, Arg194Trp, and Arg280His polymorphisms in esophageal cancer risk, a meta-analysis. *Digest Dis Sci*. 2013;58:1880–1890.
64. Shu XO, Cai Q, Gao YT, Wen W, Jin F, Zheng W. A population based case control study of the Arg399Gln polymorphism in DNA repair gene XRCC1 and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev*. 2003;12:1462–1467.
65. Abdel-rahman SZ, Soliman AS, Bondy ML, et al. Inheritance of the 194Trp and the 399Gln variant alleles of the DNA repair gene XRCC1 are associated with increased risk of early-onset colorectal carcinoma in Egypt. *Cancer Lett*. 2000;159(1):79–86. doi:10.1016/S0304-3835(00)00537-1
66. Hong YC, Lee H, Kim WC, Choi SK, ZHa W, Shin SK. Polymorphisms of XRCC1 gene, alcohol consumption and colorectal cancer. *Int J Cancer*. 2005;116:428–432. doi:10.1002/ijc.21019
67. Yin G, Morita M, Ohnaka K, et al. Genetic polymorphisms of XRCC1, alcohol consumption, and the risk of colorectal cancer in Japan. *J Epidemiol*. 2011;22:64–71. doi:10.2188/jea.JE20110059
68. Park JY, Lee SY, Jeon HS, et al. Polymorphism of the DNA repair gene XRCC1 and risk of primary lung cancer. *Cancer Epidemiol Biomarkers Prev*. 2002;11:23–27.
69. Yu H-P, Zhang X-Y, Wang X-L, et al. DNA repair gene XRCC1 polymorphisms, smoking and esophageal cancer risk. *Cancer Detect Prev*. 2004;28:194–199. doi:10.1016/j.cdp.2004.01.004
70. Huang WY, Gao YT, Rashid A, et al. Selected base excision repair gene polymorphisms and susceptibility to biliary tract cancer and biliary stones, a population-based case-control study in China. *Carcinogenesis*. 2008;29:100–105. doi:10.1093/carcin/bgm247

71. Karolina PS, Stanczyk M, Kusinska R, Kordek R, Majsterek I. Association of the Arg194Trp and the Arg399Gln polymorphisms of the XRCC1 gene with risk occurrence and the response to adjuvant therapy among Polish women with breast cancer. *Clin Breast Cancer*. 2013;13:61–68. doi:10.1016/j.clbc.2012.09.019
72. Pavithra D, Gautam M, Rama R, et al. TGFβ C-509T, TGFβ T869C, XRCC1 Arg194Trp, IKBa C642T, IL4 C-590T Genetic polymorphisms combined with socio-economic, lifestyle, diet factors and gastric cancer risk: A case control study in South Indian population. *Cancer Epidemiol*. 2018;53:21–26. doi:10.1016/j.canep.2018.01.004
73. Lunn RM, Langlois RG, Hsieh LL, Thompson CL, Bell DA. XRCC1 polymorphisms, effects on aflatoxin B-DNA adducts and glycophorin A variant frequency. *Cancer Res*. 1999;59:2557–2561.
74. Jin E-H, Kim J, Lee S-I, Hong JH. Association between polymorphisms in APE1 and XRCC1 and the risk of gastric cancer in Korean population. *Int J Clin Exp Med*. 2015;8:11484–11489.
75. Sterpone S, Mastellone V, Padua L, et al. Single-nucleotide polymorphisms in BER and HRR genes, XRCC1 haplotypes and breast cancer risk in Caucasian women. *J Cancer Res Clin Oncol*. 2010;136:631–636. doi:10.1007/s00432-010-0791-1
76. Wen -Y-Y, Pan X-F, Loh M, et al. ADPRT Val762Ala and XRCC1 Arg194Trp Polymorphisms and risk of gastric cancer in Sichuan of China. *Asian Pac J Cancer Prev*. 2012;13:2139–2144. doi:10.7314/APJCP.2012.13.5.2139
77. Bardia A, Tiwari SK, Gunisetty S, et al. Functional polymorphisms in XRCC-1 and APE-1 contribute to increased apoptosis and risk of ulcerative colitis. *Inflamm Res*. 2012;61(4):359–365. doi:10.1007/s00011-011-0418-2
78. Catts VS, Catts SV, McGrath JJ, et al. Apoptosis and schizophrenia: a pilot study based on dermal fibroblast cell lines. *Schizophr Res*. 2006;84(1):20–28. doi:10.1016/j.schres.2006.03.016
79. Fang Z, Chen F, Wang X, Yi S, Chen W, Ye G. XRCC1 Arg194Trp and Arg280His polymorphisms increase bladder cancer risk in Asian population, evidence from a meta-analysis. *PLoS One*. 2013;8:5.
80. Yan J, Wang X, Tao H, Deng Z, Yang W, Lin F. Meta-analysis of the relationship between XRCC1-Arg399Gln and Arg280His polymorphisms and the risk of prostate cancer. *Sci Rep*. 2015;5:9905. doi:10.1038/srep09905
81. Zhai XD, Mo YN, Xue XQ, et al. XRCC1 codon 280 and ERCC2 codon 751 polymorphisms and risk of esophageal squamous cell carcinoma in a Chinese population. *Bull Cancer*. 2009;96:61–65. doi:10.1684/bdc.2009.0952
82. Matullo G, Guarrera S, Sacerdote C, et al. Polymorphisms/haplotypes in DNA repair genes and smoking, a bladder cancer case-control study. *Cancer Epidemiol Biomarkers Prev*. 2005;14:2569–2578. doi:10.1158/1055-9965.EPI-05-0189
83. Tranah GJ, Giovannucci E, Ma J, Fuchs C, Hankinson SE, Hunter DJ. XRCC2 and XRCC3 polymorphisms are not associated with risk of colorectal adenoma. *Cancer Epidemiol Biomarkers Prev*. 2004;13:1090LP- 1091.
84. Han J, Hankinson SE, Hunter DJ, De Vivo I. Genetic variations in XRCC2 and XRCC3 are not associated with endometrial cancer risk. *Cancer Epidemiol Biomarkers Prev*. 2004;13:330–331. doi:10.1158/1055-9965.EPI-03-0332
85. Bastos HN, Antao MR, Silva SN, et al. Association of polymorphisms in genes of the homologous recombination DNA repair pathway and thyroid cancer risk. *Thyroid*. 2009;19(10):1067–1075. doi:10.1089/thy.2009.0099
86. Yan L, Li Q, Li X, Ji H, Zhang L. Association studies between XRCC1, XRCC2, XRCC3 polymorphisms and differentiated thyroid carcinoma. *Cell Physiol Biochem*. 2016;38:1075–1084. doi:10.1159/000443058
87. Yu KD, Chen AX, Qiu LX, Fan L, Yang C, Shao ZM. XRCC2 Arg188His polymorphism is not directly associated with breast cancer risk, evidence from 37,369 subjects. *Breast Cancer Res Treat*. 2010;123:219–225. doi:10.1007/s10549-010-0753-y
88. Loizidou MA, Michael T, Neuhausen S, et al. Genetic polymorphisms in the DNA repair genes XRCC1, XRCC2 and XRCC3 and risk of breast cancer in Cyprus. *Breast Cancer Res Treat*. 2008;112:575–579. doi:10.1007/s10549-007-9881-4
89. Singleton BK, Griffin CS, Thacker J. Clustered DNA damage leads to complex genetic changes in irradiated human cells. *Cancer Res*. 2002;62(21):6263–6269.
90. Devi KR, Ahmed J, Narain K, et al. DNA repair mechanism gene, XRCC1A (Arg194Trp) but not XRCC3 (Thr241Met) polymorphism increased the risk of breast cancer in premenopausal females A case-control study in Northeastern region of India. *Technol Cancer Res Treat*. 2017;16:1150–1159. doi:10.1177/1533034617736162
91. Qureshi Z, Mahjabeen I, Baig R, Kayani M. Correlation between Selected XRCC2, XRCC3 and RAD51 gene polymorphisms and primary breast cancer in women in Pakistan. *Asian Pac J Cancer Prev*. 2014;15:10225–10229. doi:10.7314/APJCP.2014.15.23.10225
92. Palli D, Polidoro S, D'errico M, et al. Polymorphic DNA repair and metabolic genes, a multigenic study on gastric cancer. *Mutagenesis*. 2010;25:569–575. doi:10.1093/mutage/geq042
93. Moghiti FZ, Aberkane MS, Morvan L, et al. No association between XRCC3 Thr241Met and XPD Lys751Gln polymorphisms and the risk of colorectal cancer in west Algerian population, a case-control study. *Medical Oncol*. 2014;31:942–949. doi:10.1007/s12032-014-0942-3
94. Kietthubthaw S, Sriplung H, Au WW, Ishida T. Polymorphism in DNA repair genes and oral squamous cell carcinoma in Thailand. *Int J Hyg Environ Health*. 2006;209:21–29. doi:10.1016/j.ijheh.2005.06.002
95. Huang GP, Zheng ZL, Cai L. DNA repair gene XRCC3 Thr241Met polymorphism and susceptibility to cardia and non-cardia gastric cancer, a case-control study. *Zhonghua Liu Xing Bing XueZaZhi*. 2006;27:420–423.
96. Mucha B, Przybylowska-sygut K, Dziki AJ, Dziki LS, Majsterek I. Association of Thr241Met polymorphism of XRCC3 gene with risk of colorectal cancer in the polish population. *Pol J Pathol*. 2013;64:185–190. doi:10.5114/pjp.2013.38137
97. Thiumaran RK, Bermejo JL, Rudnai P, et al. Single nucleotide polymorphisms in DNA repair genes and basal cell carcinoma of skin. *Carcinogenesis*. 2006;27:1676–1681. doi:10.1093/carcin/bgi381
98. Annamaneni S, Gorre M, Kagita S, et al. Association of XRCC1 gene polymorphisms with chronic myeloid leukemia in the population of Andhra Pradesh, India. *Hematology*. 2013;18(3):163–168. doi:10.1179/1607845412Y.00000000040
99. Chen H, Wu M, Li G, Hua L, Chen S, Huang H. Association between XRCC1 single-nucleotide polymorphism and acute radiation reaction in patients with nasopharyngeal carcinoma. *Medicine*. 2017;96:e8202. doi:10.1097/MD.00000000000008202
100. Majumder P. Ethnic populations of India as seen from an evolutionary perspective. *J Biosci*. 2001;26:533–545. doi:10.1007/BF02704750
101. Datkhile KD, Bhosale SJ, Patil MN, Khamkar TS, Durgawale PP, Kakade SV. Genetic polymorphism of DNA repair genes (XRCC1, XRCC2 & XRCC3) in breast cancer patients from rural Maharashtra. *Int J Health Sci Res*. 2017;7:281–290.
102. Nishank SS. Association of DNA damage repair gene polymorphisms hOGG1, XRCC1and p53 with sickle cell disease patients in India. *Mediterr J Hematol Infect Dis*. 2015;7:e2015046. doi:10.4084/mjhid.2015.046

Clinical and Experimental Gastroenterology

Dovepress

Publish your work in this journal

Clinical and Experimental Gastroenterology is an international, peer-reviewed, open access, online journal publishing original research, reports, editorials, reviews and commentaries on all aspects of gastroenterology in the clinic and laboratory. This journal is indexed on American Chemical Society's Chemical Abstracts Service (CAS).

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-and-experimental-gastroenterology-journal>