

A Novel Transposon, Tn65/8, Mediated Transfer of *mcr-3* Variant in ESBL-Producing *Aeromonas veronii*

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Xiaoming Wang^{1,2}
Weishuai Zhai³
Shaolin Wang² 
Zhangqi Shen²
Yang Wang²
Qidi Zhang³

¹MOE Joint International Research Laboratory of Animal Health and Food Safety, College of Veterinary Medicine, Nanjing Agricultural University, Nanjing, People's Republic of China; ²Beijing Advanced Innovation Center for Food Nutrition and Human Health, College of Veterinary Medicine, China Agriculture University, Beijing, People's Republic of China; ³College of Veterinary Medicine, Qingdao Agricultural University, Qingdao, Shandong, People's Republic of China

Purpose: The aim of this study was to determine the prevalence and transmission mechanism of *mcr-3* in *Aeromonas* spp. isolated from chicken cloaca.

Materials and Methods: *A. veronii* w55 was isolated from chicken in 2008. PCR assay was used to detect *mcr* genes and putative circular intermediate. Susceptibility testing was identified by the microdilution method. WGS was performed to obtain the whole sequence. S1-PFGE and DNA southern hybridization were used to study the location of *mcr-3.6*.

Results: PCR-based analysis indicated that 1 out of 55 *Aeromonas* spp. isolates was *mcr-3*-positive. Whole-genome sequencing revealed that the strain *A. veronii* w55 belonged to novel sequence type ST514 and had two adjacent chromosomally located *mcr* variants, *mcr-3.6* and *mcr-3*-like. The *mcr-3.6* and *mcr-3*-like genes showed 93.67% and 82.84% nucleotide sequence identity, respectively, to original *mcr-3* from *E. coli*. *A. veronii* w55 also exhibited resistance to extended-spectrum β -lactams and was positive for *bla*_{PER-3}, and this is the first time to report *bla*_{PER-3} in *A. veronii*. Genetic environment analysis revealed that the segment of *mcr-3.6-mcr-3*-like-*dgkA* was flanked by five insertion sequence elements originated from *Aeromonas* species, and the structure of ISAs2-ISAh₂-ISAs20-*mcr-3.6-mcr-3*-like-*dgkA*-ISAs2 was designated as a novel transposon Tn65/8, in which an 8405-bp circular intermediate carrying two *mcr-3* variants can be looped out.

Conclusion: This result suggested the *mcr-3* variant genes could be disseminated between various *Aeromonas* species via transposon-mediated transmission.

Keywords: *mcr-3.6*, Tn65/8, *Aeromonas veronii*

Introduction

Aeromonas species are Gram-negative, rod-shaped bacteria with a broad host range. To date, they have been shown to infect a vast number of hosts, including fish, domestic chickens, lower and higher vertebrates, and humans.¹⁻³ In animals, *Aeromonas* species cause illnesses ranging from furunculosis and diarrhea to septicemia.⁴ However, because of the wide use of broad-spectrum antibiotics in clinical settings, agriculture, and aquaculture, there has been an increase in antibiotic resistance among disease-causing *Aeromonas* species.

Colistin, a polycationic peptide, is considered as one of the last-resort drugs for the treatment of infections caused by multi-drug resistant Gram-negative bacteria.⁵ In 2016, the first plasmid-mediated colistin resistance gene *mcr-1* was characterized in Enterobacteriaceae.⁶ To date, nine mobile colistin resistance determinants (*mcr-1-mcr-9*) have been reported in Enterobacteriaceae, *Moraxella* and *Aeromonas*.⁷⁻¹⁰ Subsequently, multiple *mcr-3* variants and one *mcr-5* gene were identified in *Aeromonas* species, implying *Aeromonas* species plays an important role in the

Correspondence: Qidi Zhang
No. 700 Changcheng Road, Chengyang District, Qingdao City, Shandong Province 266109, People's Republic of China
Email zqdcu@163.com

dissemination of mobile colistin resistance genes.^{11–13} Among them *mcr-3* was first found in *Escherichia coli* isolated from pig,¹⁴ then some papers illustrated that this gene emerged in *Aeromonas* spp.¹² Subsequently, Ling et al study showed that *mcr-3.3* and *mcr-3*-like have emerged in *Aeromonas veronii* isolated from retail chicken meat,⁸ and Eichhorn et al found four *Aeromonas* isolates were positive for *mcr-3*, each of the four *mcr-3* genes showed amino acid identities of 95.0–98.0% to the original *mcr-3* protein. These variants were designated *mcr-3.6* (*Aeromonas allosaccharophila* from golden orfe), *mcr-3.7* (*Aeromonas media* from), *mcr-3.8* (*Aeromonas jandaei* from koi carp) and *mcr-3.9* (*Aeromonas hydrophila* from koi carp).¹⁴ Herein, we investigated the presence of *mcr*-type genes in *Aeromonas* spp. from chicken, and identified a novel transposon Tn6518 carrying *mcr-3* variant in ESBL-producing *Aeromonas veronii*.

Materials and Methods

Bacterial Isolates, Screening for *mcr*-Type, and Species Confirmation

Fifty-five *Aeromonas* spp. isolates were collected from commercial chickens in Sichuan Province, China, in 2008, and were screened for the presence of *mcr-1* to *mcr-9* genes using a polymerase chain reaction (PCR)-based assay with primers listed in [Table S1](#). The PCR-positive isolate was then subjected to further PCR and sequencing analysis of the 16S *rRNA*, *gyrB*, and *rpoB* loci for species identification as per Kupfer et al.¹⁵

Antimicrobial Susceptibility Testing

The minimum inhibitory concentrations (MICs) of 13 antimicrobial agents (listed in [Table S2](#)) against the *mcr-3*-positive isolate were then determined by broth microdilution according to the recommendations of the Clinical and Laboratory Standards Institute (document VET01-A4). The breakpoints for each antimicrobial agent were interpreted according to CLSI (document M45-A2) and European Committee on Antimicrobial Susceptibility Testing (EUCAST) (<http://www.eucast.org>) guidelines.¹⁶ *E. coli* ATCC 25922 was used as a quality control strain.

Genomic DNA Sequencing and Analysis of Antibiotic Resistance Genes

Genomic DNA from the *mcr-3*-positive isolate was subjected to sequence using both the Illumina Hiseq platform and the PacBio RSII System (Sinobiocore, Beijing, China). CLC Genomics Workbench 9.0 (CLC Bio, Aarhus, Denmark)

was used for assembling data from Illumina Hiseq platform, and all contigs were screened for the presence of *mcr-3* using standalone BLAST analysis. The data from PacBio RSII System were assembled using the Hierarchical Genome Assembly Process (HGAP) and Quiver as part of the SMRT analysis program (version 2.3) using the HGAP3 protocol, and then corrected using Pilon. Reference sequences for antibiotic resistance genes were obtained from the ARG-ANNOT database (<https://omictools.com/antibiotic-resistance-gene-annotation-tool>). Multilocus sequence typing (MLST) was carried out using the MLST tool from the Center for Genomic Epidemiology (<http://genomicepidemiology.org/>), and annotation of the genomic sequences was performed using ORF Finder (www.ncbi.nlm.nih.gov/gorf/orfig.cgi), Vector NTI (Invitrogen, Carlsbad, CA, USA), and ISFinder (<http://www-is.biotoul.fr>).

S1 Nuclease Pulsed-Field Gel Electrophoresis (S1-PFGE) and Southern Blotting

S1-PFGE and Southern blotting were performed to locate the *mcr-3* gene within the genome. Briefly, whole cells were embedded in agarose gel plugs and digested with S1 nuclease (TaKaRa, Dalian, China). The DNA was then separated by PFGE as described previously.¹⁷ For Southern blotting, a 450-bp digoxigenin-labelled *mcr-3* PCR amplicon was used as a probe. Genomic DNA from *Salmonella enterica* serovar Braenderup strain H9812 restricted with *Xba*I was used as the DNA marker.

Detection of Circular Intermediates

Based on the knowledge that IS30-family elements can form circular DNA intermediates, and that ISAs2 belongs to the IS30 family,^{18,19} a set of reverse primers (P-F: GGCTGGTTGTGATTGTAGAG and P-R: TCGCTCGGTCTGTTTGCTTT) were designed to investigate the potential of the ISAs2-ISAh2-ISAs20-*mcr-3.12*-*mcr-3*-like-*dgkA*-ISAs2 segment to circularize. The designed primers allowed us to obtain a complete map of the circular intermediate.

Functional Cloning of *mcr-3.6* and *mcr-3*-Like Genes in *E. coli* and *Aeromonas* Species

To determine the function of *mcr-3* variant genes, 1983-bp and 2176-bp DNA fragments, corresponding to the *mcr-3.6* and *mcr-3*-like regions along with their upstream sequences, respectively, were amplified and cloned into

pUC19, as previously reported.⁸ The resulting recombinant plasmids, pUC19-*mcr-3.6* and pUC19-*mcr-3*-like, were then separately electroporated into *E. coli* DH5 α (TaKaRa, Dalian, China) and *Aeromonas salmonicida* cells originally isolated from chicken cloaca swabs. The recombinant strains were then subjected to antibiotic susceptibility testing.

Results

Prevalence of *mcr-1* to *mcr-9* in *Aeromonas* Species Isolates

The use of colistin as animal growth enhancers has been banned in China since April, 2017. However, prior to that time, the extensive use of colistin in food animal production, especially in commercial poultry farming, increased the risk of transfer of colistin resistance determinants in bacteria. Our retrospective survey of *mcr-1* to *mcr-9* in 55 *Aeromonas* spp. isolates showed that only one ESBL-producing *A. veronii* isolate w55 was *mcr-3*-positive, indicating that the presence of *mcr-3* in *Aeromonas* in China for at least 10 years.

WGS Analysis and Functional Identification of *mcr-3.6* and *mcr-3*-Like Genes

The *mcr-3.6* variant has been deposited into NCBI with GenBank number: MF598076.1, and was first detected from *A. allosaccharophila* strain IMT43045 isolated from German *Leuciscus idus*.¹⁹ Illumina HiSeq 2500-based WGS revealed the presence of two adjacent *mcr-3* variants, *mcr-3.6* and *mcr-3*-like separated by only 66 bp, on a 36.9-kb contig of *A. veronii* w55. The two variants showed 93.7% and 82.8% nucleotide sequence identity, respectively, to the original *mcr-3* from porcine *E. coli*.²⁰ Both variants carried several missense mutations, resulting in 94.8% and 84.2% amino acid sequence similarity, respectively, to MCR-3.

In order to determine the two *mcr-3* variants function, we expressed *mcr-3.6* and *mcr-3*-like in *E. coli* DH5 α . Transformants containing pUC19-*mcr-3.6* or pUC19-*mcr-3*-like had colistin MIC of 2 μ g/mL and 0.25 μ g/mL, respectively, which was 8- and 1-fold higher MIC than that of *E. coli* DH5 α containing pUC19 alone (0.25 μ g/mL), respectively. Furthermore, to investigate the function of the two variants in *Aeromonas* species, recombinant plasmids pUC19-*mcr-3.6* and pUC19-*mcr-3*-like were separately electroporated into competent *A. salmonicida* cells, and the MICs of colistin against these two transformants were 32 μ g/mL (carrying pUC19-*mcr-3.6*) and 1 μ g/mL (carrying pUC19-*mcr-3*-like), which were 32- and 1-fold higher MIC than that

of the wild-type *A. salmonicida* strain (1 μ g/mL), respectively. These results suggested that only *mcr-3.6* could confer colistin resistance in both *E. coli* and other *Aeromonas* species, which agrees with only pUC19-*mcr-3.3* strain confers colistin resistance.⁸ Although most of the identified *mcr-3* variants in *Aeromonas* species so far derived from animals, the surveillance of these potential high-colistin resistance genes in clinical *Aeromonas* isolates are also needed.

Antimicrobial Susceptibility Profiles

Antimicrobial susceptibility testing showed that *A. veronii* w55 was resistance to amoxicillin-clavulanate (MIC, 32/16 μ g/mL), cefotaxime (8 μ g/mL), ceftriaxone (16 μ g/mL), ceftazidime (16 μ g/mL), gentamycin (16 μ g/mL), and chloramphenicol (32 μ g/mL), and was susceptible to aztreonam (2 μ g/mL), imipenem (0.03 μ g/mL), meropenem (0.03 μ g/mL), ciprofloxacin (0.5 μ g/mL) and tetracycline (4 μ g/mL) (Table S2). Interestingly, despite the presence of the two *mcr-3* variants, the MIC value for both colistin and polymyxin B against *A. veronii* w55 was 2 μ g/mL, which was determined as “susceptible” based on the EUCAST clinical breakpoints (<http://www.eucast.org>).

MLST and Other Resistance Genes

MLST analysis revealed that *A. veronii* w55 belonged to novel sequence type ST514 (<https://cge.cbs.dtu.dk/services/MLST/>). WGS analysis showed that *A. veronii* w55 not only carried colistin resistance genes *mcr-3.6* and *mcr-3*-like but also contained ESBL-producing gene *bla*_{PER-3}. *PER-3* was initially characterised from an *Aeromonas punctata* in France (GenBank accession no. AY740681.1) and recently in a medical centre in Taiwan,²¹ sharing 99% homology to *bla*_{PER-1}. Here we report the first identification of a *bla*_{PER-3}-producing *A. veronii* isolate. In addition, aminoglycoside resistance genes *strA*, *strB*, *aac(6')Ib-cr*, and *aacA4*, macrolide resistance gene *mph(A)*, phenicol resistance gene *catB3*, sulfonamide resistance gene *sul1*, and tetracycline resistance gene *tet(E)* were also detected in the genome of *A. veronii* w55. In these resistance genes, *mcr-3.6*, *mcr-3*-like, *strA*, *mph(A)*, *aacA4*, *catB3*, *sul1*, *tet(E)* located on chromosome, and others located on plasmid.

Location and Genetic Environment of *mcr-3.6*

S1-nuclease PFGE and Southern blotting using an *mcr-3*-specific probe showed that *mcr-3.6* was located on the chromosome of *A. veronii* w55 (Figure 1). The segment of




Figure 1 Location of *mcr-3.6* on the chromosome of *Aeromonas veronii* w55. (A) SI-PFGE and (B) the corresponding Southern hybridization using the *mcr-3.6* probe. Lane M, *Xba*I-digested *Salmonella enterica* serovar Braenderup strain H9812 genomic DNA marker; lane w55, *A. veronii* strain w55 genomic DNA.

mcr-3.6-mcr-3-like-dgkA in a 36.9-kb-size contig was surrounded by five IS elements (two ISAs2, one ISAs20 and two truncated ISAhv2, all of these elements were originally identified from various *Aeromonas* species, <https://www-is.bio.toul.fr/scripts/search-db.php>), which was different from that in the original *mcr-3.6*-carrying *A. allosaccharophila* IMT43045, where this segment was flanked by a truncated IS3 element (Figure 2). In detail, the upstream of *mcr-3.6-mcr-3-like-dgkA* segment contained one copy of ISAs2, ISAs20, and truncated Δ 1ISAhv2 elements. The ISAs2 and ISAs20 elements harbored 50-bp imperfect inverted repeats (IR) at both ends (as shown in Figure 2). In addition, direct target site duplications (5'-CTT-3' and 5'-AAGG-3') were also observed immediately up- and downstream of the ISAs2 and ISAs20, respectively (Figure 2). While immediate downstream of *mcr-3.6-mcr-3-like-dgkA* segment, another copy of ISAs2 and a truncated Δ 2ISAhv2 were observed. The IR elements of the downstream ISAs2 were identical to that in upstream; however, the direct target site duplications of the downstream ISAs2 (5'-AAG-3') were differed from that in the upstream (5'-CCT-3') (Figure 2).

Formation of a Circular Intermediate Containing the *mcr-3.6* Gene

Some members of the IS family, including IS3, IS30, IS110, IS26, and ISCR1 elements, utilize circular DNA intermediates containing accessory genes to undergo gene translocation via copy-and-paste mechanisms.²² ISAs2, which is first identified in *A. salmonicida*, disrupts a range of genes in *Aeromonas* species,²³ also belongs to the IS3 family and is flanked by left and right inverted repeats. Therefore, we hypothesize that *mcr-3.6* translocation could be mediated via a circular intermediate with the assistance of ISAs2 element. The reverse primers P-F and P-R targeting the Δ 1ISAhv2 and *dgkA* loci, respectively (Figure 2), were used to amplify a putative circular intermediate harboring *mcr-3.6* in *A. veronii* w55. PCR products with a size of ~2.0 kb were obtained and composed of the partial sequence of Δ 1ISAhv2 and *dgkA* and one ISAs2 element. When combined with the sequence between primers P-F and P-R, we concluded that the circular intermediate was 8405-bp in size, and contained one copy of ISAs2, Δ 1ISAhv2, ISAs20, and a segment of *mcr-3.6-mcr-3-like-dgkA* (Figure 2). This new composite transposon was thereafter designated as Tn6518 (ISAs2- Δ 1ISAhv2-ISAs20-*mcr-3.6-mcr-3-like-dgkA*-ISAs2). Nevertheless, the molecular mechanisms




Figure 2 Genetic organization of scaffolds containing *mcr-3.6* obtained from *Aeromonas veronii* strain w55 in this study, and structural comparison with *A. allosaccharophila* IMT43045 (GenBank accession no. MF598076.1). The positions and orientations of the genes are indicated by arrows, with the direction of transcription shown by the arrowhead. Grey shading indicates >99% nucleotide sequence identity. The genetic structure of transposon Tn6518 is shown in circular form.

underlying the formation of the circular intermediate form of Tn6518, and how it evolves to become a genetically stable fragment in the chromosome, require further investigation.

Discussion

Aeromonas spp. species are recognized enteric pathogens, and it is possible that the presence of *Aeromonas* isolates that carry these *mcr-3* variants in the intestine will facilitate the spread of these genes to other bacterial species living in the same habitat.^{15,24} Until now, there are several *mcr-3* variants, including *mcr-3.3* and *mcr-3.13* to *mcr-3.18*, identified among 10 *Aeromonas* isolates from different sources.¹² Shen et al reported samples from the aquatic environment had the highest prevalence (10.0%) of *mcr-3*-positive *Aeromonas* isolates,¹² which indicated that *mcr-3* have been disseminated throughout various environments, the aquatic environment might be the major reservoir. For this study, it is the first case about a novel transposon, Tn6518, mediated transfer of *mcr-3.6* in ESBL-producing *Aeromonas veronii* isolated from chicken faeces.

The first reported *mcr-3*-carrying *E. coli* WJ1 exhibited MICs of 8 ug/mL for both colistin and polymyxin B.¹⁴ However, five previously identified *Aeromonas* isolates carrying *mcr-3.3* to *mcr-3.7* exhibited colistin MICs ranging from 2 to 128 ug/mL. In the original report on *mcr-3.3* in *A. veronii*, the *mcr-3.3*-positive isolate demonstrated

borderline susceptibility to colistin (MIC of 2 mg/liter).¹¹ However, *E. coli* and *A. salmonicida* transformants carrying pUC19-*mcr-3.3* exhibited 8- and 64-fold higher MIC values (MICs of 2 mg/liter and 64 mg/liter, respectively) than the transformant carrying pUC19 alone, which was similar with our findings. These observations not only confirmed that the *mcr-3* variants can confer colistin resistance in both *Enterobacteriaceae* strains and *Aeromonads* but also suggested that the colistin resistance levels conferred by these genes may vary among different *Aeromonas* species. In the previous report, the *mcr-3* variants were surrounded by many different types of IS, like IS26, IS6100, ISKpn3 and ISAS17.^{12,15} However, in our study, the genetic environment of *mcr-3.6* was ISAs2-ISAhy2-ISAs20-*mcr-3.6*-*mcr-3-like*-*dgkA*-ISAs2, which can form a circular intermediate, and may increase the transmission of colistin resistance.

In conclusion, two *mcr-3* variants, *mcr-3.6* and *mcr-3-like*, were identified in ESBL-producing *A. veronii* isolate w55. This isolate has the novel sequence type ST514 and was isolated from a chicken cloacal sample in 2008. In addition, WGS analysis revealed that the segment of *mcr-3.6*-*mcr-3-like*-*dgkA* was surrounded by multi-copy IS elements and termed as a novel transposon Tn6518, which may be responsible for the transmission of *mcr-3.6* among various *Aeromonas* species. More worrying, *Aeromonas* species are prevalent in aquatic environments, where they can interact with different bacterial

species and genus. Overall, the presence of transposon-carrying *mcr-3* variants in *Aeromonas* species is highly concerning with regards to public health and highlights the need to investigate the prevalence of *mcr-3* in high-level colistin-resistant clinical *Aeromonas* isolates.

Nucleotide Sequence Accession No

The nucleotide sequence reported in this study has been submitted to GenBank with NCBI no. MH481281.

Acknowledgments

We thank Dr. Adam Roberts from UCL Eastman Dental Institute for assistance in registering Tn6518 in the Tn registry.

Funding

This work was supported in part by the grants from the National Key Research and Development Program of China (2016YFD0501304 and 2018YFD0500300), National Natural Science Foundation of China (81661138002), and Qingdao Science and Technology Project (No. 17-3-3-69-nsh).

Disclosure

The authors report no conflicts of interest in this work.

References

- Yang S, He T, Sun J, et al. distinct antimicrobial resistance profiling of clinically important *Aeromonas* spp. in Southwest China: a seven-year surveillance study. *Infect Drug Resist.* 2019;12:2971–2978. doi:10.2147/IDR
- Castelo-Branco DS, Silva AL, Monteiro FO, et al. *Aeromonas* and *Plesiomonas* species from scarlet ibis (*Eudocimus ruber*) and their environment: monitoring antimicrobial susceptibility and virulence. *Antonie Van Leeuwenhoek.* 2017;110(1):33–43. doi:10.1007/s10482-016-0771-9
- Dong HT, Techatanakitarnan C, Jindakittikul P, et al. *Aeromonas jandaei* and *Aeromonas veronii* caused disease and mortality in Nile tilapia, *Oreochromis niloticus* (L.). *J Fish Dis.* 2017;40:1395–1403.
- Dias C, Borges A, Saavedra MJ, et al. Biofilm formation and multidrug-resistant *Aeromonas* spp. from wild animals. *J Glob Antimicrob Resist.* 2018;12:227–234. doi:10.1016/j.jgar.2017.09.010
- Li J, Nation RL, Turnidge JD, et al. Colistin: the re-emerging antibiotic for multidrug-resistant Gram-negative bacterial infections. *Lancet Infect Dis.* 2006;6(9):589–601. doi:10.1016/S1473-3099(06)70580-1
- Liu Y, Wang Y, Walsh TR, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis.* 2016;16(2):161–168. doi:10.1016/S1473-3099(15)00424-7
- AbuOun M, Stubberfield EJ, Duggett NA, et al. *mcr-1* and *mcr-2* variant genes identified in *Moraxella* species isolated from pigs in Great Britain from 2014 to 2015. *J Antimicrob Chemother.* 2017;72(10):2745–2749. doi:10.1093/jac/dkx286
- Ling Z, Yin W, Li H, et al. Chromosome-mediated *mcr-3* variants in *Aeromonas veronii* from chicken meat. *Antimicrob Agents Chemother.* 2017;61(11):e01272–17.
- Roschanski N, Falgenhauer L, Grobbel M, et al. Retrospective survey of *mcr-1* and *mcr-2* in German pig-fattening farms, 2011–2012. *Int J Antimicrob Agents.* 2017;50(2):266–271. doi:10.1016/j.ijantimicag.2017.03.007
- Carroll LM, Gaballa A, Guldemann C, et al. Identification of novel mobilized colistin resistance gene *mcr-9* in a multidrug-resistant, colistin-susceptible *Salmonella enterica* Serotype Typhimurium Isolate. *mBio.* 2019;10(3):e00853–19.
- Wang X, Zhai W, Li J, et al. Presence of an *mcr-3* variant in *Aeromonas caviae*, *Proteus mirabilis*, and *Escherichia coli* from one domestic duck. *Antimicrob Agents Chemother.* 2018;62(2):e02106–17.
- Shen Y, Xu C, Sun Q, et al. Prevalence and genetic analysis of *mcr-3*-positive *Aeromonas* species from humans, retail meat, and environmental water samples. *Antimicrob Agents Chemother.* 2018;62(9):e00404–18.
- Ma S, Sun C, Hulth A, et al. Mobile colistin resistance gene *mcr-5* in porcine *Aeromonas hydrophila*. *J Antimicrob Chemother.* 2018;73(7):1777–1780. doi:10.1093/jac/dky110
- Eichhorn I, Feudi C, Wang Y, et al. Identification of novel variants of the colistin resistance gene *mcr-3* in *Aeromonas* spp. from the national resistance monitoring programme GERM-Vet and from diagnostic submissions. *J Antimicrob Chemother.* 2018;73(5):1217–1221. doi:10.1093/jac/dkx538
- Kupfer M, Kuhnert P, Korczak BM, Peduzzi R, Demarta A, Peduzzi R, Demarta A. Genetic relationships of *Aeromonas* strains inferred from 16S *rRNA*, *gyrB* and *rpoB* gene sequences. *Int J Syst Evol Microbiol.* 2006;56(Pt 12):2743–2751.
- Shen Z, Wang Y, Shen Y, et al. Early emergence of *mcr-1* in *Escherichia coli* from food-producing animals. *Lancet Infect Dis.* 2016;16(3):293. doi:10.1016/S1473-3099(16)00061-X
- Hu YY, Wang YL, Sun QL, et al. Colistin resistance gene *mcr-1* in gut flora of children. *Int J Antimicrob Agents.* 2017;50(4):593–597. doi:10.1016/j.ijantimicag.2017.06.011
- He S, Hickman AB, Varani AM, et al. Insertion sequence IS26 reorganizes plasmids in clinically isolated multidrug-resistant bacteria by replicative transposition. *mBio.* 2015;6(3):e762. doi:10.1128/mBio.00762-15
- Olasz F, Farkas T, Kiss J, et al. Terminal inverted repeats of insertion sequence IS30 serve as targets for transposition. *J Bacteriol.* 1997;179(23):7551–7558. doi:10.1128/JB.179.23.7551-7558.1997
- Yin W, Li H, Shen Y, et al. Novel plasmid-mediated colistin resistance gene *mcr-3* in *Escherichia coli*. *MBio.* 2017;8:e543.
- Wu CJ, Chuang YC, Lee MF, et al. Bacteremia due to extended-spectrum- β -lactamase-producing *Aeromonas* spp. at a Medical Center in Southern Taiwan. *Antimicrob Agents Chemother.* 2011;55(12):5813–5818. doi:10.1128/AAC.00634-11
- Li R, Xie M, Zhang J, et al. Genetic characterization of *mcr-1*-bearing plasmids to depict molecular mechanisms underlying dissemination of the colistin resistance determinant. *J Antimicrob Chemother.* 2017;72(2):393–401. doi:10.1093/jac/dkw411
- Gustafson CE, Chu S, Trust TJ. Mutagenesis of the paracrystalline surface protein array of *Aeromonas salmonicida* by endogenous insertion elements. *J Mol Biol.* 1994;237(4):452–463. doi:10.1006/jmbi.1994.1247
- Batra P, Mathur P, Misra MC. *Aeromonas* spp.: an emerging nosocomial pathogen. *J Lab Physicians.* 2016;8(1):1–4. doi:10.4103/0974-2727.176234

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>