Oral Maintenance Chemotherapy Using S-1/Capecitabine in Metastatic Nasopharyngeal Carcinoma Patients After Systemic Chemotherapy: A Single-Institution Experience [Corrigendum]

Guo Q, Chen M, Xu H, et al. *Cancer Manag Res.* 2020;12:1387–1396.

The authors have advised that the explanation for the # symbol is missing in Figure 2E key on page 1392. There are also missing markers for Radiotherapy to the

metastatic foci and Radiofrequency ablation in the figure. The correct figure is below.

The authors apologize for these errors.

Figure 2 Kaplan-Meier curves for (A) OS and (B) PFS in mNPC patients. (C, D) Kaplan-Meier estimates of smNPC and mmNPC patients. (E) Treatment exposure and response.

Cancer Management and Research**Dovepress****Publish your work in this journal**

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>