#### International Journal of Nanomedicine

Open Access Full Text Article

ORIGINAL RESEARCH

# Co-Delivery of Paclitaxel and Doxorubicin by pH-Responsive Prodrug Micelles for Cancer Therapy

This article was published in the following Dove Press journal: International Journal of Nanomedicine

Yanhua Jiang<sup>1</sup> Yongjian Zhou<sup>1</sup> Can Yang Zhang<sup>2</sup> Te Fang<sup>1</sup>

<sup>1</sup>Department of Anesthesiology, First Affiliated Hospital of China Medical University, Shenyang 110001, People's Republic of China; <sup>2</sup>Singapore-MIT Alliance for Research and Technology, Singapore 138602, Singapore **Background:** It is of great significance to develop intelligent co-delivery systems for cancer chemotherapy with improved therapeutic efficacy and few side-effects.

**Materials and Methods:** Here, we reported a co-delivery system based on pH-sensitive polyprodrug micelles for simultaneous delivery of doxorubicin (DOX) and paclitaxel (PTX) as a combination chemotherapy with pH-triggered drug release profiles. The physicochemical properties, drug release profiles and mechanism, and cytotoxicity of PTX/DOX-PMs have been thoroughly investigated.

**Results and Discussion:** The pH-sensitive polyprodrug was used as nanocarrier, and PTX was encapsulated into the micelles with high drug-loading content (25.6%). The critical micelle concentration (CMC) was about 3.16 mg/L, indicating the system could form the micelles at low concentration. The particle size of PTX/DOX-PMs was 110.5 nm, and increased to approximately 140 nm after incubation for 5 days which showed that the PTX/DOX-PMs had high serum stability. With decrease in pH value, the particle size first increased, and thenwas no longer detectable. Similar change trend was observed for CMC values. The zetapotential increased sharply with decrease in pH. These results demonstrated the pHsensitivity of PTX/DOX-PMs. In vitro drug release experiments and study on release mechanism showed that the drug release rate and accumulative release for PTX and DOX were dependent on the pH, showing the pH-triggered drug release profiles. Cytotoxicity assay displayed that the block copolymer showed negligible cytotoxicity, while the PTX/DOX-PMs possessed high cytotoxic effect against several tumor cell lines compared with free drugs and control.

**Conclusion:** All the results demonstrated that the co-delivery system based on pH-sensitive polyprodrug could be a potent nanomedicine for combination cancer chemotherapy. In addition, construction based on polyprodrug and chemical drug could be a useful method to prepare multifunctional nanomedicine.

**Keywords:** pH-responsive, polyprodrug micelles, co-delivery, controlled release, cancer therapy

#### Introduction

Chemotherapy is still currently the most commonly used treatment for various cancers in clinics.<sup>1–3</sup> Some efficient chemical anticancer drugs, such as doxorubicin (DOX),<sup>4</sup> paclitaxel (PTX),<sup>5</sup> and camptothecin (CPT),<sup>6</sup> have been developed for cancer chemotherapy. However, a lot of shortcomings have been reported during clinical use.<sup>7–10</sup> For example, DOX also induces damage of normal tissues, especially the heart, meanwhile killing the tumor cells.<sup>11</sup> Furthermore, small chemical

Correspondence: Te Fang Email tfang@cmu.edu.cn



© 2020 Jiang et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (http://www.dovepress.com/terms.php).

anticancer drug molecules would be cleared quickly by the reticuloendothelial system (RES) and blood/renal during circulation in the body after administration, leading to the quite low concentration at the tumor sites.<sup>12–15</sup> When the dose of drug is increased, the adverse side-effects become severe. Therefore, it is of great significance to find a method to solve this dilemma. With the rapid development of bionanotechnology recently, a nanoscale stimuli-responsive drug delivery system (DDS) has emerged and attracted more and more attention, and has been deemed to be an effective strategy to increase the drug concentration at the tumor sites and remit the unwanted side-effects of chemical anticancer drugs.<sup>16–24</sup> In the past years, various DDSs have been designed and developed for drug delivery, such as polymeric micelles (PMs),<sup>25</sup> hybrid nanoparticles (NPs),<sup>26</sup> metal-organic framework (MOF),<sup>27</sup> liposomes,<sup>28</sup> and liquid emulsion.<sup>29</sup> Generally, the chemical drugs are physically loaded in the core of the system or chemically linked on the carriers. After administration, the drug-loaded systems would remain stable and compact in the blood circulation in case the cargos are leaked. Once they arrive at the tumor sites by the enhanced permeability and retention (EPR) effect, the drug would be released by responding to the specific tumor microenvironment cues (eg, low pH, high glutathione concentration and special enzyme), followed by inducing the death of tumor cells.<sup>30–33</sup> Compared with other nanoscale carriers, PMs which are self-assembled from amphiphilic copolymers have several unique superior advantages.<sup>34–36</sup> For instance, it is easy to modify the copolymer to obtain multifunctional carriers. And the PMs possess high drug loading efficacy, high biocompatibility, and low toxicity.<sup>37</sup> Compared to the physical drug loading method, the drug molecules are chemically linked on the carriers via sensitive bonds, such as disulfide bonds and hydrazone bonds, resulting in a polyprodrug which could effectively reduce the drug burst release and efficiently protect the drugs during blood circulation.<sup>38,39</sup> Combining the advantages of both (PMs and prodrugs), some novel nanomedicines have been reported.<sup>40-42</sup> Ma et al prepared a prodrug comprised of mussel-derived biomimetic peptide with fluorescein isothiocyanate for imaging, RGD sequence for targeting, and a pH-sensitive conjugation of antitumor drugs. The prodrug could self-assemble into polyprodrug micelles which were used for active targeted delivery of anticancer drug. The in vivo experimental results showed that this system could effectively deliver drug to tumor cells and suppresstumor growth.43

Recently, several limitations of single drug cancer chemotherapy have been reported such as poor bioavailability, low therapeutic efficacy, and multidrug resistance.<sup>44</sup> Therefore, combination chemotherapy of two or more anticancer drugs has been thoroughly investigated and extensively developed in the past few years.45-48 For example, Zhu et al prepared a poly(lactic-co-glycolic acid) nanoparticle (PLGA NPs) for co-delivery of docetaxel (DTX) together with vitamin E TPGS with improved therapeutic efficacy against multidrug resistance.<sup>49</sup> Danafar et al reported drug conjugated PEG-PCL NPs for co-delivery of hydrophilic and hydrophobic anticancer drugs. The DOX-conjugated block copolymers were synthesized first, and then curcumin was physically loaded in the polymeric core during the self-assembly in aqueous. The cytotoxicity assay showed that the system had a higher cytotoxic effect against A549 cells compared with free drugs.<sup>50</sup> Therefore, co-delivery of multidrugs using intelligent nanoscale carriers could be a potent method to enhance therapeutic efficacy and overcome drug resistance.

Herein, inspired by the specific tumor microenvironment cues, we designed and prepared a co-delivery system composed of pH-sensitive polyprodrug copolymer and anticancer drug PTX. The DOX-conjugated diblock copolymer synthesized in our previous work.<sup>51</sup> Here, was these polyprodrugs were not only used as nanomedicine, but also simultaneously utilized as stimuli-responsive carrier for encapsulation and delivery of PTX with pH-triggered drug release profiles for cancer combination chemotherapy, as shown in Figure 1. The DOX-conjugated copolymer molecules could self-assemble into polymeric micelles together with hydrophobic PTX in aqueous, resulting in PTX-loaded PMs (so-called PTX/DOX-PMs). For PTX/DOX-PMs, the DOX molecules were chemically linked on the pH-sensitive diblock copolymers, and PTX molecules were physically loaded in the micellar core. The PTX/DOX-PMs showed high stability for extended circulation in the body at normal pH (pH 7.4). After deposition at the tumor sites, part of tertiary amine residues in PAE block would be protonated which led to the increase of surface charge of PTX/DOX-PMs, facilitating the cellular uptake by tumor cells. And then, the pH-sensitive bonds between polymer and DOX molecules would be cleaved by responding to the intracellular acidity. All of the tertiary amine residues in PAE block would be protonated, resulting in disassembly of PTX/DOX-PMs to release the encapsulated PTX molecules. Moreover, the drug release mechanisms of



Figure I Self-assembly of pH-responsive polyprodrug micelles for co-delivery of PTX and DOX with pH-triggered drug release profiles. **Abbreviations:** DOX, doxorubicin; PTX, paclitaxel; mPEG-*b*-PAE-*cis*-DOX, diblock copolymer poly(ethylene glycol)methyl ether-*b*-poly(β-amino esters) conjugated with DOX via acid-labile *cis*-aconityl moiety; PTX/DOX-PMs, paclitaxel and doxorubicin-loaded polymeric micelles; i.v. injection; intravenous injection; PTX/DOX, paclitaxel and doxorubicin molecules.

chemically loaded DOX and physically loaded PTX were investigated based on a comprehensive semi-empirical equation. The pH-triggered drug release profiles and cytotoxicity of systems against different tumor cell lines were investigated. In addition, some other physicochemical properties of PTX/DOX-PMs, such as particle size, zeta potential and pH -sensitivity, were also studied in this work.

## Materials and Methods

#### Materials

Diblock copolymer poly(ethylene glycol)methyl etherb-poly( $\beta$ -amino esters) conjugated with DOX via acidlabile cis-aconityl moiety (mPEG-b-PAE-cis-DOX) was synthesized as reported in our precious work.<sup>51</sup> Doxorubicin hydrochloride (DOX-HCl) was purchased from Wuhan Yuan Cheng Gong Chuang Co. Ltd (Wuhan, China). Paclitaxel (PTX) was purchased from Xi'an Helin Pharm Co., Ltd. (Xi'an, China) Methylthiazoltetrazolium (MTT) was purchased from Sigma-Aldrich. Dulbecco's modified eagle media (DMEM) growth media, fetal bovine serum (FBS), trypsin, penicillin, streptomycin, and other biological agents were all purchased from Invitrogen. A549, MDA-MB-231, A2780 and NCL-H460 cell lines were obtained from the American Type Culture Collection (ATCC). All other chemical and biological reagents were used as received.

## Preparation of PTX/DOX-PMs

The PTX/DOX-PMs were prepared using dialysis method. Briefly, polyprodrug mPEG-*b*-PAE-*cis*-DOX (20 mg) was dissolved into DMF (20 mL) with stirring for 30 min. Then PTX (5 mg, 10 mg, 20 mg) was added into the solution. The mixed solution was transferred into a preswollen cellulose membrane bag (MWCO 3500–4000), followed by placing in a beaker (1 L). The dialysis process was carried out for 48 h against deionized water at room temperature. The deionized water was replaced every 2 h in the first 12 h and then every 6 h. After filtration by 0.45  $\mu$ m filter and lyophilization, the PTX/DOX-PMs were obtained and stored at  $-20^{\circ}$ C for further studies. DOX-PMs were prepared in a similar way as reported elsewhere.<sup>51</sup>

## Characterization

The particle size of PTX/DOX-PMs and DOX-PMs was measured by dynamic light scattering (DLS, Malvern Zetasizer Nano S, Malvern, UK). As a topical experiment, PTX/DOX-PMs were re-suspended in PBS at different pH values (9.0, 8.0, 7.4, 7.0, 6.8, 6.5, 6.0, 5.5, 5.0, and 4.0) at a concentration of 1.0 mg/mL, and incubated for 2 h. The samples were measured in a 1.0 mL quartz cuvette using a diode laser of 670 nm at room temperature. The zeta potential of samples was measured in a similar method.

To evaluate the serum stability of PTX/DOX-PMs, the samples were prepared as previously mentioned. Then the samples were incubated in PBS with 20% FBS at 37°C for 5 days, the particle size and polydispersity index (PDI) of samples were recorded every day using DLS.

## Drug Loading Efficacy

The PTX loading content (LC) and encapsulation efficiency (EE) were analyzed using high- performance liquid chromatography (HPLC, Waters 2690 Separations Module system, Milford, MA) consisting of a P-900 gradient pump system and an Ultra C18 5  $\mu$ m column (250 × 4.6 mm). Briefly, 1 mg of PTX/DOX-PMs was first dissolved into dichloromethane (DCM) for 30 min. Then, the resultant solution was filtered by 0.45  $\mu$ m filter. The HPLC analysis was performed using acetonitrile/water (v/v = 4: 1) as the mobile phase at 30°C with a flow rate of 1.0 mL/min and detected at a wavelength of 227 nm (UV detector, Waters). The LC and EE were calculated according to the following equations:

$$LC (\%) = \frac{Weight of loaded drug}{Weight of drug loaded - NPs} \times 100\%$$
$$EE (\%) = \frac{Weight of loaded drug}{Weight of drug in feed} \times 100\%$$

The DOX conjugated efficiency (DCE) has already been confirmed using UV-*vis* spectrophotometer (UV-2450, Shimadzu, Japan), as reported elsewhere.<sup>51</sup>

## Critical Micelle Concentration (CMC) Measurement

The CMC values of systems were measured by the fluorescence technique using pyrene as a probe dye. In brief, the block copolymer was first dissolved into deionized water at a concentration of 0.1 mg/mL. The systems were mixed with the pre-prepared pyrene solution to prepare a series of mixed solutions with concentrations from 0.0001 to 0.1 mg/mL, and the final concentration of pyrene was  $6 \times 10^{-7}$  M. Before measurement, the obtained mixture was equilibrated in the dark for one day at 25°C. The excitation spectra of samples were tested with a fluorescence spectrophotometer (F-4500, Hitachi, Japan) at 373 nm with a bandwidth of 0.2 nm.

#### In vitro Release of PTX and DOX from PMs

To acquire sink conditions, in vitro drug release study was performed at low drug concentration. The release profiles of PTX and DOX from PTX/DOX-PMs were investigated using a dialysis method. Briefly, 2 mg of PTX/DOX-PMs was re-suspended in 2 mL of PBS at pH 7.4, 6.5 or 5.0, and then the solution was transferred into a pre-swollen cellulose membrane bag (MWCO 3500–4000) which was immersed into 38 mL of respective PBS solution in a beaker with stirring of 110 rpm at 37°C. At pre-determined time intervals, 1 mL of sample was drawn from the beaker and 1 mL fresh respective PBS was added. The PTX and DOX concentrations of every sample were analyzed using HPLC and UV-*vis* spectrophotometer as previously mentioned, respectively. The accumulative release amount of drug ( $E_r$ ) was calculated based on the following equation:

$$E_r(\%) = \frac{V_e \sum_{1}^{n-1} Ci + V_0 C_n}{m_{drug}} \times 100\%$$

where,  $m_{\rm drug}$  represented the amount of PTX or DOX in the PTX/DOX-PMs,  $V_0$  was the whole volume of the release media ( $V_0 = 40$  mL), and  $C_i$  represented the concentration of PTX or DOX in the  $i_{\rm th}$  sample.

#### Drug Release Mechanism Study

The drug release mechanism from PMs was very complex and not completely understood until now. Peppas et al established a comprehensive semi-empirical equation to roughly analyze the mechanism,<sup>52,53</sup> as follows:

$$\frac{M_t}{M_\infty} = kt^n$$

This formula has a transformation, as follows:

$$\log\left(\frac{M_t}{M_{\infty}}\right) = n\log t + \log k$$

Where,  $M_t/M_{\infty}$  was the accumulated amount of drug at time *t*. *k* was the constant incorporating structural and geometric characteristics of carrier which indicated the drug release rate, and *n* was the release exponent indicating the drug release mechanism. The value of n was 0.43 for Fickian diffusion and n < 0.43 corresponded to the combination of diffusion and erosion control, 0.43 < n < 0.85 was for anomalous transport mechanism, and *n* was 0.85 for swelling-controlled mechanism.

#### Cell Culture

The cell culture protocols for A549, MDA-MB-231, A2780, NCL-H460 and NIH 3T3 cell lines were similar as reported elsewhere.<sup>51,54</sup> The cells were cultured in the medium supplemented with 10% FBS, 100 units/mL penicillin and 100  $\mu$ g/mL streptomycin at 37°C in an incubator. After two or three times of passage, the cells were used for future studies.

#### Cytotoxicity Assay

The cytotoxic effects of diblock copolymer, free DOX, free PTX, mixture of free DOX and PTX, DOX-PMs, PTX/ DOX-PMs against these cell lines were evaluated utilizing the standard methylthiazoltetrazolium (MTT) assay. Tumor cells were seeded in a 96-well plate at a density of  $1 \times 10^4$ cells per well in 200 µL of medium. The 96-well plate was cultured in the incubator for one day. After removing the culture supernatants, the prepared diblock copolymer, free DOX, free PTX, mixed free DOX and PTX, DOX-PMs and PTX/DOX-PMs medium solution with a concentration gradient were added into the wells. Tumor cells with fresh medium were used as blank back-ground. Live tumor cells without any treatment were used as a control with no exogenous cytotoxicity. After incubation for 48 h, 20 µL of MTT solution (5 mg/mL) was added into the wells. The 96-well plate was shaken at 150 rpm for additional 10 min, followed by incubation for 4 h. The supernatants in every well were discarded, and 200 µL of DMSO was added to dissolve the formazan crystals, and the plate was shaken for 15 min. The 96-well plate was measured by a microplate reader (Multiskan Spectrum, Thermo Scientific, Finland) at 490 nm. The relative cell viability (%) was calculated using the following formula:

$$Cell \, viability = \frac{A_{sample} - A_{blank}}{A_{control} - A_{blank}} \times \, 100\%$$

where, A was the absorbance at 490 nm. The cytotoxicity test was performed in replicates of six wells.

#### Statistical Analysis

The experimental data were presented with average values, expressed as the mean  $\pm$  standard deviation (S.D.). Statistical analysis was conducted using two-sample Student's *t*-test of origin 8.5 or ANOVA analyses, and considered to be significant when p < 0.05.

#### **Results and Discussion**

## Preparation and Characterization of PTX/ DOX-PMs

The PTX/DOX-PMs were prepared using dialysis method. The particle sizes of DOX-PMs and PTX/DOX-PMs were recorded by DLS, as shown in Figure 2A. The hydrodynamic diameters of DOX-PMs and PTX/DOX-PMs were 102.3 nm and 110.5 nm, respectively. After PTX loading, the particle size was slightly increased due to the entrapped hydrophobic PTX molecules in the micellar core. As reported, the PMs with size of 50-200 nm were suitable for accumulation at tumor sites through EPR effect in cancer therapy.<sup>9</sup> In order to receive high accumulation at the site of tumor, the PTX/DOX-PMs should have high serum stability. Next, the serum stability of PTX/DOX-PMs in PBS containing 20% FBS at pH of 7.4 was evaluated, as shown in Figure 2B. After incubation for 1 day, the particle size was increased from 110.5 nm to approximately 130 nm. Even after incubation for 5 days, the particle size of PTX/DOX-PMs was still less than 150 nm, indicating that the co-delivery system, PTX/DOX-PMs, had high serum stability. In addition, the PDI of PTX/DOX-PMs after incubation for



Figure 2 (A) Hydrodynamic diameter of DOX-PMs and PTX/DOX-PMs measured by DLS. (B) Particle size and polydispersity index (PDI) of PTX/DOX-PMs after incubation in the presence of 20% FBS in PBS at pH 7.4 at 37°C for 5 days (n = 3, mean ± SD). Abbreviations: DOX-PMs, doxorubicin-loaded polymeric micelles; PTX/DOX-PMs, paclitaxel and doxorubicin-loaded polymeric micelles; DLS, dynamic light scattering; FBS, fetal bovine serum; PBS, phosphate buffer solution; PDI, polydispersity index.

different times was still less than 0.3, suggesting the good uniformity and higher stability. These findings proved that the PTX/DOX-PMs had proper particle size and high serum stability, thereby enhancing the accumulation at tumor sites via EPR effect.

The particle size, PDI, LC and EE of PTX/DOX-PMs at different mass ratios of PTX and polyprodrug copolymer were shown in Table 1. With increase of PTX in feed, the LC of PTX increased, as expected. When the PTX in feed was 5 mg (PTX: carriers= 1: 4, mass ratio), the LC and EE were 16.8 % and 80.3%, respectively. When the mass ratio was 1: 2 (PTX in feed was 10 mg), the LC markedly increased to 25.6%, while the EE was slightly decreased to 75.4%. When the PTX in feed was 20 mg (PTX: carriers= 1: 1, mass ratio), the LC slightly increased to 29.4%, but the EE was sharply decreased to 55.8%. These results showed that the 10 mg of PTX in feed (PTX: carriers= 1: 2, mass ratio) could be the optimal formulation, and these PTX/DOX-PMs were used for the following studies. Additionally, the particle size of PTX/DOX-PMs was slightly increased with increase of PTX in feed, resulting from more PTX molecules in the micellar core thereby enhancing the core of PMs. Moreover, the average number of DOX molecules conjugated on the PAE chain was and  $2.5 \pm 0.2$  per polymer molecule.<sup>51</sup>

#### pH Sensitivity of PTX/DOX-PMs

Next, the pH sensitivity of PTX/DOX-PMs was studied by measurement of particle size, zeta potential and CMC values of systems at different pH conditions. As seen in Figure 3A, when the pH was higher than 7.4, the particle size of PTX/DOX-PMs was quite stable (110 nm-120 nm). When the pH decreased from 7.4 to 6.5, the particle size of PTX/DOX-PMs sharply increased. In particular, the particle size increased to 165.8 nm at pH of 6.5. The reason could be that part of tertiary amine residues in PAE block were protonated at weakly acidic condition which induced the solubility transformation of PAE segment from hydrophobicity to hydrophilicity, resulting in swelling of PTX/DOX-PMs. When the pH decreased

sequentially (< 6.5), no particles were detectable, indicating there were no particles. The reason might be that all of the tertiary amine residues in PAE segment were ionized, leading to the hydrophilic PAE segment. Moreover, the pH-sensitive bonds cis-aconityl linker between copolymer and DOX molecules were cleaved, resulting in detachment of the hydrophobic DOX. Therefore, the original amphiphilic block copolymer mPEG-b-PAE-cis-DOX was changed to hydrophilic diblock copolymer mPEG-b-PAE, suggesting the disassembly of PTX/ DOX-PMs. Figure 3B shows the zeta potential of PTX/DOX-PMs at different pH values. As expected, the zeta potential rapidly increased (from -7.8 mV to +56.1 mV) with decrease in pH (from 9.0 to 4.0), attributed to the ionization of tertiary amine residues of PAE segment. Figure 3C shows the CMCs of DOX-PMs and PTX/DOX-PMs at different pH conditions. For DOX-PMs, the CMC was increased from 3.6 µg/mL to 45.2  $\mu$ g/mL when the pH decreased from 7.4 to 6.8. A similar change trend was observed for PTX/DOX-PMs system. The CMCs were 3.16 µg/mL (Figure S1), 18.55 µg/mL and 49.64 µg/mL, respectively, at pH 7.4, 6.8 and 6.5. This was ascribed to part of the tertiary amine residues of PAE segment being protonated which enhanced the molecular polarity, thereby requiring greater driving force for micellization to counteract the greater electrostatic repulsive force. When the pH was lower than 6.5, the CMCs of DOX-PMs and PTX/DOX-PMs were not detectable, resulting from ionized tertiary amine residues in PAE segment and cleavage of pH-sensitive bonds. The micelles were disassembled at this stage. In addition, the CMC of PTX/DOX-PMs was slightly lower than that of DOX-PMs due to the introduction of hydrophobic PTX molecules in the system. In summary, the particle size, zeta potential and CMC of PTX/DOX-PMs were dependent on the pH value. These results proved that the PTX/DOX-PMs possessed pH sensitivity.

#### pH-Triggered DOX Release in vitro

Since the pH sensitivity of PTX/DOX-PMs has been confirmed, in vitro release profiles of PTX and DOX from

**Table I** Particle Size, Polydispersity Index (PDI), Drug Loading Content (LC) and Encapsulation Efficacy (EE) of PTX/DOX-PMs at Different Mass Ratios of Drug and Carriers

Prodrug (20 mg)	PTX (mg)	Size (nm) <sup>a</sup>	PDI <sup>a</sup>	LC (%) <sup>b</sup>	EE (%) <sup>b</sup>
DOX-polymer	5	102.2±2.4	0.131±0.014	16.8±1.5	80.3±3.1
	10	110.5±3.1	0.135±0.021	25.6±1.7	75.4±2.5
	20	113.6±2.5	0.140±0.018	29.4±1.6	55.8±2.8

Note: <sup>a</sup>Measured by dynamic light scattering (DLS), <sup>b</sup>measured by high-performance liquid chromatography (HPLC).

Abbreviations: DOX, doxorubicin; PTX, paclitaxel; PTX/DOX-PMs, paclitaxel and doxorubicin-loaded polymeric micelles.



**Figure 3** Particle size (**A**) and zeta potential (**B**) of PTX/DOX-PMs incubated in PBS solutions with different pH values for 4 h (n = 3, mean  $\pm$  SD). CMC values (**C**) of DOX-PMs ad PTX/DOX-PMs in PBS solutions with different pH values (n = 3, mean  $\pm$  SD).

Abbreviations: DOX-PMs, doxorubicin-loaded polymeric micelles; PTX/DOX-PMs, paclitaxel and doxorubicin-loaded polymeric micelles; PBS, phosphate buffer solution; CMC, critical micelle concentration; ND, not detectable.

PTX/DOX-PMs at different pH values were investigated, as shown in Figure 4. Figure 4A shows the DOX release profiles from PTX/DOX-PMs at different pH conditions (7.4, 6.5 and 5.0). The drug release rate and accumulative release were obviously dependent on the pH values. At pH 7.4, the DOX release rate was quite low, and the accumulative release of DOX was less than 10% (9.8%) for 48 h. Because the DOX molecules were chemically linked on the PAE chains, and the polyprodrug molecules were stable at pH 7.4. When the pH was 6.5, the release rate rapidly accelerated. The accumulative release was about 20% for 2 h and more than 75% at 48 h, respectively, resulting from the cleavage of pH-sensitive bonds between DOX molecules and polymer. When the pH was 5.0, the

release rate was much faster in comparison to that at pH 6.5, and the accumulative release was about 30% for 2 h and higher than 95% at 48 h, respectively. The key reason might be that all of the pH-sensitive bonds between DOX and copolymer were broken. Furthermore, the PMs were disassembled, and the free DOX molecules were easily diffused into the solution. Figure 4B shows the PTX release profiles from PTX/DOX-PMs at different pH values (7.4, 6.5 and 5.0). Although the PTX release profiles were dependent on the pH values, the release rate of PTX was faster than that of DOX at pH 7.4. The accumulative release of PTX was more than 10% (12.6%) at 2 h and about 40% (39.8%) at 48 h, respectively. The reason could be that the PTX was physically



Figure 4 In vitro DOX (A) and PTX (B) release profiles of PTX/DOX-PMs in PBS at pH 7.4, pH 6.5, pH 5.0 (n = 3, mean ± SD). Abbreviations: DOX, doxorubicin; PTX, paclitaxel; PMs, polymeric micelles; PTX/DOX-PMs, paclitaxel and doxorubicin-loaded polymeric micelles; PBS, phosphate buffer solution.

loaded into the micellar core by hydrophobic interaction during the self-assembly process. Some PTX molecules could be released and diffused into the solution during incubation at pH 7.4 due to the swelling of PMs. When pH was decreased to 6.5 or 5.0, the release rate markedly accelerated compared with that at pH 7.4. The accumulative release of PTX was about 20% and 40% at pH 6.5 and 5.0 (for 2 h), and about 85% and 98% at pH 6.5 and 5.0 (for 48 h), respectively. The reason could be that the tertiary amine residues in PAE segment were gradually protonated and the pH-sensitive bonds were broken, inducing the swelling and disassembly of PTX/DOX-PMs. Collectively, the release rates of DOX and PTX were accelerated by the decrease in pH. Therefore, the drugs' (DOX and PTX) release profiles from PTX/DOX-PMs were pH-triggered.

#### Drug Release Mechanism

During preparation of co-delivery system PTX/DOX-PMs, the DOX molecules were chemically linked on the PAE segment, and the PTX molecules were physically loaded in the micellar core. The release profiles were dependent on the pH, but the release mechanism needed to be studied. Here, the drug release mechanism was investigated using a comprehensive semi-empirical model. Based on the experimental data in Figure 4, the drug release process was divided into two stages, one was from 0 h to 12 h, and the other was from 12 h to 48 h. Figure 5 showed the theoretical curves fitted to experimental release data based on the mathematical model. The release exponent (n) and rate constant (k) for two stages of DOX and PTX release profiles were tabulated in Table 2. Good linearity for two stages could be observed in Figure 5. For DOX release mechanism (Figure 5A), the n values for two stages at pH 7.4 were, respectively, 0.279 and 0.362 which were less than 0.43, indicating that the release mechanism corresponded to combination of diffusion and erosion control. Furthermore, the k values for two stages were 0.025 and 0.024, respectively, suggesting the similar drug release rate in two stages over time. At weakly acidic conditions, the n values at first stage were 0.726 and 0.625 at pH 6.5 and 5.0, respectively, which were higher than 0.43 but less than 0.85, indicating the DOX release behavior followed the anomalous transport mechanism in the first 12 h due to the cleavage of pH-sensitive bonds. At the second stage, the *n* values were 0.067 and 0.073 at pH 6.5 and 5.0, respectively, which were less than 0.43, indicating the DOX release mechanism corresponded with the combination of diffusion and erosion control. The results displayed that the free DOX molecules that detached from PAE chains due to the cleavage of pH-sensitive bonds might diffuse from PTX/DOX-PMs over time, as well as swelling and disassembly of PTX/DOX-PMs. With decrease in pH, the k value increased, which suggested the drug release rate accelerated, showing the DOX release profile was pH-dependent which was consistent with the result of in vitro experiment (Figure 4). Specifically, the k values at pH 5.0 were much higher than those at pH 6.5 and 7.4 at two stages, showing the fastest drug release rate. For PTX release mechanism (Figure 5B), the *n* value in the first 12 h at pH 7.4 was 0.431 ( $\approx$  0.43) which indicated that the PTX release from PTX/DOX-PMs corresponded to the Fickian diffusion mechanism. At the second stage, the *n* value was 0.181 (< 0.43), suggesting the PTX release mechanism was changed to the combination of diffusion and erosion control as time went on. At weakly acidic conditions



Figure 5 Plots of log ( $M_t/M_{\infty}$ ) against log t for DOX (**A**) and PTX (**B**) release from PTX/DOX-PMs at pH 7.4, 6.5 and 5.0. **Abbreviations:** DOX, doxorubicin; PTX, paclitaxel; PMs, polymeric micelles; PTX/DOX-PMs, paclitaxel and doxorubicin-loaded polymeric micelles; t, time; h, hour;  $M_t/M_{\infty}$  were the absolute cumulative amount of drug released at time t and infinite time, respectively.

(pH 6.5 and 5.0), the *n* values at the first stage were, respectively, 0.550 and 0.595 (0.43 < n < 0.85), showing the PTX release mechanism corresponded to anomalous transport mechanism during the process of swelling and disassembly of PTX/DOX-PMs. At the second stage, the *n* values were 0.155 and 0.030 (< 0.43), respectively, indicating the PTX release behaviors corresponded with the combination of diffusion and erosion control. The change trend of *k* value was similar to that of DOX release process, demonstrating the PTX release rate accelerated with pH decrease and time increase.

#### Cytotoxicity Assay

Next, the cytotoxic effects of diblock copolymer, free PTX, free DOX, mixture of free DOX/PTX, DOX-PMs and PTX/DOX-PMs for several cell lines were evaluated using MTT assay, as shown in Figures 6 and 7. As seen in Figure 6, the cytotoxicity of diblock copolymer slightly increased with the increase of copolymer concentration, but the cell viability of A549, MDA-MB-231, A2780, and NCL-H460 cells treated

with diblock copolymer was still higher than 90% even at the highest concentration (500 µg/mL). In addition, the cell viability of normal cell line (NIH 3T3, Figure S2) treated with diblock copolymer was also higher than 90% at concentration of 500 µg/mL. These results demonstrated that the diblock copolymer (nanocarrier) showed negligible cytotoxicity. Figure 7 showed the cytotoxicity of free DOX, free PTX, mixture of free DOX/PTX, DOX-PMs and PTX/DOX-PMs against four kinds of tumor cells. As for A549 (Figure 7A), the cell viabilities were, respectively, 20.2%, 12.1%, 8.5%, 9.2% and 5.5% for free PTX, free DOX, mixture of free DOX/PTX, DOX-PMs and PTX/ DOX-PMs at the highest concentration of drugs (20 µg/ mL). The IC50 values were 2.02 µg/mL, 1.21 µg/mL, 0.33 µg/mL, 0.48 µg/mL and 0.25 µg/mL, respectively, for free PTX, free DOX, mixture of free DOX/PTX, DOX-PMs and PTX/DOX-PMs, showing the PTX/DOX-PMs had the highest cytotoxicity in comparison to the other formulations. To further confirm the generality, the cytotoxicities of different formulations for MDA-MB-231 (Figure 7B), A2780

Table 2	Release	Exponent	(n),	Rate	Constant	(k)	for	PTX/DO	X-PM	s in	PBS	at	Different	pН	Values
---------	---------	----------	------	------	----------	-----	-----	--------	------	------	-----	----	-----------	----	--------

PTX/DOX-PMs	рН	n (0–12 h)	k (0–l2 h)	n (12–48 h)	k (12–48 h)
DOX	7.4	0.279±0.015	0.025±0.006	0.362±0.021	0.024±0.007
	6.5	0.726±0.036	0.084±0.011	0.067±0.012	0.609±0.038
	5.0	0.625±0.041	0.209±0.015	0.073±0.013	0.736±0.055
РТХ	7.4	0.432±0.033	0.077±0.017	0.181±0.044	0.198±0.012
	6.5	0.550±0.016	0.098±0.010	0.155±0.036	0.401±0.020
	5.0	0.595±0.022	0.238±0.023	0.030±0.008	0.887±0.039

Abbreviations: DOX, doxorubicin; PTX, paclitaxel; PTX/DOX-PMs, paclitaxel and doxorubicin-loaded polymeric micelles; PBS, phosphate buffer solution; h, hour.



Figure 6 Cell viability of block copolymer after incubating with A549 (A), MDA-MB-231 (B), A2780 (C), and NCL-H460 (D) cells for 48 h (n = 6, mean  $\pm$  SD). Abbreviations: mPEG-b-PAE, diblock copolymer poly(ethylene glycol)methyl ether-b-poly( $\beta$ -amino esters); h, hour.

(Figure 7C) and NCL-H460 (Figure 7D) cells were analyzed. For MDA-MB-231, the IC50 values were 3.32 µg/mL, 2.85 µg/mL, 1.41 µg/mL, 1.49 µg/mL and 0.91 µg/mL, respectively, for free PTX, free DOX, mixture of free DOX/PTX, DOX-PMs and PTX/DOX-PMs. For A2780, the IC50 values were 4.02 µg/mL, 3.11 µg/mL, 1.36 µg/mL, 2.01 µg/mL and 0.65 µg/mL, respectively, for free PTX, free DOX, mixture of free DOX/PTX, DOX-PMs and PTX/DOX-PMs. For NCL-H460, the IC50 values were 0.96 µg/mL, 0.75 µg/mL, 0.54 µg/mL, 0.57 µg/mL and 0.38  $\mu$ g/mL, respectively, for free PTX, free DOX, mixture of free DOX/PTX, DOX-PMs and PTX/DOX-PMs. As a result, the cell viabilities of three kinds of tumor cells were quite low at drug concentration of 20 µg/mL, suggesting the high cytotoxic effect of various drug formulations. Particularly, the cytotoxicity of different drug formulations was in order of free PTX< free DOX< DOX-PMs< free DOX/PTX< PTX/DOX-PMs. Therefore, the findings demonstrated that the PTX/DOX-PMs exhibited the highest cytotoxic effect for tumor cells. These findings showed that the co-delivery system PTX/DOX-PMs could be a promising therapeutic option for cancer treatment with high cytotoxicity.

#### Conclusion

Combination chemotherapy of multiple anticancer drugs has attracted more and more attention due to the improved therapeutic efficacy. In this work, we designed and prepared a co-delivery anticancer drug system based on pH-sensitive copolymer-based polyprodrug (mPEG*b*-PAE-*cis*-DOX) and hydrophobic chemical drug (PTX) for combination chemotherapy. The co-delivery system was able to self-assemble into polymeric micelles (PTX/ DOX-PMs) with suitable particle size (*c.a.* 110 nm) and



Figure 7 Cell viability of free DOX, free PTX, DOX-PMs and PTX/DOX-PMs after incubating with A549 (A), MDA-MB-231 (B), A2780 (C), and NCL-H460 (D) cells for 48 h (n = 6, mean ± SD).

Abbreviations: DOX, doxorubicin; PTX, paclitaxel; PMs, polymeric micelles; DOX-PMs, doxorubicin-loaded polymeric micelles; DOX+PTX, mixture of doxorubicin and paclitaxel; PTX/DOX-PMs, paclitaxel and doxorubicin-loaded polymeric micelles; h, hour.

low PDI. PTX molecules could be efficiently encapsulated into the core by hydrophobic interaction during the process of self-assembly, resulting in the high drug loading content and encapsulation efficacy. The particle size, zeta potential and CMC of PTX/DOX-PMs were increased with the decrease in pH from normal physiological environment to weakly acidic condition, especially disassembly of the system at pH 5.0, demonstrating the pH sensitivity of PTX/DOX-PMs. The PTX and DOX release profiles in vitro from PTX/DOX-PMs were obviously influenced by change in pH, indicating the pH-triggered drug release property. Furthermore, the drug release mechanism for PTX and DOX was studied based on a comprehensive semi-empirical model, proving the drug release was dependent on pH value. Furthermore, the reasonable drug loading method could be selected for "on-demand" drug

delivery. In addition, the cytotoxicity of diblock copolymer, free drugs, single drug delivery system DOX-PMs and co-delivery system PTX/DOX-PMs against different cell lines was extensively studied. The results showed that the PTX/DOX-PMs exhibited the highest cytotoxic effect for tumor cells while the carrier had negligible cytotoxicity. In short, PTX/DOX-PMs could be a promising nanomedicine for combination chemotherapy. By the way, in vivo anticancer efficacies and bio-distributions of PTX/DOX-PMs and other co-delivery systems would be systematically studied in future.

#### Acknowledgments

This work was financially supported by the Liaoning Provincial Natural science Foundation of China (NO.201 80550022).

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Sun CY, Shen S, Xu CF, et al. Tumor acidity-sensitive polymeric vector for active targeted siRNA delivery. J Am Chem Soc. 2015;137 (48):15217–15224. doi:10.1021/jacs.5b09602
- Pearce TR, Shroff K, Kokkoli E. Peptide targeted lipid nanoparticles for anticancer drug delivery. *Adv Mater.* 2012;24(28):3803–3822. doi:10.1002/adma.201200832
- Chen W, Hou Y, Tu Z, Gao L, Haag R. pH-degradable PVA-based nanogels via photo-crosslinking of thermo-preinduced nanoaggregates for controlled drug delivery. *J Control Release*. 2017;259:160–167. doi:10.1016/j.jconrel.2016.10.032
- Luo Y, Yin X, Yin X, et al. Dual pH/redox-responsive mixed polymeric micelles for anticancer drug delivery and controlled release. *Pharmaceutics*. 2019;11(4):176. doi:10.3390/pharmaceutics11040176
- Li Y, Wang H, Wang K, et al. Targeted co-delivery of PTX and TR3 siRNA by PTP peptide modified dendrimer for the treatment of pancreatic cancer. *Small*. 2017;13(2):1602697. doi:10.1002/smll.201602697
- Zhang DX, Yoshikawa C, Welch NG, Pasic P, Thissen H, Voelcker NH. Spatially controlled surface modification of porous silicon for sustained drug delivery applications. *Sci Rep.* 2019;9 (1):1367. doi:10.1038/s41598-018-37750-w
- von Roemeling C, Jiang W, Chan CK, Weissman IL, Kim BYS. Breaking down the barriers to precision cancer nanomedicine. *Trends Biotechnol.* 2017;35(2):159–171. doi:10.1016/j.tibtech.2016.07.006
- Elzoghby AO, Hemasa AL, Freag MS. Hybrid protein-inorganic nanoparticles: from tumor-targeted drug delivery to cancer imaging. *J Control Release*. 2016;243:303–322. doi:10.1016/j.jconrel.2016.10.023
- Huang X, Liao W, Zhang G, Kang S, Zhang CY. pH-sensitive micelles self-assembled from polymer brush (PAE-g-cholesterol)b-PEG-b-(PAE-g-cholesterol) for anticancer drug delivery and controlled release. *Int J Nanomed*. 2017;12:2215–2226. doi:10.2147/IJN. S130037
- Xu M, Zhang CY, Wu J, et al. PEG-detachable polymeric micelles self-assembled from amphiphilic copolymers for tumor-aciditytriggered drug delivery and controlled release. ACS Appl Mater Interfaces. 2019;11(6):5701–5713. doi:10.1021/acsami.8b13059
- Alakhov V, Klinski E, Li S, et al. Block copolymer-based formulation of doxorubicin. from cell screen to clinical trials. *Colloids Surf B*. 1999;16(1–4):113–134. doi:10.1016/S0927-7765(99)00064-8
- Zhao J, Qin Z, Wu J, Li L, Jin J, Ji J. Zwitterionic stealth peptide-protected gold nanoparticles enable long circulation without the accelerated blood clearance phenomenon. *Biomater Sci.* 2018;6 (1):200–206. doi:10.1039/C7BM00747G
- Storm G, Belliot SO, Daemen T, Lasic DD. Surface modification of nanoparticles to oppose uptake by the mononuclear phagocyte system. *Adv Drug Deliver Rev.* 1995;17(1):31–48. doi:10.1016/ 0169-409X(95)00039-A
- Sun X, Wang G, Zhang H, et al. The blood clearance kinetics and pathway of polymeric micelles in cancer drug delivery. *ACS Nano*. 2018;12(6):6179–6192. doi:10.1021/acsnano.8b02830
- Sun X, Yan X, Jacobson O, et al. Improved tumor uptake by optimizing liposome based RES blockade strategy. *Theranostics*. 2017;7 (2):319–328. doi:10.7150/thno.18078
- 16. Feng Q, Zhang Y, Zhang W, et al. Tumor-targeted and multi-stimuli responsive drug delivery system for near-infrared light induced chemo-phototherapy and photoacoustic tomography. *Acta Biomater*. 2016;38:129–142. doi:10.1016/j.actbio.2016.04.024
- Yang K, Feng L, Liu Z. Stimuli responsive drug delivery systems based on nano-graphene for cancer therapy. *Adv Drug Deliver Rev.* 2016;105:228–241. doi:10.1016/j.addr.2016.05.015

- Yin Q, Shen J, Zhang Z, Yu H, Li Y. Reversal of multidrug resistance by stimuli-responsive drug delivery systems for therapy of tumor. *Adv Drug Deliver Rev.* 2013;65(13–14):13–14. doi:10.1016/j.addr.20 13.04.011
- Mura S, Nicolas J, Couvreur P. Stimuli-responsive nanocarriers for drug delivery. *Nat Mater.* 2013;12(11):991–1003. doi:10.1038/nmat 3776
- 20. Le NTT, Nguyen TNQ, Cao VD, Hoang DT, Ngo VC, Thi TTH. Recent progress and advances of multi-stimuli-responsive dendrimers in drug delivery for cancer treatment. *Pharmaceutics*. 2019;11 (11):591. doi:10.3390/pharmaceutics11110591
- Alsuraifi A, Curtis A, Lamprou DA, Hoskins C. Stimuli responsive polymeric systems for cancer therapy. *Pharmaceutics*. 2018;10 (3):136. doi:10.3390/pharmaceutics10030136
- 22. Yang P, Li D, Jin S, et al. Stimuli-responsive biodegradable poly (methacrylic acid) based nanocapsules for ultrasound traced and triggered drug delivery system. *Biomaterials*. 2014;35(6):2079–20 88. doi:10.1016/j.biomaterials.2013.11.057
- Gulzar A, Gai S, Yang P, Li C, Ansari MB, Lin J. Stimuli responsive drug delivery application of polymer and silica in biomedicine. *J Mater Chem B.* 2015;3(44):8599–8622. doi:10.1039/C5TB00757G
- 24. Wu X, Wang Z, Zhu D, et al. pH and thermo dual-stimuli-responsive drug carrier based on mesoporous silica nanoparticles encapsulated in a copolymer–lipid bilayer. ACS Appl Mater Interfaces. 2013;5 (21):10895–10903. doi:10.1021/am403092m
- Ahmad Z, Shah A, Siddiq M, Kraatz HB. Polymeric micelles as drug delivery vehicles. *RSC Adv.* 2014;4(33):17028–17038. doi:10.1039/ C3RA47370H
- 26. Li J, Ma YJ, Wang Y, Chen BZ, Guo XD, Zhang CY. Dual redox/ pH-responsive hybrid polymer-lipid composites: synthesis, preparation, characterization and application in drug delivery with enhanced therapeutic efficacy. *Chem Eng J.* 2018;341:450–461. doi:10.1016/j. cej.2018.02.055
- 27. Cai W, Wang J, Chu C, Chen W, Wu C, Liu G. Metal-organic framework-based stimuli-responsive systems for drug delivery. *Adv Sci.* 2019;6(1):1801526. doi:10.1002/advs.201801526
- 28. Han B, Yang Y, Chen J, et al. Preparation, characterization, and pharmacokinetic study of a novel long-acting targeted paclitaxel liposome with antitumor activity. *Int J Nanomed*. 2020;15:553–571. doi:10.2147/IJN.S228715
- Sarker DK. Engineering of nanoemulsions for drug delivery. Curr Drug Deliv. 2005;2(4):297–310. doi:10.2174/156720105774370267
- Park J, Kadasala NR, Abouelmagd SA, et al. Polymer-iron oxide composite nanoparticles for EPR-independent drug delivery. *Biomaterials*. 2016;101:285–295. doi:10.1016/j.biomaterials.2016.06.007
- Cao H, Chen C, Xie D, et al. A hyperbranched amphiphilic acetal polymer for pH-sensitive drug delivery. *Polym Chem.* 2018;9 (2):169–177. doi:10.1039/C7PY01739A
- 32. Zhao J, Yang Y, Han X, et al. Redox-sensitive nanoscale coordination polymers for drug delivery and cancer theranostics. ACS Appl Mater Interfaces. 2017;9(28):23555–23563. doi:10.1021/acsami.7b 07535
- 33. Zhang CY, Gao J, Wang Z. Bioresponsive nanoparticles targeted to infectious microenvironments for sepsis management. *Adv Mater*. 2018;30(43):1803618. doi:10.1002/adma.201803618
- 34. Rafeeq T, Ana MM, Willian PG, Ghaleb AH. Drug delivery systems based on polymeric micelles and ultrasound: a review. *Curr Pharm Design*. 2016;22(19):2796–2807. doi:10.2174/1381612822666160217125215
- 35. Muley P, Kumar S, Kourati FE, Kesharwani SS, Tummala H. Hydrophobically modified inulin as an amphiphilic carbohydrate polymer for micellar delivery of paclitaxel for intravenous route. *Int J Pharm.* 2016;500(1–2):32–41. doi:10.1016/j.ijpharm.2016.01.005
- 36. Kesharwani SS, Dachineni R, Bhat GJ, Tummala H. Hydrophobically modified inulin-based micelles: transport mechanisms and drug delivery applications for breast cancer. J Drug Deliv Sci Technol. 2019;54:101254. doi:10.1016/j.jddst.2019.101254

- Zhang CY, Yang YQ, Huang TX, et al. Self-assembled pH-responsive MPEG-b-(PLA-co-PAE) block copolymer micelles for anticancer drug delivery. *Biomaterials*. 2012;33(26):6273–6283. doi:10.1016/j. biomaterials.2012.05.025
- 38. Hu X, Liu G, Li Y, Wang X, Liu S. Cell-penetrating hyperbranched polyprodrug amphiphiles for synergistic reductive milieu-triggered drug release and enhanced magnetic resonance signals. J Am Chem Soc. 2015;137(1):362–368. doi:10.1021/ja51 05848
- 39. Zhang CY, Dong X, Gao J, Lin W, Liu Z, Wang Z. Nanoparticleinduced neutrophil apoptosis increases survival in sepsis and alleviates neurological damage in stroke. *Sci Adv.* 2019;5(11):eaaz7964. doi:10.1126/sciadv.aax7964
- 40. Li Y, Lu H, Liang S, Xu S. Dual stable nanomedicines prepared by cisplatin-crosslinked camptothecin prodrug micelles for effective drug delivery. ACS Appl Mater Interfaces. 2019;11(23):20649–20659. doi:10. 1021/acsami.9b03960
- 41. Li Z, Wang H, Chen Y, et al. pH-and NIR light-responsive polymeric prodrug micelles for hyperthermia-assisted site-specific chemotherapy to reverse drug resistance in cancer treatment. *Small.* 2016;12 (20):2731–2740. doi:10.1002/smll.201600365
- Ma Y, Fan X, Li L. pH-sensitive polymeric micelles formed by doxorubicin conjugated prodrugs for co-delivery of doxorubicin and paclitaxel. *Carbohyd Polym.* 2016;137:19–29. doi:10.1016/j.carbpol. 2015.10.050
- 43. Ma Y, He P, Tian X, Liu G, Zeng X, Pan G. Mussel-derived, cancer-targeting peptide as pH-sensitive prodrug nanocarrier. ACS Appl Mater Interfaces. 2019;11(27):23948–23956. doi:10.1021/ acsami.9b09031
- 44. Zou P, Song J, Jiang B, et al. Epigallocatechin-3-gallate protects against cisplatin nephrotoxicity by inhibiting the apoptosis in mouse. *Int J Clin Exp Pathol.* 2014;7(8):4607–4616.
- 45. Duan X, Xiao J, Yin Q, et al. Smart pH-sensitive and temporal-controlled polymeric micelles for effective combination therapy of doxorubicin and disulfiram. ACS Nano. 2013;7(7):585 8–5869. doi:10.1021/nn4010796

- 46. Zhai S, Hu X, Hu Y, Wu B, Xing D. Visible light-induced crosslinking and physiological stabilization of diselenide-rich nanoparticles for redox-responsive drug release and combination chemotherapy. *Biomaterials*. 2017;121:41–54. doi:10.1016/j.biomaterials.2017.01.002
- 47. Yang J, Zhai S, Qin H, Yan H, Xing D, Hu X. NIR-controlled morphology transformation and pulsatile drug delivery based on multifunctional phototheranostic nanoparticles for photoacoustic imaging-guided photothermal-chemotherapy. *Biomaterials*. 2018;176:1–12. doi:10.10 16/j.biomaterials.2018.05.033
- 48. Hu X, Zhai S, Liu G, Xing D, Liang H, Liu S. Concurrent drug unplugging and permeabilization of polyprodrug-gated crosslinked vesicles for cancer combination chemotherapy. *Adv Mater.* 2018;30 (21):1706307. doi:10.1002/adma.201706307
- 49. Zhu H, Chen H, Zeng X, et al. Co-delivery of chemotherapeutic drugs with vitamin E TPGS by porous PLGA nanoparticles for enhanced chemotherapy against multi-drug resistance. *Biomaterials*. 2014;35(7):2391–2400. doi:10.1016/j.biomaterials.2013.11.086
- Danafar H, Rostamizadeh K, Davaran S, Hamidi M. Co-delivery of hydrophilic and hydrophobic drugs by micelles: a new approach using drug conjugated PEG–PCL nanoparticles. *Drug Dev Ind Pharm.* 2017;43 (11):1908–1918. doi:10.1080/03639045.2017.1355922
- Huang X, Liao W, Xie Z, Chen D, Zhang CY. A pH-responsive prodrug delivery system self-assembled from acid-labile doxorubicin-conjugated amphiphilic pH-sensitive block copolymers. *Mat Sci Eng C-Mater*. 2018;90:27. doi:10.1016/j.msec.2018.04.036
- He H, Markoutsa E, Zhan Y, Zhang J, Xu P. Mussel-inspired PLGA/ polydopamine core-shell nanoparticle for light induced cancer thermochemotherapy. *Acta Biomater*. 2017;59:181–191. doi:10.1016/j. actbio.2017.07.005
- Brannon-Peppas L, Blanchette JO. Nanoparticle and targeted systems for cancer therapy. *Adv Drug Deliver Rev.* 2004;56(11):1649–1659. doi:10.1016/j.addr.2004.02.014
- 54. Wang Y, Zhang H, Hao J, Li B, Li M, Wang X. Lung cancer combination therapy: co-delivery of paclitaxel and doxorubicin by nanostructured lipid carriers for synergistic effect. *Drug Deliv.* 2015;23(4):1398–1403. doi:10.3109/10717544.2015.1055619

International Journal of Nanomedicine

#### **Dove**press

Publish your work in this journal

The International Journal of Nanomedicine is an international, peerreviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch<sup>®</sup>, Current Contents<sup>®</sup>/Clinical Medicine, Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/ testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/international-journal-of-nanomedicine-journal