Drug-Related Hypertension Associated with the Efficacy of Apatinib on Hepatocellular Carcinoma [Corrigendum]

Yang X, Hou Z, Zhu K, et al. *Cancer Manag Res.* 2020;12:3163–3173.

tion in the figure caption or figure citations throughout the text.

The authors have advised there is an error with Figure 2 on page 3170. The figure parts C-H do not match the descrip-

The correct Figure 2 is shown below. The authors apologize for this error.

Figure 2 Continued.

Figure 2 Subgroup analysis of OS and PFS. **(A)** Subgroup with AFP $>400\mu\text{g/L}$, whether HTN can be a predictor of OS; **(B)** subgroup with AFP $>400\mu\text{g/L}$, whether HTN can be a predictor of PFS; **(A, B)** In patients with AFP $\leq 400\mu\text{g}$, drug-related HTN can be used as a predictor of OS ($p=0.006$) and PFS ($p=0.021$). In patients with AFP $>400\mu\text{g}$, drug-related HTN cannot be used as a predictor of OS ($p=0.081$), but can be used as a predictor of PFS ($p=0.007$). **(C)** Subgroup with Child-Pugh Score, whether HTN can be a predictor of OS; **(D)** subgroup with Child-Pugh Score, whether HTN can be a predictor of PFS; **(C, D)** in patients with Child-Pugh A, drug-related HTN can be used as a predictor of OS ($p=0.003$) and PFS ($p=0.012$). In patients with Child-Pugh B, drug-related HTN cannot be used as a predictor of OS ($p=0.267$), but can be used as a predictor of PFS ($p=0.010$). **(E)** Subgroup with macrovascular invasion, whether HTN can be a predictor of OS; **(F)** subgroup with macrovascular invasion, whether HTN can be a predictor of PFS; **(E, F)** in patients without macrovascular invasion, drug-related HTN can be used as a predictor of OS ($p=0.024$), but not as a predictor of PFS ($p=0.072$). In patients with macrovascular invasion, drug-related HTN can be used as a predictor of OS ($p=0.021$) and PFS ($p=0.001$). **(G)** Subgroup with extrahepatic spread, whether HTN can be a predictor of OS; **(H)** subgroup with extrahepatic spread, whether HTN can be a predictor of PFS. **(G, H)** In patients without extrahepatic metastases, drug-related HTN cannot be used as a predictor of OS ($p=0.078$), but can be used as a predictor of PFS ($p=0.035$). In patients with extrahepatic spread, drug-related HTN can be used as a predictor of OS ($p=0.005$) and PFS ($p=0.000$).

Cancer Management and Research

Dovepress

Publish your work in this journal

Cancer Management and Research is an international, peer-reviewed open access journal focusing on cancer research and the optimal use of preventative and integrated treatment interventions to achieve improved outcomes, enhanced survival and quality of life for the cancer patient.

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/cancer-management-and-research-journal>