Pathways of Methicillin-Resistant *Staphylococcus aureus* in Animal Model: New Insights Regarding Public Health

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Dalia A Hamza¹
Reham M Abd-Elsalam²
Sara M Nader¹
Mahmoud Elhariri³
Rehab Elhelw³
Heba S El-Mahallawy⁴

¹Department of Zoonoses, Faculty of Veterinary Medicine, Cairo University, Giza, Egypt; ²Department of Pathology, Faculty of Veterinary Medicine, Cairo University, Giza, Egypt; ³Department of Microbiology, Faculty of Veterinary Medicine, Cairo University, Giza, Egypt; ⁴Department of Animal Hygiene, Zoonoses and Animal Behaviour and Management, Faculty of Veterinary Medicine, Suez Canal University, Ismailia, Egypt

Background: *Staphylococcus aureus* is considered one of the major threats regarding food safety worldwide. Methicillin-resistant *S. aureus* (MRSA) strains in livestock, companion animals, and wild animals continue to be a potential risk to people working with them.

Aim: The current research aims to investigate the potential pathways of livestock-associated methicillin-resistant *Staphylococcus aureus* (LA-MRSA) strains in the body after oral infection using the experimental mouse model.

Methods: Seven groups of SPF male mice were purchased and housed. On day 1, six groups of mice were infected orally by the sterile gastric probe using 100 μ L/mice of LA-MRSA bacterial suspension (1×10^8 colony-forming units (CFU)/mL). The remaining group was kept as negative controls. Over 15 days, these animals have been monitored. Fresh fecal samples were screened for LA-MRSA at day 0, day 7 and day 14 following oral administration of MRSA strains. All animals were sacrificed at day 15, and internal organs (liver, lung, kidney, and intestine) were harvested aseptically and divided into two sections. The first part was histopathologically investigated, while the other half has been tested for LA-MRSA re-isolation.

Result: The oral challenge of mice by MRSA strains showed that MRSA was re-isolated from feces and intestines of all inoculated mice groups and from internal organs (liver, lung, kidney and intestine) of most mice. Results were confirmed by the detection of the bacteria in gram-stained tissue sections and changes in H&E-stained histopathological tissue sections from these organs.

Conclusion: Data from the present study indicate the possible colonization of livestock-associated methicillin-resistant *Staphylococcus aureus* (LA-MRSA) in internal organs following oral infection and thus posing a risk for food-borne infection of MRSA. Infected animals could pass LA-MRSA through feces again, resulting in increased dispersion and environmental contamination.

Keywords: *Staphylococcus aureus*, MRSA, animal model, mice, oral challenge, PCR, histopathology

Introduction

Food-borne diseases constitute a public health problem worldwide. To date, there have been more than 250 different foodborne diseases, most of which are infections caused by bacteria, viruses and parasites.

Staphylococcus aureus has a major threat regarding food safety and occupational health and is one of the most common agents incriminated in food poisoning outbreaks worldwide.¹ It is responsible for more than 10% of foodborne outbreaks associated with cheese, milk and other dairy products.²

Correspondence: Dalia A Hamza
Department of Zoonoses, Faculty of Veterinary Medicine, Cairo University, Giza 12211, Egypt
Tel +20 1 009701262
Fax +20 2 35725240
Email daliahmza@cu.edu.eg

Methicillin-resistant *S. aureus* (MRSA) are of public health importance. MRSA infections are associated with a worse prognosis than methicillin-susceptible *S. aureus* infections.^{3,4} Emergence of these resistant strains is due to the acquisition of *mecA* gene encoding Penicillin-Binding Protein 2a (PBP2a), which belongs to the family of enzymes necessary for building the bacterial cell wall.⁵ The presence of methicillin-resistant *S. aureus* strains (MRSA) in food-producing animals and its detection in retail meat samples raises the concern about the potential food-borne transmission of MRSA.⁶

Before the 1990s, the majority of MRSA cases were hospital-associated (HA-MRSA); however, the community-associated MRSA (CA-MRSA) then found to cause infections outside the healthcare environment. The third major emergent type of MRSA has been reported in live-stock animals [livestock-associated MRSA (LA-MRSA)].

This widespread of CA-MRSA and LA-MRSA has raised the question of whether MRSA is a potential foodborne pathogen or not. This prompted researches for determining the origin and pathways of LA-MRSA and its ability to cause zoonotic disease in human.⁷ Furthermore, MRSA is in need to be studied closely in an attempt to control its spread.⁸

Using animal models to study a particular disease whose features closely resemble those of disease in man are necessary in order to understand its pathogenesis and possible pathways. Numerous mouse models have been developed as substitutes for the study of infections with *S. aureus* occurring in humans. These include subcutaneous injection of staphylococci to generate skin and soft tissue infections,⁹ intravenous challenge with staphylococci to induce sepsis,¹⁰ or endocarditis¹¹ and intranasal instillation of staphylococci to induce pneumonia.¹² Our study used an oral-challenged mouse model to study the possible pathways of MRSA strains following oral infection and the understand the consequences of its sources and transmission.

Materials and Methods

Experimental design and protocols for laboratory animal housing and inoculations had been reviewed and approved by the Scientific Research Committee and Bioethics Board of Cairo University, Faculty of Veterinary Medicine, Giza, Egypt.

Bacterial Strains

MRSA strains previously obtained from milk of Mastitic animals (Cattle, buffalo and goat) were used in this study. The used strains were related to Dorgham et al.¹³

Bacterial strains were inoculated onto trypticase soy agar with 5% sheep blood and incubated for 18 to 24 hrs at 35°C.

Bacterial suspension was prepared by mixing the obtained colonies in sterile 0.9% NaCl.

MRSA cells were suspended at a concentration of 1×10^8 colony-forming units (CFU)/mL in saline using McFarland standard.¹⁴

In vivo Infectivity Assays Mice

Four weeks old, SPF male mice weighing 25 to 33 g were purchased. Mice were maintained under standard ethical conditions recommended by the Committee for the Care and Use of Laboratory Animals. Upon arrival, mice were placed and divided into 7 groups (five animals each). Experimental animal groups were individually housed in separate cages and were managed and kept at the same environmental and nutritional conditions. All animals received a common laboratory diet and water and all efforts were made to minimize the suffering of animals throughout the experiment. Fecal samples were collected from all mice and tested for the presence of MRSA strains before beginning of the experiment.

On day 1, six groups of mice were infected orally with 100 μ L of the bacterial suspension/mice using a sterile gastric probe¹⁴ while the remaining group was kept as negative controls.

Infected animals were monitored for morbidity or mortality over a period of 15 days.

Re-Isolation and Identification of MRSA Strains from the Inoculated Mice

Following oral administration of MRSA strains, fresh fecal samples were aseptically collected from mice at day 0, day 7 and day 14 post inoculations by gentle pressure on their abdomens. At day 15, all animals were sacrificed and internal organs (liver, lung, kidney and intestine) were aseptically collected from mice and divided into two parts. The first part was fixed in 10% neutral buffered formalin while the other was plated onto Columbia Agar base with 5% defibrinated sheep blood. All samples were transferred directly to the laboratory for further processing. Fresh fecal and tissue samples were plated onto Columbia Agar base with 5% defibrinated sheep blood (Oxoid, Germany). Test plates were incubated for 24–48 hrs at $37^\circ\text{C} \pm 1^\circ\text{C}$. All isolates were identified based on the colony morphology, Gram staining, coagulase plasma test,

catalase test and occurrence of hemolysis. Additionally, an API-Staph Kit (bioMérieux, Durham, N.C.) was also used for identification of *S. aureus*.

Molecular Confirmation of the Re-Isolated MRSA Strains

A total of 5–10 *S. aureus* colonies were suspended in 200 µL TE buffer. The suspension was incubated for 10 min at 56°C, and then for 10 min at 95°C before being spun at $16000 \times g$ for 2 min. After centrifugation, 5 µL of the supernatant were used as template in a 50 µL PCR reaction.

Molecular confirmation was done by amplification of the *S. aureus*-specific *nuc* gene and *mecA* gene. The primer pairs sequence used in the PCR assays are listed in Table 1.

PCR assay for the detection of *nuc* gene (encoding for the *S. aureus* specific thermonuclease) was performed as previously mentioned.¹⁵ The extracted DNA was amplified for 35 cycles consisted of 30 s at 94°C for denaturation, 30 s at 55°C for annealing and 60 s at 72°C for primer extension and a final extension for 10 min at 72°C. For amplification of the *mecA* gene (encoding for the methicillin-resistant *S. aureus*), PCR conditions included a 4 min initial denaturation at 94°C, followed by 35 cycles of 94°C for 1 min, 55°C for 1 min and 72°C for 1 min and a final extension for 10 min at 72°C.¹⁶ Twenty microliters of the obtained PCR product were then visualized and photographed after being electrophoresed on 1.5% agarose gel.

Histopathological Examination

Tissue samples from liver, kidneys, lungs and intestine collected from the control and inoculated mice groups were routinely processed for histopathological examination to obtain 5 µm sections. The sections were stained with hematoxylin and eosin and Gram's stain and examined under the microscope.¹⁷

Statistical Analysis

Data were analysed with the Chi-square (χ^2) test using PASW Statistics, Version 18.0 software (SPSS Inc., Chicago, IL, USA). A *P*-value <0.05 was considered statistically significant.

Results

Mortality Rate

The mortality rate in all mouse groups was reported throughout the experiment. None of the mice were found dead from the beginning to the end of the experiment.

Isolation, Identification and Molecular Confirmation of Re-Isolated MRSA Strains from Fecal Samples of Mice

Fecal samples from all mice under experiment were free from MRSA before the oral challenge. Following oral administration, MRSA strains were re-isolated through the study period from the collected fecal samples of mice under experiment and completely identified by morphological, biochemical characterization and by molecular amplification of *nuc* and *mecA* genes Figure 1.

Detection of MRSA in Internal Organs (Liver, Lung, Kidney and Intestine) and in Histopathological Sections

MRSA strains were re-isolated from intestine of all mouse groups after oral administration. The results of re-isolation and identification of MRSA strains from liver, lung, kidney and intestine are showed in Table 2. The proportion of LA-MRSA positive samples between different organs (liver, kidney, lung and intestine) showed no significant difference.

Histopathological Changes

No pathological changes had been detected in internal organs of the control group. However, liver of infected mice showed multiple minute focal areas to large patchy areas of hepatocellular necrosis infiltrated with neutrophils

Table 1 Primer Sequences Used for Amplification of (*Nuc*) and (*mecA*) Genes and the Suspected Product Size

Genes	Primer Sequence	Product Size	Reference
<i>nuc</i>	F. 5' GCGATTGATGGTGATACGGTT 3' R. 5'AGCCAAGCCTTGACGAACTAAAGC3'	270 bp	Al-Amery et al ¹⁵
<i>mecA</i>	F. 5' GTGAAGATATACCAAGTGATT 3' R. 5' ATGCGCTATAGATTGAAAGGAT 3'	147 bp	Zhang et al ¹⁶

Figure 1 (A) Amplified PCR products of *nuc* gene at (270 bp). Lane 1: 100 bp ladder; Lanes 2–4: positive to *Staphylococcus aureus*; (B) Amplified PCR products of *mecA* gene at (147 bp). Lane (M) 100 bp ladder; Lanes 1–6: positive to *mecA* gene.

and mononuclear cells (Figure 2B and C). Sinusoidal dilation and kupffer cells activation were also observed. The portal area revealed portal congestion, bile duct hyperplasia and leukocytic aggregations mostly, with neutrophils, macrophages and lymphocytes (Figure 2A and D). Lung tissues revealed thickening of the alveolar wall with dilated perialveolar blood capillaries, inflammatory cells mainly neutrophils and mononuclear cells (Figure 3A). In two cases, severe fibrinopurulent lobar pneumonia was observed (Figure 3B). Massive aggregation of neutrophils, macrophages and necrotic cell debris together with fibrinopurulent exudates were noticed in the alveolar lumen. Some bronchi and bronchioles showed hyperplasia of its epithelial cells with peribronchial and peribronchiolar inflammatory cell aggregation. Kidneys from infected animals showed glomerulonephritis, characterized by mesangial hypercellularity (Figure 3C), the intertubular blood vessels were dilated and the renal tubular epithelium was

vacuolated. Intestine showed sloughing and desquamation of individual enterocytes with increase lamina propria macrophages, lymphocytes and neutrophils (Figure 3D). Some enterocytes were apoptotic showing shrinkage with pyknotic nuclei.

Discussion

S. aureus is frequently colonizing most animal species worldwide; however, the emergence of MRSA strains in several food producing animals, including pigs, cattle, chicken and other animals has a serious impact regarding food safety.¹⁸ Contact with animals is recognized as a risk factor for MRSA carriage. Available data in previous literature proved that MRSA are found colonizing pigs, pig farmers and their families who are in contact with pigs in the Netherlands.¹⁹ Also, people in occupational contact with livestock, eg, farmers, veterinarians and abattoirs workers are frequently exposed and found colonized with LA-MRSA.²⁰ Furthermore, MRSA strains have been detected in different foods for human consumption including bovine milk, cheese, meat products and raw chicken meat.^{21,22}

It is important to understand the LA-MRSA source and dynamics of transmission in order to monitor and prevent the contamination of the LA-MRSA in domestic animals and retail meat.

In the present study, LA-MRSA was re-isolated from feces of all experimentally infected mice, and this may give rise to more environmental contamination. In this regard, in a recent study, LA-MRSA has been recovered from the paws of control mink groups neighboring to other minks infected with LA-MRSA spiked feed within 24 hrs following exposure to contaminated feed due to environmental dispersion.²³

Table 2 Re-Isolated MRSA Strains from Internal Organs Following Oral Challenge of Mice

Mice Groups	MRSA Recovery from Internal Organs			
	Liver + (%)	Lung + (%)	Kidney + (%)	Intestine + (%)
Group I	4 (80)	2 (40)	2 (40)	5 (100)
Group II	4 (80)	3 (60)	2 (40)	5 (100)
Group III	4 (80)	2 (40)	2 (40)	5 (100)
Group IV	5 (100)	2 (40)	1 (20)	5 (100)
Group V	5 (100)	3 (60)	3 (60)	5 (100)
Group VI	4 (80)	1 (20)	2 (40)	5 (100)
Control	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)

Note: (%) Percentage is expressed for each organ within each group.

Abbreviation: MRSA, methicillin-resistant *Staphylococcus aureus*.

Figure 2 Histopathological changes in liver of challenged mice. **(A)** Liver showing patchy area of hepatic cellular necrosis mixed with polymorph nuclear and mononuclear inflammatory cells (arrow), portal congestion, bile duct hyperplasia and leukocytic aggregations mostly, with neutrophils, macrophages and lymphocytes (H&E X400). **(B)** Liver showing multiple focal areas of hepatocellular necrosis with neutrophils (arrows) and few mononuclear cell aggregation, sinusoidal dilation and kupffer cells activation (H&E X400). **(C)** Liver showing focal area of hepatic cellular necrosis (rectangle) mixed with polymorph nuclear, mononuclear inflammatory cells and fragmented nuclei (H&E X400). **(D)** Liver showing intense periportal inflammatory cell aggregation (arrow) (H&E X400). H&E, hematoxylin and eosin stain.

In addition, the presented data from our experiment support the evidence of MRSA colonization in internal organs following oral administration.

MRSA was found colonizing the intestinal mucosa. This may be due to the presence of surface proteins “microbial surface components recognizing adhesive matrix molecules” (MSCRAMMs), which appear to play a key role in initiation of endovascular infections.^{24,25} After bacterial adherence, it will be able to grow by forming biofilm that enable it to evade the host defense mechanism.^{26,27} Deregulated barrier function of the intestinal surface epithelial lining and the ability of MRSA to invade and survive inside the epithelial and endothelial cells believed to be a key factor for mucosal bacterial invasion.²⁸ During infection, *S. aureus* produces numerous enzymes, such as proteases, lipases, and elastases that enable it to invade and destroy host tissues and metastasize to other sites.²⁹ When the bacteria gain access to the bloodstream through invasions of intestinal capillaries, it drains into the portal vein then to general circulation with

dissemination of MRSA to different organs as liver, kidney and lung. This hypothesis was confirmed in our experiment by re-isolation of MRSA and histopathological examination of liver, kidney and lung tissues. This bacterial colonization in the lung and kidney may give rise to shedding of bacteria in respiratory discharge or urinary tract.

LA-MRSA could also be re-isolated at the end of this study from internal organs (Table 2), and all examined feces, this denoted that LA-MRSA would persist in mice for 15 days after oral administration.

In a similar study in Denmark, LA-MRSA was re-isolated from paws and pharynx of minks (*Neovison vison*) after giving them LA-MRSA spiked feed. However, the infected animals were being able to get rid of being carriers after stopping of MRSA administration (Fertner et al 2019b).³⁰ On the other hand, data from our study were denied by the conclusion of Wendlandt et al⁷ who stated that although all types of MRSA may be present in/on human food, it could not be considered as a food-borne pathogen.

Figure 3 Histopathological changes in different organs of challenged mice. (A) Lung showing thickening of the alveolar wall by dilated perialveolar blood capillaries, inflammatory cells mainly neutrophils and mononuclear cells (H&E X400). (B) Lung showing severe fibrinopurulent pneumonia with giant alveoli (H&E X400). (C) Kidney showing glomerular hypertrophy with mesangial hypercellularity (H&E X400). (D) Intestine showing sloughing and desquamation of individual enterocytes with increased lamina propria macrophages, lymphocytes and neutrophils (H&E X200). H&E, hematoxylin and eosin stain.

The findings presented in this study may partly help to answer the controversial question posing a potential health risk; does isolation of MRSA from poultry, beef, and meat products linked to contamination from food handlers and poor hygiene during processing, or it comes from the animal itself.

Hence, effective control measures to prevent dispersion of MRSA begins from good hygienic practices, good manufacturing practices and hazard analysis critical control point for products from animal origin throughout the food chain production system from animal feeding and rearing in the farms to retail facilities.³¹

Generally, limiting the irresponsible use of antimicrobials in veterinary medicine in treatment and their use as growth promoters is very valuable to prevent dispersion of antimicrobial resistance.^{32,33}

Conclusion

The present study revealed that oral administration of MRSA strains in experimental mice model resulted in

their colonization into the internal organs and the infected animals could pass the resistant bacteria through feces again giving rise to more dispersion and environmental contamination. Good hygienic and manufacturing practices throughout all stages of food chain from animal husbandry to consumption of animal products are very valuable in elimination of bacterial contamination from animal feed and subsequently prevent their infection and dispersion of antimicrobial-resistant bacteria.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Hennekinne JA, De Buyser ML, Dragacci S. Staphylococcus aureus and its food poisoning toxins: characterization and outbreak investigation. *FEMS Microbiol Rev*. 2012;36(4):815–836. doi:10.1111/j.1574-6976.2011.00311.x
2. Sasidharan S, Prema B, Latha LY. Antimicrobial drug resistance of Staphylococcus aureus in dairy products. *Asian Pac J Trop Biomed*. 2011;1(2):130–132. doi:10.1016/S2221-1691(11)60010-5

3. Chen C, Fan H, Huang Y, et al. Recombinant lysostaphin protects mice from methicillin-resistant *Staphylococcus aureus* pneumonia. *Biomed Res Int*. 2014;2014.
4. Hesari MR, Salehzadeh A, Darsanaki RK. Prevalence and molecular typing of methicillin-resistant *Staphylococcus aureus* carrying Pantón–Valentine leukocidin gene. *Acta Microbiol Immunol Hung*. 2018;65(1):93–106. doi:10.1556/030.64.2017.032
5. Gajdacs M. The continuing threat of methicillin-resistant *Staphylococcus aureus*. *Antibiotics*. 2019;8(2):52. doi:10.3390/antibiotics8020052
6. Chon J, Sung K, Khan S. Methicillin-resistant *Staphylococcus aureus* (MRSA) in food-producing and companion animals and food products. *Frontiers in Staphylococcus Aureus*. 2017;8:47. doi:10.5772/66645. Available from: <https://www.intechopen.com/books/frontiers-in-staphylococcus-aureus-i-/methicillin-resistant-staphylococcus-aureus-mrsa-in-food-producing-and-companion-animals-and-food-pr>
7. Wendlandt S, Schwarz S, Silley P. Methicillin-resistant *Staphylococcus aureus*: a food-borne pathogen? *Annu Rev Food Sci Technol*. 2013;28(4):117–139. doi:10.1146/annurev-food-030212-182653
8. Gajdacs M, Zsoldiné Urbán E. Epidemiology and resistance trends of *Staphylococcus aureus* isolated from vaginal samples: a 10-year retrospective study in Hungary. *Acta Dermatovenereol Alp Pannonica Adriat*. 2019;28(4):143–147. doi:10.15570/actapa.2019.35
9. Voyich JM, Otto M, Mathema B, et al. Is Pantón–Valentine leukocidin the major virulence determinant in community-associated methicillin-resistant *Staphylococcus aureus* disease? *J Infect Dis*. 2006;194(12):1761–1770. doi:10.1086/509506
10. Kim HK, DeDent A, Cheng AG, et al. IsdA and IsdB antibodies protect mice against *Staphylococcus aureus* abscess formation and lethal challenge. *Vaccine*. 2010;28(38):6382–6392. doi:10.1016/j.vaccine.2010.02.097
11. Panizzi P, Nahrendorf M, Figueiredo JL, et al. In vivo detection of *Staphylococcus aureus* endocarditis by targeting pathogen-specific prothrombin activation. *Nat Med*. 2011;17(9):1142–1146. doi:10.1038/nm.2423
12. Wardenburg JB, Patel RJ, Schneewind O. Surface proteins and exotoxins are required for the pathogenesis of *Staphylococcus aureus* pneumonia. *Infect Immun*. 2007;75(2):1040–1044. doi:10.1128/IAI.01313-06
13. Dorgham SM, Hamza DA, Khairy EA, Hedia RH. Methicillin-resistant staphylococci in mastitic animals in Egypt. *Glob Vet*. 2013;11(6):714–720.
14. Lkhagvadorj E, Nagata S, Wada M, et al. Anti-infectious activity of synbiotics in a novel mouse model of methicillin-resistant *Staphylococcus aureus* infection. *Microbiol Immunol*. 2010;54(5):265–275. doi:10.1111/j.1348-0421.2010.00224.x
15. Al-Amery K, Elhariri M, Elsayed A, et al. Vancomycin-resistant *Staphylococcus aureus* isolated from camel meat and slaughterhouse workers in Egypt. *Antimicrob Resist Infect Control*. 2019;8(1):129. doi:10.1186/s13756-019-0585-4
16. Zhang K, McClure JA, Elsayed S, Louie T, Conly JM. Novel multiplex PCR assay for characterization and concomitant subtyping of staphylococcal cassette chromosome mec types I to V in methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol*. 2005;43(10):5026–5033. doi:10.1128/JCM.43.10.5026-5033.2005
17. Morgan A, Galal MK, Ogaly HA, et al. Tiron ameliorates oxidative stress and inflammation in titanium dioxide nanoparticles induced nephrotoxicity of male rats. *Biomed Pharmacother*. 2017;93:779–787. doi:10.1016/j.biopha.2017.07.006
18. De Neeling AJ, Van den Broek MJ, Spalburg EC, et al. High prevalence of methicillin resistant *Staphylococcus aureus* in pigs. *Vet Microbiol*. 2007;122(3–4):366–372. doi:10.1016/j.vetmic.2007.01.027
19. Van Duijkeren E, Ikawaty R, Broekhuizen-Stins MJ, et al. Transmission of methicillin-resistant *Staphylococcus aureus* strains between different kinds of pig farms. *Vet Microbiol*. 2008;126(4):383–389. doi:10.1016/j.vetmic.2007.07.021
20. European Food Safety Authority. Technical specifications on the harmonised monitoring and reporting of antimicrobial resistance in methicillin-resistant *Staphylococcus aureus* in food-producing animals and food. *EFSA J*. 2012;10(10):2897. doi:10.2903/j.efsa.2012.2897
21. Normanno G, Corrente M, La Salandra G, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) in foods of animal origin product in Italy. *Int J Food Microbiol*. 2007;117(2):219–222. doi:10.1016/j.ijfoodmicro.2007.04.006
22. Dai J, Wu S, Huang J, et al. Prevalence and characterization of *Staphylococcus aureus* isolated from pasteurized milk in China. *Front Microbiol*. 2019;10:641. doi:10.3389/fmicb.2019.00641
23. Fertner M, Pedersen K, Jensen VF, et al. Within-farm prevalence and environmental distribution of livestock-associated methicillin-resistant *Staphylococcus aureus* in farmed mink (Neovison vison). *Vet Microbiol*. 2019;231:80–86. doi:10.1016/j.vetmic.2019.02.032
24. Foster TJ. Immune evasion by staphylococci. *Nat Rev Microbiol*. 2005;3(12):948–958. doi:10.1038/nrmicro1289
25. Gordon RJ, Lowy FD. Pathogenesis of methicillin-resistant *Staphylococcus aureus* infection. *Clin Infect Dis*. 2008;46:S350–9. doi:10.1086/533591
26. Donlan RM, Costerton JW. Biofilms: survival mechanisms of clinically relevant microorganisms. *Clin Microbiol Rev*. 2002;15(2):167–193. doi:10.1128/CMR.15.2.167-193.2002
27. Salehzadeh A, Zamani H, Langeroudi MK, Mirzaie A. Molecular typing of nosocomial *Staphylococcus aureus* strains associated to biofilm based on the coagulase and protein A gene polymorphisms. *Iran J Basic Med Sci*. 2016;19(12):1325.
28. Bettenworth D, Nowacki TM, Friedrich A, Becker K, Wessling J, Heidemann J. Crohn's disease complicated by intestinal infection with methicillin-resistant *Staphylococcus aureus*. *World J Gastroenterol*. 2013;19(27):4418. doi:10.3748/wjg.v19.i27.4418
29. Lowy FD. *Staphylococcus aureus* infections. *N Engl J Med*. 1998;339:520–532. doi:10.1056/NEJM199808203390806
30. Fertner M, Pedersen K, Chriél M. Experimental exposure of farmed mink (Neovison vison) to livestock-associated methicillin-resistant *Staphylococcus aureus* contaminated feed. *Vet Microbiol*. 2019;231:45–47. doi:10.1016/j.vetmic.2019.02.033
31. European Food Safety Authority. Analysis of the baseline survey on the prevalence of methicillin-resistant *Staphylococcus aureus* (MRSA) in holdings with breeding pigs, in the EU, 2008-Part A: MRSA prevalence estimates. *EFSA J*. 2009;7(11):1376. doi:10.2903/j.efsa.2009.1376
32. Sergelidis D, Angelidis AS. Methicillin-resistant *Staphylococcus aureus*: a controversial food-borne pathogen. *Lett Appl Microbiol*. 2017;64(6):409–418. doi:10.1111/lam.12735
33. Gajdacs M, Paulik E, Szabó A. Knowledge, attitude and practice of community pharmacists regarding antibiotic use and infectious diseases: a cross-sectional survey in Hungary (KAPPhA-HU). *Antibiotics*. 2020;9(2):41. doi:10.3390/antibiotics9020041

Infection and Drug Resistance**Dovepress****Publish your work in this journal**

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>