The Role of Erastin in Ferroptosis and Its Prospects in Cancer Therapy

This article was published in the following Dove Press journal:
OncoTargets and Therapy

Yuechen Zhao¹
Yanqing Li²
Ruifeng Zhang¹
Feng Wang³
Tiejun Wang^{1,*}
Yan Jiao^{4,*}

¹Department of Radiation Oncology, The Second Hospital of Jilin University, Changchun, People's Republic of China;

²Department of Pathophysiology, College of Basic Medical Sciences, Jilin University, Changchun, People's Republic of China;

³Department of Oral and Maxillofacial Surgery, Hospital of Stomatology, Jilin University, Changchun, People's Republic of China; ⁴Department of Hepatobiliary and Pancreatic Surgery, The First Hospital of Jilin University, Changchun, People's Republic of China

*These authors contributed equally to this work

Correspondence: Tiejun Wang
Department of Radiation Oncology, The Second Hospital of Jilin University, No. 218 Ziqiang Street, Changchun 130022, People's Republic of China
Tel +8619904449098
Email drwangtiejun@yeah.net

Yan Jiao
Department of Hepatobiliary and Pancreatic Surgery, The First Hospital of Jilin University, No. 71 Xinmin Street, Changchun 130021, People's Republic of China
Tel +8613843101157
Email lagelangrill@126.com

Abstract: Erastin was initially discovered as a small molecule compound that selectively kills tumor cells expressing ST and RAS^{V12} and was later widely investigated as an inducer of ferroptosis. Ferroptosis is a recently discovered form of cell death caused by peroxidation induced by the accumulation of intracellular lipid reactive oxygen species (L-ROS) in an iron-dependent manner. Erastin can mediate ferroptosis through a variety of molecules including the cystine-glutamate transport receptor (system X_C⁻), the voltage-dependent anion channel (VDAC), and p53. Erastin is able to enhance the sensitivity of chemotherapy and radiotherapy, suggesting a promising future in cancer therapy. We hope that this review will help to better understand the role of erastin in ferroptosis and lay the foundation for further research and the development of erastin-based cancer therapies in the future.

Keywords: erastin, ferroptosis, system X_C⁻, p53, VDAC, cancer

Introduction

Ferroptosis is an iron-dependent and non-apoptotic form of cell death defined in 2012. It is characterized by excessive accumulation of lipid peroxides and reactive oxygen species (ROS).¹ Ferroptosis differs from apoptosis, necrosis, and other types of cell death in terms of morphology, genetics, metabolism and molecular biology.² Ferroptosis can occur in many organ systems, such as testes, kidneys, heart, and brain.^{3–5} Current knowledge indicates that ferroptosis occurs during various pathophysiological processes of the body, including degenerative diseases of the central nervous system, the antiviral immune response, arteriosclerosis, acute kidney injury, diabetes, and ischemia-reperfusion injury.⁶ Although ferroptosis plays a vital role in maintaining the survival of normal cells and tissues, it is increasingly recognized that some oncogenic pathways are closely related to ferroptosis, making cancer cells extremely susceptible to ferroptosis.⁷

It has been found that ferroptosis can inhibit the proliferation of malignant cells in liver cancer, pancreatic cancer, prostate cancer, breast cancer, and other cancers.^{8–11} In particular, some highly malignant cancer cells have been proved to be inherently vulnerable to ferroptosis, so inducing ferroptosis may become a new method of cancer treatment.¹² There are two main categories of ferroptosis inducers: the first type can play a role through the cystine-glutamate transporter (system X_C⁻) and includes erastin, sulfasalazine, and glutamate while the second type can directly inhibit glutathione peroxidase (GPX) activity and includes RSL3 and DP17.^{13,14} Among them, erastin differs from other ferroptosis inducers in that the latter usually trigger a single pathway, whereas erastin can trigger multiple molecules and the effect is efficient,

rapid, and lasting.¹⁵ Since naturally non-apoptotic forms are induced, erastin-based cancer treatments promise to bypass the drawbacks of traditional therapies mediated by apoptosis. In this review, we first introduce the basic characteristics of ferroptosis, and then focus in detail on the mechanism and anti-cancer characteristics of erastin in inducing ferroptosis. It is expected to provide the basis for the potential of erastin as an anti-cancer drug in the future.

The Discovery of Erastin

In 2003, Dolma et al used large-scale screening experiments to explore the killing effects of various compounds on cancer cells. They found that camptothecin (CPT) and a novel small molecule compound from combinatorial libraries could selectively kill engineered cancer cells overexpressing Small T oncoprotein (ST) and oncogenic RAS. They named this new compound eradicator of RAS and ST (erastin).¹⁶ (Figure 1) However, unlike the apoptosis induced by CPT, erastin-induced cancer cell death was found to be distinctly different. The classical apoptotic characteristics, including mitochondrial cytochrome c release, caspase 3 activation and DNA fragmentation, were not found in erastin-induced cell death, nor was erastin-induced cell death able to be inhibited by apoptotic inhibitors.^{16–18} Erastin-induced cell death was, therefore, deemed to be a novel and non-apoptotic

form of cell death.¹⁷ Erastin was shown to kill human cancer cells exclusively, rapidly and irreversibly, without affecting normal cells of the same genotype.^{15,16} Dixon et al named the cell death induced by erastin as ferroptosis.¹

Basis of Ferroptosis

As early as the 1990s, Tan et al used glutamate to act on immortalized mouse nerve cells (HT-22) to study the effect of oxidative stress on neuronal cells. It was found that glutamic acid competed for the uptake of cystine, resulting in a decrease in glutathione and eventually oxidative cell death.¹⁹ In 2008, Seiler et al identified lipid peroxidation as the key mediator of cell death in glutathione peroxidase 4 (GPX4) knockout cells. They speculated that GPX4 uses oxidative stress to activate a novel cell death pathway.²⁰ Additional research showed that this type of cell death could not be explained by either apoptosis and necrosis. In 2012, Dixon et al formally defined this mode of cell death as ferroptosis.

Iron plays an important role in many important metabolic processes in the body. Under physiological conditions, iron levels need to be properly balanced in the cell and are mainly regulated by transferrin and ferritin. Excessive ionic iron will cause “iron enrichment” and cause cell death, that is, ferroptosis.²¹ Ferroptosis is an iron-dependent form of cell death characterized by the accumulation of intracellular lipid reactive oxygen species (L-ROS). Reactive oxygen species (ROS) is a collective name for a large class of molecules. They all contain oxygen atoms and are strongly oxidizing. ROS can react with the polyunsaturated fatty acids (PUFAs) of the lipid membrane and induce lipid peroxidation to form L-ROS. High concentrations of L-ROS can trigger oxidative stress in cells, causing oxidative damage.^{22,23} Iron can contribute to the ROS pool in cells through the Fenton reaction, in which iron catalyzes the decomposition of H₂O₂ to generate hydroxyl radicals while enhancing the propagation of phospholipid oxidation and degradation of membrane lipids.²⁴ These all aggravate the formation of L-ROS and oxidative damage to cells.

Ferroptosis differs significantly from other forms of cell death (such as apoptosis, necrosis, and autophagy).^{2,25} In terms of morphology, ferroptotic cells exhibit specific mitochondrial shrinkage and increased mitochondrial membrane density, while other typical characteristics of cell death are absent.^{1,26} In terms of biochemical metabolism, the main manifestation is that ionic iron deposition causes membrane lipid peroxidation and excessive oxidative stress together

2-[1-[4-[2-(4-chlorophenoxy)acetyl]-1-piperazinyl]ethyl]-3-(2-ethoxyphenyl)-4(3H)-Quinazolinone

Figure 1 The chemical structure of erastin.

with the damaged intracellular redox homeostasis, with reduced antioxidant capacity and increased intracellular ROS, eventually lead to oxidative cell death.²⁷ Moreover, this death process can be inhibited by antioxidants and iron chelators.¹ Although many upstream pathways lead to ferroptosis, they lead directly or indirectly to an imbalance of production and degradation of intracellular L-ROS, and eventually, ferroptosis.^{1,28}

The Relevant Pathways of Ferroptosis

Ferroptosis Is Induced by Inhibiting the Cystine-Glutamate Transporter System X_C^-

System X_C^- is a reverse transporter located in the plasma membrane. It is a heterodimer composed of a light chain subunit, xCT, encoded by *SLC7A11*, and functioning as the substrate-specific subunit, and a heavy chain subunit 4F2, encoded by *SLC3A2*, which is common to other amino acid transporters. System X_C^- transfers glutamate out of cells and cystine into cells at a ratio of 1:1.^{29–31} Upon transfer into the cell, cystine is rapidly reduced to cysteine, which is then used in the synthesis of glutathione (GSH), a tripeptide composed of cysteine, glutamate, and glycine. The sulfhydryl structure contained in GSH can be oxidized and dehydrogenated, making GSH an important antioxidant and free radical scavenger in the body.³² GPX is a peroxide-degrading enzyme, and GSH is an essential cofactor in its activation.³³ GPX plays a significant role in maintaining redox homeostasis and protecting cells from lipid oxidative stress leading to death. A variety of ferroptosis inducers can inhibit cystine absorption by inhibiting system X_C^- , resulting in reduced GPX activity. The consequence of this is a reduction in the cell's antioxidant capacity and hence increased L-ROS, ultimately leading to ferroptosis.³⁴ Therefore, inhibition of the cystine-glutamate transporter system X_C^- is an important pathway to induce ferroptosis.

p53 Participates in Ferroptosis

p53 is a classic tumor suppressor that mediates tumor cell cycle arrest, aging, and apoptosis.^{35,36} With the accumulation of research on the mechanisms of cell death, it has been found that p53 not only causes apoptosis, but that activation of p53 also plays an important role in regulating ferroptosis in certain cancer cells.^{37,38} Activation of p53 was found to significantly reduce the expression of *SLC7A11* in cells, up-regulation of p53 reduced expression

of *SLC7A11* at both the protein and mRNA levels and knockdown of the p53 gene eliminated the inhibition of *SLC7A11*.^{39,40} Other studies have further demonstrated that the cell's antioxidant capacity is significantly reduced after p53 gene activation.⁴¹ Zhang et al concluded that the inhibition of *SLC7A11* expression by activation of p53 led to a decrease in system X_C^- activity, which in turn regulated ferroptosis.⁴² In addition to inhibiting the activity of system X_C^- , p53 can also mediate ferroptosis by directly targeting the diamine acetyltransferase *SAT1* and the mitochondrial glutaminase *GLS2* which is involved in the regulation of glutamine metabolism.^{43,44}

However, in some cases, p53 can also reduce cell sensitivity to ferroptosis. Studies have found that p53 activates p21 in a transcription-dependent manner and delays the onset of ferroptosis.^{45,46} In addition, Xie et al found that in colorectal cancer (CRC) cells, p53 can also inhibit ferroptosis by combining with dipeptidyl peptidase-4 (DPP4).⁴⁷ So far, it is believed that p53 is at the core of a powerful signaling network during ferroptosis. On the one hand, p53 can increase the sensitivity of cells to ferroptosis to eliminating abnormal cells and inhibiting tumorigenesis while on the other hand, p53 has another major function in protecting normal cells from various stress factors. When metabolic stress occurs, p53 can both reduce the cells' sensitivity to ferroptosis and protect them, allowing them to maintain normal physiological functions. At present, the mechanism of p53's regulation of ferroptosis under different influencing factors has not been fully studied. The role of p53 in the ferroptosis signaling regulatory network is complex. The specific mechanism of p53 in cancer treatment needs further study.

Other Pathways of Ferroptosis

GPX4 is a member of the GPX family and plays a critical role in maintaining intracellular redox homeostasis. Certain inducers of ferroptosis, such as RSL3 and DP17, have been found to act by direct inhibition of GPX4, leading to a decrease in the cellular antioxidant capacity, and eventually resulting in ferroptosis.¹ The voltage-dependent anion channel (VDAC) is an ion channel located in the outer mitochondrial membrane where it mediates and controls molecular and ion exchange between the mitochondria and the cytoplasm.^{48,49} The permeability of VDAC can be altered by drugs, causing mitochondrial metabolic disorder, ROS production, and subsequent oxidative death.⁵⁰ Under oxidative stress conditions, the transsulfuration pathway transfers a sulfur

atom from methionine to serine, yielding cysteine. The cysteine then acts as a substrate for the synthesis of GSH which assists GPXs in maintaining redox homeostasis and preventing oxidative damage. Therefore, this pathway can inhibit the occurrence of ferroptosis.⁵¹ The ferroptosis-suppressor-protein 1 (FSP1) is an oxidoreductase catalyzing the reduction of ubiquinone (also known as coenzyme Q10, CoQ10). Ubiquinone is a lipophilic free radical scavenger. FSP1 can use NAD(P)H to catalyze the regeneration of CoQ10. In this way, FSP1 can protect the ferroptosis caused by the loss of GPX4. The FSP1-CoQ10-NAD(P)H pathway is an independent parallel system, which cooperates with GPX4 to inhibit ferroptosis caused by the rise of L-ROS.⁵² Nuclear factor erythroid 2-related factor 2 (Nrf2) is also an important regulator of antioxidant response in the body. Under normal conditions, Kelch-like ECH-associated protein 1 (Keap1) promotes the ubiquitination and proteasome degradation of Nrf2. However, under oxidative stress, Keap1 is activated abnormally, which leads to the destruction of the interaction between Nrf2 and antioxidant response elements, thus participating in the regulation of ferroptosis.^{53–55} Heme oxygenase-1 and transferrin are also important sources of intracellular iron and participate in the regulation of ferroptosis.^{43,56}

Erastin, Ferroptosis, and the Mitochondria

VDAC, AIF, and MitoQ

The VDAC proteins are porins with a beta-barrel structure spanning the outer mitochondrial membrane. There are three VDAC isoforms, VDAC1, VDAC2 and VDAC3 and together they make up the most abundant proteins of the outer mitochondrial membrane. The VDAC proteins control the flow of metabolites and respiratory substrates through the outer mitochondrial membrane. These metabolites enter the mitochondrial matrix where they are used for the production of ATP which is dependent upon the maintenance of the mitochondrial membrane potential ($\Delta \Psi$).^{57,58} VDAC can alternate between the states of “open” and “closed”. In the presence of sufficient oxygen, malignant cells will still use glycolysis as a primary source of energy. This is known as the Warburg effect. After VDAC is blocked by tubulin and closed, it restricts the flow of respiratory substrates into the mitochondria. This is conducive to the aerobic glycolysis of cancer cells, leading to the Warburg effect.⁵⁹

There are many molecules involved in oxidative regulation in mitochondrial metabolism. As an important oxidoreductase in the mitochondrial inner membrane, apoptosis-inducing factor (AIF) also participates in the removal of intracellular ROS. Knocking out the expression of AIF will cause a significant increase in intracellular ROS levels.⁶⁰ In addition, mice whose AIF expression level is knocked down by 80–90% are more sensitive to oxidative stress.⁶¹ Therefore, AIF can effectively protect cells against oxidative stress. The mitochondria-targeted ROS scavenger mitoquinone (MitoQ) has powerful antioxidant properties, shown by its reduction of mitochondrial respiration and enhancement of glycolysis, thereby preventing lipid peroxidation, mitochondrial ROS production, and loss of organelle membrane potential. MitoQ is thus responsible for maintaining the integrity and function of the mitochondria. It is one of the most effective molecules preventing ferroptosis in different cell types.⁶²

Erastin as an Antagonist of Tubulin Induces the Opening of VDAC

As early as 2007, Yagoda et al found that erastin can change the permeability of the mitochondrial outer membrane and that VDAC is the target of erastin.^{17,63} Further research showed that erastin can reverse tubulin's inhibition of VDAC. Erastin can prevent and reverse the blockage of VDAC by cytoplasmic free tubulin in vivo and in vitro, allowing VDAC to open.¹⁷ This opening of VDAC leads to three main biological effects: an increase of mitochondrial metabolism (the increase of $\Delta \Psi$), a decrease in glycolysis and an increase of ROS production.⁶⁴ Since glycolysis and the inhibition of mitochondrial metabolism are metabolic characteristics of cancer cells, the promotion of VDAC opening by specific drugs and subsequent ROS production will affect most cancer cells.⁶³

Inhibiting tubulin blockage of VDAC is expected to result in two independent but simultaneous effects: increased oxidative phosphorylation and ATP synthesis with reduced glycolysis leading to a reversal of the Warburg effect (the first hit) and increased ROS formation leading to oxidative stress (the second hit).^{65,66} This anti-Warburg action can cause lethal or sub-lethal damage to cancer cells or can reduce cancer cell proliferation.⁶⁴ In addition, erastin can hyperpolarize mitochondria in cancer cells, which is followed by rapid depolarization, resulting in mitochondrial dysfunction.⁶⁴ One therapeutic advantage of erastin as a VDAC-tubulin antagonist is the specific killing of cancer cells; non-proliferating cells do

not have the high levels of free tubulin characteristic of cancer cells, so VDAC remains functional and is not regulated by free tubulin.⁶⁷

In summary, the regulation of VDAC opening by erastin will have a significant effect on mitochondrial metabolism. This will first increase oxidative phosphorylation and ROS production followed by both indirect regulation of glycolysis and reversal of the Warburg phenotype-promoting aerobic glycolysis. This will increase $\Delta\Psi$, increase mitochondrial ROS and cause oxidative stress, eventually leading to ferroptosis.⁶⁸ Therefore, erastin represents a new pharmacological target which may become a new anti-cancer drug through regulating metabolism. (Figures 2 and 3)

Erastin Inhibits Ferroptosis Induced by System X_C^-

Reina et al found that erastin can cause compensatory transcriptional upregulation of *SLC7A11*.⁶⁷ Overexpression of *SLC7A11* through gene transfection reduced erastin-induced cell death, and inhibition of *SLC7A11* expression increased erastin's anti-cancer activity.¹ Thus, it appears that that erastin can indirectly reduce cellular uptake of cystine by direct inhibition of system X_C^- . Inhibition of system X_C^- by erastin indicates that besides altering the permeability of VDAC, erastin can also activate the classic ferroptosis pathway by acting on the system X_C^- .

When system X_C^- is inhibited, the consequent absence of cysteine, as a substrate for GSH synthesis, will result in diminished levels of GSH. Biochemical and metabolomic analyses

showed that GSH was significantly depleted after erastin treatment.^{13,69} GSH is a necessary cofactor for GPX4 to catalyze the degradation of hydrogen peroxide and hydroperoxide and inhibit the production of L-ROS. Therefore, the inhibition of system X_C^- by erastin indirectly leads to the decrease of GPX4 synthesis and the subsequent decrease of cell antioxidant capacity.⁷⁰ It was found that the activity of GPX4 was decreased in a variety of cancer cells treated with erastin.^{71,72} The cell death caused by erastin inhibition of system X_C^- or GPX4 inactivation involves iron-dependent accumulation of L-ROS and consumption of PUFAs.^{13,69} However, in cancer cells treated with erastin and GPX4-deficient mouse cells, the accumulation of L-ROS, consumption of PUFAs and subsequent cell death can be prevented by treatment with small molecular antioxidants, suggesting that L-ROS-mediated cell damage is essential for ferroptosis induced by erastin.^{69,73}

In conclusion, erastin can prevent extracellular cystine from entering cells by inhibiting system X_C^- , which subsequently reduces the intracellular GSH level. GSH is an indispensable substrate for the antioxidant action of GPX4. Therefore, if the activity of GPX4 is reduced, redox homeostasis breaks down and L-ROS accumulates, leading to oxidative cell death, namely ferroptosis. (Figure 3)

Erastin Exacerbates Ferroptosis by Activating P53

Previous studies have confirmed that activation of the p53 gene can inhibit system X_C^- activity and cause ferroptosis.⁷⁴ Recent findings suggest that erastin is able to activate p53

Figure 2 Erastin induces ferroptosis by altering the permeability of VDAC.

Figure 3 The relevant pathways of ferroptosis induced by erastin.

and thus can enhance ferroptosis. After erastin treatment of lung cancer A549 cells, p53 transcription products were significantly up-regulated and ROS levels were significantly increased. After pretreatment with the ROS scavenger N-acetyl-L-cysteine (NAC), erastin exposure did not significantly affect p53 activation, suggesting that p53 activation depends on the presence of ROS induced by erastin exposure. Therefore, it is not difficult to conclude that erastin treatment results in ROS production followed by p53 activation which subsequently activates the p53 downstream pathway. More importantly, this process forms a feedback loop: erastin causes an increase in ROS, which then leads to the activation of p53, which in turn, causes increased ROS. This exacerbates the key cytotoxic and cytostatic effects of erastin on A549 cells and eventually causes ferroptosis. However, this

effect of erastin has not been found in normal lung cells, suggesting that it is specific for cancer cells⁷⁵ (Figure 3).

In 2015, Jiang et al constructed p53^{3KR} mutant cells deficient in acetylation. These cells had lost the classic p53 function of inducing cell cycle arrest and apoptosis but had retained the ability to inhibit the transcription of *SLC7A11*. When erastin was used to treat the p53^{3KR} mutant and p53-deleted cells separately, the mortality of p53^{3KR} mutant cells was very high (> 90%) in contrast to that of the p53-deleted cells ($\leq 10\%$). However, if the p53^{3KR} mutant cells overexpressed *SLC7A11*, erastin treatment resulted in a significant reduction in the cell death rate (20%).⁴⁰ Wang et al constructed the p53^{4KR98} model based on the p53^{3KR} mutant cells. The p53^{4KR98} model lost both the classical function of p53 and the ability to inhibit

SLC7A11 transcription. Erastin treatment significantly reduced both the cell death rate and tumor inhibition function of the p53^{4KR98} model.³⁸ These results suggest that the activation of p53 by erastin may play an important role in tumor inhibition by inhibiting *SLC7A11* transcription and eventually ferroptosis.

Xie et al found that p53 wild-type CRC cells were not sensitive to erastin, but the sensitivity of CRC cells to erastin recovered after the inhibition of p53 activity by drugs or gene knockout. This is different from the previously documented effects of erastin on ferroptosis in other cancer cells.⁴⁷ As discussed above, the regulatory effect of p53 on ferroptosis is related to cancer cell types. The role of erastin in the activation of p53 and in increasing the sensitivity to ferroptosis is not applicable in all cells. However, this provides a broad scope for future research: to understand the regulatory effect of erastin on ferroptosis in the p53 pathway would be helpful, firstly, in identifying specific targets for the induction of cancer cells' death and, secondly, to inhibit ferroptosis of normal cells, to reduce the side-effects of chemotherapy. At present, it is unclear to what degree the p53 gene is involved in erastin-induced ferroptosis in cancer cells, requiring further study in the future.

Regulation of Lipid Metabolism by Erastin

Lipoxygenases (LOX) are non-heme iron dioxygenase, which can catalyze diallyl site oxygenation of polyunsaturated fatty acids in cell non-bilayer phospholipids. LOX-5 is a well-studied LOX isozyme and is a key enzyme for the synthesis of many highly active oxidized lipids. LOX-5-mediated polyunsaturated fatty acid oxidation plays an important role in ferroptosis.⁷⁶ Acyl-CoA long-chain synthetases are ligases responsible for the oxidation of long-chain fatty acids. One member of this family, ACSL 4, is expressed on the endoplasmic reticulum and mitochondrial outer membrane and is mainly responsible for the catalysis of lipids to form acetyl-CoA.⁷⁷ Research shows that ACSL4 is highly involved in ferroptosis. ACSL4 is involved in the synthesis of negatively charged membrane phospholipids such as phosphatidylethanolamine and phosphatidylinositol. They play an important role in lipid metabolism by incorporating polyunsaturated fatty acids into the cell membrane.^{52,78} Therefore, ACSL4 plays an important role in the formation of ROS mediated by LOX, thus promoting ferroptosis.⁷⁶ Knockout of the GPX4 gene can lead to ferroptosis, while

the double knockout of the GPX4 and ACSL4 genes can reverse GPX4 knockout-induced ferroptosis. This indicates that ACSL4 is necessary for the process of ferroptosis.⁷⁹ In addition, the expression of ACSL4 in ferroptosis-resistant cells was lower than that in ferroptosis-sensitive cells. Therefore, ACSL4 was also used as an indicator of ferroptosis sensitivity.⁸⁰

Yuan et al found that HepG2 (human liver cancer cells) and HL60 (human promyelocytic leukemia cells) cells are highly sensitive to ferroptosis caused by erastin compared with LNCaP (human prostate cancer cells) and K562 (human chronic myeloid leukemia cells). The expression of ACSL4 mRNA and protein in HepG2 and HL60 cells were relatively high. After overexpression of ACSL4 in LNCaP and K562 cells, the cells' sensitivity to cell death induced by erastin was significantly increased.⁸⁰ This suggests that erastin regulates lipid peroxidation by regulating ACSL4, which leads to ferroptosis. In addition, an inhibitor of LOX-5, Zileuton, can inhibit erastin-induced ferroptosis by inhibiting the production of cytoplasmic ROS in HT22 cells.⁸¹ So we speculate that erastin can regulate ferroptosis by regulating pathways other than GPX4 and affecting lipid metabolism and, more importantly, because ACSL4 is overexpressed in several different cancers, such as breast cancer, prostate cancer, colon cancer, and hepatocellular carcinoma.^{82–85} This suggests that the induction of erastin is a specific anti-cancer pathway, only acting on cancer cells, and protecting normal cells from ferroptosis.

Pharmacodynamics and Safety Evaluation of Erastin

Due to its poor water solubility and unstable metabolism in the body, erastin is not suitable for direct use in vivo. Introducing other chemical groups into the aniline ring of erastin can result in compounds that are more soluble, stable, and better suited for in vivo administration. Examples of these include piperazine-erastin (PE) and imidazole ketone erastin (IKE).

Yang et al investigated the effects of PE on tumors in nude mice. They observed a significant reduction in tumor growth with no adverse effects or toxicity even at very high PE doses (60 mg/kg).¹³ A study by Zhang et al using IKE treatment of a B cell lymphoma xenograft model reported stimulation of ferroptosis and inhibition of tumor growth with no adverse effects. The use of nano-carriers to enhance efficacy and selective delivery resulted

in stronger anti-tumor effects, also with no significant toxicity.⁸⁶ A further nanoparticle study by Li et al using ferritin-bound erastin and rapamycin also observed significantly controlled tumor growth with no obvious side effects.⁸⁷

Other *in vivo* experiments have also shown that intraperitoneal injection of erastin analogs in tumor-bearing mice can significantly inhibit the growth of subcutaneous tumors in mice, and that the dose is well tolerated. Pharmacodynamic and toxicological studies have shown that according to the ratio of body surface area, erastin analogs are well tolerated at the indicated treatment dosages and thus have significant therapeutic potential.^{88,89} Zille et al also believe that not only does erastin itself not have toxic effects, but it may prevent toxicity of the tumor to the central nervous system.⁹⁰

In summary, the above studies confirm that erastin analogs can inhibit tumor growth *in vivo* and have minimal toxic and side effects. However, the use of erastin analogs alone is not enough to completely restrict the rapid growth of tumors *in vivo*. Based on current research results, combining erastin with other treatments such as radiotherapy and chemotherapy, or designing erastin analogues with higher bioavailability, greater metabolic stability, and more effective tumor invasion and accumulation rates will further optimize the therapeutic effect and reduce possible toxic and side effects.⁸⁶ It is worth noting that although current *in vivo* experiments with erastin provide very promising results, there is a need for further accurate pharmacokinetic and toxicological studies to provide a platform for further clinical trials in the future.

The Potential of Erastin in Clinical Applications In Chemotherapy

Chemotherapy is one of the three main methods for the treatment of malignant tumors. However, due to the continuous and extensive use of chemotherapeutic drugs, tumors show different degrees of drug resistance.^{91,92} This drug resistance of tumors to chemotherapy is a major factor leading to the failure of chemotherapy and poor prognosis.⁹³ Chemotherapeutic drugs eliminate cancer cells mainly by inducing apoptosis. Previous studies have confirmed that suppressed apoptosis or reduced susceptibility to apoptosis is an important mechanism of acquired drug resistance.⁹⁴ So, can we reverse drug resistance by other non-apoptotic cell death methods?

As described above, erastin can induce cancer cell death by a non-apoptotic and iron-dependent form of cell death. In addition to inducing cancer cell death itself, erastin can also be combined with chemotherapeutic drugs to enhance cancer cell sensitivity to chemotherapeutic drugs.⁹⁵ Erastin has been shown to enhance the sensitivity of lung cancer cells to cisplatin,⁹⁶ rhabdomyosarcoma cells to doxorubicin and actinomycin D,⁹⁷ glioblastoma cells to temozolomide,⁹⁸ for example. In addition, erastin can also eliminate the resistance of many types of chemotherapeutic resistant cells: it has been found to overcome the resistance of head and neck cancer cells and ovarian cancer cells to cisplatin,^{15,99} and the resistance of non-RAS-expressing acute myeloid leukemia cells to cytarabine and doxorubicin hydrochloride (Adriamycin).⁷¹ These results support the feasibility of using erastin as an anti-cancer drug in the clinic.

System X_C^- is strongly linked to drug resistance. The transport of system X_C^- causes an increase in intracellular GSH concentration, which has been confirmed to be one of the causes of chemotherapy resistance in tumor cells.¹⁰⁰ Therefore, system X_C^- can be a powerful and potential therapeutic target to overcome the drug resistance of cancer cells.^{101,102} The inhibitory effect of sulfasalazine on system X_C^- has been demonstrated in small cell lung cancer,¹⁰¹ liver cancer,¹⁰¹ genitourinary tract cancer,¹⁰³ and rectal cancer.¹⁰⁴ Erastin has a much stronger inhibitory effect on system X_C^- than other system X_C^- inhibitors such as sulfasalazine, and is effective at low concentrations, so has the potential of reversing tumor resistance.¹⁰⁵

In conclusion, there is convincing evidence for erastin's potential as an anti-cancer drug. It can be used as a new type of chemotherapeutic drug leading to cellular ferroptosis, as well as a chemotherapeutic sensitizer for various types of human cancer. It is thus an effective candidate drug.

In Radiotherapy

Radiation therapy is the second most crucial treatment for malignant tumors, second only to surgery. About 50–70% of patients with malignant tumors require radiotherapy during treatment.¹⁰⁶ However, potential organ damage is an insurmountable dose-limiting factor in radiotherapy. It is inevitable that some radiotoxic side effects may occur during or after radiotherapy.¹⁰⁷ In this context, improvement of radiotherapy efficacy as much as possible without increasing the dose has become an important way to break through these bottlenecks and is an urgent problem to be solved in the field of cancer radiotherapy.

Radiosensitizers can enhance damage to tumor tissues by promoting tumor cell apoptosis, regulating the cell cycle, accelerating DNA damage, and generating free radicals, thereby improving the efficacy of radiotherapy.¹⁰⁸ They can thus improve the therapeutic effects without increasing the dose of radiation.¹⁰⁹ Erastin increases the sensitivity of cancer cells to radiation besides its known induction of ferroptosis. Cobler et al found that erastin can increase the sensitivity of breast cancer cells to γ -rays in vivo and in vitro by inhibiting system X_C^- , and thought that erastin might prolong the duration of radiation-induced DNA damage.¹¹⁰ Other studies also found that erastin enhanced X-ray-induced cell death of cervical cancer and lung cancer, and demonstrated the same effect in tumor-bearing mice.^{111,112} More advantageously, most normal cells do not express *SLC7A11*,¹¹⁰ so erastin may specifically increase the sensitivity of cancer cells to radiation, thereby increasing the death or proliferation of cancer cells and preventing radiation damage in normal cells. Cisplatin is a known radiosensitizer widely used in the clinic.¹¹³ Erastin can increase the sensitivity of many cancer cells to cisplatin. Whether the combination of the two drugs will produce an additive effect and increase the sensitivity of cancer cells rapidly remains to be investigated. On the other hand, ionizing radiation has some effect on promoting the production of ROS mainly by destroying cellular DNA and causing cell damage.¹¹⁴ As discussed above, the most significant feature of ferroptosis induced by erastin is the increase of ROS in cells. If erastin is used as a radiosensitizer, cancer cells can produce ROS through many other pathways besides ionizing radiation. Whether this effect will lead to the rapid increase of ROS in cells leading to the aggravation of cell peroxidation and death provides us with a reasonable hypothesis.

In conclusion, erastin can be used as a novel radiosensitizer to enhance the radiosensitivity of tumors, increase the radiosensitivity of radiation-resistant tumors, or reduce the radiation dose of normal tissues. It has excellent prospects for clinical application.

Conclusion

Erastin is a small molecule compound that can specifically kill human cancer cells without affecting normal cells of the same genotype, and this process is rapid and irreversible. Erastin, as a ferroptosis inducer, is different from other ferroptosis inducers which usually trigger a single pathway. Erastin can trigger multiple pathways: inhibits the action of the cystine-glutamate transport of system X_C^- , acts on

VDAC to relieve the inhibitory effect of tubulin on VDAC, and may indirectly inhibit system X_C^- by activating p53, leading to ferroptosis. Erastin is more effective and fast-acting than other ferroptosis inducers, is effective at low concentrations and has long-lasting results. More importantly, erastin has great potential as a novel anti-cancer drug. Erastin can enhance the sensitivity of many cancer cells to various chemotherapeutic drugs and enhance the sensitivity of cancer cells to radiation. It can, therefore, be used as a new type of chemotherapy drug or chemotherapy sensitizer and radiotherapy sensitizer in cancer therapy. However, given the insufficient number of studies on erastin, further basic and clinical investigations should be conducted.

Funding

This study was funded by Special Project of Medical and Health Professionals of Jilin Province, China (NO.3D5197457429), Department of Science and Technology of Jilin Province, China (20160101119JC) and Department of Finance of Jilin Province, China (Construction Project of Difficulty Gynecologic Oncology Hierarchical Diagnosis and Treatment Medical Association and Precision Radiotherapy Training Base). The funders had no role in study design, decision to publish, or preparation of the manuscript.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Dixon SJ, Lemberg KM, Lamprecht MR, et al. Ferroptosis: an iron-dependent form of nonapoptotic cell death. *Cell*. 2012;149(5):1060–1072. doi:10.1016/j.cell.2012.03.042
2. Yu H, Guo P, Xie X, Wang Y, Chen G. Ferroptosis, a new form of cell death, and its relationships with tumorous diseases. *J Cell Mol Med*. 2017;21(4):648–657. doi:10.1111/jcmm.13008
3. Toyokuni S. Iron addiction with ferroptosis-resistance in asbestos-induced mesothelial carcinogenesis: toward the era of mesothelioma prevention. *Free Radic Biol Med*. 2019;133:206–215. doi:10.1016/j.freeradbiomed.2018.10.401
4. Weiland A, Wang Y, Wu W, et al. Ferroptosis and its role in diverse brain diseases. *Mol Neurobiol*. 2019;56(7):4880–4893. doi:10.1007/s12035-018-1403-3
5. Hu Z, Zhang H, Yang SK, et al. Emerging role of ferroptosis in acute kidney injury. *Oxid Med Cell Longev*. 2019;2019:8010614. doi:10.1155/2019/8010614
6. Hao S, Liang B, Huang Q, et al. Metabolic networks in ferroptosis. *Oncol Lett*. 2018;15(4):5405–5411. doi:10.3892/ol.2018.8066
7. Friedmann Angeli JP, Krysko DV, Conrad M. Ferroptosis at the crossroads of cancer-acquired drug resistance and immune evasion. *Nat Rev Cancer*. 2019;19(7):405–414. doi:10.1038/s41568-019-0149-1

8. Nie J, Lin B, Zhou M, Wu L, Zheng T. Role of ferroptosis in hepatocellular carcinoma. *J Cancer Res Clin Oncol*. 2018;144(12):2329–2337. doi:10.1007/s00432-018-2740-3
9. Yamaguchi Y, Kasukabe T, Kumakura S. Piperlongumine rapidly induces the death of human pancreatic cancer cells mainly through the induction of ferroptosis. *Int J Oncol*. 2018;52(3):1011–1022. doi:10.3892/ijo.2018.4259
10. Ma S, Dielschneider RF, Henson ES, et al. Ferroptosis and autophagy induced cell death occur independently after siramesine and lapatinib treatment in breast cancer cells. *PLoS One*. 2017;12(8):e0182921. doi:10.1371/journal.pone.0182921
11. Viswanathan VS, Ryan MJ, Dhruv HD, et al. Dependency of a therapy-resistant state of cancer cells on a lipid peroxidase pathway. *Nature*. 2017;547(7664):453–457. doi:10.1038/nature23007
12. Zou Y, Palte MJ, Deik AA, et al. A GPX4-dependent cancer cell state underlies the clear-cell morphology and confers sensitivity to ferroptosis. *Nat Commun*. 2019;10(1):1617. doi:10.1038/s41467-019-09277-9
13. Yang WS, SriRamaratnam R, Welsch ME, et al. Regulation of ferroptotic cancer cell death by GPX4. *Cell*. 2014;156(1–2):317–331. doi:10.1016/j.cell.2013.12.010
14. Liang C, Zhang X, Yang M, Dong X. Recent progress in ferroptosis inducers for cancer therapy. *Adv Mater*. 2019;31(51):e1904197. doi:10.1002/adma.201904197
15. Sato M, Kusumi R, Hamashima S, et al. The ferroptosis inducer erastin irreversibly inhibits system xc⁻ and synergizes with cisplatin to increase cisplatin's cytotoxicity in cancer cells. *Sci Rep*. 2018;8(1):968. doi:10.1038/s41598-018-19213-4
16. Dolma S, Lessnick SL, Hahn WC, Stockwell BR. Identification of genotype-selective antitumor agents using synthetic lethal chemical screening in engineered human tumor cells. *Cancer Cell*. 2003;3(3):285–296. doi:10.1016/S1535-6108(03)00050-3
17. Yagoda N, von Rechenberg M, Zaganjori E, et al. RAS-RAF-MEK-dependent oxidative cell death involving voltage-dependent anion channels. *Nature*. 2007;447(7146):864–868. doi:10.1038/nature05859
18. Yang WS, Stockwell BR. Synthetic lethal screening identifies compounds activating iron-dependent, nonapoptotic cell death in oncogenic-RAS-harboring cancer cells. *Chem Biol*. 2008;15(3):234–245. doi:10.1016/j.chembiol.2008.02.010
19. Tan S, Wood M, Maher P. Oxidative stress induces a form of programmed cell death with characteristics of both apoptosis and necrosis in neuronal cells. *J Neurochem*. 1998;71(1):95–105. doi:10.1046/j.1471-4159.1998.71010095.x
20. Seiler A, Schneider M, Forster H, et al. Glutathione peroxidase 4 senses and translates oxidative stress into 12/15-lipoxygenase dependent- and AIF-mediated cell death. *Cell Metab*. 2008;8(3):237–248. doi:10.1016/j.cmet.2008.07.005
21. Doll S, Conrad M. Iron and ferroptosis: a still ill-defined liaison. *IUBMB Life*. 2017;69(6):423–434. doi:10.1002/iub.1616
22. Moloney JN, Cotter TG. ROS signalling in the biology of cancer. *Semin Cell Dev Biol*. 2018;80:50–64. doi:10.1016/j.semcdb.2017.05.023
23. Zhang J, Wang X, Vikash V, et al. ROS and ROS-mediated cellular signaling. *Oxid Med Cell Longev*. 2016;2016:4350965. doi:10.1155/2016/4350965
24. He J, Yang X, Men B, Wang D. Interfacial mechanisms of heterogeneous fenton reactions catalyzed by iron-based materials: a review. *J Environ Sci (China)*. 2016;39:97–109. doi:10.1016/j.jes.2015.12.003
25. Djulbegovic MB, Uversky VN. Ferroptosis - an iron- and disorder-dependent programmed cell death. *Int J Biol Macromol*. 2019;135:1052–1069. doi:10.1016/j.ijbiomac.2019.05.221
26. Santini SJ, Cordone V, Falone S, et al. Role of mitochondria in the oxidative stress induced by electromagnetic fields: focus on reproductive systems. *Oxid Med Cell Longev*. 2018;2018:5076271. doi:10.1155/2018/5076271
27. Conrad M, Pratt DA. The chemical basis of ferroptosis. *Nat Chem Biol*. 2019;15(12):1137–1147. doi:10.1038/s41589-019-0408-1
28. Imai H, Matsuoka M, Kumagai T, Sakamoto T, Koumura T. Lipid peroxidation-dependent cell death regulated by GPX4 and ferroptosis. *Curr Top Microbiol Immunol*. 2017;403:143–170. doi:10.1007/82_2016_508
29. Lin CH, Lin PP, Lin CY, et al. Decreased mRNA expression for the two subunits of system xc⁻, SLC3A2 and SLC7A11, in WBC in patients with schizophrenia: evidence in support of the hypo-glutamatergic hypothesis of schizophrenia. *J Psychiatr Res*. 2016;72:58–63. doi:10.1016/j.jpsychires.2015.10.007
30. Koppula P, Zhang Y, Zhuang L, Gan B. Amino acid transporter SLC7A11/xCT at the crossroads of regulating redox homeostasis and nutrient dependency of cancer. *Cancer Commun (Lond)*. 2018;38(1):12. doi:10.1186/s40880-018-0288-x
31. Ruiu R, Rolih V, Bolli E, et al. Fighting breast cancer stem cells through the immune-targeting of the xCT cystine-glutamate antiporter. *Cancer Immunol Immunother*. 2019;68(1):131–141. doi:10.1007/s00262-018-2185-1
32. Paul BD, Sbodio JI, Snyder SH. Cysteine metabolism in neuronal redox homeostasis. *Trends Pharmacol Sci*. 2018;39(5):513–524. doi:10.1016/j.tips.2018.02.007
33. Ekoue DN, He C, Diamond AM, Bonini MG. Manganese superoxide dismutase and glutathione peroxidase-1 contribute to the rise and fall of mitochondrial reactive oxygen species which drive oncogenesis. *Biochim Biophys Acta Bioenerg*. 2017;1858(8):628–632. doi:10.1016/j.bbabi.2017.01.006
34. Cao JY, Dixon SJ. Mechanisms of ferroptosis. *Cell Mol Life Sci*. 2016;73(11–12):2195–2209. doi:10.1007/s00018-016-2194-1
35. Anbarasan T, Bourdon JC. The emerging landscape of p53 isoforms in physiology, cancer and degenerative diseases. *Int J Mol Sci*. 2019;20(24). doi:10.3390/ijms20246257
36. Pitolli C, Wang Y, Candi E, Shi Y, Melino G, Amelio I. p53-mediated tumor suppression: DNA-damage response and alternative mechanisms. *Cancers (Basel)*. 2019;11(12):1983. doi:10.3390/cancers11121983
37. Saint-Germain E, Mignacca L, Vernier M, Bobbala D, Ilangumaran S, Ferbeyre G. SOCS1 regulates senescence and ferroptosis by modulating the expression of p53 target genes. *Aging (Albany N Y)*. 2017;9(10):2137–2162.
38. Wang SJ, Li D, Ou Y, et al. Acetylation is crucial for p53-mediated ferroptosis and tumor suppression. *Cell Rep*. 2016;17(2):366–373. doi:10.1016/j.celrep.2016.09.022
39. Gupta AK, Bharadwaj M, Kumar A, Mehrotra R. Spiro-oxindoles as a promising class of small molecule inhibitors of p53-MDM2 interaction useful in targeted cancer therapy. *Top Curr Chem (Cham)*. 2017;375(1):3. doi:10.1007/s41061-016-0089-0
40. Jiang L, Kon N, Li T, et al. Ferroptosis as a p53-mediated activity during tumour suppression. *Nature*. 2015;520(7545):57–62. doi:10.1038/nature14344
41. Jiang L, Hickman JH, Wang SJ, Gu W. Dynamic roles of p53-mediated metabolic activities in ROS-induced stress responses. *Cell Cycle*. 2015;14(18):2881–2885. doi:10.1080/15384101.2015.1068479
42. Zhang W, Gai C, Ding D, Wang F, Li W. Targeted p53 on small-molecules-induced ferroptosis in cancers. *Front Oncol*. 2018;8:507. doi:10.3389/fonc.2018.00507
43. Kang R, Kroemer G, Tang D. The tumor suppressor protein p53 and the ferroptosis network. *Free Radic Biol Med*. 2019;133:162–168. doi:10.1016/j.freeradbiomed.2018.05.074

44. Ou Y, Wang SJ, Li D, Chu B, Gu W. Activation of SAT1 engages polyamine metabolism with p53-mediated ferroptotic responses. *Proc Natl Acad Sci U S A*. 2016;113(44):E6806–E6812. doi:10.1073/pnas.1607152113
45. Tarangelo A, Magtanong L, Biegging-Rolett KT, et al. p53 suppresses metabolic stress-induced ferroptosis in cancer cells. *Cell Rep*. 2018;22(3):569–575. doi:10.1016/j.celrep.2017.12.077
46. Tarangelo A, Dixon S. The p53-p21 pathway inhibits ferroptosis during metabolic stress. *Oncotarget*. 2018;9(37):24572–24573. doi:10.18632/oncotarget.25362
47. Xie Y, Zhu S, Song X, et al. The tumor suppressor p53 limits ferroptosis by blocking DPP4 activity. *Cell Rep*. 2017;20(7):1692–1704. doi:10.1016/j.celrep.2017.07.055
48. Becker T, Wagner R. Mitochondrial outer membrane channels: emerging diversity in transport processes. *Bioessays*. 2018;40(7):e1800013. doi:10.1002/bies.201800013
49. Mazure NM. VDAC in cancer. *Biochim Biophys Acta Bioenerg*. 2017;1858(8):665–673. doi:10.1016/j.bbabi.2017.03.002
50. Maldonado EN. VDAC-tubulin, an anti-warburg pro-oxidant switch. *Front Oncol*. 2017;7:4. doi:10.3389/fonc.2017.00004
51. McBean GJ. The transsulfuration pathway: a source of cysteine for glutathione in astrocytes. *Amino Acids*. 2012;42(1):199–205. doi:10.1007/s00726-011-0864-8
52. Doll S, Freitas FP, Shah R, et al. FSP1 is a glutathione-independent ferroptosis suppressor. *Nature*. 2019;575(7784):693–698. doi:10.1038/s41586-019-1707-0
53. Liu Z, Dong W, Yang B, et al. Tetrachlorobenzoquinone-induced Nrf2 confers neuron-like PC12 cells resistance to endoplasmic reticulum stress via regulating glutathione synthesis and protein thiol homeostasis. *Chem Res Toxicol*. 2018;31(11):1230–1239. doi:10.1021/acs.chemrestox.8b00209
54. Fan Z, Wirth AK, Chen D, et al. Nrf2-keap1 pathway promotes cell proliferation and diminishes ferroptosis. *Oncogenesis*. 2017;6(8):e371. doi:10.1038/oncsis.2017.65
55. Sun X, Ou Z, Chen R, et al. Activation of the p62-keap1-NRF2 pathway protects against ferroptosis in hepatocellular carcinoma cells. *Hepatology*. 2016;63(1):173–184. doi:10.1002/hep.28251
56. Chiang S-K, Chen S-E, Chang L-C. A dual role of heme oxygenase-1 in cancer cells. *Int J Mol Sci*. 2018;20(1):39. doi:10.3390/ijms20010039
57. Zeth K, Zachariae U. Ten years of high resolution structural research on the Voltage Dependent Anion Channel (VDAC)-recent developments and future directions. *Front Physiol*. 2018;9:108. doi:10.3389/fphys.2018.00108
58. Ponnalagu D, Singh H. Anion channels of mitochondria. *Handb Exp Pharmacol*. 2017;240:71–101.
59. Harris RA, Fenton AW. A critical review of the role of M2PYK in the warburg effect. *Biochim Biophys Acta Rev Cancer*. 2019;1871(2):225–239. doi:10.1016/j.bbcan.2019.01.004
60. Apostolova N, Cervera AM, Victor VM, et al. Loss of apoptosis-inducing factor leads to an increase in reactive oxygen species, and an impairment of respiration that can be reversed by antioxidants. *Cell Death Differ*. 2006;13(2):354–357. doi:10.1038/sj.cdd.4401776
61. Norberg E, Orrenius S, Zhivotovsky B. Mitochondrial regulation of cell death: processing of apoptosis-inducing factor (AIF). *Biochem Biophys Res Commun*. 2010;396(1):95–100. doi:10.1016/j.bbrc.2010.02.163
62. Jelinek A, Heyder L, Daude M, et al. Mitochondrial rescue prevents glutathione peroxidase-dependent ferroptosis. *Free Radic Biol Med*. 2018;117:45–57. doi:10.1016/j.freeradbiomed.2018.01.019
63. DeHart DN, Fang D, Heslop K, Li L, Lemasters JJ, Maldonado EN. Opening of voltage dependent anion channels promotes reactive oxygen species generation, mitochondrial dysfunction and cell death in cancer cells. *Biochem Pharmacol*. 2018;148:155–162. doi:10.1016/j.bcp.2017.12.022
64. Fang D, Maldonado EN. VDAC regulation: a mitochondrial target to stop cell proliferation. *Adv Cancer Res*. 2018;138:41–69.
65. Wilde L, Roche M, Domingo-Vidal M, et al. Metabolic coupling and the reverse warburg effect in cancer: implications for novel biomarker and anticancer agent development. *Semin Oncol*. 2017;44(3):198–203. doi:10.1053/j.seminoncol.2017.10.004
66. Xu XD, Shao SX, Jiang HP, et al. Warburg effect or reverse warburg effect? A review of cancer metabolism. *Oncol Res Treat*. 2015;38(3):117–122. doi:10.1159/000375435
67. Reina S, De Pinto V. Anti-cancer compounds targeted to VDAC: potential and perspectives. *Curr Med Chem*. 2017;24(40):4447–4469. doi:10.2174/0929867324666170530074039
68. Lemasters JJ. Evolution of voltage-dependent anion channel function: from molecular sieve to governor to actuator of ferroptosis. *Front Oncol*. 2017;7:303. doi:10.3389/fonc.2017.00303
69. Skouta R, Dixon SJ, Wang J, et al. Ferrostatins inhibit oxidative lipid damage and cell death in diverse disease models. *J Am Chem Soc*. 2014;136(12):4551–4556. doi:10.1021/ja411006a
70. Xie Y, Hou W, Song X, et al. Ferroptosis: process and function. *Cell Death Differ*. 2016;23(3):369–379. doi:10.1038/cdd.2015.158
71. Yu Y, Xie Y, Cao L, et al. The ferroptosis inducer erastin enhances sensitivity of acute myeloid leukemia cells to chemotherapeutic agents. *Mol Cell Oncol*. 2015;2(4):e1054549. doi:10.1080/23723556.2015.1054549
72. Shiromizu S, Yamauchi T, Kusunose N, Matsunaga N, Koyanagi S, Ohdo S. Dosing time-dependent changes in the anti-tumor effect of xCT inhibitor erastin in human breast cancer xenograft mice. *Biol Pharm Bull*. 2019;42(11):1921–1925. doi:10.1248/bpb.b19-00546
73. Seibt TM, Proneth B, Conrad M. Role of GPX4 in ferroptosis and its pharmacological implication. *Free Radic Biol Med*. 2019;133:144–152. doi:10.1016/j.freeradbiomed.2018.09.014
74. Gnanapradeepan K, Basu S, Barnoud T, Budina-Kolomets A, Kung CP, Murphy ME. The p53 tumor suppressor in the control of metabolism and ferroptosis. *Front Endocrinol (Lausanne)*. 2018;9:124. doi:10.3389/fendo.2018.00124
75. Huang C, Yang M, Deng J, Li P, Su W, Jiang R. Upregulation and activation of p53 by erastin-induced reactive oxygen species contribute to cytotoxic and cytostatic effects in A549 lung cancer cells. *Oncol Rep*. 2018;40(4):2363–2370. doi:10.3892/or.2018.6585
76. Kagan VE, Mao G, Qu F, et al. Oxidized arachidonic and adrenic PEs navigate cells to ferroptosis. *Nat Chem Biol*. 2017;13(1):81–90. doi:10.1038/nchembio.2238
77. Soupene E, Fyrst H, Kuypers FA. Mammalian acyl-CoA: lyso-phosphatidylcholineacyltransferase enzymes. *Proc Natl Acad Sci U S A*. 2008;105(1):88–93. doi:10.1073/pnas.0709737104
78. Dixon SJ, Winter GE, Musavi LS, et al. Human haploid cell genetics reveals roles for lipid metabolism genes in nonapoptotic cell death. *ACS Chem Biol*. 2015;10(7):1604–1609. doi:10.1021/acschembio.5b00245
79. Doll S, Proneth B, Tyurina YY, et al. ACSL4 dictates ferroptosis sensitivity by shaping cellular lipid composition. *Nat Chem Biol*. 2017;13(1):91–98. doi:10.1038/nchembio.2239
80. Yuan H, Li X, Zhang X, Kang R, Tang D. Identification of ACSL4 as a biomarker and contributor of ferroptosis. *Biochem Biophys Res Commun*. 2016;478(3):1338–1343. doi:10.1016/j.bbrc.2016.08.124
81. Liu Y, Wang W, Li Y, Xiao Y, Cheng J, Jia J. The 5-lipoxygenase inhibitor zileuton confers neuroprotection against glutamate oxidative damage by inhibiting ferroptosis. *Biol Pharm Bull*. 2015;38(8):1234–1239. doi:10.1248/bpb.b15-00048
82. Uyama O, Matsuyama T, Michishita H, Nakamura H, Sugita M. Protective effects of human recombinant superoxide dismutase on transient ischemic injury of CA1 neurons in gerbils. *Stroke*. 1992;23(1):75–81. doi:10.1161/01.STR.23.1.75

83. Liang YC, Wu CH, Chu JS, et al. Involvement of fatty acid-CoA ligase 4 in hepatocellular carcinoma growth: roles of cyclic AMP and p38 mitogen-activated protein kinase. *World J Gastroenterol*. 2005;11(17):2557–2563. doi:10.3748/wjg.v11.i17.2557
84. Monaco ME, Creighton CJ, Lee P, Zou X, Topham MK, Stafforini DM. Expression of long-chain fatty acyl-CoA synthetase 4 in breast and prostate cancers is associated with sex steroid hormone receptor negativity. *Transl Oncol*. 2010;3(2):91–98. doi:10.1593/tlo.09202
85. Wu X, Deng F, Li Y, et al. ACSL4 promotes prostate cancer growth, invasion and hormonal resistance. *Oncotarget*. 2015;6(42):44849–44863. doi:10.18632/oncotarget.6438
86. Zhang Y, Tan H, Daniels JD, et al. Imidazole ketone erastin induces ferroptosis and slows tumor growth in a mouse lymphoma model. *Cell Chem Biol*. 2019;26(5):623–633 e629. doi:10.1016/j.chembiol.2019.01.008
87. Li Y, Wang X, Yan J, et al. Nanoparticle ferritin-bound erastin and rapamycin: a nanodrug combining autophagy and ferroptosis for anticancer therapy. *Biomater Sci*. 2019;7(9):3779–3787. doi:10.1039/C9BM00653B
88. Sun X, Ou Z, Xie M, et al. HSPB1 as a novel regulator of ferroptotic cancer cell death. *Oncogene*. 2015;34(45):5617–5625. doi:10.1038/onc.2015.32
89. Huo H, Zhou Z, Qin J, Liu W, Wang B, Gu Y. Erastin disrupts Mitochondrial Permeability Transition Pore (mPTP) and Induces Apoptotic Death Of Colorectal Cancer Cells. *PLoS One*. 2016;11(5):e0154605. doi:10.1371/journal.pone.0154605
90. Zille M, Kumar A, Kundu N, et al. Ferroptosis in neurons and cancer cells is similar but differentially regulated by histone deacetylase inhibitors. *eNeuro*. 2019;6(1):ENEURO.0263–18.2019. doi:10.1523/ENEURO.0263-18.2019
91. Mansoori B, Mohammadi A, Davudian S, Shirjang S, Baradaran B. The different mechanisms of cancer drug resistance: a brief review. *Adv Pharm Bull*. 2017;7(3):339–348. doi:10.15171/apb.2017.041
92. Yuan R, Hou Y, Sun W, et al. Natural products to prevent drug resistance in cancer chemotherapy: a review. *Ann N Y Acad Sci*. 2017;1401(1):19–27. doi:10.1111/nyas.13387
93. Brasseur K, Gevry N, Asselin E. Chemoresistance and targeted therapies in ovarian and endometrial cancers. *Oncotarget*. 2017;8(3):4008–4042. doi:10.18632/oncotarget.14021
94. Mor G, Montagna MK, Alvero AB. Modulation of apoptosis to reverse chemoresistance. *Methods Mol Biol*. 2008;414:1–12. doi:10.1007/978-1-59745-339-4_1
95. Guo J, Xu B, Han Q, et al. Ferroptosis: a novel anti-tumor action for cisplatin. *Cancer Res Treat*. 2018;50(2):445–460. doi:10.4143/crt.2016.572
96. Yamaguchi H, Hsu JL, Chen CT, et al. Caspase-independent cell death is involved in the negative effect of EGF receptor inhibitors on cisplatin in non-small cell lung cancer cells. *Clin Cancer Res*. 2013;19(4):845–854. doi:10.1158/1078-0432.CCR-12-2621
97. Pennafort V, Queiroz MVO, Gomes ILV, Rocha MFF. Instructional therapeutic toy in the culture care of the child with diabetes type 1. *Rev Bras Enferm*. 2018;71(suppl 3):1334–1342. doi:10.1590/0034-7167-2017-0260
98. Chen L, Li X, Liu L, Yu B, Xue Y, Liu Y. Erastin sensitizes glioblastoma cells to temozolomide by restraining xCT and cystathionine-gamma-lyase function. *Oncol Rep*. 2015;33(3):1465–1474. doi:10.3892/or.2015.3712
99. Roh JL, Kim EH, Jang HJ, Park JY, Shin D. Induction of ferroptotic cell death for overcoming cisplatin resistance of head and neck cancer. *Cancer Lett*. 2016;381(1):96–103. doi:10.1016/j.canlet.2016.07.035
100. Wang SF, Chen MS, Chou YC, et al. Mitochondrial dysfunction enhances cisplatin resistance in human gastric cancer cells via the ROS-activated GCN2-eIF2alpha-ATF4-xCT pathway. *Oncotarget*. 2016;7(45):74132–74151. doi:10.18632/oncotarget.12356
101. Guo W, Zhao Y, Zhang Z, et al. Disruption of xCT inhibits cell growth via the ROS/autophagy pathway in hepatocellular carcinoma. *Cancer Lett*. 2011;312(1):55–61. doi:10.1016/j.canlet.2011.07.024
102. Ye P, Mimura J, Okada T, et al. Nrf2- and ATF4-dependent upregulation of xCT modulates the sensitivity of T24 bladder carcinoma cells to proteasome inhibition. *Mol Cell Biol*. 2014;34(18):3421–3434. doi:10.1128/MCB.00221-14
103. Takayama T, Kubo T, Morikawa A, Morita T, Nagano O, Saya H. Potential of sulfasalazine as a therapeutic sensitizer for CD44 splice variant 9-positive urogenital cancer. *Med Oncol*. 2016;33(5):45. doi:10.1007/s12032-016-0760-x
104. Ma MZ, Chen G, Wang P, et al. Xc- inhibitor sulfasalazine sensitizes colorectal cancer to cisplatin by a GSH-dependent mechanism. *Cancer Lett*. 2015;368(1):88–96. doi:10.1016/j.canlet.2015.07.031
105. Dixon SJ, Patel DN, Welsch M, et al. Pharmacological inhibition of cystine-glutamate exchange induces endoplasmic reticulum stress and ferroptosis. *Elife*. 2014;3:e02523. doi:10.7554/eLife.02523
106. Rodriguez A, Borrás JM, Lopez-Torrecilla J, et al. Demand for radiotherapy in Spain. *Clin Transl Oncol*. 2017;19(2):204–210. doi:10.1007/s12094-016-1525-x
107. Shadad AK, Sullivan FJ, Martin JD, Egan LJ. Gastrointestinal radiation injury: prevention and treatment. *World J Gastroenterol*. 2013;19(2):199–208. doi:10.3748/wjg.v19.i2.199
108. Moulder JE. Chemical radiosensitizers: the journal history. *Int J Radiat Biol*. 2019;95(7):940–944. doi:10.1080/09553002.2019.1569779
109. Wang H, Mu X, He H, Zhang XD. Cancer radiosensitizers. *Trends Pharmacol Sci*. 2018;39(1):24–48. doi:10.1016/j.tips.2017.11.003
110. Cobler L, Zhang H, Suri P, Park C, Timmerman LA. xCT inhibition sensitizes tumors to gamma-radiation via glutathione reduction. *Oncotarget*. 2018;9(64):32280–32297. doi:10.18632/oncotarget.25794
111. Pan X, Lin Z, Jiang D, et al. Erastin decreases radioresistance of NSCLC cells partially by inducing GPX4-mediated ferroptosis. *Oncol Lett*. 2019;17(3):3001–3008. doi:10.3892/ol.2019.9888
112. Shibata Y, Yasui H, Higashikawa K, Miyamoto N, Kuge Y, Hamada N. Erastin, a ferroptosis-inducing agent, sensitized cancer cells to X-ray irradiation via glutathione starvation in vitro and in vivo. *PLoS One*. 2019;14(12):e0225931. doi:10.1371/journal.pone.0225931
113. Negi P, Kingsley PA, Srivastava H, Sharma SK. Three weekly versus weekly cisplatin as radiosensitizer in head and neck cancer: a decision dilemma. *Asian Pac J Cancer Prev*. 2016;17(4):1617–1623. doi:10.7314/APJCP.2016.17.4.1617
114. Ward JF. DNA damage produced by ionizing radiation in mammalian cells: identities, mechanisms of formation, and reparability. *Prog Nucleic Acid Res Mol Biol*. 1988;35:95–125.

OncoTargets and Therapy

Dovepress

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>