Mechanism of Traditional Chinese Medicine in Treating Knee Osteoarthritis

This article was published in the following Dove Press journal:
Journal of Pain Research

Mina Wang^{1,2}

Lu Liu^{1,3}

Claire Shuiqing Zhang⁴

Zehuan Liao^{5,6}

Xianghong Jing³

Marc Fishers⁷

Luopeng Zhao^{1,8}

Xiaobai Xu¹

Bin Li¹

¹Acupuncture and Moxibustion Department, Beijing Hospital of Traditional Chinese Medicine, Capital Medical University, Beijing Key Laboratory of Acupuncture Neuromodulation, Beijing, People's Republic of China; ²Graduate School, Beijing University of Chinese Medicine, Beijing 100029, People's Republic of China; ³Institute of Acupuncture and Moxibustion, China Academy of Chinese Medical Sciences, Beijing, People's Republic of China; ⁴School of Health and Biomedical Sciences, RMIT University, Melbourne, Victoria, Australia; ⁵School of Biological Sciences, Nanyang Technological University, Singapore 637551; ⁶Department of Microbiology, Tumor and Cell Biology (MTC), Karolinska Institutet, Biomedicum, Stockholm SE-17177, Sweden; ⁷Department of Neurology, Beth Israel Deaconess Medical Centre and Harvard Medical School, Boston, MA, USA; ⁸Beijing Hospital of Traditional Chinese Medicine, Capital Medical University, Beijing Institute of Traditional Chinese Medicine, Beijing, People's Republic of China

Correspondence: Bin Li
Acupuncture and Moxibustion Department, Beijing Hospital of Traditional Chinese Medicine, Capital Medical University, Beijing Key Laboratory of Acupuncture Neuromodulation, Beijing, People's Republic of China
Tel +86 18910781852
Email libin@bjzhongyi.com

Abstract: Knee osteoarthritis (KOA) is a degenerative disease, making a unique contribution to chronic pain, edema, and limited mobility of knee joint. Traditional Chinese Medicine (TCM) is a common complementary therapy for KOA and has been found effective. The aim of this review is to consolidate the current knowledge about the mechanism of four interventions of TCM: acupuncture, moxibustion, herbs, and massage in treating KOA, and how they alleviate symptoms such as pain, swelling, and dysfunction. Furthermore, this review highlights that four therapies have different mechanisms but all of them can manage KOA through inhibiting inflammation, which indicates that alternative therapies should be considered as a viable complementary treatment for pain management in clinical practice.

Keywords: knee osteoarthritis, acupuncture, moxibustion, herbs, massage

Introduction

Osteoarthritis is a chronically degenerative joint disease which manifests as pain, swelling, stiffness, and dysfunction.¹ The incidence of osteoarthritis is high in the elder population, and an epidemiological study has also found that more than half of people who are younger than 65 years old suffer from osteoarthritis.² Knee osteoarthritis (KOA) is the most common type of osteoarthritis which reduces the quality of life among the elderly and increases economic burden in the society. Although the mechanism of KOA is still unclear, it is known that several mechanical and biological factors including age, obesity, trauma, inflammation, and genetic susceptibility have a great influence on catabolism and anabolism within chondrocytes,³ which vastly affects equilibrium of microenvironment of the knee joint. The imbalance of synthesis and degradation of chondrocytes, extracellular matrix (ECM), and subchondral bone eventually results in KOA.⁴

The current treatments, such as non-steroidal anti-inflammatory drugs, hyaluronic acid injections, and arthroplasty were proved to be helpful but also associated with certain side effects.⁵ Moreover, there is no recommended routine treatment for knee osteoarthritis. Traditional Chinese Medicine (TCM) including acupuncture, moxibustion, herbs, and massage has been used in China for thousands of years to treat many diseases.^{6,7} These methods have been practiced worldwide, mostly as an adjunctive treatment for chronic pain and diseases, such as diabetes and cancer.⁸⁻¹² Previous studies have shown that TCM is an effective therapy to relieve pain and to improve functions of knee joint for KOA patients.¹³⁻¹⁵ The mechanism of TCM to treat with KOA is intricate, which involves the promotion of the differentiation of bone marrow-derived mesenchymal stem cells into chondrocytes,¹⁶ facilitating the proliferation of

chondrocytes,¹⁷ and decelerating the inflammation of knee joint induced by inflammatory factors including interleukin (IL) and tumor necrosis factors (TNF).¹⁸ However, the precise mechanism of different types of TCM treatments remains to be fully elucidated. To the best of our knowledge, no previous reviews have summarized the mechanism of acupuncture, moxibustion, herbs, and massage in the treatment of KOA. This review aims to consolidate the mechanisms of the aforementioned four types of TCM treatments for KOA by screening published English and Chinese literature.

Mechanisms of Acupuncture

As an affordable and safe treatment, the efficacy of acupuncture in treating KOA patients has been demonstrated. Several clinical trials were conducted in USA, Germany, Spain, and China which proved high effectiveness of using acupuncture to treat KOA.^{19–21} The pathogenesis of KOA is related to inflammatory factors, cytokines, chemicals, signaling pathways, genetics, and biomechanics which mediate cartilage degradation.²² It has been demonstrated that IL-1 β and TNF- α contribute significantly to the severity and progression of KOA.²³ They can not only activate various signaling pathways such as mitogen-activated protein kinases (MAPK) signaling pathway and Toll-like receptor 4 (TLR4) signaling pathway but also promote nitric oxide (NO) and prostaglandin-E2 (PGE2) release, resulting in upregulation of matrix metalloproteinases (MMPs) and destruction of collagen II, which induce KOA.^{24,25} The mechanism of acupuncture in alleviating pain and promoting functional recovery in KOA patients involves:²⁶ 1) inhibiting the overexpression of inflammatory factors such as IL-1 β , IL-6, and TNF- α ; 2) suppressing the activity of some signaling pathways such as MAPK signaling pathway; 3) promoting the release of antioxidants; and 4) inhibiting hypertrophic differentiation of chondrocytes (Table 1).

Inhibition of Inflammatory Factors

Previous animal experiments have demonstrated that acupuncture is an effective method to inhibit the overexpression of inflammatory factors especially IL-1 β and TNF- α which exist in synovial fluid, cartilage, and subchondral bone.^{4,27} Studies have found that IL-1 β and TNF- α are crucial to increase in MMP1, MMP2, and MMP3. As the degradation of collagen II has characteristics of cartilage, the ECM degrades and synthesis of the ECM is suppressed.^{28–30} Moreover, IL-1 β promotes the expression of A Disintegrin And Metalloproteinase with

Table 1 The Mechanism of Acupuncture, Moxibustion, Herbs, and Massage in Treating Knee Osteoarthritis

Intervention	Mechanism
Acupuncture	1. inhibit the overexpression of inflammatory factors; 2. suppress the activity of some signaling pathways; 3. promote the release of antioxidants; 4. inhibit hypertrophic differentiation of chondrocytes.
Moxibustion	1. modulate NF- κ B signaling pathway; 2. suppress the activation and expression of COX-2; 3. reduce inflammatory factors directly.
Herbs	1. protect articular cartilage; 2. block Wnt/ β -catenin signaling pathway; 3. adjust p38-MAPK signaling pathway; 4. suppress inflammatory reactions.
Massage	1. change RANTES and MCP-1 expression; 2. balance musculature; 3. adjust inflammatory status.

Abbreviations: COX-2, cyclooxygenase-2; p38-MAPK, p38 mitogen-activated protein kinase; RANTES, regulated upon activation normal T cell expressed and secreted factor; MCP-1, monocyte chemoattractant protein-1.

Thrombospondin Motifs (ADAMTS)-4 and ADAMTS-5 which can also be promoted by TNF- α .³¹ Meanwhile, the levels of NO and PGE2 in knee joint increase with high level of IL-1 β and TNF- α .¹ A mouse model research confirmed that acupuncture efficiently suppresses inflammation by inhibiting expression and secretion of IL-1 β and TNF- α .³²

Inhibition of Signaling Pathway

Many signaling pathways are related to KOA, including Notch pathway, Wnt pathway, SDF-1/CXCR4 pathway, TLRs pathway, MAPK pathway, Hippo-YAP pathway, OPG-RANK-RANKL pathway, and TGF β pathway.^{33–39} MAPK signaling pathway is pivotal to pathogenesis of KOA, which induces MMP13 directly to attack the cartilage and mediate the expression of genes involving apoptosis of chondrocytes, such as Fas, Bcl-2 and Mcl-1.^{28,40,41} Acupuncture deactivates the MAPK signaling pathway, inhibiting the progression of KOA as validated in an animal experiment.⁴² Another study explored the mechanism of electroacupuncture for KOA based on Wnt/ β -catenin signaling pathway on 50 rats. The study used Western blot method to detect the levels of Wnt-4 and β -catenin in treatment and control group, and reported reduced levels of Wnt-4 and β -catenin in the treatment group.⁴³

Promoting Release of Antioxidants

Reactive oxygen species (ROS) has been hypothesized to be a crucial chemical in activating Wnt signaling pathway, mediating chondrocytes apoptosis and inducing pain.⁴⁴ Superoxide dismutase (SOD), which is a type of antioxidant that controls and removes ROS, demonstrated an increase in level in a clinical acupuncture research.^{45,46}

Inhibition of Hypertrophic Differentiation

Hypertrophic differentiation of chondrocytes causes premature bone, resulting in degradation of cartilage. Runt-related transcription factor 2 (Runx2) is the hallmark of hypertrophic differentiation of chondrocytes which can be manipulated by histone deacetylase 4 (HDAC4). A rabbit model indicated that acupuncture inhibits hypertrophic differentiation of chondrocytes in knee joint by activating HDAC4 and suppressing Runx2.⁴⁷

Mechanisms of Moxibustion

It is recognized that the meridian theory in traditional Chinese medicine is related to both acupuncture and moxibustion.⁴⁸ Moxibustion is also a well-known non-invasive procedure which involves burning moxa or mugwort (*Artemisia vulgaris*) to stimulate acupoints directly or indirectly in order to attenuate pain or accelerate recovery in many diseases, particularly in chronic diseases and refractory diseases.⁴⁹ Although moxibustion is unpopular in western countries, it is a common method in oriental regions to alleviate symptoms, such as China, Japan, and Korea because of its convenience, effectiveness, relative safety, and affordability.⁵⁰ The findings of several meta-analyses and animal experiments have indicated that moxibustion has its unique advantages for KOA patients with its analgesic and anti-inflammatory effects, which draws more attention to researchers recently.^{51,52} However, the understanding of underlying mechanism of moxibustion is still inadequate. It is believed that moxibustion plays a part in anti-nociceptive effect and functional improvement via 1) modulating NF- κ B signaling pathway; 2) suppressing the activation and expression of COX-2; and 3) reducing inflammatory factors directly such as IL-1 β and TNF- α (Table 1).

Modulation of NF- κ B Signaling Pathway

NF- κ B signaling pathway is important in regulating the expression of pro-inflammatory factors, adhesion molecules, and chemokines, such as IL-1 β , TNF- α , and inducible nitric oxide synthase (iNOS), which has a central role

in inflammation, cellular proliferation, differentiation, and apoptosis. As NF- κ B dimers are bound to I κ B proteins, once I κ B proteins degrade, NF- κ B signaling pathway is free to be activated to damage chondrocytes, leading to the occurrence of KOA.⁵³ The expression of NF- κ B p65 protein which is a sub-family member of NF- κ B transcription factors decreased, while I κ B-protein increased in soft tissue of rats knee joint in a KOA mouse model experiment, as detected by ELISA.⁵⁴

Suppression of COX-2

It is acknowledged that cyclooxygenase (COX) exists in two isoforms: COX-1 and COX-2, which have comparable sizes and structures but have different functional roles. COX-1 is constitutively expressed at high levels in platelets and gastric epithelial cells to sustain normal house-keeping functions, while COX-2 is an inducible enzyme, it is highly expressed in response to inflammation which is a major reason to cause pain and cartilage destruction of knee joint.⁵⁵ Moreover, PGE2 is produced through a COX/PGE2 synthases cascade,⁵⁶ which also plays an essential role in KOA process. The analgesic mechanism of moxibustion was investigated in a clinical research by detecting peripheral blood concentrations of COX-2 in KOA patients, and it displayed reduction on COX-2 concentration in the moxibustion groups.⁵⁷

Reduction of Inflammatory Factors

Similarly, moxibustion can also cause reduction of inflammatory factors like IL-1 β and TNF- α as the mechanisms of acupuncture, which was validated in several studies.^{18,58}

Mechanisms of Herbs

Herbal therapy is based on the syndrome differentiation in order to satisfy personalized needs of the patients.⁵⁹ Compared to single herb, physicians prefer herbal formulae which are complex mixtures of numerous herbs that contain abundant therapeutic compounds so that the curative effects are maximized and the toxicity or adverse effects are minimized by interplay of different herbs.⁶⁰ Recently, a large number of progress has been made in the research of treating KOA, some promising ingredients and herbal formulae have been identified that can delay the progression of KOA, promote cartilage repair, ameliorate knee joint disorders.⁶¹ The potential mechanism of herbal therapy in treating with KOA patients is related to 1) protecting articular cartilage including cartilage cells and matrix; 2) blocking Wnt/ β -catenin signaling pathway; 3)

adjusting p38-MAPK signaling pathway; and 4) suppressing inflammatory reactions (Table 1).

Protection of Articular Cartilage

The critical pathological characteristic of KOA is the degradation of chondrocytes and cartilage matrix, leading to degeneration of articular cartilage. Research found that herbal therapy could prevent the losses of proteoglycans and collagen II as well as accelerate the proliferation of chondrocytes.⁶² Cartilage oligomeric matrix protein (COMP), which is a tissue-specific protein, binds to collagen II fiber and cross-links C-telopeptides of type II collagen (CTX-II), a degradation product of collagen II.⁶³ They impact on stabilizing the collagen fiber network of articular cartilage, and once collagen II degrades, they are released into the synovial fluid and absorbed into serum as biochemical markers to reflect the destruction of articular cartilage.⁶⁴ Several studies used the concentrations of COMP and CTX-II in serum to inspect the mechanism of herbal therapy for KOA patients, and they showed that the concentrations of COMP and CTX-II in serum have a significant decrease in the herbal treatment group,^{65,66} which proved that certain herbal therapy has the ability to protect articular cartilage.

Blockage of Wnt/ β -Catenin Signaling Pathway

Wnt/ β -catenin signaling pathway is served as a pivotal regulator of bone remodeling through working independently or interacting with the other signaling pathways.⁶⁷ The Wnt signaling pathway is categorized by canonical and non-canonical Wnt signaling pathways. In the canonical Wnt signaling pathway, soluble Wnt proteins bind to receptor *fz1*, resulting in nuclear translocation of β -catenin protein and activation of signaling pathway. Accumulation of β -catenin leads to inflammatory reactions and overexpression of MMPs and ADAMTS5, which contributes to the degradation of cartilage, eventually deteriorates the progression of KOA.⁶⁸ Herbal therapy can block Wnt/ β -catenin signaling pathway, and mitigate dysfunction of knee joint in KOA patients, evidenced in a clinical research.⁶⁹

Adjustment of P38-MAPK Signaling Pathway

p38-MAPK signaling pathway is one of the critical members of MAPK signaling pathways, which has emerged as a key mediator of pro-inflammatory cytokines, chemokines, COX-2,

and MMPs to control cell proliferation, differentiation, senescence, and apoptosis.⁷⁰ Therefore, p38-MAPK signaling pathway adjusts the destruction of cartilage and degradation of matrix-associated proteins and collagen II,⁷¹ which is also the central role of pathogenesis of KOA. A rabbit model experiment was carried out to demonstrate the mechanism of herbal formula in treating degenerative chondrocytes. The experiment revealed that the herbal formula has impact on adjusting p38-MAPK signaling pathway to inhibit the apoptosis of chondrocytes.⁷²

Suppression of Inflammatory Reactions

As mentioned before, inflammatory reactions play an important role in the development of KOA, which bring about pain, swelling, and incompetence of knee joint.⁷³ Various pro-inflammatory factors, chemicals, and proteins, involving IL-1 β , TNF- α , MMPs, hyaluronic acid (HA), SOD, NO, and PGE2, participate in inducing and worsening inflammatory reactions.^{74,75} Remarkably, they do not only damage cartilage directly but activate multiple signaling pathways to cause degradation of ECM and collagen II at the same time, resulting in accelerating the failure of knee joint.⁷⁶ Animal experiments or clinical researches all confirmed that herbal therapy tremendously affects the level of multifarious inflammation-related factors,^{77–80} of which, is the mechanism of herbs in treating with KOA patients.

Mechanisms of Massage

Massage, an easy, cost-efficient, safe, and self-sufficient therapy, is one of the most popular treatments of complementary and alternative medicine used to reduce stress, relieve pain, and other conditions.⁸¹ Furthermore, its effectiveness has been reported in relieving the symptoms of KOA, including pain, edema, and weakness of muscles. It has also been emphasized in previous studies that massage therapy is beneficial to blood circulation, muscle tone and flexibility, especially for quadriceps, and inhibition of inflammatory factors.⁸² Even if there is not as much evidence as that of acupuncture, the recent explorations have made some conclusions that the underlying mechanism is associated with: 1) changing in RANTES and MCP-1 expression; 2) balancing musculature; 3) adjusting inflammatory status (Table 1).

Changes in RANTES and MCP-1 Expression

It has been identified that a plethora of chemokines and their receptors are expressed by cells within the knee joint,

and they play a central role in the KOA pathogenesis.⁸³ Chemokines are small chemotactic proteins and composed of more than 50 members that are relevant to inflammatory processes via providing a chemotactic gradient to recruit leucocytes directly, activating integrins, promoting mediator expression and regulating vascularization.⁸⁴ Within this complicated inflammatory milieu, regulated upon activation normal T cell expressed and secreted factor (RANTES) and monocyte chemotactic protein-1 (MCP-1) (two kinds of chemokines, also named CCL5 and CCL2) have broad responsibility to monocyte and macrophage recruitment to induce pain and physical disability of knee joint as they and their receptors are detected in KOA synovial tissue and synovial fluids.⁸⁵ Therefore, the less amount of RANTES and MCP-1 leads to less recruitment of monocyte and macrophage, eventually alleviating the progression of KOA.⁸⁶ A patient model in China demonstrated that massage has the best ability in modulating the expression of RANTES and MCP-1 compared to acupuncture and moxibustion, thus contributing more to patient recovery.⁸⁷

Balance of Musculature

Lower extremity muscle strength is significantly associated with functional capabilities of our daily life, such as walking and climbing stairs.⁸⁸ It has been reported that generating knee joint actions and alleviating loading requires normal quadriceps muscle strength.⁸⁹ In contrast to normal strength, quadriceps muscle fatigue sets KOA patients at high risk of pain and functional limitation, which also means these symptoms may be prevented by muscle exercise.⁹⁰ As we put forward above, massage is an effective therapy to passively exercise muscle and relax mind, which is due to improved endurance and flexibility of muscle. Therefore, it is beneficial for reducing pain and difficulty in performing functional tasks by the knee joint. Several studies have confirmed that massage is conducive in repairing the imbalance of musculature.^{91,92} Moreover, two different clinical researches compared the effect of massage with Celebrex and Fenbid, respectively, in increasing the strength of quadriceps (they used Biodex system 3-type isokinetic tester (Biodex Company, US) to test peak torque (PT), total work (TW), and average power (AP) before and after massage), and they both found that massage was superior in improving the strength of quadriceps for KOA patients.^{13,93}

Adjustment of Inflammatory Status

Inflammatory cytokines that drive the progression of KOA are chronic and low-grade inflammation, and they are extensively studied in the pathogenesis of KOA.⁹⁴ There is a wealth of evidence that inflammatory cytokines induce cartilage catabolism and halt anabolism, as a consequence of gradual structural changes within knee joint tissues and contributing to joint pain and dysfunction.⁹⁵ Although life-style changes such as exercise are convenient and effective, the regimens are difficult to implement in those people who are of advanced age or disability of mobility with KOA.⁹⁶ The regular complementary therapy —massage has its advantages in helping those individuals to slowdown the progression of inflammation reactions, preventing KOA from deteriorating. A rabbit model was established to assess the effect of massage in the expression of IL-1 and MMP-9, of which the promising result suggested that massage does have effect in adjusting inflammatory status of rabbit knee joint.⁹⁷

Conclusions

This review focuses on the mechanism of four classical types of Traditional Chinese Medicine in treating with KOA: acupuncture, moxibustion, herbs, and massage have different mechanism but all of them are associated with mediating inflammatory factors to alleviate KOA. Acupuncture inhibits KOA-related signaling pathways (MAPK, Wnt, Notch, SDF-1/CXCR4, TLRs, Hippo-YAP OPG-RANK-RANKL, and TGF β pathway) and hypertrophic differentiation of chondrocytes, but promotes the release of antioxidants. Moxibustion affects NF- κ B signaling pathway and the activation and expression of COX-2. Herbs modulate Wnt/ β -catenin signaling pathway and p38-MAPK signaling pathway as well as protect articular cartilage through preventing losses of proteoglycans and collagen II and promoting proliferation of chondrocytes. Massage changes the expression of RANTES and MCP-1 and harmonizes musculature. It has been confirmed in clinical doctors or scientists that acupuncture, moxibustion, herbs, and massage are beneficial to KOA, but most studies pay attention to individual side that TCM affects. The potential mechanism is more complicated because of interaction among diverse factors within knee joint. Therefore, the crosstalk between various factors should not be neglected in future studies. Overall, different interventions of TCM have their own aspects in treating with KOA, but they all

suppress inflammation of knee joint to relieve pain and make up for the abuse of acesodyne.

Funding

This study was funded by the National Key Research and Development Plan (2019YFC1709703), 2019 National Administration of Traditional Chinese-Project of building evidence based practice capacity for TCM-Project BEBPC-TCM (NO. 2019XZZX-ZJ002), Beijing Traditional Chinese Medicine Science and Technology Project (Grant No. JJ2018-53), China Association for Science and Technology Young Talent Lifting Project (2019-2021ZGZJXH-QNRC001), Capital Health Development Scientific Research Project Excellent Young Talents (Capital development 2020-4-2236), Beijing Dongcheng District Excellent Talent Development Funding (2019WJGW-10-05), China Postdoctoral Science Foundation Funded Project (2018M630261), Beijing Hospitals Authority Youth Programme (QML20181001), and Beijing Traditional Chinese Medicine Administration-Innovative and Inheritable “Fist” Construction Projects of Yanjing School.

Disclosure

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

- Herrero-Baumont G, Perez-Baos S, Sanchez-Pernaute O, Roman-Blas JA, Lamuedra A, Largo R. Targeting chronic innate inflammatory pathways, the main road to prevention of osteoarthritis progression. *Biochem Pharmacol*. 2019.
- Vina ER, Kwoh CK. Epidemiology of osteoarthritis: literature update. *Curr Opin Rheumatol*. 2018;30(2):160–167. doi:10.1097/BOR.0000000000000479
- Sacitharan PK, Vincent TL. Cellular ageing mechanisms in osteoarthritis. *Mamm Genome*. 2016;27(7–8):421–429. doi:10.1007/s00335-016-9641-z
- Wojdasiewicz P, Poniatowski AA, Szukiewicz D. The Role of Inflammatory and Anti-Inflammatory Cytokines in the Pathogenesis of Osteoarthritis. *Mediators Inflamm*. 2014;2014(2):561459. doi:10.1155/2014/561459
- Guo J, Chen Y, Li Z, et al. The cerebral mechanism of acupuncture for treating knee osteoarthritis: study protocol for a randomized controlled trial. *Trials*. 2019;20(1):126. doi:10.1186/s13063-019-3233-7
- Sai-bo L, Yao-chi W. Influence of traditional Chinese medicine on histomorphology and cytokine changes in knee osteoarthritis. *J Clin Rehabil Tissue Eng Res*. 2008;12(07):1313–1317.
- Liao Z, Zhao Y. Effect of Heat-Producing Needling Technique on the Local Skin Temperature: clinical Dataset. *Data*. 2018;3(4):65. doi:10.3390/data3040065
- Wang Z, Wang J, Chan P. Treating type 2 diabetes mellitus with traditional Chinese and Indian medicinal herbs. *Evidence Based Complementary Altern Med*. 2013;2013.
- Khan RMM, Chua ZJY, Tan JC, Yang Y, Liao Z, Zhao Y. From pre-diabetes to diabetes: diagnosis, treatments and translational research. *Medicina*. 2019;55(9):546. doi:10.3390/medicina55090546
- Phua W, Wong M, Liao Z, Tan N. An aPPAR α Functional Consequence in Skeletal Muscle Physiology via Peroxisome Proliferator-Activated Receptors. *Int J Mol Sci*. 2018;19(5):1425. doi:10.3390/ijms19051425
- Yang Y, Yang T, Liu S, et al. Concentrated ambient PM_{2.5} exposure affects mice sperm quality and testosterone biosynthesis. *PeerJ*. 2019;7:e8109. doi:10.7717/peerj.8109
- Xiang Y, Guo Z, Zhu P, Chen J, Huang Y. Traditional Chinese medicine as a cancer treatment: modern perspectives of ancient but advanced science. *Cancer Med*. 2019;8(5):1958–1975. doi:10.1002/cam4.2108
- Yuan-zhi F, Li G, Jun-tao Y, Min F, Yao-chi W. Effects of acupuncture and Chinese massage on the functions of knee flexors and extensors in patients with knee osteoarthritis: randomized controlled trial. *World J Acupuncture Moxibustion*. 2010;20(4):29–36.
- Si Y, Ma Y, Guo Y, et al. Efficacy and safety of Shaoyang Xibi decoction in patients with knee osteoarthritis: a multi-center, single-blind, randomized controlled trial. *J Tradit Chin Med*. 2018;38(5):733–739.
- Xu YK, Zhu FG, Feng EY, et al. [External therapies of traditional Chinese medicine combined with sodium hyaluronate injected in articular cavity therapy on knee osteoarthritis: meta-analysis]. *Zhongguo Zhong Yao Za Zhi*. 2018;43(9):1934–1939. doi:10.19540/j.cnki.cjcm.20180307.021. Chinese.
- Lin J, Wu G, Chen J, et al. Electroacupuncture inhibits sodium nitropruside-mediated chondrocyte apoptosis through the mitochondrial pathway. *Mol Med Rep*. 2018;18(6):4922–4930. doi:10.3892/mmr.2018.9498
- Xiaofang XI, Xuezhi LI, Liu F, et al. Repair effects of close-to-bone needling combined with electroacupuncture on extracellular matrix of cartilage in rabbits with knee osteoarthritis. *Zhongguo Zhen Jiu*. 2016;36(12):1288.
- Fu Y, Chen S, Mao J, et al. Mechanism of thermosensitive moxibustion on knee osteoarthritis in rabbit models. *Zhongguo Zhen Jiu*. 2018;38:291–296.
- Uryu N, Okada K, Kawakita K. Analgesic effects of indirect moxibustion on an experimental rat model of osteoarthritis in the knee. *Acupuncture Med*. 2007;25(4):175–183.
- Jianguo W, Lijuan H. Observation on the therapeutic effect of warming needle moxibustion on knee osteoarthritis. *Zhongguo Zhen Jiu*. 2007;03:191–192.
- Changdu L, Xinyong H, Xuguang Y, Qiongfeng W, Siqin H. Observation on therapeutic effect of warming needle moxibustion on knee osteoarthritis of deficiency-cold type. *Zhongguo Zhen Jiu*. 2006;03:189–191.
- Shi-quan S, Tao T. Discussion on the mechanism of acupuncture and moxibustion in the treatment of knee osteoarthritis. *Zhongguoyiyadaobao*. 2017;4(10):127–130.
- Xiao-cui Y, Qiang W, Wen S, et al. Electroacupuncture potentiates peripheral cB2 receptor-inhibited chronic pain in a mouse model of knee osteoarthritis. *J Pain Res*. 2018;11:2797–2808. doi:10.2147/JPR.S171664
- HOUHUANG C, XIANG S, LI L, et al. Electroacupuncture serum inhibits TNF- α -mediated chondrocyte inflammation via the Ras-Raf-MEK1/2-ERK1/2 signaling pathway. *Mol Med Rep*. 2017;16(5):5807–5814. doi:10.3892/mmr.2017.7366
- Gonzalez-Molina J, Gramolelli S, Liao Z, Carlson JW, Ojala PM, Lehti K. MMP14 in sarcoma: a regulator of tumor microenvironment communication in connective tissues. *Cells*. 2019;8(9):991. doi:10.3390/cells8020137
- Lida Z, Fanjing W, Zhen W, Guoe J, Xianghua C. Clinical application and mechanism of acupuncture in the treatment of knee osteoarthritis in the past 5 years. *Shanxizhongyixueyuanxuebao*. 2017;18(2):76–79.
- Yue P, Gao L, Chen M, Guo W, Xin SC. [Effect of Warm-needle-moxibustion on Behavior Reactions and TNF- α and MMP-3 Contents in Knee Cartilage of Rabbits with Knee Osteoarthritis]. *Zhen Ci Yan Jiu*. 2016;41(3):235–239. Chinese.

28. Huang ZM, Du SH, Huang LG, Li JH, Xiao L, Tong P. Regulation of senescence associated signaling mechanisms in chondrocytes for cartilage tissue regeneration. *Osteoarthritis Cartilage*. 2016;24(2):1246–1262. doi:10.1016/j.joca.2016.02.009
29. Li L, Li Z, Li Y, Hu X, Zhang Y, Fan P. Profiling of inflammatory mediators in the synovial fluid related to pain in knee osteoarthritis. *BMC Musculoskelet Disord*. 2020;21(1):99. doi:10.1186/s12891-020-3120-0
30. Paiva KBS, Granjeiro JM. Matrix Metalloproteinases in Bone Resorption, Remodeling, and Repair. *Prog Mol Biol Transl Sci*. 2017;148:203–303.
31. Kim J-R, Yoo JJ, Kim HA. Therapeutics in Osteoarthritis Based on an Understanding of Its Molecular Pathogenesis. *Int J Mol Sci*. 2018;19(3):674. doi:10.3390/ijms19030674
32. Hai WD, Fei B, Hong WZ, Hua S, Xiang. ZY. Influence of acupuncture on IL-1 β and TNF- α expression in the cartilage of rats with knee osteoarthritis. *China J Orthop Trauma*. 2011;24(9):775–778.
33. Kwon H, Paschos NK, Hu JC, Athanasiou K. Articular cartilage tissue engineering: the role of signaling molecules. *Cell Mol Life Sci*. 2016;73(6):1173–1194. doi:10.1007/s00018-015-2115-8
34. Zhou Y, Wang T, Hamilton JL, Chen D. Wnt/ β -catenin Signaling in Osteoarthritis and in Other Forms of Arthritis. *Curr Rheumatol Rep*. 2017;19(9):53. doi:10.1007/s11926-017-0679-z
35. Li X, Li J, Cheng K, et al. Effect of low-intensity pulsed ultrasound on MMP-13 and MAPKs signaling pathway in rabbit knee osteoarthritis. *Cell Biochem Biophys*. 2011;61(2):427–434. doi:10.1007/s12013-011-9206-4
36. Liu YX, Wang GD, Wang X, Zhang YL, Zhang TL. Effects of TLR-2/NF- κ B signaling pathway on the occurrence of degenerative knee osteoarthritis: an in vivo and in vitro study. *Oncotarget*. 2017;8(24):38602–38617. doi:10.18632/oncotarget.16199
37. Wang K, Li Y, Han R, et al. T140 blocks the SDF-1/CXCR4 signaling pathway and prevents cartilage degeneration in an osteoarthritis disease model. *PLoS One*. 2017;12(4):e0176048. doi:10.1371/journal.pone.0176048
38. Liu X, Du M, Wang Y, Liu S, Liu X. BMP9 overexpressing adipose-derived mesenchymal stem cells promote cartilage repair in osteoarthritis-affected knee joint via the Notch1/Jagged1 signaling pathway. *Exp Ther Med*. 2018;16(6):4623–4631. doi:10.3892/etm.2018.6754
39. Thomas M, Fronk Z, Gross A, et al. Losartan attenuates progression of osteoarthritis in the synovial temporomandibular and knee joints of a chondrodysplasia mouse model through inhibition of TGF- β 1 signaling pathway. *Osteoarthritis Cartilage*. 2019;27(4):676–686. doi:10.1016/j.joca.2018.12.016
40. Yun-Tao G, Jian C, Zhu-Long M, et al. Research progress on osteoarthritis treatment mechanisms. *Biomed Pharmacother*. 2017;93:1246–1252. doi:10.1016/j.biopha.2017.07.034
41. Shi-chao G, Hai-bo Y, Hong-xiao L, Yun-hui S. Research progress on MAPK signal pathway in the pathogenesis of osteoarthritis. *zhongguo gu shang/China J Orthop Trauma*. 2014;27(5):441–444.
42. Jian-jun G, Hong-mei Q, Cai-ru Z. Effect of dragon - tiger contending acupuncture on the mitogen - activated protein kinase signaling pathway in rats with knee osteoarthritis. *HUNAN J TRADITIONAL CHIN MED*. 2018;34(11):138–141.
43. Zhang YY, Li XH, Wu MX. [Effect of electroacupuncture at Wnt/ β -catenin signaling pathway on inhibiting cartilage degeneration in rats with knee osteoarthritis]. *Zhongguo Zhen Jiu*. 2019;39(10):1081–1086. doi:10.13703/j.0255-2930.2019.10.013. Chinese.
44. Liao Z, Chua D, Tan NS. Reactive oxygen species: a volatile driver of field cancerization and metastasis. *Mol Cancer*. 2019;18(1):65. doi:10.1186/s12943-019-0961-y
45. Wen-ji C, Qun W, Qun-wei Q, Yu-jie S. The Effect of Electrothermal Needle Acupuncture on Oxidative Stress Levels in Knee Osteoarthritis Patients. *Shanghai J Acu-mox*. 2017;36(2):218–221.
46. Liao Z, Tan ZW, Zhu P, Tan NS. Cancer-associated fibroblasts in tumor microenvironment—Accomplices in tumor malignancy. *Cell Immunol*. 2018;343:103729.
47. Fei L, Xue-zhi L, Ni-ni F, et al. Effect of electroacupuncture combined with close-to-bone needling on expression of cartilage cell HDAC4 and Runx2. in rabbits with knee osteoarthritis. *J Chongqing Med Univ*. 2016;41(2):179–184.
48. Liu Y, Chen W, Tan Y, et al. Analysis of the Registration Information on Interventions of Acupuncture and Moxibustion Trials in the International Clinical Trials Registry Platform. *Evid Based Complement Alternat Med*. 2018;2018:1054629. doi:10.1155/2018/9567061
49. Song GM, Tian X, Jin YH, et al. Moxibustion is an Alternative in Treating Knee Osteoarthritis: the Evidence From Systematic Review and Meta-Analysis. *Medicine*. 2016;95(6):e2790. doi:10.1097/MD.00000000000004864
50. Uryu N, Okada K, Kawakita K. Analgesic effects of indirect moxibustion on an experimental rat model of osteoarthritis in the knee. *Acupuncture Med*. 2007;25(4):175–183. doi:10.1136/aim.25.4.175
51. Li A, Wei ZJ, Liu Y, Li B, Guo X, Feng SQ. Moxibustion Treatment for Knee Osteoarthritis: a Systematic Review and Meta-Analysis. *Medicine (Baltimore)*. 2016;95(14):e3244. doi:10.1097/MD.00000000000003244
52. Xiu-mei R, Jin-jin C, Xue-yong S, et al. Knee osteoarthritis treated with moxibustion: a randomized controlled trial. *zhongguo gu shang/China J Orthop Trauma*. 2011;31(12):1057–1061.
53. Lepetosa P, Papavassiliou KA, Papavassiliou AG. Redox and NF- κ B signaling in osteoarthritis. *Free Radic Biol Med*. 2019;132:90–100. doi:10.1016/j.freeradbiomed.2018.09.025
54. Lu H, Rong-lin C, Ling H, et al. The Effect of Moxibustion on NF-kb β 5 Protein and IKB-protein in Knee Joint Soft Tissue of Rats with Knee Osteoarthritis. *Clin j traditional chin med*. 2017;29(6):856–858.
55. Norregaard R, Kwon TH, Frøkiær J. Physiology and pathophysiology of cyclooxygenase-2 and prostaglandin E2 in the kidney. *Kidney Res Clin Pract*. 2015;34(4):194–200. doi:10.1016/j.krcp.2015.10.004
56. Jia Z, Zhang Y, Ding G, Heiney KM, Huang S, Zhang A. Role of COX-2/mPGES-1/prostaglandin E2 cascade in kidney injury. *Mediators Inflamm*. 2015;2015:147894. doi:10.1155/2015/147894
57. Xiaoqiu Y, Jianmin D, Zhenghe C. Comparison of the Analgesic Effect of Different Needle Warming Moxibustion Methodson Syndrome of Yang-deficiency with Congealing Cold in KOA and their Mechanism. *World Chin Med*. 2015;10(9):1402–1405.
58. Xiu-bin L, Zheng-xiang L, Li-xin W. The Effect of Indirect Gentle Moxibustion on Serum IL-1 and TNF-Contents in Patients with Knee Osteoarthritis. *Shanghai J Acu-mox*. 2016;35(12):1459–1461.
59. Bauer BA, Tilburt JC, Sood A, et al. Complementary and alternative medicine therapies for chronic pain. *Chin J Integr Med*. 2016;22(6):403–411. doi:10.1007/s11655-016-2258-y
60. Liu L, Wang LP, He S, Immune Homeostasis: MY. Effects of Chinese Herbal Formulae and Herb-Derived Compounds on Allergic Asthma in Different Experimental Models. *Chin J Integr Med*. 2018;24(5):390–398. doi:10.1007/s11655-018-2836-2
61. Wang HM, Liu JN, Zhao Y. Progress on integrated Chinese and Western medicine in the treatment of osteoarthritis. *Chin J Integr Med*. 2010;16(4):378–384. doi:10.1007/s11655-010-0512-2
62. Xiang-ming N, Hong L, Ji-hong Z, Hong-min Z. Study on Histology of Herbs Invigorating Kidney Cure Knee Osteoarthritis. *J Clin Med*. 2014;1(12):2080–2081.
63. Das Gupta E, Ng WR, Wong SF, Bhurhanudeen AK, Yeap SS. Correlation of serum cartilage oligomeric matrix protein (COMP) and interleukin-16 (IL-16) levels with disease severity in primary knee osteoarthritis: a pilot study in a Malaysian population. *PLoS One*. 2017;12(9):e0184802.
64. Bai B, Li Y. Combined detection of serum CTX-II and COMP concentrations in osteoarthritis model rabbits: an effective technique for early diagnosis and estimation of disease severity. *J Orthop Surg Res*. 2016;11(1):149.

65. Feng W, Dingrui X, Wenwu L, Yuanfang D. Effect of Xiaoyu Jiegu powder on COMP, serum CTX- II in rabbit model with knee osteoarthritis. *Clin j traditional chin med*. 2015;27(12):1723–1725.
66. Peng Z, Guan-tong S, Yue-long C. Effect of Invigorating Kidney Formula, Softening Liver Formula and Dissipating Phlegm Formula on Synovocyte Proliferation and COMP Expression in Knee Osteoarthritis. *Chin J Traditional Med Traumatol Orthopedics*. 2008;15(12):18–21.
67. Mohammed MK, Shao C, Wang J, et al. Wnt/ β -catenin signaling plays an ever-expanding role in stem cell self-renewal, tumorigenesis and cancer chemoresistance. *Genes Dis*. 2016;3(1):11–40. doi:10.1016/j.gendis.2015.12.004
68. Zhang Y, Pizzute T, Pei M. A review of crosstalk between MAPK and Wnt signals and its impact on cartilage regeneration. *Cell Tissue Res*. 2014;358(3):633–649. doi:10.1007/s00441-014-2010-x
69. Meng-ya T, Hui-ying N, Xuemin Z, Lifang S. Clinical study on the role of reinforcing kidney activating blood drugs(TCD) in the regulation of classic Wnt/ β -catenin pathway in patients with osteoarthritis. *Zhong Guo Zheng Gu*. 2014;26(8):12–17.
70. Joos H, Albrecht W, Laufer S, Brenner RE. Differential effects of p38MAP kinase inhibitors on the expression of inflammation-associated genes in primary, interleukin-1 β -stimulated human chondrocytes. *Br J Pharmacol*. 2010;160(5):1252–1262. doi:10.1111/j.1476-5381.2010.00760.x
71. Chen Y, Shou K, Gong C, Yang H, Yang Y, Bao T. Anti-Inflammatory Effect of Geniposide on Osteoarthritis by Suppressing the Activation of p38 MAPK Signaling Pathway. *Biomed Res Int*. 2018;2018:8384576.
72. Fei-tai L, Yu L, Yi-yuan Z, et al. Effects of Serum Medicated with Duhuo Jisheng Decoction on the Induced Degeneration of Articular Chondrocytes in Vitro Culture Based upon “caveolin-p38MAPK” Signal Pathway. *Rheumatism Arthritis*. 2015;4(4):5–9.
73. Shun L, Jun-jie Z, Xiao-tao X, Wen-wu G, Xian-qi C, Cheng-hui L. The clinical use and its mechanisms research on angelicae pubescentis and loranthe decoction in KOA treatment. *Global Traditional Chin Med*. 2015;8(9):1149–1152.
74. Kim K-S, Cho H-S, Lee S-D, et al. Inhibitory effect of Buthus martensi Karsch extracts on interleukin-1 β -induced expression of nitric oxide (NO) synthase and production of NO in human chondrocytes and LPS-induced NO and prostaglandin E2 production in mouse peritoneal macrophages. *Toxicol vitro*. 2005;19(6):757–769. doi:10.1016/j.tiv.2005.04.008
75. Dong R, Tan Y, Fan A, Liao Z, Liu H, Wei P. Molecular Dynamics of the Recruitment of Immunoreceptor Signaling Module DAP12 Homodimer to Lipid Raft Boundary Regulated by PIP2. *J Phys Chem B*. 2019.
76. Philp AM, Davis ET, Jones SW. Developing anti-inflammatory therapeutics for patients with osteoarthritis. *Rheumatology*. 2017;56(6):869–881. doi:10.1093/rheumatology/kew278
77. Jianqin S, Hongying C, Zhonghong Y, Jingping M, Lihong G. Experimental Study on Effects of Electroacupuncture Combined with Weilingxian Intra-articular Injection on MMPs and HA in Rats with Knee Osteoarthritis. *J Emergency Traditional Chin Med*. 2016;25(5):791–794.
78. Feifei L, Shuguang Y, Haibo J, Xicheng L, Suting D. Effect of Gutong Powder on the Expression of IL-1 β , TNF- α and MMP-3 in the Patients with Knee Osteoarthritis. *Clin j traditional chin med*. 2017;29(11):1912–1915.
79. Lianfeng J, Shanshan L. Drynaria on Synovial Cytokine TNF - α in Rabbit Model of Knee Osteoarthritis. *Chin Arch Traditional Chin Med*. 2014;32(6):1398–1401.
80. Li H, Peng Y, Wang X, et al. Astragaloside inhibits IL-1 β -induced inflammatory response in human osteoarthritis chondrocytes and ameliorates the progression of osteoarthritis in mice. *Immunopharmacol Immunotoxicol*. 2019;41(4):497–503. doi:10.1080/08923973.2019.1637890
81. Atkins V. The Effects of Self-Massage on Osteoarthritis of the Knee: a Randomized, Controlled Trial. *Int j ther massage bodywork*. 2013;6:4.
82. Godoy Cortés V, Gallego-Izquierdo T, Lázaro Navas I, Pecos Mart ND. Effectiveness of massage therapy as co-adjuvant treatment to exercise in osteoarthritis of the knee: a randomized control trial. *J back Musculoskeletal Rehabil*. 2014;27:521–529.
83. Haringman JJ, Smeets TJ, Reinders-Blankert P, Tak PP. Chemokine and chemokine receptor expression in paired peripheral blood mononuclear cells and synovial tissue of patients with rheumatoid arthritis, osteoarthritis, and reactive arthritis. *Ann Rheum Dis*. 2006;65(3):294–300. doi:10.1136/ard.2005.037176
84. Endres M, Andreas K, Kalwitz G, et al. Chemokine profile of synovial fluid from normal, osteoarthritis and rheumatoid arthritis patients: CCL25, CXCL10 and XCL1 recruit human subchondral mesenchymal progenitor cells. *Osteoarthritis Cartilage*. 2010;18(11):1458–1466. doi:10.1016/j.joca.2010.08.003
85. Raghu H, Lepus CM, Wang Q, et al. CCL2/CCR2, but not CCL5/CCR5, mediates monocyte recruitment, inflammation and cartilage destruction in osteoarthritis. *Ann Rheum Dis*. 2017;76(5):914–922. doi:10.1136/annrheumdis-2016-210426
86. Scanzello CR. Chemokines and inflammation in osteoarthritis: Insights from patients and animal models. *J Orthop Res*. 2017;35(4):735–739. doi:10.1002/jor.23471
87. Chun-ri L, Yu H, Bao-qiang D. Research of RANTES and MCP-1 Expression in Knee Osteoarthritis after Acupuncture Moxibustion and Massage Treatment. *J Liaoning Univ TCM*. 2013;15(10):79–81.
88. Luc-Harkey BA, Safran-Norton CE, Mandl LA, Katz JN, Losina E. Associations among knee muscle strength, structural damage, and pain and mobility in individuals with osteoarthritis and symptomatic meniscal tear. *BMC Musculoskelet Disord*. 2018;19(1):258. doi:10.1186/s12891-018-2182-8
89. Muraki S, Akune T, Teraguchi M, et al. Quadriceps muscle strength, radiographic knee osteoarthritis and knee pain: the ROAD study. *BMC Musculoskelet Disord*. 2015;16(1):305. doi:10.1186/s12891-015-0737-5
90. Elboim-Gabyzon M, Rozen N, Laufer Y. Quadriceps femoris muscle fatigue in patients with knee osteoarthritis. *Clin Interv Aging*. 2013;8:1071–1077. doi:10.2147/CIA.S42094
91. Jia-nan D, Li G, Jian-hua L, et al. Effect of Pain Point Manipulation on Isokinetic Myodynamia in Early and Mid-stage Knee Osteoarthritis. *Chin J Traditional Med Traumatol Orthopedics*. 2015;23(6):35–38+41.
92. Hua LJ, Li G, Min F, Zhe W, Yang L. Effects of Tuina on muscular tension of flexor and extensor in patients with knee osteoarthritis. *Chin J Orthopaedics Traumatol*. 2011;24(575–577):213–217.
93. Yuan-zhi F, Min F, Jun-tao Y, Li G, Wu-quan S, Yao-chi W. Clinical Study on Tuina Therapy for Knee Osteoarthritis Patients Musculus Quadriceps Fexoris Strength. *J Chin Med*. 2010;25(5):1007–1010.
94. Maldonado M, Nam J. The role of changes in extracellular matrix of cartilage in the presence of inflammation on the pathology of osteoarthritis. *Biomed Res Int*. 2013;2013:284873.
95. Scanzello CR. Role of low-grade inflammation in osteoarthritis. *Curr Opin Rheumatol*. 2017;29(1):79–85. doi:10.1097/BOR.0000000000000353
96. Robinson WH, Lepus CM, Wang Q, et al. Low-grade inflammation as a key mediator of the pathogenesis of osteoarthritis. *Nat Rev Rheumatol*. 2016;12(10):580–592. doi:10.1038/nrrheum.2016.136
97. Ying C, Mingxin X, Jianjun L, Shuai Z, Shilong S. Effect of Qiang Huo Guizhi manipulation combined with Tuina on IL-1 and MMP-9 in rabbit knee osteoarthritis. *J Pract Med*. 2019;35(2):181–185.

Journal of Pain Research**Dovepress****Publish your work in this journal**

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication. The manuscript

management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>