Prevalence and Predictors of CD4+ T-Lymphocytopenia Among HIV-Negative Tuberculosis Patients in Uganda

This article was published in the following Dove Press journal:
Research and Reports in Tropical Medicine

Joseph Baruch Baluku^{1,3}
Joseph Musazi⁴
Rose Mulwana¹
Araali Robert Mugabo¹
Felix Bongomin⁵
Wincelous Katagira³

¹Mulago National Referral Hospital, Pulmonology Division, Kampala, Uganda; ²Mildmay Uganda, Kampala, Uganda; ³Makerere University Lung Institute, Kampala, Uganda; ⁴Makerere University Infectious Disease Institute, Kampala, Uganda; ⁵Department of Medical Microbiology & Immunology, Faculty of Medicine, Gulu University, Gulu, Uganda

Purpose: CD4+ T-lymphocytopenia is a risk for tuberculosis (TB) infection, reactivation and severe disease. We sought to determine the prevalence and predictors of CD4+ T-lymphocytopenia among HIV-negative patients with bacteriologically confirmed TB in Uganda.

Patients and Methods: Eligible participants were adult HIV-negative patients with bacteriologically confirmed TB at the National TB Treatment Centre in Uganda. CD4+ and CD8+ T-lymphocyte counts were determined by flow cytometry. We defined CD4+ T-lymphocytopenia as a CD4+ T-lymphocyte count of $<418 \text{ cells/mm}^3$ as per the population estimate for Ugandans. We performed logistic regression analysis to determine predictors of CD4+ T-lymphocytopenia.

Results: We enrolled 216 participants whose mean age (standard deviation ($\pm$ SD)) was 32.5 (± 12.1) years, of whom 146 (67.6%) were males. The prevalence of CD4+ T-lymphocytopenia was 25% (54/216) (95% confidence interval (CI): 19.6–31.2%). Patients with anaemia (adjusted odds ratio (aOR): 3.83, 95% CI: 1.59–9.23, $p = 0.003$), weight loss (aOR: 3.61, 95% CI: 1.07–12.23, $p = 0.039$) and a low CD8+ T-cell count (aOR: 6.10, 95% CI: 2.68–13.89, $p < 0.001$) were more likely to have CD4+ T-lymphocytopenia while those with monocytosis (aOR: 0.35, 95% CI: 0.14–0.89, $p = 0.028$) were less likely to have CD4+ T-lymphocytopenia.

Conclusion: There was a high prevalence of CD4+ T-lymphocytopenia among HIV-negative TB patients. Patients with weight loss, anaemia and a low CD8+ count were more likely to have CD4+ T-lymphocytopenia while those with monocytosis were less likely to have CD4+ lymphocytopenia. The findings suggest that CD4+ lymphocytopenia is indicative of severe disease and globally impaired cell-mediated immune responses against TB.

Keywords: HIV negative, tuberculosis, CD4+, CD8+, lymphocytopenia, monocytosis, weight loss, anaemia

Plain Language Summary

CD4+ T-cells are a crucial component of the immune response against tuberculosis (TB). Individuals with low CD4+ T-cell counts are at risk of TB infection and more likely to have severe forms of TB. We determined the prevalence and predictors of low CD4+ T-cell counts among HIV-negative TB patients at a national TB treatment centre in Uganda. We found that a quarter of HIV-negative TB patients in this setting have a low CD4+ T-cell count (less than 418 cells per mm^3). Patients with anaemia, weight loss, and low CD8+ T-cell counts were more likely to have low CD4+ T-cell counts while those with high monocyte counts were

Correspondence: Joseph Baruch Baluku
Mulago National Referral Hospital,
Pulmonology Division, P.O. Box 7051,
Kampala, Uganda
Tel +256706327972
Email bbjoe18@gmail.com

less likely to have low CD4+ T-cell counts. The results indicate that HIV-negative TB patients with low CD4+ T-cell counts also have impairment in the function of other cells of the immune system and severe TB disease.

Background

Globally, tuberculosis (TB) contributed 1.2 million deaths among HIV-negative individuals in 2018.¹ The control of TB infection depends on intact innate and adaptive immune responses that are mainly effected by monocytes (as macrophages) and cluster of differentiation (CD) 4+ T-lymphocytes, respectively.² CD4+ T-lymphocytes produce interferon gamma (IFN γ) that contributes to the recruitment and activation of innate immune cells that ultimately clear TB infection by phagocytosis and autophagy.^{3,4} IFN γ derived from CD4+ T-lymphocytes is particularly important for the control of mycobacterial burden, abating intracellular multiplication of *Mycobacterium tuberculosis* (Mtb), preventing dissemination from pulmonary infection and promoting long-term survival of the infected host.^{5,6}

Among HIV-positive individuals, it has been unequivocally demonstrated that CD4+ T-lymphocytopenia increases the risk for TB infection and mortality among TB/HIV co-infected individuals.^{7,8} Although lower CD4+ T-lymphocyte counts are reported among HIV-negative TB patients than controls, the prevalence and predictors of the CD4+ T-lymphocytopenia are not well characterised.^{9,10} The few studies that have evaluated the prevalence of CD4+ T-lymphocytopenia among HIV-negative TB patients report it to be 14.4–83%.^{11–14} However, these studies are limited by the heterogeneity of the study population whereby participants without bacteriological confirmation of TB were included. Further, predictors of CD4+ T-lymphocytopenia among HIV-negative TB patients are not widely reported. We aimed to determine the prevalence and predictors of CD4+ T-lymphocytopenia among HIV-negative bacteriologically confirmed TB patients at a national tuberculosis treatment centre in Uganda.

Patients and Methods

Study Measurements

We analysed data from a cross-sectional study that enrolled adult bacteriologically confirmed TB patients at the National Tuberculosis Treatment Centre (NTTC) in Uganda between August 2017 and March 2018.¹⁵ Demographic data, clinical symptoms and signs, full hemogram parameters, CD4+ and CD8+ T-lymphocyte counts were extracted from the master

dataset. A full description of methods employed in determination of these variables is provided elsewhere.¹⁵ CD4+ and CD8+ T-lymphocyte counts were determined by flow cytometry using a flow cytometer (BD FACSCalibur™). We defined CD4+ T-lymphocytopenia as a CD4+ count of <418 cells/mm³ considering a normal range estimate of 418–2105 cells/mm³ for Ugandans.¹⁶ Full hemogram parameters were determined by a hemoanalyser (Sysmex® Automated haematology analyser XN series – XN 1000) at Mulago Hospital Haematology laboratory. The HIV status was determined by immunochromatographic rapid tests according to the Uganda national HIV testing guidelines.¹⁷

Study Setting and Population

The NTTC is a unit of Mulago National Referral Hospital and is a tertiary referral centre for diagnosis and treatment of drug susceptible and drug resistant TB. The centre is located in Kampala city, the capital city of Uganda, and treats 25% of the country's TB cases. In this analysis, we included bacteriologically confirmed adult (≥ 18 years) TB patients who were HIV negative and had a CD4+ T-lymphocyte count result. There were no exclusion criteria. In the primary study, participants were consecutively sampled from the outpatient and inpatient facilities of the NTTC until the desired sample size was obtained.¹⁵

Statistical Analysis and Sample Size Estimation

Considering a prevalence of CD4+ T-lymphocytopenia of 14.4%¹³ reported in Senegal, a sample size of at least 211 is sufficient to determine the prevalence of CD4+ T-lymphocytopenia at the 95% confidence interval (CI) and a possible 10% occurrence of missing data. Data were analysed using STATA 15.1 (StataCorp, College Station, TX, USA). Categorical data were summarised as frequencies and percentages while continuous variables were summarised as means with the corresponding standard deviations (SD). The prevalence of CD4+ T-lymphocytopenia was calculated as a proportion of participants with a CD4+ count of <418 cells/mm³ to the total number of study participants. To determine factors that are independently associated with CD4+ T-lymphocytopenia, we performed bi-variable logistic regression analysis and fitted factors that had a $p \leq 0.2$ into a multivariable logistic regression model. A $p \leq 0.05$ was considered statistically significant considering a 95% CI. Missing data on monocyte counts were imputed using

multiple imputations with chained equations (MICE) using age, gender and platelet count, with 30 imputations. Since modelling involved haematological parameters, multi-collinearity would be expected. Prior to fitting statistical models, we checked for multi-collinearity between covariates to determine if there existed substantial correlation (Pearson's correlation coefficient > 0.3). The haematocrit, mean corpuscular haemoglobin (MCH), and red cell count were highly correlated with the haemoglobin level and adding them in the adjusted model induced multi-collinearity. Thus, these were dropped from the final model in favour of the haemoglobin level.

Ethical Approvals and Consent to Participate

Study participants provided written informed consent prior to study enrolment. The study was approved by the School of Medicine Research and Ethics Committee of Makerere University College of Health Sciences (REC REF 2017–087).

Results

The analysis was performed on 216 HIV-negative adult patients with bacteriologically confirmed TB.

Baseline Characteristics of Study Participants

The mean age ($\pm$ SD) of the study participants was 32.54 ($\pm$ 12.08) years and 146 (67.6%) were males. Other baseline characteristics are shown in Table 1. The frequencies of the parameters of the full hemogram, CD8+ and CD4/CD8 ratio of the study participants are shown in Table 2.

Prevalence of CD4+ T-Lymphocytopenia Among HIV-Negative TB Patients

The prevalence of CD4+ T-lymphocytopenia was 25% (54/216) (95% CI: 19.6% - 31.2%). For the entire study population, the mean ($\pm$ SD) CD4+, CD8+ T-cell counts and CD4+/CD8+ ratio were 636.62 ($\pm$ 304.91) cells/mm³, 429.88 ($\pm$ 264.74) cells/mm³ and 1.77 ($\pm$ 0.89) respectively.

Predictors of CD4+ T-Lymphocytopenia Among HIV-Negative Pulmonary TB Patients

At multivariable logistic regression analysis, HIV-negative TB patients with anaemia (adjusted odds ratio (aOR): 3.83,

95% CI: 1.59–9.23, $p = 0.003$), weight loss (aOR: 3.61, 95% CI: 1.07–12.23, $p = 0.039$) and a low CD8+ T-cell count (aOR: 6.10, 95% CI: 2.68–13.89, $p < 0.001$) were more likely to have CD4+ T-lymphocytopenia while those with monocytosis (aOR: 0.35, 95% CI: 0.14–0.89, $p = 0.028$) were less likely to have CD4+ T-lymphocytopenia. The multivariable logistic regression model is shown in Table 3.

Discussion

In this study, we evaluated the prevalence and predictors of CD4+ T-lymphocytopenia among HIV-negative TB patients at a national tuberculosis treatment centre in Uganda. We found a high prevalence of CD4+ T-lymphocytopenia of 25%. Patients with anaemia, weight loss and low CD8+ T-cell counts were more likely to have CD4+ T-lymphocytopenia while those with monocytosis were less likely to have CD4+ T-lymphocytopenia.

The high prevalence of CD4+ T-lymphocytopenia indicates that a high proportion of HIV-negative TB patients in this setting may have impaired immune responses to *Mtb* infection. The consequences are higher rates of disseminated disease, absence of smear positivity and characteristic imaging findings that could delay diagnosis and superimposed opportunistic infections, all of which could potentially influence treatment outcomes.¹³ CD4+ T-lymphocytopenia among patients with tuberculosis could be explained by homing of T-lymphocytes in peripheral blood to heavily infected sites in the lung parenchyma and lung vasculature.¹⁸ It follows that response to TB treatment would correlate with improvement of the CD4+ T-lymphocyte counts in peripheral blood. Resolution of CD4+ T-lymphocytopenia upon treatment of TB has been reported.^{9,12} This suggests that TB is the cause of the T-lymphocytopenia as opposed to the CD4+ T-lymphocytopenia being a risk factor for TB. On the contrary, Zonios et al have shown that an idiopathic form of CD4+ T-lymphocytopenia is a risk factor for mycobacterial infections through a prospective study design.¹⁹ The cross-sectional nature of our study does not establish whether CD4+ T-lymphocytopenia was caused by TB or was a risk factor for TB among HIV-negative individuals in our study population.

CD4+ T-lymphocytopenia was associated with low CD8+ T-lymphocyte counts which would otherwise have enhanced TB immune responses.²⁰ CD4+ T-lymphocytes are required for proliferation of CD8+ T-lymphocytes that modulate cytotoxic effects on *Mtb* infected macrophages and participate in memory immune responses against TB.^{21–23} Interestingly, the majority of the participants

Table 1 Baseline Characteristics of Study Participants

Baseline Characteristic	n (%)
Education level	
None	12 (5.6)
Primary school	64 (29.6)
Secondary school	98 (45.4)
Tertiary	42 (19.4)
Residence	
Urban	149 (69.0)
Rural	67 (31.0)
Alcohol use	105 (48.4)
Cigarette smoking	63 (29.2)
Symptoms	
Cough	213 (98.6)
Weight loss	170 (78.7)
Night sweats	145 (67.1)
Fever	134 (62.0)
Chills	89 (41.2)
Anorexia	61 (28.2)
Number of symptoms	
<4	82 (38.0)
≥4	134 (62.0)
Symptoms duration days (mean (± SD))	
Cough	92.23 (± 116.54)
Fever	32.53 (± 45.55)
Night sweats	78.15 (± 497.43)
Anorexia	29.15 (± 36.79)
Weight loss	44.34 (± 46.75)
Body mass index (kilograms per metres ²), mean (± SD)	18.78 (±2.84)
Rifampicin resistant tuberculosis	15 (11.5)

Abbreviation: SD, standard deviation.

had a normal CD4/CD8 ratio, although the clinical utility of the CD4/CD8 ratio is not well established among HIV-negative TB patients. One meta-analysis indicates that the CD4/CD8 ratio in peripheral blood is significantly decreased among newly diagnosed patients with TB, although the bronchoalveolar CD4/CD8 ratio was increased on the contrary.²⁷

The central role of the CD4+ T-lymphocytes in Mtb immune responses is further supported by our finding of lower odds of CD4+ T-lymphocytopenia among patients with monocytosis. Mtb-specific-CD4+ Th1 cell responses induce monocytosis by production of IFN- γ or tumour necrotic factor alpha (TNF- α) that recruits monocytes/macrophages to effect TB phagocytosis.^{24,25} Further, there is cell to cell communication between CD4+ T-lymphocytes and

Table 2 Haematological and Immunological Characteristics of the Study Participants

Characteristics ^a	<Lower Limit of Normal n (%)	Normal n (%)	>Upper Limit of Normal n (%)
White blood cell count (n=212)	1 (0.5)	125 (59.0)	86 (40.5)
Red blood cell count (n=213)	12 (5.6)	183 (85.9)	18 (8.5)
Hemoglobin (n=213)	51 (23.9)	161 (75.6)	1 (0.5)
Hematocrit (n=213)	37 (17.4)	161 (75.6)	15 (7.0)
Mean corpuscular volume (n=213)	58 (27.2)	147 (69.0)	8 (3.8)
Mean corpuscular hemoglobin (n=213)	59 (27.7)	152 (71.4)	2 (0.9)
Mean corpuscular hemoglobin concentration (n=213)	129 (60.6)	29 (13.6)	55 (25.8)
Platelet count (n=212)	5 (2.4)	107 (50.5)	100 (47.2)
Mean platelet volume (n=209)	1 (0.5)	157 (75.1)	51 (24.4)
Neutrophil count (n=155)	1 (0.7)	50 (32.3)	104 (67.1)
Lymphocyte count (n=155)	53 (34.2)	102 (65.8)	0(0.0)
Monocyte count (n=155)	4 (2.6)	75 (48.4)	76 (49.0)
Eosinophil count (n=155)	40 (25.8)	115 (74.2)	0 (0.0)
Basophil count (n=155)	1 (0.7)	149 (96.1)	5 (3.2)
CD8 count ² (n=216)	59 (27.3)	156 (72.2)	1(0.5)
CD4/CD8 ratio ^b	11 (5.1%)	203(94.0)	2 (0.9)

Notes: ^aFull hemogram ranges are for adult Ugandans.³⁹ ^bNormal ranges are for adult Ugandans.¹⁶

monocytes that could influence the cytokine repertoire for TB containment.²⁶ Therefore, CD4+ T-lymphocytopenia could be indicative of global impairment of cell-mediated immune responses against TB.

We also found that anaemia and weight loss were independently associated with CD4+ T-lymphocytopenia. Anaemia and weight loss are prognostic of delayed sputum conversion and mortality in TB disease and the association with CD4+ T-lymphocytopenia may be indicative of severe TB disease.²⁸⁻³¹ Therefore, CD4+ T-lymphocytopenia is likely to have prognostic importance even among HIV-negative TB patients.³² In support of this, baseline levels of some subsets of CD4+ T-lymphocytes have been shown to predict TB culture conversion.³³ Weight loss in tuberculosis is mediated by appetite mediators, leptin and ghrelin, whose levels correlate with CD4+ T-lymphocyte-derived IFN- γ .³⁴ Our observed association of weight loss and anaemia with CD4+ T-lymphocytopenia may be consequent to malnutrition observed among patients with longstanding infections.^{35,36} This is a plausible explanation in our study population whose symptoms had lasted over 90 days.

Skogmar et al also found a prevalence of CD4+ T-lymphocytopenia of 25% and 10% among HIV-negative

Table 3 Predictors of CD4+ T-Lymphocytopenia Among HIV-Negative TB Patients

Characteristics	Crude Odds Ratio		Adjusted Odds Ratio	
	cOR (95% CI)	p-value	aOR (95% CI)	p-value
Sex				
Male	1		1	
Female	0.58 (0.29–1.18)	0.133	0.72 (0.29–1.77)	0.475
Age (years)	1.02 (0.99–1.04)	0.152	1.01 (0.98–1.04)	0.479
BMI				
≥18.5	1		1	
<18.5	2.05 (1.08–3.90)	0.029	1.09 (0.47–2.51)	0.848
Anorexia				
No	1		1	
Yes	1.74 (0.90–3.35)	0.099	1.72 (0.75–3.93)	0.202
Weight loss				
No	1		1	
Yes	4.38 (1.49–12.85)	0.007	3.61 (1.07–12.23)	0.039 [†]
Hemoglobin (g/dl) ^a				
<10.8	3.59 (1.82–7.08)	<0.001	3.83 (1.59–9.23)	0.003 [†]
Normal	1		1	
>17.1	–		–	
Platelet count (cells/μL) ^a				
<106,000	1.49 (0.24–9.37)	0.668	0.85 (0.08–9.61)	0.899
Normal	1		1	
>384,000	0.49 (0.26–0.95)	0.034	0.47 (0.20–1.07)	0.073
Monocyte count (cells/μL) ^a				
<200	0.67 (0.07–6.04)	0.722	1.04 (0.10–10.85)	0.976
Normal	1		1	
>700	0.46 (0.23–0.92)	0.028	0.35 (0.14–0.89)	0.028 [†]
CD8+ Count ^b				
<256	6.74 (3.41–13.33)	<0.001	6.10 (2.68–13.89)	<0.001 [†]
Normal	1		1	
>1619	–		–	

Notes: ^aNormal ranges are for adult Ugandans.³⁹ ^bNormal ranges are for adult Ugandans.¹⁶ [†]Statistically significant.

Abbreviations: LLN, lower limit of normal; ULN, upper limit of normal; BMI, body mass index; cOR, crude odds ratio; aOR, adjusted odds ratio.

TB patients when considering a cut off of CD4+ count of <500 cells/mm³ and 350 cells/mm³ respectively in Ethiopia.¹² In contrast, Mhmoud et al found the proportion of HIV-negative TB patients with severe CD4+ T-lymphocytopenia (CD4+ count <300 cells/mm³) to be 83% in Sudan.¹⁴ Notably, these studies included participants without bacteriological confirmation of TB and different cut offs of the CD4+ count are employed. Dissimilar results can thus be expected.

Other studies have found CD4+ T-lymphocytopenia to be associated with extensive TB disease, low serum albumin level, low BMI, low haematocrit, smear positive TB,

wasting and being bed ridden.^{11,12,37} These studies also invariably included patients without bacteriological confirmation of TB. Nonetheless, similar to Jones et al (1997), we found a low BMI to be significantly associated with CD4+ T-lymphocytopenia at bivariable logistic regression analysis.¹¹ Also, since the haematocrit correlates with the haemoglobin levels,³⁸ our association of CD4+ T-lymphocytopenia with anaemia is similar to the association of a low haematocrit with CD4+ T-lymphocytopenia reported by Jones et al. Moreover, the haematocrit demonstrated a high correlation with the haemoglobin levels in our study participants (Pearson's correlation co-efficient of

0.803) hence its exclusion in the final logistic regression model in favour of the haemoglobin level.

One of the study limitations was missing data on the monocyte counts that necessitated imputation of missing data. We also did not evaluate the qualitative function of the CD4 T-lymphocytes to further characterise the immune responses among HIV-negative patients with CD4+ T-lymphocytopenia. Lastly, there might have been a referral bias of TB patients with severe disease as our study population was drawn from a national referral facility thus affecting generalizability of study findings. Nevertheless, the strength of our study lies in its utilisation of a homogenous population of bacteriologically confirmed TB cases.

Conclusion

The prevalence of CD4+ T-lymphocytopenia was high among HIV-negative TB patients. HIV-negative TB patients with weight loss, anaemia and a low CD8+ T-cell count were more likely to have CD4+ T-lymphocytopenia while those with monocytosis were less likely to have CD4+ T-lymphocytopenia. These findings suggest that HIV-negative TB patients with CD4+ T-lymphocytopenia have severe TB disease and immune dysfunction. We recommend further evaluation of the prognostic value CD4+ T-lymphocytopenia among HIV-negative TB patients.

Data Sharing Statement

Datasets generated and/or analysed during this study are available from the corresponding author upon reasonable request.

Funding

None.

Disclosure

The authors declare no conflicts of interest.

References

- Global tuberculosis report 2019. Available from: <https://www.who.int/tb/global-report-2019>. Accessed November 8, 2019
- de Martino M, Lodi L, Galli L, Chiappini E. Immune response to Mycobacterium tuberculosis: a narrative review. *Front Pediatr*. 2019;7. doi:10.3389/fped.2019.00350.
- Prezzemolo T, Guggino G, La Manna MP, Di Liberto D, Dieli F, Caccamo N. Functional signatures of human CD4 and CD8 T cell responses to Mycobacterium tuberculosis. *Front Immunol*. 2014;5. doi:10.3389/fimmu.2014.00180.
- Kumar P. IFN γ -producing CD4+ T lymphocytes: the double-edged swords in tuberculosis. *Clin Transl Med*. 2017;6(1):21. doi:10.1186/s40169-017-0151-8
- Sakai S, Kauffman KD, Sallin MA, et al. CD4 T cell-derived IFN- γ plays a minimal role in control of pulmonary Mycobacterium tuberculosis infection and must be actively repressed by PD-1 to prevent lethal disease. *PLoS Pathog*. 2016;12(5):e1005667. doi:10.1371/journal.ppat.1005667
- Yao S, Huang D, Chen CY, Halliday L, Wang RC, Chen ZW. CD4+ T cells contain early extrapulmonary tuberculosis (TB) dissemination and rapid TB progression and sustain multi-effector functions of CD8+ T and CD3- lymphocytes: mechanisms of CD4+ T cell immunity. *J Immunol*. 2014;192(5):2120–2132. doi:10.4049/jimmunol.1301373
- Ellis PK, Martin WJ, Dodd PJ. CD4 count and tuberculosis risk in HIV-positive adults not on ART: a systematic review and meta-analysis. *PeerJ*. 2017;5:e4165. doi:10.7717/peerj.4165
- Aung ZZ, Saw YM, Saw TN, et al. Survival rate and mortality risk factors among TB-HIV co-infected patients at an HIV-specialist hospital in Myanmar: a 12-year retrospective follow-up study. *Int J Infect Dis*. 2019;80:10–15. doi:10.1016/j.ijid.2018.12.008
- Al-Aska A, Al-Anazi A, Al-Subaei S, et al. CD4+ T-lymphopenia in HIV negative tuberculous patients at King Khalid University Hospital in Riyadh, Saudi Arabia. *Eur J Med Res*. 2011;16(6):285–288. doi:10.1186/2047-783X-16-6-285
- Uppal SS, Tewari SC, Verma S, Dhot PS. Comparison of CD4 and CD8 lymphocyte counts in HIV-negative pulmonary TB patients with those in normal blood donors and the effect of antitubercular treatment: hospital-based flow cytometric study. *Cytometry B Clin Cytom*. 2004;61B(1):20–26. doi:10.1002/cyto.b.20018
- Jones BE, Oo MM, Taikwel EK, et al. CD4 cell counts in human immunodeficiency virus-negative patients with tuberculosis. *Clin Infect Dis*. 1997;24(5):988–991. doi:10.1093/clinids/24.5.988
- Skogmar S, Schön T, Balcha TT, et al. CD4 cell levels during treatment for tuberculosis (TB) in Ethiopian adults and clinical markers associated with CD4 lymphocytopenia. *PLoS One*. 2013;8(12):e83270. doi:10.1371/journal.pone.0083270
- Kony SJ, Hane AA, Larouze B, et al. Tuberculosis-associated severe CD4+ T-lymphocytopenia in HIV-seronegative patients from Dakar. *J Infect*. 2000;41(2):167–171. doi:10.1053/jinf.2000.0721
- Mhmoud NA, Fahal AH, van de Sande WWJ. CD4+ T-lymphocytopenia in HIV-negative tuberculosis patients in Sudan. *J Infect*. 2012;65(4):370–372. doi:10.1016/j.jinf.2012.06.003
- Baluku JB, Nassozi S, Gyagenda B, et al. Prevalence of malaria and TB coinfection at a national tuberculosis treatment centre in Uganda. *J Trop Med*. 2019. doi:10.1155/2019/3741294
- Nanzigu S, Waako P, Petzold M, et al. CD4-T-lymphocyte reference ranges in Uganda and its influencing factors. *Lab Med*. 2011;42(2):94–101. doi:10.1309/LMFT0VCE1UGO9YGD
- Ministry of Health. National HIV Testing Services Policy and Implementation Guidelines. Published online 2016.
- Jacques MK, Jayaraman P, Kuchroo VK, Behar SM. Defining the contribution of T cell exhaustion to failed immunity during TB. *J Immunol*. 2016;196(1 Supplement):65.17.
- Zonios DI, Falloon J, Bennett JE, et al. Idiopathic CD4+ lymphocytopenia: natural history and prognostic factors. *Blood*. 2008;112(2):287–294. doi:10.1182/blood-2007-12-127878
- Lin PL, Flynn JL. CD8 T cells and Mycobacterium tuberculosis infection. *Semin Immunopathol*. 2015;37(3):239–249. doi:10.1007/s00281-015-0490-8
- Serbina NV, Flynn JL, Mansfield JM. CD8+ T cells participate in the memory immune response to Mycobacterium tuberculosis. *Infect Immun*. 2001;69(7):4320–4328. doi:10.1128/IAI.69.7.4320-4328.2001
- Canaday DH, Wilkinson RJ, Li Q, Harding CV, Silver RF, Boom WH. CD4+ and CD8+ T cells kill intracellular Mycobacterium tuberculosis by a perforin and Fas/Fas ligand-independent mechanism. *J Immunol*. 2001;167(5):2734–2742. doi:10.4049/jimmunol.167.5.2734
- Serbina NV, Lazarevic V, Flynn JL. CD4+ T cells are required for the development of cytotoxic CD8+ T cells during Mycobacterium tuberculosis infection. *J Immunol*. 2001;167(12):6991–7000. doi:10.4049/jimmunol.167.12.6991

24. Hertoghe T, Wajja A, Ntambi L, et al. T cell activation, apoptosis and cytokine dysregulation in the (co)pathogenesis of HIV and pulmonary tuberculosis (TB). *Clin Exp Immunol.* 2000;122(3):350–357. doi:10.1046/j.1365-2249.2000.01385.x
25. Sampath P, Moideen K, Ranganathan UD, Bethunaickan R. Monocyte subsets: phenotypes and function in tuberculosis infection. *Front Immunol.* 2018;9. doi:10.3389/fimmu.2018.01726.
26. Schrier SB, Hill AS, Plana D, Lauffenburger DA. Synergistic communication between CD4+ T cells and monocytes impacts the cytokine environment. *Sci Rep.* 2016;6(1):1–11. doi:10.1038/srep34942
27. YIN Y, QIN J, DAI Y, ZENG F, PEI H, WANG J. The CD4+/CD8+ ratio in pulmonary tuberculosis: systematic and meta-analysis article. *Iran J Public Health.* 2015;44(2):185–193.
28. Nagu TJ, Spiegelman D, Hertzmark E, et al. Anemia at the initiation of tuberculosis therapy is associated with delayed sputum conversion among pulmonary tuberculosis patients in Dar-es-Salaam, Tanzania. *PLoS One.* 2014;9(3):e91229. doi:10.1371/journal.pone.0091229
29. Kerkhoff AD, Wood R, Cobelens FG, Gupta-Wright A, Bekker L-G, Lawn SD. The predictive value of current haemoglobin levels for incident tuberculosis and/or mortality during long-term antiretroviral therapy in South Africa: a cohort study. *BMC Med.* 2015;13(1):1–13. doi:10.1186/s12916-015-0320-9
30. García-García M, de L, Ponce-de-León A, et al. Tuberculosis-related deaths within a well-functioning DOTS control program. *Emerg Infect Dis.* 2002;8(11). doi:10.3201/eid0811.020021
31. Filate M, Mehari Z, Alemu YM. Longitudinal body weight and sputum conversion in patients with tuberculosis, Southwest Ethiopia: a retrospective follow-up study. *BMJ Open.* 2018;8(9):e019076. doi:10.1136/bmjopen-2017-019076
32. Pilheu JA, De Salvo MC, Gonzalez J, Rey D, Elias MC, Ruppi MC. CD4+ T-lymphocytopenia in severe pulmonary tuberculosis without evidence of human immunodeficiency virus infection. *Int J Tuberc Lung Dis.* 1997;1(5):422–426.
33. Riou C, Gray CM, Lugongolo M, et al. A subset of circulating blood mycobacteria-specific CD4 T cells can predict the time to Mycobacterium tuberculosis sputum culture conversion. *PLoS One.* 2014;9(7):e102178. doi:10.1371/journal.pone.0102178
34. Zheng Y, Ma A, Wang Q, et al. Relation of leptin, ghrelin and inflammatory cytokines with body mass index in pulmonary tuberculosis patients with and without type 2 diabetes mellitus. *PLoS One.* 2013;8(11):e80122. doi:10.1371/journal.pone.0080122
35. Nájera O, González C, Toledo G, López L, Ortiz R. Flow cytometry study of lymphocyte subsets in malnourished and well-nourished children with bacterial infections. *Clin Diagn Lab Immunol.* 2004;11(3):577–580. doi:10.1128/CDLI.11.3.577-580.2004
36. Hughes S, Kelly P. Interactions of malnutrition and immune impairment, with specific reference to immunity against parasites. *Parasite Immunol.* 2006;28(11):577–588. doi:10.1111/j.1365-3024.2006.00897.x
37. Davoudi S, Rasoolinegad M, Younesian M, et al. CD4+ cell counts in patients with different clinical manifestations of tuberculosis. *Braz J Infect Dis.* 2008;12(6):483–486. doi:10.1590/S1413-86702008000600008
38. Quintó L, Aponte JJ, Menéndez C, et al. Relationship between haemoglobin and haematocrit in the definition of anaemia. *Trop Med Int Health.* 2006;11(8):1295–1302. doi:10.1111/j.1365-3156.2006.01679.x
39. Eller LA, Eller MA, Ouma B, et al. Reference intervals in healthy adult Ugandan blood donors and their impact on conducting international vaccine trials. *PLoS One.* 2008;3(12):e3919. doi:10.1371/journal.pone.0003919

Research and Reports in Tropical Medicine

Publish your work in this journal

Research and Reports in Tropical Medicine is an international, peer-reviewed, open access journal publishing original research, case reports, editorials, reviews and commentaries on all areas of tropical medicine, including: Diseases and medicine in tropical regions; Entomology; Epidemiology; Health economics issues; Infectious disease; Laboratory science and new technology in tropical medicine;

Parasitology; Public health medicine/health care policy in tropical regions; and Microbiology. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/research-and-reports-in-tropical-medicine-journal>

Dovepress