The association between adjustment disorder diagnosed at psychiatric treatment facilities and completed suicide

Jaimie L Gradus^{1,2}

Ping Qin³

Alisa K Lincoln⁴

Matthew Miller⁵

Elizabeth Lawler⁶

Timothy L Lash^{2,7}

¹National Center for PTSD VA Boston Healthcare System, Boston, MA, USA;

²Department of Epidemiology, Boston University School of Public Health, Boston, MA, USA; ³National Centre for Register-based Research, Aarhus University, Aarhus, Denmark; ⁴Health Sciences and Sociology Departments, Northeastern University, Boston, MA, USA; ⁵Department of Health Policy and Management, Harvard University School of Public Health, Boston, MA, USA; ⁶Department of Medicine, Harvard University, Boston, MA, USA; ⁷Department of Clinical Epidemiology, Aarhus University Hospital, Aarhus, Denmark

Abstract: Adjustment disorder is a diagnosis given following a significant psychosocial stressor from which an individual has difficulty recovering. The individual's reaction to this event must exceed what would be observed among similar people experiencing the same stressor. Adjustment disorder is associated with suicidal ideation and suicide attempt. However the association between adjustment disorder and completed suicide has yet to be examined. The current study is a population-based case control study examining this association in the population of Denmark aged 15 to 90 years. All suicides in Denmark from 1994 to 2006 were included, resulting in 9,612 cases. For each case, up to 30 controls were matched on gender, exact date of birth, and calendar time, yielding 199,306 controls. Adjustment disorder diagnosis was found in 7.6% of suicide cases and 0.52% of controls. Conditional logistic regression analyses revealed that those diagnosed with adjustment disorder had 12 times the rate of suicide as those without an adjustment disorder diagnosis, after controlling for history of depression diagnosis, marital status, income, and the matched factors.

Keywords: adjustment disorder, suicide, case-control study

Adjustment disorder is a diagnosis given following a significant psychosocial stressor from which an individual has difficulty recovering (eg, death of a loved one, failure to attain an important goal, termination from employment, or other similar events). To meet diagnostic criteria for this disorder, this reaction must exceed what would be observed among similar people experiencing the same stressor, and the stressor must occur no more than one month before symptom onset.¹

Suicidal behavior is more common among people diagnosed with adjustment disorder than people without this diagnosis. Patients hospitalized in a psychiatric department with an adjustment disorder diagnosis have a higher prevalence of preceding suicide attempts at admission (96%) than patients without adjustment disorders (78.6%).² Further, among adolescents admitted to a psychiatric hospital, adjustment disorder was more common among those hospitalized for a suicide attempt than among youths with no history of suicide attempt.³ This difference has been observed in other populations as well. Among patients hospitalized in Switzerland, those brought to the psychiatric emergency room for a suicide attempt were more likely to have had an adjustment disorder diagnosis (47.9%) than those brought to the psychiatric emergency room for other reasons (22.3%).⁴ Additionally, adjustment disorders were the most common diagnosis among patients under the age of 21 years in a retrospective study of patients who had made a suicide attempt in Singapore.⁵

While there is evidence of an association between adjustment disorders and attempts, the association between adjustment disorders and completed suicide has yet to

Correspondence: Jaimie L Gradus
National Center for Posttraumatic Stress Disorder, Women's Health Sciences Division, VA Boston Healthcare System, 150 South Huntington Avenue (116B-3), Boston, MA 02130, USA
Tel +1 857 364 6688
Fax +1 857 364 4515
Email jaimie.gradus@va.gov

be examined. Studies of the association between adjustment disorder and suicidal behavior are an imperfect substitute for studies of the association with suicidal behavior, because there are important differences between people who think about or attempt suicide, and people who die by suicide.⁶ Therefore, the current study aimed to examine adjustment disorder as a predictor of completed suicide. We hypothesized that people diagnosed with adjustment disorder would have a higher rate of completed suicide than people without this diagnosis.

Methods

Study population

The source population for the current study was the entire population of Denmark aged 15 to 90 years at risk for suicide between January 1st, 1994 and December 31st, 2006. Participants had to reside in Denmark for the entire calendar year before suicide (or the match date in control participants) to allow us to collect complete data on social variables. The primary criterion for identification as a case in the study was having completed suicide according to the definition below, which yielded 9,612 cases. We matched up to 30 controls from a 25% representative sample of the Danish population to suicide cases on gender, date of birth, and calendar time, yielding 199,306 controls. If 30 or fewer matched controls were available, we selected all available controls. If more than 30 controls matched a case, then we randomly selected 30 from those available.

Data collection

We obtained data for the study from the Danish national medical and administrative registries. The Cause-of-Death Register contains information on cause and date of death, and was used to identify suicide cases.⁷ The Danish Psychiatric Central Register contains dates on outpatient and inpatient psychiatric treatment in Denmark, as well as validated diagnoses, and was used to obtain data on adjustment disorder and psychiatric covariate diagnoses.⁸ Finally, we used the Civil Registration System for control participant selection,⁹ and we used the Integrated Database for Labour Market Research to obtain data on social variables.⁷ We linked these registries using the Civil Registration Number, which is assigned to all citizens and residents of Denmark.

Variable definitions

Appendix 1 lists the variables involved in the current analyses and their International Classification of Disease codes.^{10,11} Suicide is classified as a cause of death in Denmark as

follows. After any unnatural death, independent forensic medical doctors conduct an inquest. This inquest involves collecting information about the death itself and the deceased from the deceased's physician, family and friends, and the person(s) who discovered the body. Information is collected about place of death, method of dying, and presence of a suicide note. If the cause of death is still uncertain following this inquest, an autopsy is conducted. The final determination of death from suicide is made by the independent forensic doctor based on information from all of these sources. This process has been described in more detail elsewhere.¹²

We included only psychiatric diagnoses that occurred before the first adjustment disorder diagnosis as potential confounders and modifiers, as it is possible that these disorders are on the causal pathway between adjustment disorder and suicide. In addition, we used only primary diagnoses to identify both adjustment disorder and psychiatric covariates. Psychiatric covariates included depression, substance abuse, and anxiety disorders. For demographic covariates, we assessed marital status, income, and geographic location. Marital status was recorded as of November in the calendar year before suicide date or match date for control participants. Income was recorded in the calendar year before suicide date or match date for control participants, and was categorized into quartiles within strata of sex and age groups.

Statistical analyses

We calculated the frequency and proportion of cases and controls within categories of the exposure and covariates. We used conditional logistic regression used to estimate the association between adjustment disorder diagnosis and completed suicide. As risk set sampling was used to sample control participants, the odds ratios obtained from these conditional logistic regression analyses provide unbiased estimates of the corresponding rate ratios.¹³ We used change-in-estimate methods as a model building strategy to select confounders from the candidate psychiatric and demographic covariates listed above.¹⁴

To address potential misclassification of psychiatric disorders, we conducted a bias analysis to assess the impact of residual confounding by depression, substance abuse, and anxiety disorders.¹⁵ We calculated the corrected estimates using a sensitivity of 50%, 70%, and 90% and assumed that misclassification was nondifferential and independent for cases and controls because the data were recorded prospectively. In addition, we assumed that specificity would be perfect, given the extensive process for psychiatric disorder validation that occurs in Denmark. Finally, because cases may

have been treated in a psychiatric facility more than controls before committing suicide, and therefore would have more accurate psychiatric disorder confounder classification, we conducted a bias analysis to examine differential imperfect sensitivity of confounder classification among cases and controls. For this analysis, we used a sensitivity for cases of 0.92 (the minimum sensitivity that resulted in nonnegative cells in the corrected tables) and a sensitivity for controls of 0.5. We used SAS software (version 9.1.3; SAS Institute, Cary, NC) to conduct all analyses.

Results

The descriptive characteristics of the cases and controls are displayed in Table 1. Suicide cases were primarily male (71%), single (58%), and in the lowest income quartile (42%). The initial conditional logistic regression analysis, adjusting for the control-to-case matching on gender, date of birth, and calendar time, revealed that those diagnosed with adjustment disorder had 19 times the rate of suicide of people without this preceding diagnosis (95% confidence interval [CI]: 17–21). Using the change-in-estimate model

building method, we identified depression, marital status, and income as confounders of this association. Those with adjustment disorder had 12 times the rate of suicide as those without a preceding adjustment disorder, when controlling for these variables in addition to the matched factors (95% CI: 11–14).

We conducted a bias analysis to evaluate whether residual confounding by psychiatric covariates would affect the observed associations between adjustment disorder and suicide, assuming a valid bias model and the accuracy of the values assigned to the bias parameters. The results of this analysis are displayed in Table 2. At all levels of sensitivity tested, an association between adjustment disorder and suicide remained. Finally, we conducted a bias analysis to explore differential misclassification of psychiatric covariates among cases and controls, given that cases may have been treated in a psychiatric facility more than controls before committing suicide, and may therefore have a more accurately recorded psychiatric history (Table 3). Assuming a valid bias model, and accurate values assigned to parameters, it does not appear from these results that dif-

Table 1 Characteristics of suicide cases and controls, Denmark 1994–2006

	Cases (n = 9612)	Controls (n = 199,306)
Gender		
Male	6837 (71%)	142,865 (72%)
Female	2775 (29%)	56,441 (28%)
Age group		
15–34 years	1701 (18%)	42,627 (21%)
35–54 years	3608 (38%)	93,703 (47%)
55–74 years	2818 (29%)	51,176 (26%)
75–90 years	1485 (15%)	11,800 (5.9%)
Marital status		
Married	3121 (32%)	107,758 (54%)
Cohabiting	823 (8.0%)	25,130 (13%)
Single	5569 (58%)	64,471 (32%)
Unmarried minor	99 (1.0%)	1947 (1.0%)
Income		
Lowest income quartile	4014 (42%)	49,322 (25%)
Second income quartile	2242 (23%)	50,011 (25%)
Third income quartile	1786 (19%)	49,952 (25%)
Highest income quartile	1570 (16%)	50,021 (25%)
Adjustment disorder	728 (7.6%)	1040 (0.52%)
Other psychiatric diagnoses		
Depression	1713 (18%)	2542 (1.3%)
Substance abuse	1024 (11%)	2686 (1.4%)
Anxiety disorders	352 (3.7%)	1309 (0.66%)

Table 2 Bias analysis exploring the effects of imperfect nondifferential sensitivity of psychiatric disorders on the association between adjustment disorder and suicide^a

	Adjustment disorder Odds ratio
Adjusted for depression (uncorrected)	7.5
Depression sensitivity = 0.5	4.9
Depression specificity = 1.0	
Depression sensitivity = 0.7	6.2
Depression specificity = 1.0	
Depression sensitivity = 0.9	7.1
Depression specificity = 1.0	
Adjusted for substance abuse (uncorrected)	11
Substance abuse sensitivity = 0.5	7.8
Substance abuse specificity = 1.0	
Substance abuse sensitivity = 0.7	9.2
Substance abuse specificity = 1.0	
Substance abuse sensitivity = 0.9	10
Substance abuse specificity = 1.0	
Adjusted for anxiety disorders (uncorrected)	14
Anxiety disorders sensitivity = 0.5	12
Anxiety disorders specificity = 1.0	
Anxiety disorders sensitivity = 0.7	13
Anxiety disorders specificity = 1.0	
Anxiety disorders sensitivity = 0.9	14
Anxiety disorders specificity = 1.0	

Notes: ^aWe set all specificity equal to 1.0, because the probability that a person truly without these disorders would be treated at a psychiatric facility with a diagnosis of these disorders is near zero.

Table 3 Bias analysis exploring the effects of differential imperfect sensitivity of psychiatric disorders on association between adjustment disorder and suicide^a

	Adjustment disorder Odds ratio
Adjusted for depression (uncorrected)	7.5
Depression sensitivity cases = 0.92; controls = 0.5	4.9
Depression specificity = 1.0	
Adjusted for substance abuse (uncorrected)	11
Substance abuse sensitivity cases = 0.92; controls = 0.5	7.8
Substance abuse specificity = 1.0	
Adjusted for anxiety disorders (uncorrected)	14
Anxiety disorder sensitivity cases = 0.92; controls = 0.5	12
Anxiety disorder specificity = 1.0	

Notes: ^aWe set all specificity equal to 1.0, because the probability that a person truly without these disorders would be treated at a psychiatric facility with a diagnosis of these disorders is near zero.

ferential sensitivity of classification of psychiatric disorder confounders among cases and controls would have a significant impact on the results.

Discussion

Consistent with our hypotheses, we found that adjustment disorder was associated with an increased rate of completed suicide, controlling for history of depression, income, marital status, and the matched factors. Possible mechanisms that could account for this observed association include subsyndromal psychiatric symptoms, particularly depressed mood, which are associated with suicidal ideation among adolescents and with adjustment disorder.¹⁶ In addition, stressful events that may result in an adjustment disorder diagnosis, which were not assessed in the current study, are associated with suicidal behavior among adolescents and young adults.¹⁷

The current investigation has limitations worth noting. First, we restricted psychiatric covariates to the period before adjustment disorder diagnosis and only used primary diagnoses from the Psychiatric Central Register (and not secondary diagnoses). It is possible that there is residual confounding of our observed adjustment disorder and suicide associations due to underascertainment of comorbid psychiatric illness. Because of this concern, we conducted a bias analysis, and found that even with a sensitivity of confounder classification of 50%, the associations between adjustment disorder and suicide could not be explained by this type of misclassification. Second, it is possible that cases were treated in a psychiatric facility more frequently before committing suicide, and therefore would have better psychiatric confounder documentation than controls. We conducted a bias analysis to

address this concern. Assuming a valid bias model, it appears that this differential misclassification would not have had a substantial impact on the results of the current study. Finally, we only included psychiatric diagnoses that were obtained at a psychiatric setting in Denmark. It is possible to be diagnosed with a psychiatric disorder at a somatic hospital in Denmark, and therefore the exposed participants in the current study may represent more severe cases of adjustment disorder, which may in turn be more strongly associated with suicide. We examined the number of adjustment disorder cases with a primary discharge code from somatic hospitals in the same time period, and there were too few to conduct a separate set of analyses on this group to see if the pattern of results was different from those presented here.

Despite these limitations, the current study is the first to report on the relation between adjustment disorder and completed suicide. Research to date has focused on the association between adjustment disorder and suicidal ideation or suicidal attempts. The current study examined the association between adjustment disorder and completed suicide using a population-based sample and prospectively collected data. The results of this study highlight completed suicide as an important potential sequelae of adjustment disorder diagnosis – one that warrants increased attention from researchers and clinician's alike.

Acknowledgments

We would like to thank Preben Bo Mortensen for enabling access to the data and for providing research resources. The authors report no conflicts of interest in this work.

References

1. World Health Organization. *The ICD-10 Classification of Mental and Behavioural Disorders, Diagnostic Criteria for Research*. Geneva, Switzerland: World Health Organization; 1993.
2. Kryzhanovskaya L, Canterbury R. Suicidal behavior in patients with adjustment disorders. *Crisis*. 2001;22:125–131.
3. Goldston DB, Daniel SS, Reboussin BA, Reboussin DM, Kelley AE, Frazier PH. Psychiatric diagnoses of previous suicide attempters, first-time attempters, and repeat attempters on an adolescent inpatient psychiatry unit. *J Am Acad Child Adolesc Psychiatry*. 1998;37:924–932.
4. Schnyder U, Valach L. Suicide attempters in a psychiatric emergency room population. *Gen Hosp Psychiatry*. 1997;19:119–129.
5. Wai BHK, Hong C, Heok KE. Suicidal behavior among young people in Singapore. *Gen Hosp Psychiatry*. 1999;21:128–133.
6. Maris RW, Berman AL, Silverman MM. Introduction to the study of suicide. In: Maris RW, Berman AL, Silverman MM, editors. *Comprehensive Textbook of Suicidology*. New York, NY: The Guilford Press; 2000. p. 3–25.
7. Qin P, Agerbo E, Mortensen PB. Suicide risk in relation to socioeconomic, demographic, psychiatric, and familial factors: A national register-based study of all suicides in Denmark, 1981–1997. *Am J Psychiatry*. 2003;160:765–772.

8. Munk-Jørgensen P, Mortensen PB. The Danish Psychiatric Central Register. *Dan Med Bull.* 1997;44:82–84.
9. Pedersen CB, Gøtzsche H, Møller J, Mortensen PB. The Danish civil registration system. A cohort of eight million persons. *Dan Med Bull.* 2006;53:441–449.
10. World Health Organization. *Manual of the International Classification of Diseases (ICD-8)*. Geneva, Switzerland: World Health Organization; 1967.
11. Janca A, Ustun TB, van Drimmelen J, Dittmann V, Isaac M. *ICD-10 Symptom Checklist for Mental Disorders, Version 1.1*. Geneva: World Health Organization, 1994.
12. Statens Institute for Folkesundhed [National Institute for Public Health]. *Selv mord i Danmark – Markant fald i selvmord, men stigende antal selvmordsforsøg. Hvorfor? [Suicide in Denmark – a decrease in suicide completion but a increase in suicide attempt. Why?]* København: Kailow Graphic A/S; 2005.
13. Rothman KJ, Greenland S, Lash TL. Case-control studies. In: Rothman KJ, Greenland S, Lash TL, editors. *Modern Epidemiology*. 3rd ed. Philadelphia, PA: Lippincott, Williams & Wilkins; 2008. p. 111–127.
14. Greenland S. Introduction to regression modeling. In: Rothman KJ, Greenland S, Lash TL, editors. *Modern Epidemiology*. 3rd ed. Philadelphia, PA: Lippincott, Williams & Wilkins; 2008. p. 418–458.
15. Greenland S, Lash TL. Bias analyses. In: Rothman KJ, Greenland S, Lash TL, editors. *Modern Epidemiology*. 3rd ed. Philadelphia, PA: Lippincott, Williams & Wilkins; 2008:345–380.
16. Pelkonen M, Marttunen M, Henriksson M, Lönnqvist J. Suicidality in adjustment disorder: Clinical characteristics of adolescent outpatients. *Eur J Child Adolesc Psychiatry.* 2005;14:174–180.
17. Beautrais AL, Joyce P, Mulder RT. Precipitating factors and life events in serious suicide attempts among youths aged 13 through 24 years. *J Am Acad Child Adolesc Psychiatry.* 1997;36(11):1543–1551.

Appendix I Definition of variables used in the current study

	ICD-8 Code (1981–1993)	ICD-10 Code (1994–2006)
Adjustment disorder	–	F43.2
Depression		
Depressive episodes	296, 300.4	F32
Recurrent depressive disorder	296, 300.4	F33
Dysthymia	300.4	F34.1
Substance abuse		
Mental/behavioral disorders due to use of alcohol	291, 303	F10
Mental/behavioral disorders due to use of drugs	294, 304	F11–F19
Anxiety disorders		
Other anxiety disorders	300	F40
Phobic anxiety disorders	300.2	F41
Obsessive–compulsive disorder	300.3	F42
Completed suicide	–	X60–X84

Abbreviation: ICD, International Classification of Disease.

Clinical Epidemiology

Publish your work in this journal

Clinical Epidemiology is an international, peer-reviewed, open access journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification, systematic

Submit your manuscript here: <http://www.dovepress.com/clinical-epidemiology-journal>

reviews, risk & safety of medical interventions, epidemiology & biostatistical methods, evaluation of guidelines, translational medicine, health policies & economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.

Dovepress