Novel Carbapenem-Resistant *Klebsiella pneumoniae* ST147 Coharboring *bla*_{NDM-1}, *bla*_{OXA-48} and Extended-Spectrum β -Lactamases from Pakistan

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Aamir Jamal Gondal^{1,2}
Sidrah Saleem¹
Shah Jahan³
Nakhshab Choudhry⁴
Nighat Yasmin²

¹Department of Microbiology, University of Health Sciences, Lahore, Pakistan;

²Department of Biomedical Sciences, King Edward Medical University, Lahore, Pakistan; ³Department of Immunology, University of Health Sciences, Lahore, Pakistan; ⁴Department of Biochemistry, King Edward Medical University, Lahore, Pakistan

Purpose: The emergence of multidrug-resistant *Klebsiella pneumoniae* (*K. pneumoniae*) is associated with the acquisition of multiple carbapenemases. Their clonal spread is a worldwide concern due to their critical role in nosocomial infections. Therefore, the identification of high-risk clones with antibiotic resistance genes is very crucial for controlling its global spread.

Materials and Methods: A total of 227 *K. pneumoniae* strains collected during April 2018 to November 2019 were confirmed by PCR. Carbapenemases and extended-spectrum β -lactamases (ESBL) were detected phenotypically. Confirmation of carbapenemases was carried out by PCR and Sanger sequencing. The clonal lineages were assigned to selected isolates by multilocus sequence typing (MLST), and the plasmid analysis was done by PCR-based detection of the plasmid replicon typing.

Results: Of the total *K. pneumoniae*, 117 (51.5%) were carbapenem resistant (CRKP) and 140 (61.7%) were identified as ESBL producers. Intermediate to high resistance was detected in the tested β -lactam drugs while polymyxin-B and tigecycline were found to be susceptible. Among CRKP, 91 (77.8%) isolates were detected as carbapenemase producing, while 55 (47%) were positive for *bla*_{NDM-1} 23.9% (n=28), *bla*_{OXA-48} 22.2% (n=26) and *bla*_{VIM} 0.85% (n=1) while 12.7% (n=7) carried both *bla*_{NDM-1} and *bla*_{OXA-48} genes. The CRKP coharboring *bla*_{NDM-1} and *bla*_{OXA-48} genes (n=7) were positive for *bla*_{CTX-M} *bla*_{SHV} (n=3), *bla*_{SHV} (n=1) and *bla*_{CTX-M} (n=3). The novel CRKP with the coexistence of *bla*_{NDM-1}, *bla*_{OXA-48}, *bla*_{CTX-M} and *bla*_{SHV} genes were associated with the high-risk clone ST147 (n=5) and ST11 (n=2). The assigned replicon types were IncL/M, IncFII, IncA/C and IncHI1.

Conclusion: This is the first report of the coexistence of *bla*_{NDM-1}, *bla*_{OXA-48}, *bla*_{CTX-M} and *bla*_{SHV} genes on a high-risk lineage ST147 from Pakistan. This study highlights the successful dissemination of carbapenemase resistance genes in the high-risk clones that emphasizes the importance of monitoring and controlling the spread of these diverse clones globally.

Keywords: high-risk clone, New Delhi metallo- β -lactamase, MLST, *K. pneumoniae*, carbapenem resistance

Correspondence: Aamir Jamal Gondal
Department of Biomedical Sciences, King Edward Medical University, Nelagumbad Chowk, Anarkali Bazar, Lahore 54000, Pakistan
Tel +92-300-8822055
Email ajgondal119@gmail.com

Introduction

Accelerated emergence and effective propagation of carbapenem-resistant *Klebsiella pneumoniae* (CRKP) across the world have become a prominent public health challenge due to high mortality rate in healthcare-associated nosocomial

infections.^{1,2} The CRKP has a unique ability to acquire multiple resistance encoding genes through horizontal gene transfer interceded by broad-host-range plasmids, transposons and insertion sequences thereby turning out as one of the most successful nosocomial pathogen.³ Lack of stewardship and irrational use of carbapenems for the infections of ESBL producing *K. pneumoniae* has led to the evolution of transmissible plasmid-encoding resistance genes that supported the selection of high-risk clones of CRKP involving diverse geographic regions and populations.⁴ During 2014, the World Health Organization declared CRKP as the third most critical nosocomial pathogen for future concern.⁵ As carbapenemases and ESBL/AmpC β -lactamases are critical in the acquisition of multidrug resistance, the identification of such broad-spectrum resistance genes is required for the development of new intervention strategies.⁶

Clinically important carbapenemase genes encompass *bla*_{KPC-2}, *bla*_{VIM}, *bla*_{IMB}, *bla*_{NDM-1} and *bla*_{OXA-48}.⁷ Global dissemination of such plasmid-encoded carbapenemases has increased alarmingly yet their geographic prevalence varies significantly.^{8,9} Clinical literature remains expressive about the linkage of NDM-1, most common MBL-type carbapenemase and its transmission to other parts of the world from the endemic areas of Indian subcontinent since the first isolation of *bla*_{NDM-1} producing *K. pneumoniae* ST14 and *Escherichia coli* (*E. coli*) in a patient treated in India and later shifted to Sweden in 2009.¹⁰ Frequent reports of NDM-1, KPC-2 and OXA-48 type carbapenemase are available from Pakistan during the past decade.^{11,12} Similarly, the OXA-48 is endemic in several countries since its first identification from Turkey in 2001.¹³ CRKP co-harboring at least two carbapenemases were reported globally such as KPC-3 and VIM-2 in Italy, NDM-1 and KPC-2 from Brazil⁹ and Pakistan,¹² NDM-1 and OXA-48 in Morocco,¹⁴ Switzerland,¹⁵ China¹⁶ and Sultanate of Oman.¹⁷ However, the understanding of molecular and genetic context of the carbapenemases is scarce especially in the developing countries.

Carbapenemases have spread worldwide through evolution of high-risk clones by acquiring, retaining and efficiently transmitting resistance genes. Such globally identified high-risk *K. pneumoniae* clones for the dissemination of carbapenemases include ST258, ST11 and ST147 co-harboring broad range of plasmids. Several STs were found to be associated with *bla*_{NDM-1} producing *K. pneumoniae* such as ST258, ST340, ST512 and ST147 along with different plasmids IncF, IncA/C

and IncL/M.⁹ Therefore, careful detection and treatment strategies are required especially in developing countries where carbapenemase-producing strains have diverse opportunities.

However, insufficient data are available from Pakistan that describes molecular versatility of resistance genes in relation to genetic analysis and prevalence of high-risk clones. Hence, it is imperative to promptly detect and examine these successful clones to get insights into the global spread of antimicrobial drug resistance. Therefore, the current study aimed to ascertain the prevalence of carbapenemases and to analyze their clonal relatedness.

Materials and Methods

Bacterial Collection and Identification

Clinical strains were collected during the course of routine diagnostic bacterial cultures from tertiary care hospitals of Lahore, Pakistan. A total of 227 clinical strains of *K. pneumoniae* were included from different sample types from April 2018 to November 2019. The isolates were characterized phenotypically by colony morphology, Gram's staining and biochemical characteristics by using API-20E according to the manufacturer's instructions (BioMerieux, France). The study was sanctioned by institutional review board of the University of Health Sciences, Lahore, Pakistan.

Antimicrobial Susceptibility Testing (AST)

AST was carried out by standard disc diffusion method according to the CLSI guidelines using the following antibiotic discs: imipenem (IPM), meropenem (MEM), ertapenem (ETP), ceftazidime (CAZ), ampicillin (AMP), amoxicillin-clavulanic acid (AMC), cefepime (FEP), cefotaxime (CTP), aztreonam (ATM), gentamicin (CN), amikacin (AK), ciprofloxacin (CIP), doxycycline (DO), polymyxin-B (PB), tigecycline (TGC), cefotaxime (CTX), trimethoprim-sulfamethoxazole (SXT) and piperacillin-tazobactam (TZP) (Oxoid, UK). *E. coli* ATCC 25922 and *Pseudomonas aeruginosa* ATCC 27853 were used as quality control strains.¹⁸

Minimal Inhibitory Concentrations (MICs)

MIC of antibiotics was determined by standard broth micro-dilution method using cation-adjusted Mueller-Hinton broth in accordance with CLSI guidelines with antibiotic concentrations ranging from 0.5 to 1024 μ g/mL.¹⁸

Phenotypic Characterization

Carbapenemases were identified phenotypically by carbapenem inactivation method (CIM) while the detection of ESBLs was carried out by double-disc synergy test (DDST) using amoxicillin-clavulanic acid alone and in combination with ceftazidime as per the guidelines of CLSI.¹⁸

DNA Isolation from Bacterial Strains

Genomic DNA was prepared from pure bacterial culture plates by heat lysis method as reported previously and stored at -20°C for onward processing.¹⁹

Molecular Profile Analysis by Polymerase Chain Reaction (PCR)

Klebsiella species, carbapenemase resistance genes (*bla*_{NDM-1}, *bla*_{VIM}, *bla*_{IMP} and *bla*_{OXA-48}) and ESBL encoding genes (*bla*_{TEM}, *bla*_{CTX-M} and *bla*_{SHV}) were detected through PCR using specific primers as given in Table 1. PCR reaction mixture of 50 μL consisting of 25 μL of 2x PCR Master Mix, 1 μL of each primer, 2 μL of DNA and dH₂O upto 50 μL . Amplification was carried out in thermal cycler (Proflex, ABI) with different annealing temperatures given in Table 1. Agarose gel (1–1.5%) was used to resolve and analyze the PCR products. *bla*_{NDM-1} genes were further analyzed for allelic discrimination by Sanger's sequencing method. *K. pneumoniae* ATCC BAA-2146 was used as NDM positive control.

Multilocus Sequence Typing (MLST) of *Klebsiella* Species

MLST of CRKP strains coharboring *bla*_{NDM-1} and *bla*_{OXA-48} was performed using seven housekeeping genes (*gapA*, *infB*, *mdh*, *pgi*, *phoE*, *rpoB* and *tonB*) as described by *K. pneumoniae* MLST website.²⁰ The mutation analysis of NDM-1 was carried out by cycle sequencing using specific primers given in Table 1. The cycle sequencing was performed by BigDye terminator v3.1 kit on Proflex thermal cycler while the sequencing products were analyzed through capillary electrophoresis on Genetic Analyzer AB-3500 (Life Technologies by Thermofisher, USA) as per the kit instructions. Data were analyzed by using the sequencing analysis software v6.1 and checked on basic local alignment (BLAST) at NCBI for allele identification. CRKP STs were assigned using the MLST database (<http://bigsdB.pasteur.fr/klebsiella/klebsiella.html>).

Plasmid Analysis

CRKP coharboring *bla*_{NDM-1} and *bla*_{OXA-48} were further analyzed for the presence of plasmids. Plasmid DNA was extracted from single colony of CRKP by using the plasmid isolation kit (Thermofisher Scientific) and DNA was stored at -20°C . The samples were run on 0.8% agarose gel for the detection of plasmids. Plasmids were classified according to their incompatibility groups by using the PCR-based replicon typing method as described before.²¹

Results

During the 19-month study period, 227 clinical strains of *Klebsiella* were identified by phenotypic and genotypic methods. Of the total, 129 (56.8%) were isolated from males while remaining 98 (43.2%) were from females. These isolates were collected from wound 29.5% (n=67), pus 17.6% (n=40), blood 15.5% (n=35), tracheal secretion 13.2% (n=30), sputum 11% (n=25), urine 7.04% (n=16) and tissue 6.16% (n=14). Isolates originated from different sections of the hospital such as general surgery 26.4% (n=60), SICU 18.5% (n=42), general medicine 15.8% (n=36), dermatology 5.72% (n=13), nephrology 4.40% (n=10), chest medicine 8.37% (n=19), cardiology 3.96% (n=9), pediatric medicine 7.04% (n=16), oncology 5.28% (n=12) and orthopedic surgery 4.40% (n=10).

Antimicrobial Susceptibility Testing and Phenotypic Confirmatory Tests

As high as 51.5% (n=117) clinical strains of *K. pneumoniae* were carbapenem resistant (CRKP) while remaining 48.5% (n=110) were susceptible (CS). Out of 117 CRKP, 77.7% (n=91) were detected as carbapenemase-producing strains (CPKP). Most of the CRKP exhibited resistant to intermediate resistant profile for the β -lactam combination agents, carbapenems, fluoroquinolones, aminoglycosides and trimethoprim/sulfamethoxazole. Antimicrobial resistance pattern of CRKP strains was as follows: meropenem (96.9%), imipenem (98%), ertapenem (90%), amoxicillin/clavulanic acid (93.5%), ceftazidime (91.2%), ceftriaxone (96%), cefotaxime (95%), aztreonam (90.3%), ciprofloxacin (87%), amikacin (37.1%), tigecycline (21.1%) and polymyxin-B (13.7%). The MIC values of the tested β -lactam antibiotics were as follows: 4 to >1024 mg/L for ertapenem, 8 to >1024 mg/L for meropenem, 8 to >1024 mg/L for imipenem in all tested strains. All of the isolates were recognized as MDR (72%) or XDR (28%). The MIC results of the selected strains are given in Table 2.

Table 1 Primer Sequences Used for PCR

Gene	Primer Sequence (5'–3')	Annealing (°C)	Amplicon Size (bp)	Reference
<i>rpoB</i> (<i>K. pneumoniae</i>)	F: CAA CGG TGT GGT TAC TGA CG R: TCT ACG AAG TGG CCG TTT TC	55	108	53
<i>pehX</i> (<i>K. oxytoca</i>)	F: GAT ACG GAG TAT GCC TTT ACG GTG R: TAG CCT TTA TCA AGC GGA TAC TGG	55	343	53
<i>gyrA</i> (<i>Klebsiella</i> genus)	F: CGC GTA CTA TAC GCC ATG AAC GTA R: ACC GTT GAT CAC TTC GGT CAG G	55	441	53
<i>bla_{SHV}</i>	F: CTT TAT CGG CCC TCA CTC AA R: AGG TGC TCA TCA TGG GAA AG	55	237	54
<i>bla_{TEM}</i>	F: CGC CGC ATA CAC TAT TCT CAG AAT GA R: ACG CTC ACC GGC TCC AGA TTT AT	55	445	54
<i>bla_{CTX-M}</i>	F: ATG TGC AGY ACC AGT AAR GTK ATG GC R: TGG GTR AAR TAR GTS ACC AGA AYC AGC GG	55	593	54
<i>bla_{NDM-I}</i>	F: ATG GAA TTG CCC AAT ATT ATG CAC R: TCA GCG CAG CTT GTC GGC	52	813	55
<i>bla_{VIM}</i>	F: GAT GGT GTT TGG TCG CAT A R: CGA ATG CGC AGC ACC AG	52	390	56
<i>bla_{OXA-48}</i>	F: GCG TGG TTA AGG ATG AAC AC R: CAT CAA GTT CAA CCC AAC CG	52	438	56
<i>bla_{IMP}</i>	F: GGA ATA GAG TGG CTT AAY TCT C R: GGT TTA AYA AAA CAA CCA CC	52	232	56
<i>bla_{NDM-I}</i>	F: TGGCTTTTGAACTGTCGCACC R: CTGTCACATCGAAATCGCGCGA	60	1000	57
<i>gapA</i>	F: TGA AGT ATG ACT CCA CTC ACG G R: AAC GCC TTT CAT TGC GCC TTC GGA A	60	662	20
<i>infB</i>	F: CTC TCT GCT GGA CTA CAT TCG R: CGC TTT CAG CTC CAG AAC TTC	52	462	20
<i>mdh</i>	F: CCC AAC TGC CTT CAG GTT CAG R: CCT TCC ACG TAG GCG CAT TCC	52	756	20
<i>pgi</i>	F: GAG AAA AAC CTG CCG GTG CTG CTG R: CGG TTA ATC AGG CCG TTA GTG GAG C	52	566	20
<i>phoE</i>	F: ACC TGG CGC AAC ACC GAT TTC TTC R: TTC AGC TGG TTG ATT TTG TAA TCC AC	52	602	20
<i>rpoB</i>	F: GGC GAA ATG GCG GAA AAC CA R: GAG TCT TCG AAG TTG TAA CC	52	1075	20
<i>tonB</i>	F: CTC TAT ACT TCG GTA CAT CAG GTT R: CCT GTT TGG CGG CCA GCA CCT GGT	48	539	20

Abbreviations: *rpoB*, RNA polymerase beta-subunit gene; *pehX*, polygalacturonase gene; *gyrA*, DNA gyrase subunit A gene; *bla_{SHV}*, beta-lactamase sulfhydryl reagent variable gene; *bla_{NDM-I}*, New Delhi metallo-beta-lactamase gene; *bla_{VIM}*, metallo-beta-lactamase verona integron gene; *bla_{TEM}*, beta-lactamase temoneira gene; *bla_{OXA-48}*, beta-lactamase oxacillinase 48 gene; *bla_{CTX-M}*, beta-lactamase cefotaxime munich gene; *bla_{IMP}*, beta-lactamase imipenemase gene; *gapA*, glyceraldehyde-3-phosphate dehydrogenase A gene; *infB*, translation initiation factor IF-2 gene; *mdh*, malate dehydrogenase gene; *pgi*, phosphoglucose isomerase gene; *phoE*, phosphoprotein E gene; *tonB*, periplasmic energy transducer gene; *rpoB*, beta-subunit of RNA polymerase gene.

Table 2 MIC Values of Selected Carbapenem-Resistant *K. pneumoniae* Strains

Strain ID	Resistance Profile	MIC (µg/mL)											
		MEM	CAZ	SXT	AMP	AMC	CPT	CIP	ATM	AK	PB	DO	CTX
KP-17	MDR	4	8	2/38	64	32/16	0.5	1	16	16	2	4	1
KP-97	XDR	512	32	8/152	8	8/4	16	8	32	8	8	16	8
KP-104	MDR	64	32	2/38	8	16/8	0.5	1	4	64	2	4	8
KP-188	MDR	32	64	2/38	16	32/16	0.5	1	4	16	2	4	32
KP-191	MDR	16	4	16/304	8	32/16	0.5	1	8	16	2	2	16
KP-194	MDR	16	4	4/76	16	8/4	0.5	1	4	32	2	2	8
KP-199	XDR	>1024	128	32/608	512	256/128	0.5	64	128	128	8	32	256
KP-222	MDR	256	128	2/38	64	32/16	0.5	1	4	8	2	4	32
KP-246	XDR	128	256	8/152	32	64/32	16	1	16	64	16	64	64
KP-268	XDR	512	256	16/304	128	16/8	16	64	64	1024	4	128	256
KP-272	MDR	256	64	2/38	8	8/4	0.5	1	32	64	2	4	32
KP-284	MDR	64	32	2/38	32	8/4	0.5	16	4	16	2	4	32
KP-289	XDR	64	16	8/152	32	256/128	32	128	128	512	8	512	8
KP-315	XDR	>1024	256	8/152	64	128/64	16	4	512	512	8	256	4
KP-326	MDR	128	64	2/38	16	8/4	0.5	0.5	32	16	2	4	16
KP-333	MDR	32	16	8/152	8	8/4	0.5	0.5	16	16	2	4	2
KP-426	MDR	8	4	2/38	16	16/8	0.5	1	16	32	2	4	0.5
KP-443	MDR	64	32	4/76	8	128/64	0.5	1	16	8	2	4	0.5
KP-465	MDR	128	128	2/38	16	8/4	0.5	1	128	16	32	4	0.5
KP-494	MDR	64	128	2/38	8	8/4	4	32	64	16	2	4	0.5
KP-544	XDR	32	64	32/608	32	128/64	0.5	1	256	512	16	128	256
KP-562	MDR	64	256	2/38	64	8/4	0.5	1	32	128	2	4	0.5
KP-611	XDR	256	64	64/1216	64	128/64	0.5	16	16	64	2	4	64
KP-663	MDR	128	512	32/608	64	8/4	0.5	1	64	8	2	4	0.5
KP-675	MDR	32	32	2/38	256	8/4	0.5	1	64	8	8	4	0.5
KP-668	MDR	64	32	2/38	8	64/32	2	1	128	16	1	4	0.5
KP-687	MDR	32	64	2/38	128	8/4	8	1	32	16	1	4	0.5
KP-704	XDR	512	128	64/1216	128	256/128	0.5	256	32	1024	16	512	32

Abbreviations: MDR, multidrug resistant; XDR, extensively drug resistant.

Significantly higher frequency of CPKP was observed in wound samples 49.4% (n=45; $p=0.002$), pus samples 27.4% (n=25; $p<0.026$) and tracheal secretion samples 23.2% (n=21; $p=0.029$). Clinical strains of CRKP from wound and pus samples were significantly associated with the general surgery ($p<0.001$) while those from tracheal secretion samples were significantly associated with the SICU ($p=0.008$) as compared to the other samples obtained from the general surgery and SICU. Out of 227 *K. pneumoniae* strains, 61.6% (n=140) were ESBL producers and 38.3% (n=87) were non-ESBL producers. Among the 140 strains of ESBL producing *K. pneumoniae*, 9.28% (n=13) isolates were resistant to one of the third-generation cephalosporins (3GCs), 28.5% (n=40) were resistant against 2 of the 3GCs and 62.1% (n=87) were resistant to all the 3GCs. Association analysis demonstrated that 80% (n=112) ESBL producers were collected

from the samples of wound, pus and tracheal secretions ($p=0.003$).

Antibiotic Resistance Genes

Out of 117 CRKP, 47% (n=55) were positive for the carbapenemase resistance genes by PCR including *bla*_{NDM-1} 23.9% (n=28), *bla*_{OXA-48} 22.2% (n=26) and *bla*_{VIM} 0.85% (n=1); *bla*_{IMP} was not detected. However, 12.7% (n=7) of CPKP harbored *bla*_{NDM-1} and *bla*_{OXA-48} genes. *bla*_{NDM-1} positive strains were further confirmed by DNA sequencing. The presence of the β -lactamase-encoding genes *bla*_{CTX-M}, *bla*_{TEM} and *bla*_{SHV} was detected in the ESBL-producing *K. pneumoniae* strains (n=140). Single ESBL gene was detected in 38.5% (n=54): *bla*_{CTX-M} 10.7% (n=15), *bla*_{SHV} 22.8% (n=32), *bla*_{TEM} 5% (n=7) and double ESBL genes were detected in 61.4% (n=86): *bla*_{CTX-M}, *bla*_{SHV} 32.1% (n=45), *bla*_{TEM}, *bla*_{SHV} 12.1% (n=17), *bla*_{TEM}, *bla*_{CTX-M}

Table 3 Resistance Profile of Carbapenem-Resistant *K. pneumoniae*

Strain ID	Ward	Sample Type	MIC (µg/mL) Imipenem	Resistance Profile	Profile of Resistance Genes	Replicon and Sequence Type
KP-17	SICU	Wound	256	MDR	<i>bla</i> _{NDM-1} (MT312213)*, <i>bla</i> _{OXA-48} , <i>bla</i> _{CTX-M} , <i>bla</i> _{SHV}	ST147, IncL/M, IncFII, IncA/C, IncFHI
KP-97	SICU	Pus	64	XDR	<i>bla</i> _{NDM-1} (MT320894)*, <i>bla</i> _{CTX-M} , <i>bla</i> _{SHV}	-
KP-104	OPD	Wound	32	MDR	<i>bla</i> _{NDM-1} (MT320895)*, <i>bla</i> _{CTX-M} , <i>bla</i> _{SHV}	-
KP-188	SICU	Tracheal secretion	64	MDR	<i>bla</i> _{NDM-1} (MT320896)*, <i>bla</i> _{SHV}	-
KP-191	GS	Wound	16	MDR	<i>bla</i> _{NDM-1} (MT320897)*, <i>bla</i> _{CTX-M}	-
KP-194	GS	Wound		MDR	<i>bla</i> _{NDM-1} (MT320898)*, <i>bla</i> _{SHV}	-
KP-199	OPD	Pus	>1024	XDR	<i>bla</i> _{NDM-1} (MT320899)*, <i>bla</i> _{OXA-48} , <i>bla</i> _{CTX-M} , <i>bla</i> _{SHV}	ST147, IncL/M, IncFII, IncA/C
KP-222	GS	Sputum	128	MDR	<i>bla</i> _{NDM-1} (MT320900)*, <i>bla</i> _{OXA-48} , <i>bla</i> _{CTX-M}	ST147, IncL/M, IncFII, IncA/C, IncFHI
KP-246	GS	Wound	32	XDR	<i>bla</i> _{NDM-1} (MT320901)*, <i>bla</i> _{CTX-M}	-
KP-268	SICU	Tip cells	64	XDR	<i>bla</i> _{NDM-1} (MT320902)*, <i>bla</i> _{TEM} , <i>bla</i> _{CTX-M} , <i>bla</i> _{SHV}	-
KP-272	GS	Wound	32	MDR	<i>bla</i> _{NDM-1} (MT320903)*, <i>bla</i> _{CTX-M} , <i>bla</i> _{SHV}	-
KP-284	GS	Wound	8	MDR	<i>bla</i> _{NDM-1} (MT320904)*, <i>bla</i> _{CTX-M} , <i>bla</i> _{SHV}	-
KP-289	GS	Wound	16	XDR	<i>bla</i> _{NDM-1} (MT320905)*, <i>bla</i> _{TEM}	-
KP-315	SICU	Tracheal secretion	32	XDR	<i>bla</i> _{NDM-1} (MT320906)*, <i>bla</i> _{SHV}	-
KP-326	OPD	Blood	128	MDR	<i>bla</i> _{NDM-1} (MT320907)*, <i>bla</i> _{OXA-48} , <i>bla</i> _{CTX-M}	ST147, IncL/M, IncFII, IncA/C, IncFHI
KP-333	GS	Pus	32	MDR	<i>bla</i> _{NDM-1} (MT320908)*, <i>bla</i> _{SHV}	-
KP-426	BURN	Tracheal secretion	64	MDR	<i>bla</i> _{NDM-1} (MT320909)*, <i>bla</i> _{CTX-M}	-
KP-443	GS	Wound	64	MDR	<i>bla</i> _{NDM-1} (MT320910)*	-
KP-465	WSW	Wound	128	MDR	<i>bla</i> _{NDM-1} (MT320911)*, <i>bla</i> _{OXA-48} , <i>bla</i> _{SHV}	ST11, IncL/M, IncFII, IncA/C, IncHI
KP-494	SICU	Tracheal secretion	64	MDR	<i>bla</i> _{NDM-1} (MT320912)*, <i>bla</i> _{OXA-48} , <i>bla</i> _{CTX-M} , <i>bla</i> _{SHV}	ST147, IncL/M, IncFII, IncA/C
KP-544	GS	Blood	32	XDR	<i>bla</i> _{NDM-1} (MT320913)*	-
KP-562	SICU	Tracheal secretion	64	MDR	<i>bla</i> _{NDM-1} (MT320914)*, <i>bla</i> _{CTX-M}	-
KP-611	PS	Pus	128	XDR	<i>bla</i> _{NDM-1} (MT320915)*, <i>bla</i> _{CTX-M}	-
KP-663	GS	Wound	64	MDR	<i>bla</i> _{NDM-1} (MT320916)*, <i>bla</i> _{SHV} , <i>bla</i> _{CTX-M}	-
KP-675	SICU	Tracheal secretion	128	MDR	<i>bla</i> _{NDM-1} (MT320917)*, <i>bla</i> _{OXA-48} , <i>bla</i> _{CTX-M}	ST11, IncL/M, IncFII, IncA/C

(Continued)

Table 3 (Continued).

Strain ID	Ward	Sample Type	MIC (μg/mL) Imipenem	Resistance Profile	Profile of Resistance Genes	Replicon and Sequence Type
KP-668	GS	Pus	64	MDR	<i>bla</i> _{NDM-1} (MT320918)*, <i>bla</i> _{TEM-1} , <i>bla</i> _{SHV}	-
KP-687	SICU	Tracheal secretion	32	MDR	<i>bla</i> _{NDM-1} (MT320919)*, <i>bla</i> _{SHV}	-
KP-704	CM	Sputum	32	XDR	<i>bla</i> _{NDM-1} (MT320920)*, <i>bla</i> _{CTX-M}	-

Note: *GenBank Accession Number.

Abbreviations: GS, general surgery; PS, plastic surgery; SICU, surgical ICU; CM, chest medicine.

15% (n=21) and *bla*_{TEM}, *bla*_{SHV}, *bla*_{CTX-M} 2.14% (n=3). The CRKP strains coharboring *bla*_{NDM-1} and *bla*_{OXA-48} genes (n=7) were positive for *bla*_{CTX-M}, *bla*_{SHV} (n=3), *bla*_{SHV} (n=1) and *bla*_{CTX-M} (n=3). The results are shown in Table 3.

Sequence Type Analysis and Plasmid Detection of NDM-I Producing Isolates

CRKP coharboring *bla*_{NDM-1} and *bla*_{OXA-48} (n=7) were further analyzed for sequence typing. High-risk *K. pneumoniae* clones ST147 coharbored *bla*_{NDM-1}, *bla*_{OXA-48}, *bla*_{CTX-M}, *bla*_{SHV} (n=3), *bla*_{NDM-1}, *bla*_{OXA-48}, *bla*_{CTX-M} (n=2), while ST11 coharbored *bla*_{NDM-1}, *bla*_{OXA-48}, *bla*_{SHV} (n=1) and *bla*_{NDM-1}, *bla*_{OXA-48}, *bla*_{CTX-M} (n=1). Plasmid analysis of CRKP coharboring *bla*_{NDM-1} and *bla*_{OXA-48} identified the following replicon types: IncL/M, IncFII, IncA/C and IncH1.²¹

Discussion

The emergence of CRKP has resulted in limited effective treatment strategies thus posing a major healthcare threat worldwide.¹ Global dissemination of transmissible carbapenemases by virtue of horizontal gene transfer involving certain high-risk clones⁹ has become alarming especially in developing countries in the backdrop of inconsistent antibiotic policies. *K. pneumoniae* is among one of the most commonly detected multidrug-resistant member of the Enterobacteriaceae family.²²

In the present study, we identified 51.1% CRKP strains that consisted of 77.7% carbapenemase producers. A large-scale study conducted in Turkey has detected only 3.1% (n=45/1452) CRKP isolates,² while from the European cohort study, 55% (n=944/1717) isolates were carbapenem resistant and 39.84% (n=684/1717) were carbapenemase producers.²³ Similarly, 10.69% (n=247/2310) CRKP strains were reported previously.²⁴ However, the highest percentages of carbapenem resistant and carbapenemase producers

are reported from Pakistan such as another study identified 88% carbapenemase producers.²⁵ Due to the presence of high carbapenem resistance among *K. pneumoniae* in Pakistan, it is tempting to speculate that *K. pneumoniae* strains have the ability to retain diverse resistance determinants especially in a situation of uncontrolled use of high amounts of antibiotics. Wound samples (49.4%) were the major source of the CRKP infection that were significantly associated with the general surgery ward. In line with our study, 40% of wound samples with carbapenemase production were reported recently in association with the emergency department.²⁶ However, blood, urine, sputum, tracheal secretion and pus were the major source of CRKP in other studies.^{11,24,25,27–29} The identification of CRKP strains from different anatomical sites highlights the importance of diverse set of sampling sites for the surveillance studies.

In consistent with the previous studies,^{2,11,30–32} the most effective antibiotics against the isolates were polymyxin-B (13.7%) and tigecycline (21.1%). However, intermediate to high resistance levels were observed against carbapenemases (meropenem, imipenem and ertapenem) 90% to 98%, cephalosporins 86% to 92%, aztreonam 90.3%, ciprofloxacin 87% and amikacin 37.1% that counts for 72% MDR and 28% XDR isolates. Sattar et al¹² have reported 45% MDR *K. pneumoniae* strains with 85% to 90% resistance to cephalosporins and 30% resistance to imipenem. Another study from Pakistan reported 22.5% MDR *K. pneumoniae* strains among the study population in 2013.³³ The detailed analysis of antibiotic resistance among the *K. pneumoniae* from Pakistan suggested that the resistance has been increasing.

The most frequently detected carbapenemases among *K. pneumoniae* are *bla*_{KPC} enzymes followed by *bla*_{NDM-1}, *bla*_{OXA-48}-like and *bla*_{VIM} in *K. pneumoniae*.²² In our study, the detailed resistome analysis revealed the presence of carbapenemase resistance genes in 55 out of 117 CRKP strains. The most commonly detected carbapenemase

genes were *bla*_{NDM-1} (23.9%; n=28/55) and *bla*_{OXA-48} (22.2%; n=26/55) while *bla*_{VIM} was identified in only 1 isolate and *bla*_{IMP} was not detected. In consistent with our results, several studies from Pakistan reported that the most prevalent carbapenemase genes in Enterobacteriaceae/*K. pneumoniae* are *bla*_{NDM-1} 83.3% (n=30/37),²⁵ 70% (n=10),²⁸ 14.6% (n=13/82)³⁴ followed by *bla*_{OXA-48} 86% (n=49/57),³⁵ 50% (n=5/10)²⁸ and *bla*_{VIM} 13.4% (n=11/82),³⁴ 3.5% (n=2/57).³⁵ The results of our study are also in line with the observations that India, Bangladesh and Pakistan are the major reservoir countries for the widespread dissemination of carbapenemase genes such as *bla*_{NDM-1} and *bla*_{OXA-48}.³⁶ Since the first report of *bla*_{NDM-1} detection in Pakistan in 2010,³⁷ carbapenemase genes have spread significantly. Moreover, in the present study, the *bla*_{NDM-1} and *bla*_{OXA-48} coproduction was detected in 7 out of 55 CRKP. The co-occurrence of *bla*_{NDM-1} and *bla*_{OXA-48} has been reported previously in Asian and European countries.^{16,24,38–40} However, in Pakistan the coexistence of carbapenem-resistant genes is not commonly detected in *K. pneumoniae*. In clinical isolates of *K. pneumoniae*, *bla*_{KPC-2}, *bla*_{NDM-1} (n=2/20),¹² *bla*_{NDM-1}, *bla*_{OXA-48} (n=2/10)²⁸ and *bla*_{VIM}, *bla*_{NDM-1} (n=4/28) encoding genes in community-based *E. coli* isolates⁴¹ have been identified.

The CRKP strains coharboring *bla*_{NDM-1} and *bla*_{OXA-48} examined in our study belonged to either ST11; single locus variant of ST258⁴² or emerging ST147 high-risk CRKP clone with resistance genes located on different plasmids.⁹ NDM-type carbapenemases have been described associated with ST11, ST14, ST147, ST340, ST149 and ST231.² The ST11 is typically associated with the acquisition of multidrug resistance due to its ability to capture multiple plasmids⁴² and *K. pneumoniae* strains with multiple resistance genes have been reported previously.⁴³ In concurrence with this study, our data also revealed that different antimicrobial resistance and replicon type exist within the identified ST11 isolates. One of the ST11 isolate was polymyxin-B resistant while other was susceptible. Our results are in line with the previously reported study from Pakistan where out of 3 ST11 strains, 2 strains were colistin resistant and 1 strain was colistin susceptible.²⁸ The ST11 isolates identified in our study coharbored *bla*_{NDM-1} and *bla*_{OXA-48} genes, whereas ST11 isolates positive for *bla*_{NDM-1} (n=7), *bla*_{NDM-7} (n=2) and *bla*_{NDM-5} (n=1) were recently reported from Pakistan.²⁵

Previously, the pandemic lineage ST147 in *K. pneumoniae* has been correlated with the spread of

carbapenemase resistant genes such as *bla*_{CTX-M}, *bla*_{VIM}, *bla*_{OXA-48}, *bla*_{KPC} and *bla*_{NDM-1}.^{44,45} ST147 has also been associated with *bla*_{NDM-1}, *bla*_{CTX-M}, *bla*_{SHV},³⁰ *bla*_{CMY-4}, *bla*_{OXA-48},⁴⁶ *bla*_{NDM-1}, *bla*_{OXA-48},³⁸ *bla*_{NDM-1}⁴⁷ and *bla*_{OXA-48}.²⁴ Antecedently, two studies are available from Pakistan that reported the existence of ST147 *K. pneumoniae* with *bla*_{OXA181} resistant gene¹¹ and ST147 *K. pneumoniae* isolate with *bla*_{NDM-1}, *bla*_{NDM-5}.²⁵ The coexistence of *bla*_{NDM-1}, *bla*_{OXA-48} has also been detected in ST307 from China¹⁶ and *bla*_{NDM-1}, *bla*_{OXA-232} in ST231 from Pakistan.²⁷ However, in our study we have identified the co-emergence of *bla*_{NDM-1}, *bla*_{OXA-48}, *bla*_{CTX-M}, *bla*_{SHV} among ST147 (n=5), a globally spread high-risk clone. The identification of S147 with *bla*_{NDM-1}, *bla*_{OXA-48}, *bla*_{CTX-M}, *bla*_{SHV} is alarming as it indicates that strong selection has occurred towards the resistance in these clinical isolates from Pakistan.

Subsequently, four replicon types IncL/M, IncFII, IncA/C and IncHI were detected in the present study. Previous studies have shown that IncL/M-type plasmid was related to the OXA-48-type carbapenemases and responsible for the *bla*_{OXA-48} gene dissemination.⁴⁸ The molecular studies have reported that the most frequent replicon type identified in *K. pneumoniae* species is IncFII replicon⁹ while IncA/C type replicons are responsible for the horizontal spread of NDM-type carbapenemase along with IncFIIK, IncL/M and IncHI.⁴⁹ Moreover, among the typed resistant plasmids, IncL/M and IncFII plasmids may be regarded as epidemic as they have been detected in different countries with different origins and sources.⁵⁰ On the other hand, IncR, IncFIIK-type and IncA/C type replicons have been identified in OXA-48-type carbapenemases and IncR type replicons in NDM-type carbapenemases.⁵¹ Previously reported replicon types among the NDM-producing *K. pneumoniae* from Pakistan include IncN, IncA/C⁵² and IncFII, IncR.¹¹ In our study, the identified replicon types (IncL/M, IncFII, IncA/C and IncHI) are reported to be responsible for OXA-48-type and NDM-type carbapenemases dissemination.

Conclusion

We reported the first identification of high-risk CRKP clone ST147 coharboring several carbapenem resistance genes *bla*_{NDM-1}, *bla*_{OXA-48}, *bla*_{CTX-M}, *bla*_{SHV} from Pakistan. Taken into account the presence of highest genetic diversity among *K. pneumoniae* worldwide, the identification of high-risk clone with multidrug resistance and coexistence of different classes of β -lactamases in the same strain highlight the severity of health challenges

posed by *K. pneumoniae* worldwide. Our findings suggested that the high antimicrobial resistance existed among study isolates that can also be associated with the presence of several β -lactams genes in a high-risk clone. Therefore, the continuous monitoring of carbapenemases is necessary to prevent the national and transnational spread of these powerful isolates especially in case when the healthcare facilities are inadequate.

Acknowledgments

We are thankful to the University of Health Sciences, Lahore, and King Edward Medical University, Lahore for the provision of samples and research facilities.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published and agreed to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- Pitout JD, Nordmann P, Poirel L. Carbapenemase-producing *Klebsiella pneumoniae*, a key pathogen set for global nosocomial dominance. *Antimicrob Agents Chemother*. 2015;59(10):5873–5884. doi:10.1128/AAC.01019-15
- Davarcı İ, Şenbayrak S, Aksaray S, Koçoğlu ME, Kuşkucu MA, Samastı M. Molecular epidemiology of carbapenem-resistant *Klebsiella pneumoniae* isolates. *Anatol Clin*. 2019;24(1):1–7.
- Ye Y, Xu L, Han Y, Chen Z, Liu C, Ming L. Mechanism for carbapenem resistance of clinical Enterobacteriaceae isolates. *Exp Ther Med*. 2018;15(1):1143–1149. doi:10.3892/etm.2017.5485
- Durdu B, Hakyemez IN, Bolukcu S, Okay G, Gultepe B, Aslan T. Mortality markers in nosocomial *Klebsiella pneumoniae* bloodstream infection. *Springerplus*. 2016;5(1):1892. doi:10.1186/s40064-016-3580-8
- Organization WH. *Antimicrobial Resistance: Global Report on Surveillance*. World Health Organization; 2014.
- Jia X, Dai W, Ma W, et al. Carbapenem-resistant *E. cloacae* in Southwest China: molecular analysis of resistance and risk factors for infections caused by NDM-1-producers. *Front Microbiol*. 2018;9:658. doi:10.3389/fmicb.2018.00658
- Aguirre-Quinonero A, Martínez-Martínez L. Non-molecular detection of carbapenemases in Enterobacteriaceae clinical isolates. *J Infect Chemother*. 2017;23(1):1–11. doi:10.1016/j.jiac.2016.09.008
- van Duin D, Doi Y. The global epidemiology of carbapenemase-producing Enterobacteriaceae. *Virulence*. 2017;8(4):460–469. doi:10.1080/21505594.2016.1222343
- Lee C-R, Lee JH, Park KS, Kim YB, Jeong BC, Lee SH. Global dissemination of carbapenemase-producing *Klebsiella pneumoniae*: epidemiology, genetic context, treatment options, and detection methods. *Front Microbiol*. 2016;7:895.
- Bonomo RA, Burd EM, Conly J, et al. Carbapenemase-producing organisms: a global scourge. *Clin Infect Dis*. 2018;66(8):1290–1297. doi:10.1093/cid/cix893
- Nahid F, Zahra R, Sandegren L. A blaOXA-181-harboring multi-resistant ST147 *Klebsiella pneumoniae* isolate from Pakistan that represent an intermediate stage towards pan-drug resistance. *PLoS One*. 2017;12(12):e0189438. doi:10.1371/journal.pone.0189438
- Sattar H, Toleman M, Nahid F, Zahra R. Co-existence of blaNDM-1 and blaKPC-2 in clinical isolates of *Klebsiella pneumoniae* from Pakistan. *J Infect Chemother*. 2016;28(4):346–349. doi:10.1179/1973947814Y.00000000223
- Willemsen I, van Esser J, Kluytmans-van den Bergh M, Kluytmans-van den Bergh M, et al. Retrospective identification of a previously undetected clinical case of OXA-48-producing *K. pneumoniae* and *E. coli*: the importance of adequate detection guidelines. *Infection*. 2016;44(1):107–110. doi:10.1007/s15010-015-0805-7
- Duman Y, Ersoy Y, Gursoy N, Altunışık Toplu S, Otlı B. A silent outbreak due to *Klebsiella pneumoniae* that co-produced NDM-1 and OXA-48 carbapenemases, and infection control measures. *Iran J Basic Med Sci*. 2020;23(1):46–50. doi:10.22038/IJBMS.2019.35269.8400
- Seiffert SN, Marshall J, Perreten V, Carattoli A, Furrer H, Endimiani A. Emergence of *Klebsiella pneumoniae* co-producing NDM-1, OXA-48, CTX-M-15, CMY-16, QnrA and ArmA in Switzerland. *Int J Antimicrob Agents*. 2014;44(3):260–262. doi:10.1016/j.ijantimicag.2014.05.008
- Xie L, Dou Y, Zhou K, et al. Coexistence of blaOXA-48 and truncated blaNDM-1 on different plasmids in a *Klebsiella pneumoniae* isolate in China. *Front Microbiol*. 2017;8:133. doi:10.3389/fmicb.2017.00133
- Dortet L, Poirel L, Al Yaqoubi F, Nordmann P. NDM-1, OXA-48 and OXA-181 carbapenemase-producing enterobacteriaceae in sultanate of Oman. *Clin Microbiol Infect*. 2012;18(5):E144–E148. (). doi:10.1111/j.1469-0691.2012.03796.x
- Wayne P. *CLSI Performance Standards for Antimicrobial Susceptibility Testing. CLSI Supplement M100*. Clinical and Laboratory Standards Institute; 2018.
- Martzy R, Bica-Schröder K, Pálvölgyi ÁM, et al. Simple lysis of bacterial cells for DNA-based diagnostics using hydrophilic ionic liquids. *Sci Rep*. 2019;9(1):1–10. doi:10.1038/s41598-019-50246-5
- Brolund A, Rajer F, Giske CG, Melefors Ö, Titelman E, Sandegren L. Dynamics of resistance plasmids in extended-spectrum- β -lactamase-producing Enterobacteriaceae during postinfection colonization. *Antimicrob Agents Chemother*. 2019;63(4):e02201–e02218. doi:10.1128/AAC.02201-18
- Carlioni E, Andreoni F, Omiccioli E, Villa L, Magnani M, Carattoli A. Comparative analysis of the standard PCR-Based Replicon Typing (PBRT) with the commercial PBRT-KIT. *Plasmid*. 2017;90:10–14. doi:10.1016/j.plasmid.2017.01.005
- Grundmann H, Glasner C, Albiger B, et al. Occurrence of carbapenemase-producing *Klebsiella pneumoniae* and *Escherichia coli* in the European survey of carbapenemase-producing Enterobacteriaceae (EuSCAPE): a prospective, multinational study. *Lancet Infect Dis*. 2017;17(2):153–163. doi:10.1016/S1473-3099(16)30257-2
- David S, Reuter S, Harris SR, et al. Epidemic of carbapenem-resistant *Klebsiella pneumoniae* in Europe is driven by nosocomial spread. *Nat Microbiol*. 2019;4(11):1919–1929. doi:10.1038/s41564-019-0492-8
- Guo L, An J, Ma Y, et al. Nosocomial outbreak of OXA-48-producing *Klebsiella pneumoniae* in a Chinese hospital: clonal transmission of ST147 and ST383. *PLoS One*. 2016;11(8).
- Qamar MU, Walsh TR, Toleman MA, et al. Dissemination of genetically diverse NDM-1, -5, -7 producing-Gram-negative pathogens isolated from pediatric patients in Pakistan. *Future Microbiol*. 2019;14(8):691–704. doi:10.2217/fmb-2019-0012
- Tekeli A, Dolapci İ, Evren E, Oguzman E, Karahan ZC. Characterization of *Klebsiella pneumoniae* Coproducing KPC and NDM-1 Carbapenemases from Turkey. *Microb Drug Resist*. 2019;26:118–125.

27. Masseron A, Poirel L, Ali BJ, Syed M, Nordmann P. Molecular characterization of multidrug-resistance in Gram-negative bacteria from the Peshawar teaching hospital, Pakistan. *New Microbes New Infect.* 2019;32:100605. doi:10.1016/j.nmni.2019.100605
28. Lomonaco S, Crawford MA, Lascols C, et al. Resistome of carbapenem- and colistin-resistant *Klebsiella pneumoniae* clinical isolates. *PLoS One.* 2018;13(6):e0198526. doi:10.1371/journal.pone.0198526
29. Ejaz H, Wang N, Wilksch JJ, et al. Phylogenetic analysis of *Klebsiella pneumoniae* from hospitalized children, Pakistan. *Emerg Infect Dis.* 2017;23(11):1872. doi:10.3201/eid2311.170833
30. Kiaei S, Moradi M, Hosseini-Nave H, Ziasistani M, Kalantar-Neyestanaki D. Endemic dissemination of different sequence types of carbapenem-resistant *Klebsiella pneumoniae* strains harboring bla_{NDM} and 16S rRNA methylase genes in Kerman hospitals, Iran, from 2015 to 2017. *Infect Drug Resist.* 2019;12:45. doi:10.2147/IDR.S186994
31. Gamal D, Fernández-Martínez M, Salem D, et al. Carbapenem-resistant *Klebsiella pneumoniae* isolates from Egypt containing bla_{NDM}-1 on IncR plasmids and its association with rmtF. *Int J Infect Dis.* 2016;43:17–20. doi:10.1016/j.ijid.2015.12.003
32. Samonis G, Maraki S, Karageorgopoulos D, Vouloumanou E, Falagas M. Synergy of fosfomycin with carbapenems, colistin, netilmicin, and tigecycline against multidrug-resistant *Klebsiella pneumoniae*, *Escherichia coli*, and *Pseudomonas aeruginosa* clinical isolates. *Eur J Clin Microbiol Infect Dis.* 2012;31(5):695–701. doi:10.1007/s10096-011-1360-5
33. Sabir R, Alvi SFD, Fawwad A. Antimicrobial susceptibility pattern of aerobic microbial isolates in a clinical laboratory in Karachi-Pakistan. *Pak J Med Sci.* 2013;29(3):851. doi:10.12669/pjms.293.3187
34. Nahid F, Khan AA, Rehman S, Zahra R. Prevalence of metallo- β -lactamase NDM-1-producing multi-drug resistant bacteria at two Pakistani hospitals and implications for public health. *J Infect Public Health.* 2013;6(6):487–493. doi:10.1016/j.jiph.2013.06.006
35. Demir Y, Zer Y, Karaoglan I. Investigation of VIM, IMP, NDM-1, KPC AND OXA-48 enzymes in Enterobacteriaceae strains. *Pak J Pharm Sci.* 2015;28.
36. Wailan A, Paterson D. The spread and acquisition of NDM-1: a multifactorial problem. *Expert Rev Anti Infect Ther.* 2014;12(1):91–115. doi:10.1586/14787210.2014.856756
37. Kumarasamy KK, Toleman MA, Walsh TR, et al. Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and epidemiological study. *Lancet Infect Dis.* 2010;10(9):597–602. doi:10.1016/S1473-3099(10)70143-2
38. Solgi H, Badmasti F, Giske CG, Aghamohammad S, Shahcheraghi F. Molecular epidemiology of NDM-1 and OXA-48-producing *Klebsiella pneumoniae* in an Iranian hospital: clonal dissemination of ST11 and ST893. *J Antimicrob Chemother.* 2018;73(6):1517–1524. doi:10.1093/jac/dky081
39. Lixandru BE, Cotar AI, Straut M, et al. Carbapenemase-producing *Klebsiella pneumoniae* in Romania: a six-month survey. *PLoS One.* 2015;10(11):e0143214. doi:10.1371/journal.pone.0143214
40. Jamal WY, Albert MJ, Khodakhast F, Poirel L, Rotimi VO. Emergence of new sequence type OXA-48 carbapenemase-producing Enterobacteriaceae in Kuwait. *Microb Drug Resist.* 2015;21(3):329–334. doi:10.1089/mdr.2014.0123
41. Rahman S, Hussain MA, Murtaza A, et al. How ESBL- and carbapenemase-producing genes are mobilized-analysis of *Escherichia coli* isolates recovered from poultry retail meat in district mardan KPK Pakistan. Paper presented at: 2019 16th International Bhurban Conference on Applied Sciences and Technology (IBCAST) 2019.
42. Navon-Venezia S, Kondratyeva K, Carattoli A. *Klebsiella pneumoniae*: a major worldwide source and shuttle for antibiotic resistance. *FEMS Microbiol Rev.* 2017;41(3):252–275. doi:10.1093/femsre/fux013
43. Souza RD, Pinto NA, Hwang I, et al. Molecular Epidemiology and Resistome Analysis of Multidrug-Resistant ST11 *Klebsiella pneumoniae* Strain Containing Multiple Copies of Extended-Spectrum β -Lactamase Genes Using Whole-Genome Sequencing. 2017.
44. Damjanova I, Toth A, Paszti J, et al. Expansion and countrywide dissemination of ST11, ST15 and ST147 ciprofloxacin-resistant CTX-M-15-type β -lactamase-producing *Klebsiella pneumoniae* epidemic clones in Hungary in 2005—the new ‘MRSA’s’? *J Antimicrob Chemother.* 2008;62(5):978–985. doi:10.1093/jac/dkn287
45. Samuelson Ø, Toleman M, Hasseltvedt V, et al. Molecular characterization of VIM-producing *Klebsiella pneumoniae* from Scandinavia reveals genetic relatedness with international clonal complexes encoding transferable multidrug resistance. *Clin Microbiol Infect.* 2011;17(12):1811–1816. doi:10.1111/j.1469-0691.2011.03532.x
46. Grami R, Mansour W, Ben Haj Khalifa A, et al. Emergence of ST147 *Klebsiella pneumoniae* producing OXA-204 carbapenemase in a University Hospital, Tunisia. *Microb Drug Resist.* 2016;22(2):137–140. doi:10.1089/mdr.2014.0278
47. Wang X, Xu X, Li Z, et al. An outbreak of a nosocomial NDM-1-producing *Klebsiella pneumoniae* ST147 at a teaching hospital in mainland China. *Microb Drug Resist.* 2014;20(2):144–149. doi:10.1089/mdr.2013.0100
48. Poirel L, Bonnin RA, Nordmann P. Genetic features of the widespread plasmid coding for the carbapenemase OXA-48. *Antimicrob Agents Chemother.* 2012;56(1):559–562. doi:10.1128/AAC.05289-11
49. Bonnin RA, Poirel L, Carattoli A, Nordmann P. Characterization of an IncFII plasmid encoding NDM-1 from *Escherichia coli* ST131. *PLoS One.* 2012;7(4):e34752. doi:10.1371/journal.pone.0034752
50. Carattoli A. Resistance plasmid families in Enterobacteriaceae. *Antimicrob Agents Chemother.* 2009;53(6):2227–2238. doi:10.1128/AAC.01707-08
51. Erdem F, Oncul O, Aktas Z. Characterization of resistance genes and polymerase chain reaction-based replicon typing in carbapenem-resistant *Klebsiella pneumoniae*. *Microb Drug Resist.* 2019;25(4):551–557. doi:10.1089/mdr.2018.0231
52. Sartor AL, Raza MW, Abbasi SA, et al. Molecular epidemiology of NDM-1-producing Enterobacteriaceae and *Acinetobacter baumannii* isolates from Pakistan. *Antimicrob Agents Chemother.* 2014;58(9):5589–5593. doi:10.1128/AAC.02425-14
53. Chander Y, Ramakrishnan M, Jindal N, Hanson K, Goyal SM. Differentiation of *Klebsiella pneumoniae* and *K. oxytoca* by multiplex polymerase chain reaction. *Int J Appl Res Vet M.* 2011;9(2):138.
54. Rajivgandhi G, Maruthupandy M, Ramachandran G, Priyanga M, Manoharan N. Detection of ESBL genes from ciprofloxacin resistant Gram negative bacteria isolated from urinary tract infections (UTIs). *Front Lab Med.* 2018;2(1):5–13. doi:10.1016/j.flm.2018.01.001
55. Tada T, Shrestha B, Miyoshi-Akiyama T, et al. NDM-12, a novel New Delhi metallo- β -lactamase variant from a carbapenem-resistant *Escherichia coli* clinical isolate in Nepal. *Antimicrob Agents Chemother.* 2014;58(10):6302–6305. doi:10.1128/AAC.03355-14
56. Candan ED, Aksöz N. *Escherichia coli*: characteristics of carbapenem resistance and virulence factors. *Braz Arch Biol Technol.* 2017;60.
57. Toleman MA, Bugert JJ, Nizam SA. Extensively drug-resistant New Delhi metallo- β -lactamase-encoding bacteria in the environment, Dhaka, Bangladesh, 2012. *Emerg Infect Dis.* 2015;21(6):1027. doi:10.3201/eid2106.141578

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/infection-and-drug-resistance-journal>