Investigating the Effect of Nano-Curcumin on the Expression of Biofilm Regulatory Genes of *Pseudomonas aeruginosa*

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Parastoo Sharifian¹
Somayeh Yaslianifard²
Parviz Fallah³
Siavash Aynesazi⁴
Mahmood Bakhtiyari⁵
Mohammad
Mohammadzadeh²

¹Department of Microbiology, School of Medicine, Alborz University of Medical Sciences, Karaj, Iran; ²Department of Microbiology, School of Medicine, Dietary Supplements and Probiotic Research Center, Alborz University of Medical Sciences, Karaj, Iran; ³Department of Laboratory Science, Faculty of Allied Medicine, Alborz University of Medical Sciences, Karaj, Iran; ⁴Department of Microbiology, Faculty of Science, North Branch, Islamic Azad, Tehran, Iran; ⁵Department of Community Medicine and Epidemiology, School of Medicine, Non-Communicable Diseases Research Center, Alborz University of Medical Sciences, Karaj, Iran

Background: *Pseudomonas aeruginosa* is an opportunistic pathogen that causes serious nosocomial infections, especially in immunodeficient patients and cystic fibrosis, cancer, and burned individuals. The biofilm that plays an important role in the virulence of *P. aeruginosa* is under the regulation of quorum sensing and two-component regulatory systems of bacteria. Curcumin, an active phenolic extract of turmeric has shown an inhibitory effect on the biofilm formation of some pathogenic bacteria. Thus, the present study aims to evaluate the effect of Nano-Curcumin on the expression of major regulatory genes involved in biofilm formation of *P. aeruginosa*.

Materials and Methods: The biofilm formation of *P. aeruginosa* ATCC 10145 was assessed in the presence of 15, 20, and 25 µg/mL concentrations of Nano-Curcumin using the microplate titer method. The effect of Nano-Curcumin on the expression level of regulatory genes were determined by relative reverse transcriptase-realtime PCR.

Results: In the absence of Nano-Curcumin, *P. aeruginosa* strain ATCC 10145 strongly produced biofilm (3+) and in the presence of 15 and 20 µg/mL, biofilm formation was reduced to moderate (2+) and weak biofilm producer (1+), respectively. Nano-Curcumin at a concentration of 25µg/mL inhibited biofilm formation in *P. aeruginosa*. The expression of regulatory genes was not affected by biofilm inhibitory concentrations of Nano-Curcumin.

Conclusion: The antibiofilm mechanism of Curcumin is not related to the downregulation of regulatory systems of *P. aeruginosa* and probably it prevents the formation of a complete biofilm structure.

Keywords: *Pseudomonas aeruginosa*, biofilm formation, Nano-Curcumin

Introduction

Pseudomonas aeruginosa is a gram-negative opportunistic pathogen found everywhere and able to infect immunocompromised patients including those with cystic fibrosis and extensive burn wounds.¹ Because of more antibiotic resistance of bacterial cells living as biofilms, the relevant infections with biofilm producer strains are very hard to eradicate.²

The biofilm matrix of *Pseudomonas aeruginosa* is formed from extracellular polymeric materials, including polysaccharides, extracellular DNA, and proteins.³ The complex regulation of biofilm formation involves multiple bacterial types of machinery, including QS systems and two-component regulatory systems that both interact mainly with extracellular polysaccharide (EPS) production.⁴ Two-component regulatory systems such as GacS/GacA and RetS/LadS play the main role in

Correspondence: Mohammad
Mohammadzadeh
Tel +989126767995
Email M.mohammadzadeh84@yahoo.com

controlling factors associated with biofilm formation. The GacS/GacA system acts as a super-regulator of the QS system and is involved in the production of multiple virulence factors as well as in biofilm formation.⁵ The Gac system consists of a transmembrane sensor kinase (GacS) and a transcriptional regulator (GacA) which has a positive effect on the upregulation of the QS systems involved in biofilm formation. It has also been reported that RetS interacts with the GacS/GacA system and inhibits the phosphorylation of GacA.⁶ Furthermore, *P. aeruginosa* possesses two main QS systems (Las and Rhl) which drive the production of autoinducer signaling acyl-homoserine lactone molecules.⁷ The QS regulator LasR and RhlR can bind to the promoter region of the *psl* and *pel* operons and lead to enhancement of biofilm formation.⁸

Because of the emergence of multidrug-resistant (MDR) *P. aeruginosa* in developing countries,⁹ complementary therapy for control of infections associated with these bacteria can be very beneficial. Recently, dietary phytochemicals with a long history of medical use in humans have been investigated to control biofilms as they are non-toxic.^{1,10} It has been shown that the phenolic compound *Curcuma longa* (Curcumin) can inhibit biofilm formation in several bacteria such as *P. aeruginosa*.^{11–13} Different mechanisms have been suggested for the anti-biofilm activity of Curcumin by some studies, but the molecular mechanism of Curcumin for inhibition of biofilm is until unclear. Curcumin may affect biofilm formation by changing the expression of regulatory genes associated with *P. aeruginosa* signaling pathways. Curcumin has low solubility in aqueous solutions that lead to lower tissue diffusion. The Curcumin nanoparticle formulation (Nano-Curcumin) is more soluble than Curcumin and more resistant to enzymatic hydrolysis. Moreover, Nano-Curcumin has increased tissue uptake and also has a longer half-life than Curcumin that makes it more appropriate to investigate its effects on the biofilm formation.¹⁴ According to the evidences, this study aimed to evaluate the effect of Nano-Curcumin on the expression of major genes involved in the regulation of biofilm in *P. aeruginosa*.

Materials and Methods

Bacterial Strains and Culture Conditions

Pseudomonas aeruginosa ATCC 10145 was used as a positive biofilm producer strain and *P. aeruginosa* ATCC 27853 was used as a biofilm negative control.

Bacteria were cultured in Trypticase soy broth (TSB) medium (Merck, Germany). The cultures were incubated at 37 °C overnight, then were preserved at –70 °C for long term storage after adding 20% glycerol.

Biofilm Formation Assay

Biofilm Assay Optimization

The microtiter plate assay was performed for optimizing biofilm formation conditions.¹⁵ Initial 1:100 dilution was prepared from fresh 18 hours culture of bacterial strains in Brain heart infusion (BHI) (Merck, Germany). Then, 100 µL was added in 96-well plate and incubated for 48 hours at 37°C. After 48 hours, planktonic bacteria were removed and the wells were washed two times with tap water with the plate left at room temperature for 3 hours. The walls were stained with 125 µL of 0.1% (w/v) crystal violet for 10 min at room temperature, then washed with tap water. After drying, 200 µL of 30% (v/v) acetic acid was added for 10 min, and finally, OD of positive and negative controls was measured at 570nm using Microplate Readers (Awareness, USA). Biofilm formation results were obtained regarding OD of controls as follows: ODco+ ≤ ODco- = no biofilm producer; ODco+ > ODco-: ≤2×ODco- = weak biofilm producer; ODco+ > ODco-: 2–4×ODco- = moderate biofilm producer; ODco+ > ODco-: >4×ODco- = strong biofilm producer. OD values were measured based on the mean OD of four assessments.

Biofilm Inhibition Assay

For increasing the solubility of Curcumin, the polymeric nanoparticles of Curcumin (Nano-Curcumin) from source Curcumin (Merck, Germany) were used in this study. The Planetary Ball Mill technology was performed to synthesize Nano-Curcumin by the reference laboratory of Amirkabir University of Technology. Biofilm formation of *P. aeruginosa* ATCC 10145 strain was assessed at the final 15, 20, and 25µg/mL concentrations of Nano-Curcumin. Nano-Curcumin was dissolved into Brain-Heart Infusion Broth (Merck, Germany) containing 10% of Dimethyl Sulfoxide (DMSO). Bacterial culture in BHI with 10% DMSO without Nano-Curcumin was used as a control.

Bacterial Growth Inhibition Assay

The effect of Nano-Curcumin on bacterial growth was assessed through measuring OD at 600nm. *P. aeruginosa* ATCC 10145 cultures at the initial concentration of 0.5 McFarland was incubated for 48 hours in the absence and

presence of different concentrations of Nano-Curcumin (15, 20, 25 µg/mL) in BHI with 10% DMSO. Then, OD values of different samples with Nano-Curcumin were compared with the control sample.

RNA Extraction

RNA was extracted from three samples; *Pseudomonas aeruginosa* ATCC 10145 strain in the presence and the absence of Nano-Curcumin after 48 hours of incubation under optimal biofilm formation conditions and one 16-hour fresh planktonic bacteria. All samples were centrifuged and the supernatant was discarded. All samples were diluted with sterile normal saline to 3.0 McFarland. The total RNA was extracted using the High Pure RNA Isolation Kit (Roche, Switzerland) according to the kit protocol. The quantity of RNA samples was analyzed through OD measurement by NanoDrop (Boeco, Germany). RNA was adjusted to the same concentration for all samples. The purity of RNA was assessed by the negative results of a house-keeping *tufB* gene using traditional PCR.

cDNA Synthesis

Reverse transcriptase reaction was accomplished for cDNA synthesis using Transcriptor First Strand cDNA Synthesis Kit (Roche, Switzerland) regarding kit protocol. The quantity and quality of cDNA samples were assessed by OD measurement and positive results of *tufB* gene in PCR, respectively. cDNA was adjusted as the same concentration for all samples. The primer sequences of *tufB* are listed in Table 1.

Polymerase Chain Reaction (PCR)

PCR was carried out for controlling the specific primers and optimizing annealing temperatures for amplification of

gacA, *retS*, *lasR*, and *rhlR* genes. Amplifications were performed in reactions containing 12 µL of 2X ready to use Master Mix (Fermentase, Germany), 1 µL of each 20 pmol Forward and reverse primers, 10 µL of sterile distilled water, and 1 µL of 1 µg adjusted cDNA. Electrophoresis on 1.5% agarose gel was done for evaluating the PCR amplicons. The primer sequences and PCR conditions are mentioned in Table 1.

Real-Time PCR

Relative Real-time PCR was used to investigate the effect of Nano-Curcumin on the expression of genes *gacA*, *retS*, *lasR*, and *rhlR*. The comparison of gene expression was performed in five cDNAs prepared from samples with different culture conditions: a fresh culture (bacterial planktonic form), a sample of bacteria isolated from the optimal biofilm formation condition, and three samples of bacteria isolated from bacterial biofilm conditions in the presence of 15, 20, and 25 µg/mL of Nano-Curcumin. Amplifications were performed in double reaction in final volume 20 µL including 10 µL of 2X ready to use SYBR Green Master Mix (TaKaRa, Japan), 1 µL of each 20 pmol Forward and Reverse primers, 7 µL double-distilled water, and 1 µL of adjusted cDNA (1 µg/µL). Reactions were carried out in 35 cycles via Rotor-Gene Q 5plex (Qiagen, USA) according to thermal conditions as mentioned above. For normalization of the expression, the house-keeping *tufB* gene was used as an internal control. No template controls were used for each reaction for the screening of unwanted amplifications. The differences in gene expression were calculated through the $2^{-\Delta\Delta Ct}$ method.

Table 1 Sequences of Primers and PCR Conditions of Evaluating Genes

Target	Primer (5' to 3')	Amplicon Sizes	PCR Thermal Conditions
<i>tufB</i>	F: GTGAGAACGTTGGTATCCTG R: GTGGTACGGAAGTAGAACTG	193 bp	{95 °C (20 Sec), 60 °C (15 Sec), 72 °C (20 Sec)}×35
<i>gacA</i>	F: AGGTGCAGCGTGATTAAG R: GCATCTTCACGTCCATCAG	181 bp	{95 °C (20 Sec), 57 °C (15 Sec), 72 °C (20 Sec)}×35
<i>retS</i>	F: TCCCTATTTCCGCCAGAC R: CCAGCTTGACCACGATTG	143 bp	{95 °C (20 Sec), 58 °C (15 Sec), 72 °C (20 Sec)}×35
<i>lasR</i>	F: TGGATGCTCAAGGACTACG R: TTCACATTGGCTTCCGAG	173 bp	{95 °C (20 Sec), 56 °C (15 Sec), 72 °C (20 Sec)}×35
<i>rhlR</i>	F: CTGGAAAAGGAAGTGCGG R: GTAGTTCTGCATCTGGTATCG	141 bp	{95 °C (20 Sec), 58 °C (15 Sec), 72 °C (20 Sec)}×35

Statistical Analysis

The experiments were performed in five times. The means and standard deviations of ΔC_t values were analyzed using GraphPad Prism 8.0 Software (San Diego, Canada). The ANOVA test was carried out to calculate the significance of the results. A P value of less than 0.05 was considered significant.

Results

Determination of Optimum Condition of Biofilm Formation

Based on the microtiter plate assay, the control positive *P. aeruginosa* ATCC 10145 strain was confirmed as a strong biofilm producer strain. The OD ratios for biofilm quantification have been mentioned in Table 2.

Nano-Curcumin Blocks Biofilm Formation in *P. aeruginosa* in a Concentration-Dependent Way, but Does Not Affect Bacterial Growth

Nano-Curcumin was dissolved at a maximum concentration of 25 $\mu\text{g/mL}$ in BHI containing 10% DMSO. Bacterial growth was not inhibited by Nano-Curcumin, but biofilm formation was inhibited completely. In the presence of 15 and 20 $\mu\text{g/mL}$, biofilm formation levels were reduced to moderate (2+) and weak (1+), respectively (Figures 1 and 2). The OD values of bacterial growth in the presence of different concentrations of Nano-Curcumin are listed in Table 3.

Quality Control of RNA and cDNA

The extracted RNA samples were adjusted to 250 ng/ μL , and cDNA samples were provided as 1 $\mu\text{g}/\mu\text{L}$ for PCR. The purity of RNA as well as the quality of cDNA and

primers was confirmed by negative results for RNA and positive PCR results for cDNA samples. The results of the PCR of *tufB* gene in RNA samples were negative, suggesting the purity of RNA. Further, positive PCR results of *tufB*, *gacA*, *retS*, *lasR*, and *rhlR* genes showed the reliability of primers and cDNA of samples.

Nano-Curcumin Does Not Affect Gene Expression of Biofilm Regulatory Factors

The relative expression of genes *gacA*, *retS*, *lasR*, and *rhlR* was determined with C_t values. The mean amounts of expression of target genes were normalized to the mean amount of the *tufB* reference gene in the same sample. The level of expression of *gacA*, *lasR*, and *rhlR* regulators showed a significant increase in samples from biofilm conditions in the absence and presence of different concentrations of Nano-Curcumin in comparison to the sample of the fresh planktonic form of bacteria. However, there were no significant differences between the expression level of *retS* gene in all samples. The results showed that while biofilm inhibited with Nano-Curcumin, it did not affect on the expression of regulator genes involved in biofilm formation (Figure 3).

Discussion

P. aeruginosa is an important human opportunistic pathogen causing acute and chronic infections, especially in immunocompromised patients. In recent decades, the excessive use of antibiotics has led to the emergence of resistant strains in these bacteria.¹⁶

Biofilm formation is one of the mechanisms of these bacteria that reduce the effect of antibiotic therapy. *P. aeruginosa* biofilm is a structure composed of different biomolecules that evolve as these components come together. Bacterial attachment factors, extracellular polysaccharides (EPS), and extracellular DNA (eDNA) are involved in the formation of this structure.⁶ Regulation of the expression of essential genes in the process of secretion and formation of biofilm components is under the direct and indirect control of two-component regulatory systems and bacterial quorum sensing. Failure in these systems results in halting biofilm production or it's defective structure.¹⁷

The natural compounds with antimicrobial effects as an alternative or complementary therapy has been recently considered. The antimicrobial activity of essential oils of some plants on multi-drug-resistant *P. aeruginosa* isolates in vitro

Table 2 OD Values and Biofilm Formation Level of *P. aeruginosa* ATCC 10145 Strain in 570 nm

OD of Positive Control	OD of Negative Control	Ratios	Biofilm Level
0.79	0.15	5.26	3+ (strong)
0.83	0.19	4.36	3+ (strong)
0.61	0.23	2.65	2+ (Moderate)
0.74	0.17	4.35	3+ (strong)
0.67	0.16	4.18	3+ (strong)
Mean: 0.73	Mean: 0.18	Mean: 4.16	Total: 3+ (strong)

Figure 1 Nano-Curcumin inhibits the Biofilm in a concentration-dependent manner. The biofilm formation was reduced to 2+, 1+ and negative in 15, 20 and 25 µg/mL concentrations of Nano-Curcumin.

Figure 2 Inhibition of biofilm formation in *P. aeruginosa* 10145 in the absence and presence of 15, 20, and 25 µg/mL Nano-Curcumin. (**P value < 0.001). The error bar represents SD in five repeated experiments.

has been shown by some previous studies.^{18,19} Curcumin is an active phenolic component of *Curcuma longa* that has been investigated due to its pharmaceutical attributes.^{6,20}

Some studies confirming the effect of Curcumin on biofilm production in *P. aeruginosa* suggest that this effect is likely due to the QS of the bacterium and the downregulation of the regulatory genes involved in this process. In a study, Roudashti et al²¹ using RT-PCR showed that C12-HSL and C4-HSL signals of *P. aeruginosa* PAO1 strain significantly

diminished in the presence of Curcumin. Rudrappa et al¹ hypothesized that Curcumin may target QS responses in *P. aeruginosa* PAO1 through delays in the construction of the HSL or by disabling the receiving QS signals. Nevertheless, other studies showed it affects the expression of the genes directly involved in the biofilm production.²²

In the present study, the inhibitory effect of Curcumin was surveyed on the biofilm formation and QS of *P. aeruginosa*.

Table 3 Bacterial Growth in the Absence and the Presence of Different Concentrations of Nano-Curcumin According to OD Values of 600 nm After 48 Hours

Culture Conditions	BHI +10% DMSO	Nano-Curcumin Concentrations		
		BHI+10% DMSO+ 25µg/mL	BHI+10% DMSO+ 20µg/mL	BHI+10% DMSO+ 15µg/mL
OD values	0.71	0.69	0.76	0.68
	0.73	0.63	0.67	0.63
	0.65	0.59	0.69	0.75
	0.69	0.61	0.63	0.66
	0.69	0.68	0.70	0.65
Mean	0.69	0.64	0.69	0.67

The results of this study indicated that although Nano-Curcumin has an inhibitory effect on biofilm formation in *P. aeruginosa* in a concentration-dependent manner, it has no significant effect on the expression of major regulatory genes that play an important role in biofilm formation.

This may suggest that Nano-Curcumin has a different role other than changing the expression of regulatory genes that are involved in bacterial biofilm synthesis.

A research indicated that Curcumin at concentrations ≥ 25 µM had an inhibitory effect on bacterial growth, causing leakage on the bacterial membranes, that lead to the death of some bacterial species such as *Staphylococcus aureus*, *Escherichia coli*, and *P. aeruginosa*.²³ The blockade of the proteolytic activity of autotransporter proteins of Enterococcal and Enteropathogenic *Escherichia coli* by Curcumin was mentioned in a more recent study. It was also observed that Curcumin does not affect bacterial growth and gene expression of those auto-transporters, but it prevents their release through binding to their cleavage site.²⁴

Degradation of bacterial biofilms by Curcumin through its interaction with biofilm matrix proteins of some bacterial species was reported by a study.²⁰ Furthermore, a novel protein CdrA was shown in the biofilm structure. This protein is an auto-transporter which mediates bacterial aggregation and static biofilm formation with interaction with Psl; a major EPS of *P. aeruginosa* biofilm.²⁵ Curcumin may destabilize the biofilm structure and reduce bacterial aggregation by inhibiting the release of this protein to the extracellular matrix or biofilm degradation resulting in interaction with this type of biofilm component proteins.

Figure 3 The expression level of genes involved in the regulation of biofilm in *P. aeruginosa* in planktonic phase, biofilm formation condition in the absence and presence of Nano-Curcumin; expression of *gacA*, *lasR* and *rhlR* genes increased by between 10 to 14 times in bacteria in biofilm conditions in the absence and presence of Nano-Curcumin compared to fresh planktonic bacteria. The *retS* gene showed no changes in the expression under biofilm conditions compared to the bacterial planktonic phase.

The results of some other studies indicated the inhibitory effect of Curcumin on bacterial cell division^{26,27} as well as its synergistic effect on some antibiotics in the treatment of antibiotic-resistant bacteria.^{28,29} Despite the remarkable effects of Curcumin on many bacteria, there are some limitations to using it as an antibacterial agent. Curcumin is poorly water-soluble while water is the most important constituent of the human body. Further, it has been shown that Curcumin may be toxic to eukaryotic cells at the concentration above 10 µg/mL.³⁰ In the present study, we used Nano-Curcumin that has more solubility than exact Curcumin in water. However, Nano-Curcumin was dissolved in a maximum at a concentration of 25 µg/mL in the presence of 10% DMSO. This concentration inhibited biofilm formation in the strong biofilm producer strain of *P. aeruginosa*, while it did not affect bacterial growth.

The results of this study revealed that non-toxic concentrations of Curcumin in eukaryotic cells were not able to inhibit bacterial growth. For this reason, the systemically use of Curcumin to treat *P. aeruginosa* infections seem unexpected.

The cytotoxic effect of plant oils in high-concentrations also has been reported in some studies, which lead to the limited application of these compounds as a therapeutic agent in *In vivo* conditions.^{31,32}

According to the results of this study and similar studies, it seems that the anti-biofilm effect of Curcumin in *P. aeruginosa* is more related to inhibition of secretion of different components of the biofilm to extracellular space of bacteria or direct disruption of biofilm stability due to reaction with its structural protein component rather than affecting the expression of regulatory genes related to QS and two-component regulatory systems that control biofilm formation in this bacterium. Further studies seem to be necessary to determine the exact mechanism of biofilm inhibition. Further, due to the low solubility of Curcumin and its minor inhibitory effect on bacterial growth at its soluble concentrations, its topical use can be considered as a complementary therapy in superficial infection with *P. aeruginosa*.

Acknowledgments

We wish to thank the Deputy of Research and Technology and Department of Microbiology, Alborz University of Medical Sciences for supporting this study.

Disclosure

The authors report no conflicts of interest in this work.

References

- Rudrappa T, Bais HP. Curcumin, a known phenolic from *Curcuma longa*, attenuates the virulence of *Pseudomonas aeruginosa* PAO1 in whole plant and animal pathogenicity models. *J Agric Food Chem*. 2008;56(6):1955–1962. doi:10.1021/jf072591j
- Skindersoe ME, Alhede M, Phipps R, et al. Effects of antibiotics on quorum sensing in *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother*. 2008;52(10):3648–3663. doi:10.1128/AAC.01230-07
- Taghadosi R, Shakibaie MR, Ghanbarpour R, Hosseini-Nave H. Role of antigen-43 on biofilm formation and horizontal antibiotic resistance gene transfer in non-O157 Shiga toxin producing *Escherichia coli* strains. *Iran J Microbiol*. 2017;9(2):89.
- De Kievit T. Quorum sensing in *Pseudomonas aeruginosa* biofilms. *Environ Microbiol*. 2009;11(2):279–288. doi:10.1111/j.1462-2920.2008.01792.x
- Parkins MD, Ceri H, Storey DG. *Pseudomonas aeruginosa* GacA, a factor in multihost virulence, is also essential for biofilm formation. *Mol Microbiol*. 2001;40(5):1215–1226. doi:10.1046/j.1365-2958.2001.02469.x
- Rasamiravaka T, Labtani Q, Duez P, El Jaziri M. The formation of biofilms by *Pseudomonas aeruginosa*: a review of the natural and synthetic compounds interfering with control mechanisms. *Biomed Res Int*. 2015;2015:1–17. doi:10.1155/2015/759348
- Jimenez PN, Koch G, Thompson JA, Xavier KB, Cool RH, Quax WJ. The multiple signaling systems regulating virulence in *Pseudomonas aeruginosa*. *Microbiol Mol Biol Rev*. 2012;76(1):46–65. doi:10.1128/MMBR.05007-11
- Sakuragi Y, Kolter R. Quorum-sensing regulation of the biofilm matrix genes (*pel*) of *Pseudomonas aeruginosa*. *J Bacteriol*. 2007;189(14):5383–5386. doi:10.1128/JB.00137-07
- Tuon FF, Gortz LW, Rocha JL. Risk factors for pan-resistant *Pseudomonas aeruginosa* bacteremia and the adequacy of antibiotic therapy. *Braz J Infect Dis*. 2012;16(4):351–356. doi:10.1016/j.bjid.2012.06.009
- Brackman G, Defoirdt T, Miyamoto C, et al. Cinnamaldehyde and cinnamaldehyde derivatives reduce virulence in *Vibrio* spp. by decreasing the DNA-binding activity of the quorum sensing response regulator LuxR. *BMC Microbiol*. 2008;8(1):149. doi:10.1186/1471-2180-8-149
- Li B, Li X, Lin H, Zhou Y. Curcumin as a promising antibacterial agent: effects on metabolism and biofilm formation in *S. mutans*. *Biomed Res Int*. 2018;2018.
- Raorane CJ, Lee J-H, Kim Y-G, Rajasekharan SK, García-Contreras R, Lee J. Antibiofilm and antivirulence efficacies of flavonoids and curcumin against *Acinetobacter baumannii*. *Front Microbiol*. 2019;10:990. doi:10.3389/fmicb.2019.00990
- Vaughn AR, Haas KN, Burney W, et al. Potential role of curcumin against biofilm-producing organisms on the skin: a review. *Phytother Res*. 2017;31(12):1807–1816. doi:10.1002/ptr.5912
- Kurita T, Makino Y. Novel curcumin oral delivery systems. *Anticancer Res*. 2013;33(7):2807–2821.
- O'Toole GA. Microtiter dish biofilm formation assay. *J Vis Exp*. 2011;47:e2437.
- Tielen P, Kuhn H, Rosenau F, Jaeger K-E, Flemming H-C, Wingender J. Interaction between extracellular lipase LipA and the polysaccharide alginate of *Pseudomonas aeruginosa*. *BMC Microbiol*. 2013;13(1):159. doi:10.1186/1471-2180-13-159
- Ganesh PS, Rai VR. Attenuation of quorum-sensing-dependent virulence factors and biofilm formation by medicinal plants against antibiotic resistant *Pseudomonas aeruginosa*. *J Tradit Complement Med*. 2018;8(1):170–177. doi:10.1016/j.jtcme.2017.05.008
- Donadu M, Usai D, Pinna A, et al. In vitro activity of hybrid lavender essential oils against multidrug resistant strains of *Pseudomonas aeruginosa*. *J Infect Dev Ctries*. 2018;12(01):009–014. doi:10.3855/jidc.9920

19. Trong Le N, Viet Ho D, Quoc Doan T, et al. In vitro antimicrobial activity of essential oil extracted from leaves of *Leodeo domatio-phorus* Chaowasku, DT Ngo and HT Le in Vietnam. *Plants*. 2020;9(4):453. doi:10.3390/plants9040453
20. Singh AK, Prakash P, Singh R, et al. Curcumin quantum dots mediated degradation of bacterial biofilms. *Front Microbiol*. 2017;8:1517. doi:10.3389/fmicb.2017.01517
21. Roudashti S, Zeighami H, Mirshahabi H, Bahari S, Soltani A, Haghi F. Synergistic activity of sub-inhibitory concentrations of curcumin with ceftazidime and ciprofloxacin against *Pseudomonas aeruginosa* quorum sensing related genes and virulence traits. *World J Microbiol Biotechnol*. 2017;33(3):50. doi:10.1007/s11274-016-2195-0
22. Shariati A, Asadian E, Fallah F, et al. Evaluation of Nano-curcumin effects on expression levels of virulence genes and biofilm production of multidrug-resistant *Pseudomonas aeruginosa* isolated from burn wound infection in Tehran, Iran. *Infect Drug Resist*. 2019;12:2223. doi:10.2147/IDR.S213200
23. Tyagi P, Singh M, Kumari H, Kumari A, Mukhopadhyay K. Bactericidal activity of curcumin I is associated with damaging of bacterial membrane. *PLoS One*. 2015;10(3):e0121313. doi:10.1371/journal.pone.0121313
24. Sanchez-Villamil JI, Navarro-Garcia F, Castillo-Romero A, Gutierrez-Gutierrez F, Tapia D, Tapia-Pastrana G. Curcumin blocks cytotoxicity of enteroaggregative and enteropathogenic *Escherichia coli* by blocking Pet and EspC proteolytic release from bacterial outer membrane. *Front Cell Infect Microbiol*. 2019;9:334. doi:10.3389/fcimb.2019.00334
25. Reichhardt C, Wong C, da Silva DP, Wozniak DJ, Parsek MR. CdrA interactions within the *Pseudomonas aeruginosa* biofilm matrix safeguard it from proteolysis and promote cellular packing. *MBio*. 2018;9(5):e01376–01318. doi:10.1128/mBio.01376-18
26. Kaur S, Modi NH, Panda D, Roy N. Probing the binding site of curcumin in *Escherichia coli* and *Bacillus subtilis* FtsZ—a structural insight to unveil antibacterial activity of curcumin. *Eur J Med Chem*. 2010;45(9):4209–4214. doi:10.1016/j.ejmech.2010.06.015
27. Rai D, Singh JK, Roy N, Panda D. Curcumin inhibits FtsZ assembly: an attractive mechanism for its antibacterial activity. *Biochem J*. 2008;410(1):147–155. doi:10.1042/BJ20070891
28. Kali A, Devaraj Bhuvaneshwar P, Charles M, Seetha KS. Antibacterial synergy of curcumin with antibiotics against biofilm producing clinical bacterial isolates. *J Basic Clin Pharmacol*. 2016;7(3):93. doi:10.4103/0976-0105.183265
29. Sasidharan NK, Sreekala SR, Jacob J, Nambisan B. In vitro synergistic effect of curcumin in combination with third generation cephalosporins against bacteria associated with infectious diarrhea. *Biomed Res Int*. 2014;2014:1–8. doi:10.1155/2014/561456
30. Damarla SR, Komma R, Bhatnagar U, Rajesh N, Mulla SMA. An evaluation of the genotoxicity and subchronic oral toxicity of synthetic curcumin. *J Toxicol*. 2018;2018:1–27. doi:10.1155/2018/6872753
31. Amorese V, Donadu M, Usai D, et al. In vitro activity of essential oils against *Pseudomonas aeruginosa* isolated from infected hip implants. *J Infect Dev Ctries*. 2018;12(11):996–1001. doi:10.3855/jidc.10988
32. Bua A, Usai D, Donadu M, et al. Antimicrobial activity of *Austroepatorium inulaefolium* (HBK) against intracellular and extracellular organisms. *Nat Prod Res*. 2018;32(23):2869–2871. doi:10.1080/14786419.2017.1385014

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>