Low-Intensity Continuous Ultrasound for the Symptomatic Treatment of Upper Shoulder and Neck Pain: A Randomized, Double-Blind Placebo-Controlled Clinical Trial [Corrigendum]

Pettersen S, Plancher K, Klyve D, Draper D, Ortiz R. *J Pain Res.* 2020;13:1277–1287.

Page 1277, Abstract, Results, second sentence, the text “Participants treated with active therapy observed a significant mean pain reduction from baseline of 2.61 points for active ($p<0.001$), compared to 1.58 points decrease from baseline for placebo ($p=0.087$), resulting in a 1.03 points significant decrease in the active group over placebo ($p=0.003$)” should read “Participants treated with active therapy observed a significant mean pain reduction from baseline of 2.61 points for active ($p<0.001$), compared to 1.58 points decrease from baseline for placebo ($p=0.087$), resulting in a 1.03 points significant decrease in the active group over placebo ($p=0.009$)”.

Page 1281, Results, Pain Change from Baseline section, last sentence, the text “While this could indicate a placebo

effect, the difference between groups for change from baseline was significant for all 4 weeks assessed ($p<0.001$ for weeks 1–3, $p=0.003$ for week 4), showing a more significant change from baseline for the active LICUS group compared to placebo” should read “While this could indicate a placebo effect, the difference between groups for change from baseline was significant for all 4 weeks assessed ($p<0.001$ for weeks 1–2, $p<0.01$ for weeks 3–4), showing a more significant change from baseline for the active LICUS group compared to placebo”.

Page 1282, Table 2 is incorrect. The correct table is shown below.

The authors apologize for these errors and advise they do not affect the results of the paper.

Table 2 Primary Outcomes. The Pain Reported at Baseline and Weekly Averages. Change in Pain from Baseline and Comparison Between Groups Evaluated Using Independent *t*-tests

Pain diary data (NRS)				
Time	Active	Placebo	Between Group Mean 95% CI	P-value
Baseline	5.60 ± 1.58	5.44 ± 1.95	0.163 (-1.22 to 1.55)	0.81
Week 1	3.54 ± 1.64	4.58 ± 2.20	-1.03 (-1.65 to -0.42)	0.001
Week 2	3.26 ± 1.58	4.74 ± 2.39	-1.48 (-2.12 to -0.84)	<0.001
Week 3	3.09 ± 1.54	3.92 ± 2.10	-0.827 (-1.46 to -0.19)	0.011
Week 4	2.98 ± 1.62	3.86 ± 2.30	-0.87 (-1.64 to -0.106)	0.026
NRS mean change from baseline 95% CI				
Week 1	-2.05 ± 1.64 -2.76 to -1.35 $p<0.001$	-0.860 ± 2.21 -2.53 to 0.81 $p=0.306$	-1.20 (-1.81 to -0.579)	<0.001
Week 2	-2.34 ± 1.59 -3.03 to -1.65 $p<0.001$	-0.693 ± 2.38 -2.49 to 1.11 $p=0.443$	-1.65 (-2.28 to -1.00)	<0.001

(Continued)

(Continued).

Pain diary data (NRS)				
Time	Active	Placebo	Between Group Mean 95% CI	P-value
Week 3	-2.51 ± 1.55 -3.18 to -1.83 p<0.001	-1.52 ± 2.11 -3.14 to 0.110 p=0.07	-0.989 (-1.62 to -0.351)	0.003
Week 4	-2.61 ± 1.63 -3.34 to -1.90 p<0.001	-1.58 ± 2.30 -3.40 to 0.24 p=0.087	-1.03 (-1.80 to -0.264)	0.009

Journal of Pain Research

Dovepress

Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication. The manuscript

management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>