Cannabidiol as a Novel Therapeutic for Immune Modulation

This article was published in the following Dove Press journal:
ImmunoTargets and Therapy

Nadia Peyravian^{1,2}

Sapna Deo^{1,2}

Sylvia Daunert¹⁻³

Joaquin J Jimenez^{1,2,4}

¹Department of Biochemistry and Molecular Biology, University of Miami Miller School of Medicine, Miami, FL, USA; ²Dr. JT Macdonald Foundation Biomedical Nanotechnology Institute of the University of Miami, Miami, FL, USA; ³University of Miami Clinical and Translational Science Institute, Miami, FL, USA; ⁴Department of Dermatology & Cutaneous Surgery, University of Miami Miller School of Medicine, Miami, FL, USA

Abstract: The immune-suppressive effects of cannabidiol (CBD) are attributed to the modulation of essential immunological signaling pathways and receptors. Mechanistic understanding of the pharmacological effects of CBD emphasizes the therapeutic potential of CBD as a novel immune modulator. Studies have observed that the antagonists of CB₁ and CB₂ receptors and transient receptor potential vanilloid 1 reverse the immunomodulatory effects of CBD. CBD also inhibits critical activators of the Janus kinase/signal transducer and activator of transcription signaling pathway, as well as the nucleotide-binding oligomerization domain-like receptor signaling pathway, in turn decreasing pro-inflammatory cytokine production. Furthermore, CBD protects against cellular damage incurred during immune responses by modulating adenosine signaling. Ultimately, the data overwhelmingly support the immunosuppressive effects of CBD and this timely review draws attention to the prospective development of CBD as an effective immune modulatory therapeutic.

Keywords: cannabidiol, CBD, immune modulation, CB₁ and CB₂ receptors, TRPV1, JAK/STAT, inflammasome

Cannabidiol (CBD)

Cannabidiol (CBD) is one of the biochemical compounds, referred to as cannabinoids, isolated from the *Cannabis* plant (Figure 1).^{1,2} CBD is the second most prominent cannabinoid within the plant and comprises nearly 40% of the plant extract.^{3,4} In recent years, CBD has become one of the most widely studied cannabinoids for its anti-inflammatory and immunomodulatory effects.^{5,6} These anti-inflammatory and immunomodulatory pharmacological properties of CBD have set the precedent for its use as a natural compound for autoimmune disorders such as multiple sclerosis (MS), type 1 diabetes, and rheumatoid arthritis.^{7,8} To date, however, the FDA has approved only one CBD prescription drug, Epidiolex, for treatment of two rare and otherwise drug-resistant forms of epilepsy, Dravet Syndrome and Lennox-Gastaut Syndrome.^{9,10} An overview of elemental receptor and signaling pathway followed by a critical review of CBD's involvement in the modulation of the receptors and immunological signaling cascades encourage further development of CBD as a state-of-the-art immunomodulatory therapy.^{8,11}

CB₁ and CB₂ Receptors

CB₁ and CB₂ are members of the G-protein coupled receptor family and are encoded by the *CNR1* and *CNR2* genes, respectively.¹² CB₁ receptors have a prominent presence in the central nervous system (CNS) while CB₂ receptors are more localized in immune

Correspondence: Joaquin J Jimenez; Sylvia Daunert
Department of Biochemistry and Molecular Biology, University of Miami Miller School of Medicine, 1011 NW 15th Street, Miami, FL 33136, USA
Tel +1 305 243 6586
Email j.jimenez@med.miami.edu; sdaunert@med.miami.edu

Figure 1 Chemical structure of cannabidiol (CBD).

cells.^{13–15} CB₁ receptor activation initiates down-stream modulation of three main signaling pathways (Figure 2).¹² Activated CB₁ receptor coupled to G_{i/o} inhibits adenylyl cyclase (AC) activity, in turn inhibiting cyclic adenosine monophosphate (cAMP) formation. As a result, intracellular signal transduction, and protein kinase A (PKA) activity is hindered.^{16–18} The cAMP-PKA signal transduction is crucial for many cellular responses including enzyme activation and gene expression.¹⁹ Studies have also shown that activation of the CB₁ receptor also leads to regulation of mitogen-activated protein kinases (MAPKs) that are vital to inflammatory and immune cellular responses, including extracellular signal-regulated protein kinase 1/2 (ERK1/2), p38, and c-jun terminal kinase (JNK).^{20,21} The phosphoinositide 3-kinase (PI3K)/

protein kinase B (Akt) is a third pathway involved in cell growth and survival that may also be activated by CB₁ receptor.^{22,23} Similarly, CB₂ receptor agonism modulates and inhibits AC and cAMP leading to the inactivation of PKA by G_{i/o} protein coupling.^{24,25}

CBD and CB₁ and CB₂ Receptors

The low affinity of CBD for both CB₁ and CB₂ receptors ($K_i > 2000$ nM) suggests the pharmacodynamics of CBD to be independent of these receptors.^{26–29} For example, CBD suppressed cytokine production in CB₁ and CB₂ receptor knock-out (*CNRI^{-/-}/CNR2^{-/-}*) mice, suggesting the effects of CBD are independent of CB₁ and CB₂ receptors.³⁰ Specifically, Interleukin-2 (IL-2), a T-cell-derived cytokine, was significantly reduced in splenocytes of *CNRI^{-/-}/CNR2^{-/-}* mice upon treatment with CBD (1 μ M). CBD (10 μ M) also significantly decreased adaptive immune response cytokine production of Interferon gamma (IFN- γ). However, CBD has been shown to function as an inverse agonist of CB₁ and CB₂ receptors and can antagonize the effects of CB₁ and CB₂ receptor agonists.⁴ The use of CB₁ and CB₂ receptor antagonists reverses the anti-inflammatory and immune-suppressive effects of CBD.^{31,32} For instance, while treatment with CBD (10 mg/kg) was shown to reduce gastrointestinal motility in a bacterial lipopolysaccharide (LPS) sepsis mouse model, the effects were reversed upon treatment with CB₁ receptor antagonist, AM251 (1 mg/kg).³¹ Similar reversal of CBD-induced neuroprotective effects was observed in a hypoxic-ischemic brain injury model in newborn pigs.³² In this model, CBD (1 mg/kg) inhibited inflammatory cytokine interleukin-1 (IL-1) production in hypoxic-ischemic pigs, and co-administration with CB₂ receptor antagonist, AM630 (1 mg/kg), reversed this CBD-mediated neuroprotective effect. Conclusively, the immune-protective effects of CBD are pharmacologically supported to be independent of CB₁ and CB₂ receptors as well as by inverse agonism of CB₁ and CB₂.

Figure 2 Signaling pathway activity downstream of CB₁. Activated CB₁ receptor couples with G_{i/o} to inhibit adenylyl cyclase (AC) activity, thus inhibiting cyclic adenosine monophosphate (cAMP) production and protein kinase A (PKA) activity. CB₁ receptor activation also regulates mitogen-activated protein kinases (MAPK) and phosphoinositide 3-kinase (PI3K)/protein kinase B (Akt) pathways.

Transient Receptor Potential Vanilloid 1 (TRPV1)

The transient receptor potential vanilloid 1 (TRPV1) is a homotetrameric membrane protein and is the most well-studied nonselective ion channel of the transient receptor potential cationic family.³³ TRPV1 is expressed in various immune cells including lymphocytes, microglia, and astrocytes and is involved in pathophysiological responses of the immune system.³⁴ Numerous exogenous and endogenous stimuli activate TRPV1 with capsaicin as the most notable

agonist.³⁵ TRPV1 is responsible for detecting stimuli to maintain and regulate cellular homeostasis by promoting Ca^{2+} influx as a secondary messenger for the induction of pro-inflammatory cytokines and chemokines.³⁶ Similar to CB_1 and CB_2 receptors described above, activation of TRPV1 may lead to varying downstream signaling such as PI3K/Akt and ERK1/2 signaling pathways (Figure 3).³⁷ TRPV1 may also lead to transforming growth factor-activated kinase 1 (TAK-1) dependent JNK/MAPK cascade that leads to pro-inflammatory nuclear factor- κB (NF- κB) activation.³⁸ Janus kinase/signal transducers and activators of transcription (JAK/STAT) pathway may also be activated downstream of TRPV1 activation leading to increased expression of pro-inflammatory cytokines.³⁹

CBD and TRPV1

As TRPV1 activation induces pro-inflammatory and immune mediators, TRPV1 channels are a potential target for novel immune modulatory therapeutics.^{40–42} Vanilloid receptor knockout (TRPV1 $^{-/-}$) mice treated with CBD dose-dependently (5 mg/kg – 50 mg/kg) inhibited induction of myeloid-derived suppressor cells (MDSCs), suggesting that TRPV1 receptors are critical for induction of MDSCs by CBD.⁴³ CBD (10 mg/kg) was also shown to be an effective therapeutic in

a rodent model of thermal hyperalgesia which is associated with immune activation.^{44,45} In contrast, co-administration of CBD (10 mg/kg) with TRPV1 antagonist, capsazepine (CPZ) (2 mg/kg), partially reversed the anti-hyperalgesia effects of CBD. Treatment with CPZ (10 mg/kg) fully reversed the anti-hyperalgesia effects, signifying that CBD modulates TRPV1 for immune suppression.⁴⁵ Similarly, CBD (10 μM) increased phagocytosis of CNS immune response microglial cells and treatment with CPZ (10 μM) abolished this enhanced phagocytosis, attributing the increased phagocytic effects to modulation of TRPV1.⁴⁶ The role of TRPV1 in the immunomodulatory effects of CBD was also apparent in CBD-induced endometrial cancer cell apoptosis. The apoptotic effects of CBD (5 μM) on the endometrial cancer cell line were dependent on TRPV1 modulation as these effects were reversed upon co-administration with TRPV1 antagonist, iRTX (20 nM), hence signifying that CBD may be a potential candidate for immune modulation of TRPV1.⁴⁷

Janus Kinase (JAK) and Signal Transducer and Activator of Transcription 3 (STAT3)

Janus kinase/signal transducers and activators of transcription (JAK/STAT) signaling pathway is an evolutionary conserved cascade fundamental to T cell-mediated immunity and adaptive immune responses.^{48–50} Broadly, cytokines or growth factors activate their respective transmembrane receptors, facilitating activation of receptor-bound JAKS^{51,52} (Figure 4). Activated JAKs (JAK1, JAK2, JAK3, TYK2) proceed to phosphorylate latent STAT monomers (STAT1, STAT2, STAT3, STAT4, STAT5a, STAT5b, STAT6) leading to STAT dimerization, translocation to the nucleus, binding to specific DNA promoter sequences, and ultimately transcription of the target gene.^{53,54} Signal transducer and activator of transcription 3 (STAT3) is one of the seven STAT proteins involved in the regulation of critical cellular gene transcriptions including cell differentiation, proliferation, metastasis, as well as immune responses.⁵⁵ Constitutive activation of JAK-STAT3 signaling is the basis for the development and progression of many cancers; thus, it is not surprising that JAK-STAT3 remains a principal anticancer target.⁵⁶ However, as STAT3 is a transcription factor that remains generally confined to the nucleus, the drug-ability of STAT3

Figure 3 Transient receptor potential vanilloid 1 (TRPV1) signaling. TRPV1 activation initiates down-stream signaling of three major pathways including PI3K/AKT. Transforming growth factor-activated kinase 1 (TAK-1) dependent c-jun terminal kinase (JNK)/MAPK and Janus kinase/signal transducers and activators of transcription (JAK/STAT) signaling cascades may also be activated as a result of TRPV1 activation leading to nuclear factor- κB (NF- κB) activation within the nucleus and transcription of target genes.

Figure 4 Janus kinase/signal transducers and activators of transcription (JAK/STAT) signaling pathway. Cytokines facilitate activation of receptor-bound JAKs leading to phosphorylation, dimerization, and translocation of STAT to the nucleus to promote transcription of target genes.

remains low, posing an unresolved scientific and clinical challenge.⁵⁷

CBD and JAK/STAT

The use of CBD, to modulate of JAK/STAT is one novel approach towards immunosuppression. Transcriptomic profiles of CBD-treated human Gingival Mesenchymal Stem Cells (hGMSCs) revealed down-regulation of JAK/STAT pathways, particularly complete down-regulation of Interleukin-6 (IL-6), a potent but not exclusive activator of JAK/STAT3 signaling.^{58,59} Similarly, treatment with CBD (ranging from 0.5 μ M to 20 μ M) has been shown to significantly inhibit IL-2 and IFN- γ production in mouse splenocytes.^{30,60} Both IL-2 and IFN- γ are activators of JAK/STAT. IL-2 is also a stimulator of PI3K-Akt and MAPK signaling cascades critical in regulating lymphocyte proliferation. CBD also inhibits another anti-inflammatory and immune regulatory cytokine involved in the modulation of JAK/STAT, Interleukin-10 (IL-10), in human leukemic T-cells.^{61,62}

CBD also has been shown to inhibit JAK/STAT activators, including IFN- γ and inflammatory and immune response tumor necrosis factor (TNF) cytokine in a rheumatoid arthritis model.^{63–65} This study revealed that CBD (5 mg/kg per day intraperitoneally (ip)) reduced TNF and IFN- γ production in arthritic mice. The anti-arthritic and immunosuppressive effects of CBD were further underlined by concentration-dependent (0–10 μ g/mL) inhibition of lymphocyte

proliferation.⁶⁵ CBD (5 mg/kg) is also effective in the suppression of T-cell proliferation and microglial activation, as well as reduction of axon damage within the spinal cord of in an encephalomyelitis murine model of Multiple Sclerosis (MS) by suppression of fundamental JAK/STAT initiators including IL-2, TNF- α , and IFN- γ .¹¹ Notably, T cells express many IL receptors that can lead to activation of JAK/STAT; therefore, suppression of T cells by CBD may inhibit activation of this signaling pathway. High concentration of CBD (20 μ g/mL) also attenuates TNF- α and IFN- γ expression in the intestinal lymphatic system as quantified in rat mesenteric lymph node and spleen isolated T cells and human lymphocytes.⁶⁶

The immunomodulatory effects of CBD have also been observed in Type 1 diabetes mellitus. Type 1 diabetes mellitus is an autoimmune disease leading to T lymphocyte-mediated destruction of insulin-producing pancreatic β cell.⁶⁷ Administration of CBD (5 mg/kg/day ip) has shown to significantly prevent the onset of autoimmune diabetes in non-obese diabetes (NOD) prone mice in comparison to mice that did not receive CBD.^{68,69} These mice displayed significant reduction in Th1 inflammatory cytokine production of IFN- γ and TNF- α , and an increase in Th2 cytokine production of IL-4 and IL-10, suggesting that CBD may exert an immunomodulatory mechanism resulting in a Th1 to Th2 immune response shift.⁷⁰ Similarly, CBD (5 mg/kg/day) was shown to significantly reduce pro-inflammatory cytokine IFN- γ and TNF- α in the plasma of NOD mice treated with CBD.⁶⁹ These mice also exhibited a decrease in Th1 cytokine production and increase in the production of Th2-associated cytokines.⁶⁹ These results suggest that CBD may have an immunomodulatory role, shifting immune responses from Th1 to Th2.⁶⁹ Expression of other initiators and activators of JAK/STAT including IL-6, TNF- α , and IL-12 were also reduced in the CBD-treated splenocytes derived from these non-obese diabetes-prone (NOD) female mice.

CBD has been shown to reduce proinflammatory signaling by modulation of IFN β /STAT pathway⁷¹ such that CBD (10 μ M) reduced IL-1 β , IL-6, and IFN β expression in LPS activated microglial cells. IL-1 β activates myeloid differentiation factor 88 (MyD88)-adaptor protein-dependent pathway that leads to activation of NF- κ B-dependent transcription. The NF- κ B pathway is primarily responsible for regulating the expression of many pro-inflammatory genes including cytokines TNF α , IL-1, and IL-6. IFN β , on the other hand, activates chemokines such as CXCL10, CCL5, and CCL2 by binding to IFN receptor and inducing phosphorylation of JAK leading to STAT pathway activation. CBD also suppresses NF- κ B mediated transcription, in turn enhancing the anti-

inflammatory effects. This process is mediated via STAT3 by increasing anti-inflammatory STAT3 phosphorylation while reducing pro-inflammatory STAT1 phosphorylation.⁷¹ Collectively, these studies further illuminate the pharmacodynamics of CBD and underline the immunomodulatory properties of CBD by down-regulation of JAK/STAT signaling cascade, thus supporting the development of CBD as a novel immune modulation therapeutic.

Nucleotide-Binding Oligomerization Domain-Like Receptor (NLR) Signaling Pathway

Nucleotide-binding oligomerization domain-like receptors (NLRs) serve as cellular receptors that detect and orchestrate innate immune system protection in response to cellular stress and insult, such as pathogen-associated molecular patterns (PAMPs) and danger-associated molecular patterns (DAMPs).^{72,73} When dangerous cellular stimuli or stress is detected, the NLR Family Pyrin Domain Containing 3 gene (NLRP3) consisting of a central nucleotide-binding and oligomerization (NACHT) domain and leucine-rich repeat (LRR) forms an inflammasome complex with an adaptor protein apoptosis-associated speck-like protein (ASC). The ASC consists of an N-terminal pyrin domain (PYD) and C-terminal caspase-recruitment domain (CARD) and cysteine protease caspase 1 (Figure 5).⁷⁴ This activated NLRP3 inflammasome complex leads to activation of caspase 1, in turn, regulating pro-inflammatory interleukin-1 β (IL-1 β) and interleukin-18 (IL-18) cytokine production that play significant roles in NF- κ B, MAPK and IFN pro-inflammatory pathways.⁷⁵

CBD and NLR

An alternative pharmacological mechanism by which CBD induces immunomodulatory effects is by inhibiting activation of the NLR signaling cascade. Next-generation sequencing and gene ontology data reveal that inflammasome activation is inhibited in mesenchymal stem cells treated with CBD (5 μ M) in comparison to stem cells not treated with CBD.⁷⁶ In this study, CBD prevented NLRP3-inflammasome pathway activation by suppressing the expression of key genes belonging to NLRP3-inflammasome pathway including NLRP3 and caspase 1 and inhibited downstream production of IL-18 in Human Gingival Mesenchymal Stem Cells (hGMSCs).⁷⁶ Down-regulation of NLRP3 genes in CBD-treated hGMSCs was also confirmed using immunocytochemical and protein extraction. Gene expression profiling also indicated CBD-

Figure 5 Nucleotide-binding oligomerization domain-like receptor (NLR) signaling pathway. Pathogen-associated molecular patterns (PAMPs) and danger-associated molecular patterns (DAMPs) activate NLR Family Pyrin Domain Containing 3 (NLRP3) consisting of a central nucleotide-binding and oligomerization (NACHT) domain and leucine-rich repeat (LRR). NLRP3 forms an inflammasome complex with adaptor protein apoptosis-associated speck-like protein (ASC) consisting of a protein pyrin domain (PYD) caspase-recruitment domain (CARD) and cysteine protease caspase 1. Activated NLRP3 inflammasome complex activates caspase 1, leading to regulation of pro-inflammatory interleukin-1 β (IL-1 β) and interleukin-18 (IL-18) cytokine production.

induced down-regulation of encoding genes belonging, but not limited to the NLRP3-inflammasome pathway, including pro-inflammatory cytokines (IL6ST, IL-1 β), Interleukin receptors or subunits (IL1R1, IL11RA, IL13RA), Toll-like receptor adaptor myeloid differentiation 88 (MYD88), Interferon Gamma Receptors (IFNGR1 and IFNGR2), Mitogen-Activated Protein Kinases (MAPK1, MAPK12, and MAPK14), transcription factors (STAT3 and STAT6), NF- κ B complex (NFKB2, NFKB3/RELA), and the Matrix Metalloproteinase 3 (MMP3).⁷⁶

To further understand the immunosuppressive effects and modulation of NLRP3 inflammasome pathway, CBD has also been compared to known NLRP3 inflammasome inhibitors, oridonin and MCC950, for their respective inhibition of NLRP3-inflammasome activity.⁷⁷ Before evaluating anti-inflammasome activity of CBD, in vitro toxicity studies were conducted to ensure cellular viability of human monocytes treated with CBD. No significant cytotoxicity was observed among cells treated with CBD (0.1, 1, 10 and 100 μ M), suggesting CBD as a safe and non-toxic potential phyto-therapeutic. LPS was used to significantly increase the production of pro-inflammatory cytokine IL-1 β in monocytes, and CBD (0.1, 1, and 10 μ M) successfully attenuated this IL-1 β production.⁷⁷ As the NLRP3 inflammasome complex is responsible for IL-1 β production, specific CBD-induced modulation of NLRP3 was evaluated by

comparing IL-1 β concentration in LPS-stimulated monocytes treated with CBD, oridonin, and MCC950. CBD and these NLRP3 inhibitors exhibited comparable IL-1 β cytokine inhibitory effects, thus stressing the immunomodulatory effects of CBD by inhibition of the NLRP3 inflammasome pathway, hence, further emphasizing CBD as a prospective immunomodulatory agent.

Adenosine Signaling

Adenosine is a purine nucleoside that protects against excessive cellular damage during stress and inflammation.^{78,79} The protective effects of adenosine are mediated by G protein-coupled A_{2A} receptor, which attenuates inflammation via activation of cAMP-mediated pathway, and ultimately leading to inhibition of T cell differentiation, downregulation of neutrophil superoxide production, and inhibition of proinflammatory cytokines expression including TNF- α (Figure 6).^{80,81} However, rapid intracellular uptake of adenosine, facilitated by the equilibrative nucleoside transporter (ENT), prevents adenosine interaction with their respective immune cell adenosine receptors, and, consequently, blocking these protective effects.^{82,83} One method for hindering adenosine reuptake and promoting immune modulation is to prevent intracellular transport of adenosine by use of an ENT inhibitor.⁸⁴

Figure 6 Adenosine signaling. Equilibrative nucleoside transporter proteins (ENT) facilitate diffusion of adenosine across the cell membrane. Adenosine activates G protein-coupled adenosine receptor (A_{2A}), leading to cAMP formation mediated by adenylate cyclase activity and subsequent activation of protein kinase A (PKA).

CBD and Adenosine Signaling

CBD is a competitive ENT inhibitor ($K_i < 250$ nM) resulting in subsequent anti-inflammatory and immunosuppressive effects by inhibiting adenosine uptake and enhancing adenosine signaling.⁸⁵ In vitro data highlight properties of CBD similar to adenosine receptor agonists such as enhancing endogenous adenosine signaling and inhibiting adenosine uptake in murine microglia and macrophages pretreated with CBD.⁸⁶ Moreover, in vivo treatment with CBD (1 mg/kg) decreased TNF- α as measured in the serum of LPS-treated mice.⁸⁶ However, this effect was eliminated in an A_{2A} receptor knockout mouse model and annulled in the presence A_{2A} adenosine receptor antagonist.⁸⁶ In a separate study, adenosine uptake inhibition was compared in rat retinal microglial cells treated with CBD and the ENT inhibitor S-(4-nitrobenzyl)-6-thioinosine (NBMPR) to understand if the ENT is involved in the pharmacodynamic role of CBD.⁸⁷ In this study, CBD (0.5 μ M) alone was shown to inhibit adenosine uptake in retinal microglial cells and treatment with NBMPR alone inhibited adenosine uptake in a dose-dependent manner. However, adenosine uptake inhibition was not enhanced in CBD-treated cells that were also treated with NBMPR, which lead to the suggestion that the immunosuppressive effects are mediated by competitive binding to ENT.⁸⁷ Other studies have reported that TNF- α in LPS-treated mice is reduced in the presence of adenosine uptake inhibitors.⁸⁸ As a result, endogenous adenosine is readily available to bind to the A_{2A} receptor. To study if CBD inhibits TNF- α by A_{2A} receptor mediation, rats were treated with CBD (1 mg/kg) and A_{2A} receptor antagonist ZM241385 (10 mg/kg). In these experiments, CBD-treated rats showed significant reduction of TNF- α , while treatment with the ZM241385 alone or treatment with CBD and ZM241385 did not inhibit TNF- α concentration, suggesting that CBD inhibits adenosine uptake and reduces TNF- α production by A_{2A} receptor modulation.⁸⁷

The immunosuppressive effects of CBD by inhibition of adenosine reuptake and regulation of A_{2A} receptor have also been evaluated in a murine model of Multiple Sclerosis. For that, mice were infected with Theiler's Murine Encephalomyelitis Virus (TMEV) RNA to induce the deleterious inflammatory effects, such as encephalomyelitis, and then treated with CBD.⁸⁹ Treatment with CBD (5 mg/kg ip) inhibited VCAM-1 production, a mediator of lymphocyte adhesion and leukocyte infiltration produced in response to inflammation. CBD was also found to reduce CCL2 and CCL5 chemokine expression as well as pro-inflammatory cytokines TNF- α and IL-1 β .⁸⁹ CCL2 and CCL5 are associated with macrophage,

T cell, and leukocyte recruitment during inflammation.⁹⁰ These pharmacological effects of CBD were also observed to be dependent on A_{2A} receptor mediation in vivo. ZM241385 was administered to TMEV-induced mice and then treated with CBD shortly after. CBD attenuated VCAM-1 expression and immune cell infiltration. However, these effects were not observed in mice receiving ZM241385 alone, suggesting that the A_{2A} receptor is involved in the mechanism of action of CBD.⁸⁹ These studies further solidify the immunosuppressive effects of CBD and link the modulatory properties of CBD to inhibition of adenosine uptake and its enhancement of adenosine signaling.

Conclusions

Mechanistic understanding of fundamental pharmacological pathways and receptors followed by a survey and analytical review of promising work involving CBD in the context of the aforementioned receptors and signaling cascades validate CBD as a prospective immune modulatory therapeutic agent. CBD exerts immunomodulatory effects in CB₁ and CB₂ receptor knockout mice, suggesting that the significant reduction of immune response by CBD is independent of CB₁ and CB₂ receptors. However, receptor antagonists of CB₁ and CB₂ receptors have shown to reverse the immune modulatory effects of CBD, indicating that CBD is also an inverse agonist of CB₁ and CB₂ receptors. Co-administration of CBD with a TRPV1 antagonist reversed CBD inhibition of immune activation, further indicating the involvement of TRPV1 in the immunomodulatory effects of CBD. Moreover, CBD also down-regulates JAK/STAT signaling by inhibiting immune regulatory cytokines, such as TNF- α and IFN- γ , involved in the modulation of JAK/STAT. CBD also prevents NLRP3-inflammasome pathway activation as well as suppresses gene expression of downstream proteins of the NLRP3-inflammasome pathway. These immunosuppressive effects result in decreased production of pro-inflammatory cytokines, including IL-1 β and IL-18. CBD has also been shown to inhibit cellular adenosine uptake, resulting in enhanced adenosine signaling and protection against tissue injury during inflammation and immune response. Adenosine receptor, A_{2A}, inhibition with an antagonist or in A_{2A} receptor knockout mouse, was shown to reverse these immune modulatory effects of CBD. Ultimately, the findings emphasize the immunosuppressive properties of CBD and its potential use as an attractive therapeutic strategy for immunomodulation.

However, to enhance CBD as a promising therapeutic option, additional mechanistic studies should be conducted in the context of other autoimmune disorders, using various immune system models and stimuli, to generate new knowledge of the immune modulatory effects of CBD in vitro and in vivo. In addition, long-term studies are needed to determine CBD's therapeutic efficacy in chronic immune diseases. Safety and toxicity should be evaluated in these long-term studies to identify any limitations associated with using CBD as a recurring immune modulator. Furthermore, as CBD is a highly hydrophobic molecule and subject to extensive first pass metabolism, its dosages are relatively high. Therefore, there are limitations on how varying dosage affects exhibited immunomodulatory properties. Conclusively, further studies should be conducted to attain a comprehensive understanding of CBD pharmacokinetics and dosages required for therapeutic immune modulation with the ultimate purpose of enhancing CBD as a prospective immune modulatory therapeutic.

Acknowledgments

This work was supported by the National Institutes of Health (R01GM114321, R01GM127706, R01MH104656, R01MH110415, HL126559, DA039576, DA040537, and DA044579) and the National Science Foundation (CHE-1506740, CBET-1841419). SD thanks the Miller School of Medicine of the University of Miami for the Lucille P. Markey Chair in Biochemistry and Molecular Biology.

Disclosure

Nadia Peyravian, Sapna Deo, Sylvia Daunert, and Joaquin J Jimenez report a patent pending: 62/985,529. The authors report no other potential conflicts of interest in this work.

References

1. VanDolah HJ, Bauer BA, Mauck KF. Clinicians' Guide to Cannabidiol and Hemp Oils. *Mayo Clin Proc.* 2019;94(9):1840–1851. doi:10.1016/j.mayocp.2019.01.003
2. Berman P, Futoran K, Lewitus GM, et al. A new ESI-LC/MS approach for comprehensive metabolic profiling of phytocannabinoids in Cannabis. *Sci Rep.* 2018;8(1):14280. doi:10.1038/s41598-018-32651-4
3. Lafaye G, Karila L, Blecha L, Benyamina A. Cannabis, cannabinoids, and health. *Dialogues Clin Neurosci.* 2017;19(3):309–316.
4. Peres FF, Lima AC, Hallak JEC, Crippa JA, Silva RH, Abilio VC. Cannabidiol as a promising strategy to treat and prevent movement disorders? *Front Pharmacol.* 2018;9:482. doi:10.3389/fphar.2018.00482
5. McAllister SD, Soroceanu L, Desprez PY. The antitumor activity of plant-derived non-psychoactive cannabinoids. *J Neuroimmune Pharmacol.* 2015;10(2):255–267. doi:10.1007/s11481-015-9608-y
6. Schonhofen P, de Medeiros LM, Bristol IJ, et al. Cannabidiol Exposure During Neuronal Differentiation Sensitizes Cells Against Redox-Active Neurotoxins. *Mol Neurobiol.* 2015;52(1):26–37. doi:10.1007/s12035-014-8843-1

7. Katchan V, David P, Shoenfeld Y. Cannabinoids and autoimmune diseases: A systematic review. *Autoimmun Rev*. 2016;15(6):513–528. doi:10.1016/j.autrev.2016.02.008
8. Pisanti S, Malfitano AM, Ciaglia E, et al. Cannabidiol: state of the art and new challenges for therapeutic applications. *Pharmacol Ther*. 2017;175:133–150. doi:10.1016/j.pharmthera.2017.02.041
9. Devinsky O, Cross JH, Wright S. Trial of cannabidiol for drug-resistant seizures in the dravet syndrome. *N Engl J Med*. 2017;377(7):699–700.
10. Thiele EA, Marsh ED, French JA, et al. Cannabidiol in patients with seizures associated with Lennox-Gastaut syndrome (GWPCARE4): a randomised, double-blind, placebo-controlled Phase 3 trial. *Lancet*. 2018;391(10125):1085–1096. doi:10.1016/S0140-6736(18)30136-3
11. Kozela E, Lev N, Kaushansky N, et al. Cannabidiol inhibits pathogenic T cells, decreases spinal microglial activation and ameliorates multiple sclerosis-like disease in C57BL/6 mice. *Br J Pharmacol*. 2011;163(7):1507–1519. doi:10.1111/j.1476-5381.2011.01379.x
12. Zou S, Kumar U. Cannabinoid Receptors and the Endocannabinoid System: signaling and Function in the Central Nervous System. *Int J Mol Sci*. 2018;19(3). doi:10.3390/ijms19030833.
13. Turcotte C, Blanchet MR, Laviolette M, Flamand N. The CB2 receptor and its role as a regulator of inflammation. *Cell Mol Life Sci*. 2016;73(23):4449–4470. doi:10.1007/s00018-016-2300-4
14. Basu S, Dittel BN. Unraveling the complexities of cannabinoid receptor 2 (CB2) immune regulation in health and disease. *Immunol Res*. 2011;51(1):26–38. doi:10.1007/s12026-011-8210-5
15. Busquets-Garcia A, Bains J, Marsicano G. CB1 Receptor Signaling in the Brain: extracting Specificity from Ubiquity. *Neuropsychopharmacology*. 2018;43(1):4–20. doi:10.1038/npp.2017.206
16. Bellone C, Luscher C, Mameli M. Mechanisms of synaptic depression triggered by metabotropic glutamate receptors. *Cell Mol Life Sci*. 2008;65(18):2913–2923. doi:10.1007/s00018-008-8263-3
17. Saroz Y, Kho DT, Glass M, Graham ES, Grimsey NL. Cannabinoid Receptor 2 (CB2) Signals via G-alpha-s and Induces IL-6 and IL-10 Cytokine Secretion in Human Primary Leukocytes. *ACS Pharmacol Transl Sci*. 2019;2(6):414–428. doi:10.1021/acspsci.9b00049
18. Haushalter KJ, Casteel DE, Raffaeiner A, Stefan E, Patel HH, Taylor SS. Phosphorylation of protein kinase A (PKA) regulatory subunit RIalpha by protein kinase G (PKG) primes PKA for catalytic activity in cells. *J Biol Chem*. 2018;293(12):4411–4421. doi:10.1074/jbc.M117.809988
19. Sassone-Corsi P. The cyclic AMP pathway. *Cold Spring Harb Perspect Biol*. 2012;4(12):a011148–a011148. doi:10.1101/cshperspect.a011148
20. Dobovisek L, Hojnik M, Ferik P. Overlapping molecular pathways between cannabinoid receptors type 1 and 2 and estrogens/androgens on the periphery and their involvement in the pathogenesis of common diseases (Review). *Int J Mol Med*. 2016;38(6):1642–1651. doi:10.3892/ijmm.2016.2779
21. Dalton GD, Howlett AC. Cannabinoid CB1 receptors transactivate multiple receptor tyrosine kinases and regulate serine/threonine kinases to activate ERK in neuronal cells. *Br J Pharmacol*. 2012;165(8):2497–2511. doi:10.1111/j.1476-5381.2011.01455.x
22. Ozaita A, Puighermanal E, Maldonado R. Regulation of PI3K/Akt/GSK-3 pathway by cannabinoids in the brain. *J Neurochem*. 2007;102(4):1105–1114. doi:10.1111/j.1471-4159.2007.04642.x
23. Sanchez MG, Ruiz-Llorente L, Sanchez AM, Diaz-Laviada I. Activation of phosphoinositide 3-kinase/PKB pathway by CB(1) and CB(2) cannabinoid receptors expressed in prostate PC-3 cells. Involvement in Raf-1 stimulation and NGF induction. *Cell Signal*. 2003;15(9):851–859. doi:10.1016/s0898-6568(03)00036-6
24. Demuth DG, Molleman A. Cannabinoid signalling. *Life Sci*. 2006;78(6):549–563. doi:10.1016/j.lfs.2005.05.055
25. Khan MI, Sobocinska AA, Czarnecka AM, Krol M, Botta B, Szczylik C. The therapeutic aspects of the endocannabinoid system (ECS) for Cancer and their development: from nature to laboratory. *Curr Pharm Des*. 2016;22(12):1756–1766. doi:10.2174/1381612822666151211094901
26. Zlebnik NE, Cheer JF. Beyond the CB1 receptor: is cannabidiol the answer for disorders of motivation? *Annu Rev Neurosci*. 2016;39:1–17. doi:10.1146/annurev-neuro-070815-014038
27. Pertwee RG. The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *Br J Pharmacol*. 2008;153(2):199–215. doi:10.1038/sj.bjp.0707442
28. Bouron A. Phyto and endocannabinoids exert complex actions on calcium and zinc signaling in mouse cortical neurons. *Biochem Pharmacol*. 2018;152:244–251. doi:10.1016/j.bcp.2018.04.003
29. Juknat A, Rimmerman N, Levy R, Vogel Z, Kozela E. Cannabidiol affects the expression of genes involved in zinc homeostasis in BV-2 microglial cells. *Neurochem Int*. 2012;61(6):923–930. doi:10.1016/j.neuint.2011.12.002
30. Kaplan BL, Springs AE, Kaminski NE. The profile of immune modulation by cannabidiol (CBD) involves deregulation of nuclear factor of activated T cells (NFAT). *Biochem Pharmacol*. 2008;76(6):726–737. doi:10.1016/j.bcp.2008.06.022
31. de Filippis D, Iuvone T, d'amico A, et al. Effect of cannabidiol on sepsis-induced motility disturbances in mice: involvement of CB receptors and fatty acid amide hydrolase. *Neurogastroenterol Motil*. 2008;20(8):919–927. doi:10.1111/j.1365-2982.2008.01114.x
32. Pazos MR, Mohammed N, Lafuente H, et al. Mechanisms of cannabidiol neuroprotection in hypoxic-ischemic newborn pigs: role of 5HT(1A) and CB2 receptors. *Neuropharmacology*. 2013;71:282–291. doi:10.1016/j.neuropharm.2013.03.027
33. Assas BM, Miyan JA, Pennock JL. Cross-talk between neural and immune receptors provides a potential mechanism of homeostatic regulation in the gut mucosa. *Mucosal Immunol*. 2014;7(6):1283–1289. doi:10.1038/mi.2014.80
34. Huang WX, Yu F, Sanchez RM, et al. TRPV1 promotes repetitive febrile seizures by pro-inflammatory cytokines in immature brain. *Brain Behav Immun*. 2015;48:68–77. doi:10.1016/j.bbi.2015.01.017
35. Kong WL, Peng YY, Peng BW. Modulation of neuroinflammation: role and therapeutic potential of TRPV1 in the neuro-immune axis. *Brain Behav Immun*. 2017;64:354–366. doi:10.1016/j.bbi.2017.03.007
36. Elokely K, Velisetty P, Delemotte L, et al. Understanding TRPV1 activation by ligands: insights from the binding modes of capsaicin and resiniferatoxin. *Proc Natl Acad Sci U S A*. 2016;113(2):E137–45. doi:10.1073/pnas.1517288113
37. Jiang XX, Liu GY, Lei H, Li ZL, Feng QP, Huang W. Activation of transient receptor potential vanilloid 1 protects the heart against apoptosis in ischemia/reperfusion injury through upregulating the PI3K/Akt signaling pathway. *Int J Mol Med*. 2018;41(3):1724–1730. doi:10.3892/ijmm.2017.3338
38. Wang Z, Yang Y, Yang H, et al. NF-kappaB feedback control of JNK1 activation modulates TRPV1-induced increases in IL-6 and IL-8 release by human corneal epithelial cells. *Mol Vis*. 2011;17:3137–3146.
39. Yang XL, Wang X, Shao L, et al. TRPV1 mediates astrocyte activation and interleukin-1beta release induced by hypoxic ischemia (HI). *J Neuroinflammation*. 2019;16(1):114. doi:10.1186/s12974-019-1487-3
40. Du Q, Liao Q, Chen C, Yang X, Xie R, Xu J. The role of transient receptor potential vanilloid 1 in common diseases of the digestive tract and the cardiovascular and respiratory system. *Front Physiol*. 2019;10:1064. doi:10.3389/fphys.2019.01064
41. Lowin T, Straub RH. Cannabinoid-based drugs targeting CB1 and TRPV1, the sympathetic nervous system, and arthritis. *Arthritis Res Ther*. 2015;17:226. doi:10.1186/s13075-015-0743-x

42. Gouin O, L'Herondelle K, Lebonvallet N, et al. TRPV1 and TRPA1 in cutaneous neurogenic and chronic inflammation: pro-inflammatory response induced by their activation and their sensitization. *Protein Cell*. 2017;8(9):644–661. doi:10.1007/s13238-017-0395-5
43. Hegde VL, Nagarkatti PS, Nagarkatti M. Role of myeloid-derived suppressor cells in amelioration of experimental autoimmune hepatitis following activation of TRPV1 receptors by cannabidiol. *PLoS One*. 2011;6(4):e18281. doi:10.1371/journal.pone.0018281
44. Baral P, Udit S, Chiu IM. Pain and immunity: implications for host defence. *Nat Rev Immunol*. 2019;19(7):433–447. doi:10.1038/s41577-019-0147-2
45. Costa B, Giagnoni G, Franke C, Trovato AE, Colleoni M. Vanilloid TRPV1 receptor mediates the antihyperalgesic effect of the nonpsychoactive cannabinoid, cannabidiol, in a rat model of acute inflammation. *Br J Pharmacol*. 2004;143(2):247–250. doi:10.1038/sj.bjp.0705920
46. Hassan S, Eldeeb K, Millns PJ, Bennett AJ, Alexander SP, Kendall DA. Cannabidiol enhances microglial phagocytosis via transient receptor potential (TRP) channel activation. *Br J Pharmacol*. 2014;171(9):2426–2439. doi:10.1111/bph.12615
47. Fonseca BM, Correia-da-Silva G, Teixeira NA. Cannabinoid-induced cell death in endometrial cancer cells: involvement of TRPV1 receptors in apoptosis. *J Physiol Biochem*. 2018;74(2):261–272.
48. Harrison DA. The Jak/STAT pathway. *Cold Spring Harb Perspect Biol*. 2012;4(3):a011205–a011205. doi:10.1101/cshperspect.a011205
49. Souza-Neto JA, Sim S, Dimopoulos G. An evolutionary conserved function of the JAK-STAT pathway in anti-dengue defense. *Proc Natl Acad Sci U S A*. 2009;106(42):17841–17846. doi:10.1073/pnas.0905006106
50. Waldmann TA, Chen J. Disorders of the JAK/STAT Pathway in T Cell Lymphoma Pathogenesis: implications for Immunotherapy. *Annu Rev Immunol*. 2017;35:533–550. doi:10.1146/annurev-immunol-110416-120628
51. Bousoik E, Montazeri Aliabadi H. “Do We Know Jack” About JAK? A Closer Look at JAK/STAT Signaling Pathway. *Front Oncol*. 2018;8:287. doi:10.3389/fonc.2018.00287
52. Babon JJ, Lucet IS, Murphy JM, Nicola NA, Varghese LN. The molecular regulation of Janus kinase (JAK) activation. *Biochem J*. 2014;462(1):1–13. doi:10.1042/BJ20140712
53. Villarino AV, Kanno Y, Ferdinand JR, O'Shea JJ. Mechanisms of Jak/STAT signaling in immunity and disease. *J Immunol*. 2015;194(1):21–27. doi:10.4049/jimmunol.1401867
54. Darnell JE, Kerr IM, Stark GR. Jak-STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science*. 1994;264(5164):1415–1421. doi:10.1126/science.8197455
55. Johnston PA, Grandis JR. STAT3 signaling: anticancer strategies and challenges. *Mol Interv*. 2011;11(1):18–26. doi:10.1124/mi.11.1.4
56. Yu H, Lee H, Herrmann A, Buettner R, Jove R. Revisiting STAT3 signalling in cancer: new and unexpected biological functions. *Nat Rev Cancer*. 2014;14(11):736–746. doi:10.1038/nrc3818
57. Siveen KS, Sikka S, Surana R, et al. Targeting the STAT3 signaling pathway in cancer: role of synthetic and natural inhibitors. *Biochim Biophys Acta*. 2014;1845(2):136–154. doi:10.1016/j.bbcan.2013.12.005
58. Chiricosta L, Silvestro S, Pizzicannella J, et al. Transcriptomic analysis of stem cells treated with moringin or cannabidiol: analogies and differences in inflammation pathways. *Int J Mol Sci*. 2019;20(23):6039. doi:10.3390/ijms20236039
59. Murakami M, Kamimura D, Hirano T. Pleiotropy and Specificity: insights from the Interleukin 6 Family of Cytokines. *Immunity*. 2019;50(4):812–831. doi:10.1016/j.immuni.2019.03.027
60. Kaplan BL, Rockwell CE, Kaminski NE. Evidence for cannabinoid receptor-dependent and -independent mechanisms of action in leukocytes. *J Pharmacol Exp Ther*. 2003;306(3):1077–1085. doi:10.1124/jpet.103.051961
61. Verma R, Balakrishnan L, Sharma K, et al. A network map of Interleukin-10 signaling pathway. *J Cell Commun Signal*. 2016;10(1):61–67. doi:10.1007/s12079-015-0302-x
62. Srivastava MD, Srivastava BI, Brouhard B. Delta9 tetrahydrocannabinol and cannabidiol alter cytokine production by human immune cells. *Immunopharmacology*. 1998;40(3):179–185. doi:10.1016/S0162-3109(98)00041-1
63. Miscia S, Marchisio M, Grilli A, et al. Tumor necrosis factor alpha (TNF-alpha) activates Jak1/Stat3-Stat5B signaling through TNFR-1 in human B cells. *Cell Growth Differ*. 2002;13(1):13–18.
64. Horvath CM. The Jak-STAT pathway stimulated by interferon gamma. *Sci STKE*. 2004;2004(260):tr8. doi:10.1126/stke.2602004tr8
65. Malfait AM, Gallily R, Sumariwalla PF, et al. The nonpsychoactive cannabis constituent cannabidiol is an oral anti-arthritis therapeutic in murine collagen-induced arthritis. *Proc Natl Acad Sci U S A*. 2000;97(17):9561–9566. doi:10.1073/pnas.160105897
66. Zgair A, Lee JB, Wong JCM, et al. Oral administration of cannabis with lipids leads to high levels of cannabinoids in the intestinal lymphatic system and prominent immunomodulation. *Sci Rep*. 2017;7(1):14542. doi:10.1038/s41598-017-15026-z
67. Burrack AL, Martinov T, Fife BT. T cell-mediated beta cell destruction: autoimmunity and alloimmunity in the context of type 1 diabetes. *Front Endocrinol (Lausanne)*. 2017;8:343. doi:10.3389/fendo.2017.00343
68. Weiss L, Zeira M, Reich S, Slavin S, Raz I, Mechoulam R. Cannabidiol arrests onset of autoimmune diabetes in NOD mice. *Neuropharmacology*. 2008;54(1):244–249. doi:10.1016/j.neuropharm.2007.06.029
69. Weiss L, Zeira M, Reich S, et al. Cannabidiol lowers incidence of diabetes in non-obese diabetic mice. *Autoimmunity*. 2006;39(2):143–151. doi:10.1080/08916930500356674
70. Mechoulam R, Peters M, Murillo-Rodriguez E, Hanus LO. Cannabidiol—recent advances. *Chem Biodivers*. 2007;4(8):1678–1692. doi:10.1002/cbdv.200790147
71. Kozela E, Pietr M, Juknat A, Rimmerman N, Levy R, Vogel Z. Cannabinoids Delta(9)-tetrahydrocannabinol and cannabidiol differentially inhibit the lipopolysaccharide-activated NF-kappaB and interferon-beta/STAT proinflammatory pathways in BV-2 microglial cells. *J Biol Chem*. 2010;285(3):1616–1626. doi:10.1074/jbc.M109.069294
72. Tschopp J, Schroder K. NLRP3 inflammasome activation: the convergence of multiple signalling pathways on ROS production? *Nat Rev Immunol*. 2010;10(3):210–215. doi:10.1038/nri2725
73. Schroder K, Tschopp J. The inflammasomes. *Cell*. 2010;140(6):821–832. doi:10.1016/j.cell.2010.01.040
74. Ruland J. Inflammasome: putting the pieces together. *Cell*. 2014;156(6):1127–1129. doi:10.1016/j.cell.2014.02.038
75. Kolly L, Busso N, Palmer G, Talabot-Ayer D, Chobaz V, So A. Expression and function of the NALP3 inflammasome in rheumatoid synovium. *Immunology*. 2010;129(2):178–185. doi:10.1111/j.1365-2567.2009.03174.x
76. Libro R, Scionti D, Diomedea F, et al. Cannabidiol modulates the immunophenotype and inhibits the activation of the inflammasome in human gingival mesenchymal stem cells. *Front Physiol*. 2016;7:559. doi:10.3389/fphys.2016.00559
77. Liu C, Ma H, Slitt AL, Seeram NP. Inhibitory effect of cannabidiol on the activation of nlrp3 inflammasome is associated with its modulation of the p2x7 receptor in human monocytes. *J Nat Prod*. 2020. doi:10.1021/acs.jnatprod.0c00138
78. Hasko G, Cronstein BN. Adenosine: an endogenous regulator of innate immunity. *Trends Immunol*. 2004;25(1):33–39. doi:10.1016/j.it.2003.11.003
79. Noji T, Karasawa A, Kusaka H. Adenosine uptake inhibitors. *Eur J Pharmacol*. 2004;495(1):1–16. doi:10.1016/j.ejphar.2004.05.003
80. Borea PA, Gessi S, Merighi S, Vincenzi F, Varani K. Pharmacology of adenosine receptors: the state of the art. *Physiol Rev*. 2018;98(3):1591–1625. doi:10.1152/physrev.00049.2017

81. Cronstein BN, Sitkovsky M. Adenosine and adenosine receptors in the pathogenesis and treatment of rheumatic diseases. *Nat Rev Rheumatol*. 2017;13(1):41–51. doi:10.1038/nrrheum.2016.178
82. Jacobson KA, Gao ZG. Adenosine receptors as therapeutic targets. *Nat Rev Drug Discov*. 2006;5(3):247–264. doi:10.1038/nrd1983
83. Pastor-Anglada M, Perez-Torras S. Who is who in adenosine transport. *Front Pharmacol*. 2018;9:627. doi:10.3389/fphar.2018.00627
84. Chen JF, Eltzschig HK, Fredholm BB. Adenosine receptors as drug targets—what are the challenges? *Nat Rev Drug Discov*. 2013;12(4):265–286. doi:10.1038/nrd3955
85. Ibeas Bih C, Chen T, Nunn AV, et al. Molecular Targets of Cannabidiol in Neurological Disorders. *Neurotherapeutics*. 2015;12(4):699–730. doi:10.1007/s13311-015-0377-3
86. Carrier EJ, Auchampach JA, Hillard CJ. Inhibition of an equilibrative nucleoside transporter by cannabidiol: a mechanism of cannabinoid immunosuppression. *Proc Natl Acad Sci U S A*. 2006;103(20):7895–7900. doi:10.1073/pnas.0511232103
87. Liou GI, Auchampach JA, Hillard CJ, et al. Mediation of cannabidiol anti-inflammation in the retina by equilibrative nucleoside transporter and A2A adenosine receptor. *Invest Ophthalmol Vis Sci*. 2008;49(12):5526–5531. doi:10.1167/iovs.08-2196
88. Noji T, Takayama M, Mizutani M, et al. KF24345, an adenosine uptake inhibitor, suppresses lipopolysaccharide-induced tumor necrosis factor- α production and leukopenia via endogenous adenosine in mice. *J Pharmacol Exp Ther*. 2002;300(1):200–205. doi:10.1124/jpet.300.1.200
89. Mecha M, Feliu A, Inigo PM, et al. Cannabidiol provides long-lasting protection against the deleterious effects of inflammation in a viral model of multiple sclerosis: a role for A2A receptors. *Neurobiol Dis*. 2013;59:141–150. doi:10.1016/j.nbd.2013.06.016
90. Owen JL, Criscitiello MF, Libreros S, et al. Expression of the inflammatory chemokines CCL2, CCL5 and CXCL2 and the receptors CCR1-3 and CXCR2 in T lymphocytes from mammary tumor-bearing mice. *Cell Immunol*. 2011;270(2):172–182. doi:10.1016/j.cellimm.2011.05.004

ImmunoTargets and Therapy

Dovepress

Publish your work in this journal

ImmunoTargets and Therapy is an international, peer-reviewed open access journal focusing on the immunological basis of diseases, potential targets for immune based therapy and treatment protocols employed to improve patient management. Basic immunology and physiology of the immune system in health, and disease will be also covered. In addition, the journal will focus on the impact of management

programs and new therapeutic agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/immunotargets-and-therapy-journal>