Awareness of Physicians in Yemen Toward High Blood Pressure Management According to the Eighth Joint National Committee (JNC 8) Guideline

This article was published in the following Dove Press journal:
International Journal of General Medicine

Abdulsalam M Halboup ¹
Gamil Q Othman¹
Mohammed M Battah ¹
Karem H Alzoubi ²
Hebah Sallom³

¹Departments of Clinical Pharmacy and Pharmacy Practice, Faculty of Pharmacy, University of Science and Technology, Sana'a, Yemen; ²Department of Clinical Pharmacy, Faculty of Pharmacy, Jordan University of Science and Technology, Irbid, Jordan; ³Department of Clinical Pharmacy, Faculty of Pharmacy, Near East University, Nicosia, Cyprus, Turkey

Background: Hypertension is a common public health problem that impacts more than one-third of the world population. Awareness of physicians towards the guidelines of high blood pressure management is an essential step to reduce the consequences of high blood pressure.

Objective: This study was aimed to assess the awareness of physicians towards high blood pressure treatment according to the recent report of the Joint National Committee (JNC8) guideline.

Methods: A self-administered questionnaire was distributed to 400 physicians during the period from February to April 2017. Physicians were recruited from public and private hospitals as well as clinics. A validated questionnaire that incorporated the changes seen in JNC 8, as well as the specific modality of hypertension management based on other guidelines, was administered to the participating physicians.

Results: Three hundred and eighty-nine physicians completed the questionnaire; with all the interviewed physicians have ever heard about JNC 8. The practice of general practitioners (GPs) was significantly deviated from the recommended guideline of blood pressure management as compared to consultants, specialists, and residents. Additionally, certain variations were found among consultants, specialists, and residents with slight superiority of consultants towards most aspects.

Conclusion: The finding of this study highlights an inadequate knowledge of GPs in Yemen towards high blood pressure management guidelines. The findings of this study emphasize the necessity for continuous medical education programs that are specially designed to target GPs. Continued update of medical curricula in Yemeni universities is also needed.

Keywords: hypertension, practitioners, physicians, guidelines, Yemen

Introduction

Hypertension is a common public-health challenge worldwide. By 2025, it is estimated that over 1.56 Billion of the world population will have hypertension.¹ Long-standing hypertension is considered a major risk factor for coronary artery disease, stroke, heart failure, peripheral vascular disease, vision loss, and chronic kidney disease.²⁻⁵ The Joint National Committee (JNC) on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure issues hypertension-related guidelines after reviewing emerging therapies and outcomes of ongoing clinical trials. The JNC 8 was the latest guideline released in 2014. It includes major

Correspondence: Mohammed M Battah
Tel +967-777404880
Email mmalbattah@gmail.com

differences from the previous reports regarding the threshold of starting antihypertensive medications, the therapeutic goal of blood pressure (JNC 8 defined similar therapeutic goals for the majority of hypertensive patients except elderly), and the first line drugs category. In JNC 8, the first line drug can be equally selected from four specific classes rather than five classes in previous versions of the guideline. As per JNC 8, β -blockers were not recommended as first line therapy for patients without compelling indication. Additionally, one major difference is that the evidence of JNC 8 was limited to a systematic review of randomized clinical trials.⁶ However, several other points about hypertension were not addressed in the JNC 8 including the definition of hypertension, the use of standard medications in a hypertensive patient with heart failure, and the potential benefits of lifestyle changes. Therefore, these issues were incorporated from the guidelines of JNC 7, the American College of Cardiology Foundation/American Heart Association (ACCF/AHA), the lifestyle workgroup, and the American College of Obstetrician and Gynecologist.^{7–10}

The JNC reports have served as a valuable source for making a clinical decision. The recommended guidelines help in decreasing the variations between practitioners and improving the quality of health care.¹¹ Many studies have connected poor control of blood pressure with inadequate adherence to hypertension guidelines.^{12,13} Implementation of hypertension guidelines in a community with a high prevalence of hypertension might help in reducing untreated/inappropriately treated patients and later might help in reducing the long-term consequences associated with high blood pressure.

In Yemen, the estimated prevalence of hypertension for the age group between 30 and 64 is 17.1% and the crude prevalence was 13.5%.¹⁴ Moreover, among elderly patients who were seeking cataract surgery in eye camps, hypertension was found to be the most prevalent systemic disease.¹⁵ However, no study has been conducted to evaluate the quality of the provided healthcare recommendations for the management of hypertension in Yemen. Hence, this study was aimed to assess the awareness of Yemeni physicians towards hypertension definition, diagnosis, and choice of reasonable therapeutics modalities. This study is also aimed to reveal the knowledge and practice of high blood pressure management among Yemeni physicians with different specialties.

Materials and Methods

Study Design

A self-administered questionnaire was used ([appendix](#)). The final questionnaire was developed in English language, and it consisted of two parts, the first part included four questions related to the demographic data of the participants, including age, gender, physician's category/rank, and type of professional practice setting where they work. The second part included 17 questions, incorporated from JNC 8, JNC 7, ACCF/AHA, American College of Obstetrician and Gynecologist, and the lifestyle workgroup in order to have a comprehensive view about the management of hypertension. Each question has only one correct answer. The questionnaire consisted of 17 items. Five items dealt with classification and goal of therapy, another five items dealt with hypertension with compelling indications, four more items dealt with the lifestyle modifications, one question about pregnancy hypertension, one question about the combined use of Angiotensin Converting Enzyme Inhibitors (ACEi) along with Angiotensin Receptor Blockers (ARBs), and one question about the use of β -blockers in a hypertensive patient without compelling indications.

Validation

The questionnaire has been distributed among four academics with expertise in community medicine, clinical pharmacy, pharmacy practice, and pharmacology, and two internal medicine consultants. This is to ensure the accuracy, appropriateness, and relevance of the included items. The final questionnaire was designed as multiple-choice and the correct answer was one of three options; namely, agree, disagree, or I do not know. In order to ensure reliability and consistency of the questions, questionnaires were distributed among 30 physicians that were not included in the study sample. The calculated Cronbach's alpha was found to be 0.651.

Location and Time of the Study

This study was conducted in Sana'a, the capital of Yemen. The study included 389 physicians; who were interviewed in three public hospitals, two military hospitals, thirty-two private hospitals, and seven clinics. These institutions were chosen to be representative of all hospitals in Yemen since these institutions serve as the referral hospitals for most of the patients in Yemen. The questionnaires were administered and collected within a three-month period from February to

April 2017. The interviewed physicians took an average of 10 minutes to fill the questionnaire.

The study included all physicians who dealt with hypertension patients during their practice, such as general practitioners, cardiologists, nephrologists, gynecologists, and internal medicine physicians.

Sample Size Calculation

The last annual health statistics report that was published in 2014 reported the total number of physicians in Sana'a was 1732 physicians. Of them, 1081 were general prescribers and 651 were specialized physicians.¹⁶ OpenEpi website (http://www.openepi.com/Menu/OE_Menu.htm) was used for the sample size calculation. Participants (n=315) were determined as representative samples with 95% confidence level and 5% accuracy. With expected non-response rate of 20%, the final sample size was approximately 380.

Data Collection

The questionnaire was handed to physicians after explaining the purpose of the study to them. Five students in the fifth year of their pharmacy course at university of Science and Technology were trained on the study protocol before collecting data. Physicians were asked to complete the questionnaire and handed it back immediately after responding to the questions. Those physicians were divided into four groups: general practitioners (those who finished bachelor of medicine without doing specialty in any medical field), residents (those who have enrolled in four or-five-year residency program), specialists (those who have finished internal medicine residency), and consultants (those who have a subspecialty in a specific medical field, such as cardiologists or five-year elapsed after being specialist).¹⁷ The data obtained was checked and prepared for entry into statistical software for subsequent data analysis.

Data Analysis

All data were analyzed using IBM SPSS Statistics version 21.0 for Windows® (IBM Corp., Armonk, NY, USA). Descriptive statistics were used to express the demographic data of the participants. Chi-squared test was used to express the association between the participants' specialties and the questions. $P < 0.05$ was considered statistically significant.

Results

Demographic Data

The questionnaire was completed by 389 physicians; thus, the response rate of the study was 97%. The demographic characteristics of the participants are shown in Table 1. The participants of the study were mainly specialists (38%, n=146) followed by GPs (23%, n=91), residents (20%, n=79), and consultants (19%, n=73). Participants were mainly working in public hospitals (44.73%, n=174) followed by private hospitals (35.73%, n=139), and private clinics (19.54%, n=76). Participants were mostly male physicians (71%, n=276), and the age of the majority (36%, n=140) was more than 40 years (Table 1). All participants reported that they have ever heard about JNC 8.

Lifestyle Modifications

The majority (86%, n=336) of participants agreed that DASH eating plan is recommended as a part of lifestyle changes for hypertension. Moderate physical exercise, reduction in sodium consumption, and weight loss were also recommended by 93.5%, n=364; 69.75%, n=270; 96.75%, n=376 of physicians, respectively (Table 2).

Hypertension Classification and Goal of Therapy

The majority of the participants (78.7%, n=306) knew the correct definition of the prehypertension. The majority also 78.9% (n= 307) agreed that blood pressure higher than

Table 1 Demographics of Physicians Involved in the Study (n=389)

Characteristics		Frequency	Percent (%)
Age (Years)	Under 30	111	29%
	From 30 to 40	138	35%
	Above 40	140	36%
Gender	Male	276	71%
	Female	113	29%
Physician category	Consultant	73	19%
	Specialist	146	38%
	Resident	79	20%
	GPs	91	23%
Professional practice	Public hospital	174	44.73%
	Private hospital	139	35.73%
	Private clinic	76	19.54%

Abbreviation: GPs, general practitioners.

Table 2 Correct Answers Regarding Life Style Modifications (n=389)

Lifestyle Modifications	Consultant		Specialist		Resident		GPs		Total	Chi-Square	P-value
	Freq.	%	Freq.	%	Freq.	%	Freq.	%			
DASH	65	89	131	90	68	86	72	79	86%	10.369	0.110
Moderate exercise	70	96	140	96	75	95	79	87	93.5%	12.858	0.045*
Reduce sodium intake	58	79	102	70	54	68	56	62	69.75%	25.993	0.000*
Weight loss	72	99	142	97	78	99	84	92	96.75%	10.321	0.112

Note: *Significant (P<0.05).

Abbreviations: DASH, dietary approaches to stop hypertension; GPs, general practitioners.

160/100 mmHg is classified as stage 2 hypertension. Yet, only 64% (n=249) of physicians agreed that lowering blood pressure in patients aged 60 years or older without compelling indication to a level of less than 150/90 mm Hg is the target of their treatment. Moreover, 52.25% (n=205) of the physicians agreed that stage two hypertensive patients need antihypertensive drugs combination if the target cannot be achieved by a single drug (Table 3).

Hypertension with Compelling Indications

According to JNC 8, specific medication classes were recommended based on compelling indications, such as chronic kidney disease and diabetes mellitus. Assessment of physician awareness based on these indications as well as other indications from JNC 7, such as post-myocardial infarction, stage C heart failure, and cerebrovascular accident. About 58% (n= 223) of the participants agreed to use ACE inhibitors or ARBs as a first line for the management of hypertension with renal impairment in a patient without renal artery stenosis. For management of hypertension in a patient with DM, about 78.75% (n= 304) of the physicians agreed to use one of the four medication classes (ACEi/ARB, calcium channel blockers, or diuretics) as a first line for this group. In a patient with stage C heart failure,

about 77.25% (n= 305) of the physicians agreed to use standard regimen that consisted of ACEi/ARB, β -blockers, diuretics, and spironolactone as a first line. For the management of hypertension in ischemic heart disease, about 75% (n= 295) of the physicians agreed to use β -blockers with the ACE inhibitor as a first line. For recurrent stroke prevention in a hypertensive patient, approximately 62.75% (n= 243) of physicians agreed to use either ACE inhibitors or diuretics as first line. Moreover, only 46% (n=179) of the participants agreed to use labetalol as a first line agent in a pregnant women with severe hypertension (Table 4).

β -Blockers in a Patient Without Compelling Indication

Using β -blockers in a patient without compiling indications is not recommended in JNC 8 owing to the increased risk of cardiovascular death. However, in JNC 7 β -blockers were used as a first line. Approximately, 64% (n= 247) of the physicians agreed not to use β -blockers as a first line in a hypertensive patient without compelling indications (Table 5). However, GPs were associated with lower disagreement rates with this as compared to consultants, specialists, and residents.

Table 3 Correct Answers Regarding Hypertension Classification, Goal, and Drug Combination (n=389)

HTN Classifications and Goals	Consultant		Specialist		Resident		GPs		Total	Chi-Square	P-value
	Freq.	%	Freq.	%	Freq.	%	Freq.	%			
Prehypertension	67	91.8	109	74.7	63	79.7	67	73.6	78.7%	15.355	0.018*
Stage2 HTN	63	86.3	125	85.6	59	74.7	60	65.9	78.9%	23.286	0.001*
Goal of BP in elderly (≥ 60 year)	54	74.0	96	65.8	51	64.6	48	52.7	64.0%	21.321	0.002*
Need drug combination in stage 2 HTN	38	52	86	59	51	65	30	33	52.25%	35.856	0.000*

Note: *Significant (P<0.05).

Abbreviations: HTN, hypertension; BP, blood pressure; GPs, general practitioners.

Table 4 Correct Answers Regarding Hypertensive Patients with Compelling Indications (n=389)

Compelling Indications	Consultant		Specialist		Resident		GPs		Total	Chi-Square	P-value
	Freq.	%	Freq.	%	Freq.	%	Freq.	%			
HF (stage C)	57	78	125	86	61	77	62	68	77.25%	11.215	0.082
CKD	51	70	84	58	45	57	43	47	58%	16.588	0.011*
First line during pregnancy	35	48	71	49	36	46	37	41	46%	7.694	0.261
CVA	55	75	96	66	51	65	41	45	62.75%	26.301	0.000*
DM	62	85	112	77	62	78	68	75	78.75%	26.705	0.000*
Post MI	58	79	118	81	57	72	62	68	75%	8.466	0.206

Note: *Significant (P<0.05).

Abbreviations: HF, heart failure; CKD, chronic kidney disease; CVA, cerebrovascular accident; DM, diabetes mellitus; MI, myocardial infarction; GPs, general practitioners.

Table 5 Correct Answer Regarding ACE and ARBs Combination as Well as β -Blockers Without Compelling Indications (n=389)

Drug Recommendation Question	Consultant		Specialist		Resident		GPs		Total	Chi-Square	P-value
	Freq.	%	Freq.	%	Freq.	%	Freq.	%			
β -blockers without compelling indications	53	73	96	66	57	72	41	45	64%	24.5	0.000*
ACEIs and ARBs should not be used as combination	54	74	100	68	57	72	46	51	66.25%	28.0	0.000*

Note: *Significant (P<0.05).

Abbreviations: ACEIs, angiotensin converting enzyme inhibitors; ARBs, angiotensin receptor blockers; GPs, general practitioners.

ACEI and ARBs Combination

About 66.25% (n= 257) of the physicians agreed not to combine ACE inhibitors with ARBs for hypertension management (Table 5). Again, lower rate of correct practice was associated with GPs as compared to consultants, specialists, and residents.

Discussion

The results of this survey, which included 389 physicians representing different settings in Sana'a, Yemen, are the first evaluating awareness of physicians about the management of hypertension in Yemen. The results of this survey showed that all participating physicians have ever heard about the guidelines of JNC 8 without necessarily reflecting on their practice. The majority of the participants (96.75%, 93.5%, 86%, and 69.75%) recommended weight loss, moderate physical exercise, DASH, and reduction of sodium intake, respectively, as parts of lifestyle modifications for a patient with hypertension. Similar findings were obtained from another study conducted in Saudi Arabia.¹⁸

Regarding hypertension classification, goal, and drug combinations, the majority (78.7, n=306) of the participants in this study knew the correct definition of the prehypertension state, and agreed that blood pressure greater than 160/100 mm Hg is classified as stage-2

hypertension. Moreover, about half of the physicians' responses demonstrated that a patient with stage two hypertension needs antihypertensive drugs combination if the first drug cannot achieve blood pressure target.

The JNC 8 guideline recommended reducing blood pressure to less than 150/90 mm Hg in elderly patients.⁶ In this study, the majority (64%, n=249) of physicians agreed that lowering blood pressure to less than 150/90 mm Hg is the target of their practice. Compared to this finding, 75% of internal and family medicine residents in Saudi Arabia correctly reported blood pressure targets in the elderly.¹⁹ The higher awareness among Saudi physicians regarding this point may be attributed, in part, to their updated knowledge about the current guidelines since they have more continuous education opportunities and the guidelines resources are routinely available in their clinic. It's worthy to mention also that the recommendation of blood pressure treatment goal of less than 150/90 mm Hg in elderly patients is based on expert opinion (grade E), and has remained a point of controversy among experts. In this study, awareness of physicians regarding the definition of prehypertensive stage was higher (78.7, n=305) than the study that was conducted in Jordan by Al-Azzam and his colleague (36.55%). Saudi physicians, on the other hand, exhibited higher awareness regarding combination therapy

in stage 2 hypertension (69.7%) compared with the Yemeni physicians (52.25%, n=205).

In hypertension with compelling indications, such as chronic kidney disease, diabetes mellitus, heart failure, ischemic heart disease, and cerebrovascular accident, specific antihypertensive medications classes are recommended in order to ameliorate disease progression and to improve patient's outcome.^{9,20,21} For example, in a hypertensive patient with chronic kidney disease, antihypertensive agents that reduce proteinuria have an advantage; therefore, ACE inhibitors or ARBs have been recommended to be more advantageous in such conditions since they improve kidney-related outcomes.^{22–26} However, in the current study, only 58% (n=226) of the physicians agreed to use ACE inhibitors or ARBs as the first line therapy for the management of hypertension with renal impairment in cases without renal artery stenosis. Physicians' awareness in this study was higher than that of Jordanian physicians (40.7%) and lower than that of Saudi physicians (73.4%).^{17,19} The discrepancy in the awareness level among physicians from different Middle Eastern countries may be attributed to many reasons, such as the date of releasing the guidelines, the strength of the recommendation, the practicing area, and the specialties of the studied groups.

It is well documented that high blood pressure is the most important risk factor for cerebral vascular accidents, and lowering the blood pressure in a hypertensive patient has been confirmed to reduce the relative risk of stroke by 35% to 44%.²⁷ Other studies have proven that a combination of ACE inhibitor with thiazide diuretics has extended benefit in recurrent stroke reduction beyond the reduction in blood pressure.^{28,29} The finding of this study showed that 62.75% (n=244) of physicians agreed to use either ACE inhibitors, diuretics or combination of both drugs as first line for recurrent stroke prevention in hypertensive patients. Nevertheless, Al-Azzam and his colleague demonstrated in their study that only 21.4% of the physicians knew the benefits of this combination and recommended its use for their patients.¹⁷ Again, the variation in the awareness between studies may be attributed, in part, to the time of the releasing JNC 7 guidelines. In this study, the guidelines have been extensively disseminated and adopted by Yemeni physicians since the JNC 7 report was released in 2003 compared with the Jordanian study, which was conducted two years after the release of the JNC 7 guidelines.

Regarding diabetes patients with hypertension, the majority of the Yemeni physicians (78.75%, n=306) agreed to use ACE inhibitors, ARBs, thiazide diuretics or calcium channel blockers as the first line therapy for this group of patients. Compared with this finding, 74.3% of physicians in Saudi Arabia agreed to use any of the above medications as first line in a hypertensive patient with diabetes,¹⁹ meaning that the recommendations of JNC 8 guidelines were adopted quickly by Yemeni physicians.

The use of ACE inhibitor, β -blockers, diuretics, and spironolactone as a standard combination for patients with heart failure in stage C has advantages such as slowing the progression of heart failure and remodeling, decreased mortality, and improved patient's quality of life.^{8,30,31} In this study, the majority (77.25%, n=300) of Yemeni physicians agreed to use this standard therapy in hypertensive patients with stage C heart failure. Previous studies showed that 40.1% of Jordanian physicians and 43% of Kuwaiti physicians correctly selected ACE inhibitors and ARBs, respectively, as the first line in a hypertensive patient with heart failure.^{17,32}

Regarding hypertension complicated by ischemic heart disease, studies have shown that both selective and non-selective Beta-1 blockers reduce mortality and the risk of recurrence of myocardial infarction as well as decreasing cardiac dysfunctions.^{33,34} In this study, the majority (75%, n=292) of the physicians agreed to use β -blockers with ACE inhibitors as a first line therapy in a patient who has myocardial infarction and hypertension. Only, 59.3% of the Jordanian physicians correctly selected β -blockers as a first line in a hypertensive patient with ischemic heart disease.¹⁷

Labetalol is an effective antihypertensive agent that blocks both beta-and alpha-adrenergic receptors. It is recommended as a first line agent in pregnant women with severe hypertension.³⁵ Unfortunately, only 46% (n=179) of the participants in our study agreed to use labetalol as a first line agent in pregnant women with severe hypertension. Several studies revealed that labetalol is a preferred antihypertensive agent in chronic hypertension during pregnancy, especially in severe cases since methyldopa has a mild effect on hypertension with slow onset and nifedipine has many adverse effects.^{10,36,37}

β -blockers, according to JNC 8, are no longer recommended as first line agents to treat hypertension in patients without compelling indication owing to their association with greater risk of cardiovascular death when compared with amlodipine-based regimen.³⁸ In

this study, the adherence to this recommendation was reported by 64% (n=249) of the Yemeni physicians. Moreover, it was noticed that the consultants, specialists and residents quickly adopted to the updated guideline compared with GPs. Furthermore, combination therapy of ACE inhibitors and ARBs during the management of hypertension patients should be avoided according to JNC 8. Unfortunately, only 66.25% (n=258) of the Yemeni physicians agreed not to combine ACE inhibitors with ARBs in a patient with hypertension. A comparable result was obtained from family and internal medicine residents in Saudi Arabia where 67.9% of them agreed not to use ACE inhibitors and ARBs together.¹⁹

The finding of this study showed small variations among the responses of consultants, specialists, and residents in different areas of questioning, with slight superiority to consultants in some of the questions. However, GPs demonstrated significantly lower knowledge and awareness regarding guidelines of hypertension management.

This study has some potential limitations; the recruitment of physicians was only from Sana'a city, the capital of Yemen, although other cities may have been considered in order to generalize the awareness level. The self-administered questionnaire is a simple and low-cost tool of measuring awareness. However, it has inherited social desirability, gender, and recall bias, which could lead to overestimation.

Conclusion

This study highlights the inadequate knowledge of GPs in Yemen towards high blood pressure management. The findings of this study emphasize the necessity of continuous medical educations for physicians that are specially designed to target GPs and to continually update the curricula of medical students in Yemen.

Ethical Approval and Consent to Participate

The study protocol was approved by the Ethical Committee of the Medical Research at the University of Science and Technology (MECA No: ECA/UST191). Verbal informed consent from participants was acceptable and approved by the Ethical Committee of the Medical Research at the University of Science and Technology. All participants were informed about the objectives of the study and consented to participate.

Acknowledgments

The authors want to thank 5th year pharmacy students (Dalia Al- Qadi, Tasbeeh Al-Sunidar, Wafa'a Al-Wasabi, Rua'a Ockba, Areej Al-Hitary, Anwar Qadi, Kwther Sary) at University of Science and Technology, Girl Branch for helping in data collection.

Author Contributions

All authors contributed to data analysis, drafting and revising the article, gave final approval of version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors declare that there is no conflicts of interest.

References

1. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *Lancet*. 2005;365(9455):217–223. doi:10.1016/S0140-6736(05)17741-1.
2. Lackland DT, Weber MA. Global burden of cardiovascular disease and stroke: hypertension at the core. *Can J Cardiol*. 2015;31(5):569–571. doi:10.1016/j.cjca.2015.01.009.
3. Wang H, Naghavi M, Allen C, et al. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016;388(10053):1459–1544. doi:10.1016/S0140-6736(16)31012-1
4. Singh R, Suh I, Singh V, et al. Hypertension and stroke in Asia: prevalence, control and strategies in developing countries for prevention. *J Hum Hypertens*. 2000;14(10):749–763. doi:10.1038/sj.jhh.1001057
5. Wong TY, Mitchell P. The eye in hypertension. *Lancet*. 2007;369(9559):425–435. doi:10.1016/S0140-6736(07)60198-6.
6. James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA*. 2014;311(5):507–520.
7. Chobanian AV. National heart, lung, and blood institute joint national committee on prevention, detection, evaluation, and treatment of high blood pressure; national high blood pressure education program coordinating committee: the seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure: the JNC 7 report. *JAMA*. 2003;289:2560–2572. doi:10.1001/jama.289.19.2560
8. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol*. 2013;62(16):1495–1539. doi:10.1016/j.jacc.2013.05.020
9. Eckel RH, Jakicic JM, Ard JD, et al. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2014;63(25Part B):2960–2984. doi:10.1016/j.jacc.2013.11.003
10. ACOG. ACOG Practice Bulletin No. 203: chronic hypertension in pregnancy. *Obstet Gynecol*. 2019;133(1):e26–e50.

11. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: executive summary: a report of the American College of Cardiology/American Heart Association task force on clinical practice guidelines. *J Am Coll Cardiol*. 2019;74(10):1376–1414. doi:10.1016/j.jacc.2019.03.009
12. Hill MN, Miller NH, Degeest S, et al. Adherence and persistence with taking medication to control high blood pressure. *J Am Soc Hypertens*. 2011;5(1):56–63. doi:10.1016/j.jash.2011.01.001
13. Ardery G, Carter BL, Milchak JL, et al. Explicit and implicit evaluation of physician adherence to hypertension guidelines. *J Clin Hypertens (Greenwich)*. 2007;9(2):113–119. doi:10.1111/j.1524-6175.2007.06112.x
14. Motlagh B, O'Donnell M, Yusuf S. Prevalence of cardiovascular risk factors in the Middle East: a systematic review. *Eur J Cardiovasc Prev Rehabil*. 2009;16(3):268–280. doi:10.1097/HJR.0b013e328322ca1b.
15. Al-Shaer MY, Bamashmus MA, Al-Akily SA, Al-Akhlee HA. Prevalence of hypertension, diabetes mellitus and Hepatitis B and C among people seeking cataract surgery in eye camps in Yemen. *Yemeni J Med Health Res*. 2019;8:1&2.
16. MOPHP. *Annual Statistical Health Report*. mopph website 2014. Available from: <http://www.mopph-ye.org/arabic/docs/Report2011.pdf>. Accessed March 1, 2019..
17. Al-Azzam SI, Najjar RB, Khader YS. Awareness of physicians in Jordan about the treatment of high blood pressure according to the seventh report of the Joint National Committee (JNC VII). *Eur J Cardiovasc Nurs*. 2007;6(3):223–232. doi:10.1016/J.EJCNURSE.2006.10.001.
18. Al-Gelban KS, Khan MY, Al-Khaldi YM, et al. Adherence of primary health care physicians to hypertension management guidelines in the Aseer region of Saudi Arabia. *Saudi J Kidney Dis Transpl*. 2011;22(5):941–948.
19. Alshehri NA, Almigbal TH, Alodhayani A, Batais MA. Family and internal medicine resident's awareness of and adherence to the Joint National Committee 8 (JNC 8) hypertension guidelines: a cross sectional study. *Biomed Res*. 2017;28:12.
20. Haller H, Ito S, JL I Jr, et al. Olmesartan for the delay or prevention of microalbuminuria in type 2 diabetes. *N Engl J Med*. 2011;364(10):907–917. doi:10.1056/NEJMoa1007994
21. Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2018;71(19). e127–e248.
22. Strippoli GF, Craig M, Deeks JJ, Schena FP, Craig JC. Effects of angiotensin converting enzyme inhibitors and angiotensin II receptor antagonists on mortality and renal outcomes in diabetic nephropathy: systematic review. *BMJ*. 2004;329(7470):828. doi:10.1136/bmj.38237.585000.7C.
23. Contreras G, Greene T, Agodoa LY, et al. Blood pressure control, drug therapy, and kidney disease. *Hypertension*. 2005;46(1):44–50. doi:10.1161/01.HYP.0000166746.04472.60
24. Silvarino R, Rios P, Baldovinos G, et al. Is chronic kidney disease progression influenced by the type of renin-angiotensin-system blocker used? *Nephron*. 2019;143(2):100–107. doi:10.1159/000500925
25. Zheng CM, Wang JY, Chen TT, et al. Angiotensin-converting enzyme inhibitors or angiotensin receptor blocker monotherapy retard deterioration of renal function in Taiwanese chronic kidney disease population. *Sci Rep*. 2019;9(1):2694. doi:10.1038/s41598-019-38991-z
26. DeJong C, Grant RW. Continuation of angiotensin-converting enzyme inhibitors and angiotensin receptor blockers in the face of kidney disease progression-safe and possibly life saving. *JAMA Intern Med*. 2020;180(5):727. doi:10.1001/jamainternmed.2020.0300.
27. Neal B, MacMahon S, Chapman N; for Blood Pressure Lowering Treatment Trialists' Collaboration. Effects of ACE inhibitors, calcium antagonists, and other blood-pressure-lowering drugs: results of prospectively designed overviews of randomised trials. Blood Pressure Lowering Treatment Trialists' Collaboration. *Lancet*. 2000;356(9246):1955–1964. doi:10.1016/s0140-6736(00)03307-9.
28. Harrow C, Fisher M. Strokes and diabetes. *Pract Diabetes Int*. 2005;22(6):215–221. doi:10.1002/pdi.820.
29. Tzourio C, Anderson C, Chapman N, et al. for PROGRESS Collaborative Group. Effects of blood pressure lowering with perindopril and indapamide therapy on dementia and cognitive decline in patients with cerebrovascular disease. *Arch Intern Med*. 2003;163:1069–1075.
30. Pitt B, Pfeffer MA, Assmann SF, et al. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med*. 2014;370(15):1383–1392. doi:10.1056/NEJMoa1313731
31. Edelmann F, Musial-Bright L, Gelbrich G, et al. Tolerability and feasibility of beta-blocker titration in HFpEF versus HFrEF: insights from the CIBIS-ELD trial. *JACC Heart Fail*. 2016;4(2):140–149. doi:10.1016/j.jchf.2015.10.008
32. Al-Ali KA, Al-Ghanim FA, Al-Furaih AM, Al-Otaibi N, Makboul G, El-Shazly MK. Awareness of hypertension guidelines among family physicians in primary health care. *Alexandria J Med*. 2013;49(1):81–87. doi:10.1016/j.ajme.2012.07.003.
33. Himmelmann A. New information on the role of beta-blockers in cardiac therapy. *Cardiovasc Drugs Ther*. 1999;13(6):469–477. doi:10.1023/A:1007893301243.
34. Khan NA, McAlister FA, Rabkin SW, et al. The 2006 Canadian Hypertension Education Program recommendations for the management of hypertension: part II - Therapy. *Can J Cardiol*. 2006;22(7):583–593. doi:10.1016/S0828-282X(06)70280-X
35. Raio L, Bolla D, Baumann M. Hypertension in pregnancy. *Curr Opin Cardiol*. 2015;30(4):411–415. doi:10.1097/HCO.0000000000000190.
36. Battarbee AN, Sinkey RG, Harper LM, Oparil S, Tita ATN. Chronic hypertension in pregnancy. *Am J Obstet Gynecol*. 2020;222(6):532–541. doi:10.1016/j.ajog.2019.11.1243.
37. Shah S. *Hypertensive Disorders in Pregnancy*. *Obstetric and Gynecologic Nephrology*. Springer; 2020:11–23.
38. Dahlöf B, Sever PS, Poulter NR, et al. Prevention of cardiovascular events with an antihypertensive regimen of amlodipine adding perindopril as required versus atenolol adding bendroflumethiazide as required, in the Anglo-Scandinavian Cardiac Outcomes Trial-Blood Pressure Lowering Arm (ASCOT-BPLA): a multicentre randomised controlled trial. *Lancet*. 2005;366(9489):895–906. doi:10.1016/S0140-6736(05)67185-1

International Journal of General Medicine**Dovepress****Publish your work in this journal**

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>