Emodin Retarded Renal Fibrosis Through Regulating HGF and TGF β –Smad Signaling Pathway

This article was published in the following Dove Press journal:
Drug Design, Development and Therapy

Fan Yang¹
Lu Deng²
JinPeng Li³
MuHu Chen¹
Ying Liu¹
YingChun Hu¹
Wu Zhong¹

¹Department of Emergency Medicine, Affiliated Hospital of Southwest Medical University, Luzhou, Sichuan 646000, People's Republic of China; ²Department of Thyroid Surgery, Affiliated Hospital of Southwest Medical University, Luzhou, Sichuan 646000, People's Republic of China; ³Department of Thyroid and Breast Surgery, Wuhan University Zhongnan Hospital, Wuhan, Hubei 430071, People's Republic of China

Background: Renal fibrosis is a frequently occurring type of chronic kidney disease that can cause end-stage renal disease. It has been verified that emodin or HGF can inhibit the development of renal fibrosis. However, the antifibrotic effect of emodin in combination with HGF remains unclear.

Methods: Cell viability was detected with CCK8. Gene and protein expression in HK2 cells was detected by qRT-PCR and Western blot, respectively. Moreover, a unilateral ureteral obstruction-induced mouse model of renal fibrosis was established for investigating the antifibrotic effect of emodin in combination with HGF in vivo.

Results: HGF notably increased the expression of collagen II in TGF β -treated HK2 cells. In addition, HGF-induced increase in collagen II expression was further enhanced by emodin. In contrast, fibronectin, α SMA and Smad2 expression in TGF β -stimulated HK2 cells was significantly inhibited by HGF and further decreased by combination treatment (emodin plus HGF). Moreover, we found that combination treatment exhibited better antifibrotic effects compared with emodin or HGF in vivo.

Conclusion: These data demonstrated that emodin plus HGF exhibited better antifibrotic effects compared with emodin or HGF. As such, emodin in combination with HGF may serve as a new possibility for treatment of renal fibrosis.

Keywords: fibrosis, combination, emodin, TGF β

Introduction

Renal fibrosis is considered to originate from activation of myofibroblasts in the kidney and infiltration of inflammatory cells.^{1,2} A number of patients with chronic kidney disease will deteriorate to end-stage renal disease (ESRD) once renal fibrosis occurs. The only effective method is transplantation in that situation.³ Furthermore, it has been confirmed that chronic kidney disease patients account for 10% of the world's total population.^{4,5} As such, a new method for cure of renal fibrosis is of great significance.

HGF) contains an α -chain and a β -chain. These chains are composed of a serine protease-like structure and four kringle domains.^{6,7} Upregulation of the HGF–Met pathway initiates dynamic biological responses that help morphogenesis (eg, epithelial tubulogenesis) and cells survival.⁸ Recent reports have indicated that HGF has a significant antifibrotic effect.^{9,10} In addition, *Rheum officinale* has been confirmed to have some bioactivity, including antifibrotic activity.^{11,12} Emodin is the major bioactive compound of *R. officinale* and has strong anti-

Correspondence: Wu Zhong
Department of Emergency Medicine,
Affiliated Hospital of Southwest Medical
University, 25 Taiping Street, Luzhou,
Sichuan 646000, People's Republic of
China
Email zhongwu3987@126.com

inflammatory and antioxidant properties.¹³ It has been reported that emodin notably suppresses the development of multiple diseases.^{14,15} Additionally, it has previously been verified that emodin can relieve the progression of renal fibrosis.¹⁶ However, the antifibrotic effect of emodin in combination with HGF has not been confirmed yet. TGF β participates in many cellular processes, including cell growth, differentiation, apoptosis, and homeostasis.^{17,18} Many studies have indicated that TGF β plays a crucial role in renal fibrosis.^{19,20} Yasumura et al verified that TGF β contributes to the occurrence of renal fibrosis.²¹ In this study, we sought to confirm whether emodin can increase the antifibrotic effect of HGF, with the purpose of providing new strategies for patients.

Methods

Cell Culture

Human renal tubular endothelial cells (HK2; ATCC, Manassas, VA, USA) were cultured in RPMI 1640 medium (Thermo Fisher Scientific, Waltham, MA, USA) supplemented with 10% FBS and 1% penicillin (Thermo Fisher Scientific) in an incubator (37°C, 5% CO₂). To mimic renal fibrosis in vitro, HK2 cells were stimulated with TGF β ₁ (5 ng/mL; PeproTech, Rocky Hill, NJ, USA) for 48 hours.

CCK8 Assays

HK2 cells (5×10³ cells/well) were seeded at 37°C overnight. Then, cells were treated with emodin (10, 20, 40, and 60 μ M) for 2 days. After that, 10 μ L CCK8 reagent (Beyotime) per well was added for 2 hours. Finally, cell viability was measured with a microplate reader (Bio-Rad Laboratories, Benicia, CA, USA) at 450 nm absorbance. Standard emodin products (95% purity) were purchased from Sigma-Aldrich (St Louis, MO, USA).

Immunofluorescence Staining

HK2 cells were seeded overnight. After 72 hours' treatment, cells were blocked with 10% FBS for 30 minutes and then incubated with antiSmad4 antibody (Abcam, Cambridge, MA, USA; 1:1,000) overnight. Then, cells were incubated with goat antirabbit IgG (Abcam; 1:5,000) for 1 hour. After that, the nuclei were stained with DAPI (Beyotime) for 5 minutes. Finally, cells were observed under microscopy (Olympus CX23, Tokyo, Japan).

Reverse-Transcription Quantitative Polymerase Chain Reaction

Total RNA was extracted using Trizol reagent (Thermo Fisher Scientific). RNA integrity was measured by agarose-gel electrophoresis. Then, cDNA was obtained by reverse transcription (PrimeScript first-strand cDNA-synthesis kit; Takara, Tokyo, Japan). PCR reactions were carried out by SYBR Premix Ex Taq II (Takara) with an Applied Biosystems 7500 real-time PCR system (Thermo Fisher Scientific). Primers were: collagen II — forward 5'-AACCAAGGGAAGCATTACTAC-3', reverse 5'-GAGCAAATATTCCATAAGCC-3'; fibronectin — forward 5'-CTTGAAGCCCAGCAACCTAC-3', reverse 5'-ACAGCCCACAGACCACACAT-3'; α SMA — forward 5'-TTTGAGACGAGGGAGATGGT-3', reverse 5'-GTTCTTGGTTGGAGCGGAT-3'; and GAPDH — forward 5'-GTCCACCGCAAATGCTTCTA-3', reverse 5'-TGCTGTCACCTTCACCGTTC-3'. Amplification conditions were 95°C for 30 seconds, followed by 40 cycles of 95°C for 5 seconds and 60°C for 34 seconds. The 2^{- $\Delta\Delta$ Ct} method was used to quantify the data. β -actin was used as an internal control.

Western Blot

Cells or renal tissue were lysed using RIPA lysis buffer (Beyotime). Protein concentration was quantified using a BCA protein-assay kit (Beyotime). Protein (30 μ g) was separated with 10% SDS-PAGE and then transferred into PVDF membranes (Beyotime). After blocking with 5% skim milk, the membranes were incubated with primary antibodies (anti-collagen II 1:1,000, anti- α SMA 1:1,000; anti-fibronectin 1:1,000; anti-Smad2 1:1,000; and anti-GAPDH 1:1,000) at 4°C overnight. All primary antibodies were purchased from Abcam. Then, membranes were incubated with secondary antibodies (goat antirabbit IgG; Abcam, 1:5,000) at room temperature for 1 hour. Protein bands were visualized using BeyoECL Plus (Beyotime) and measured using IPP Image-Pro Plus software.

Animal Study

The experimental protocol was approved by the Ethics Committee for Animal Experimentation of Southwest Medical University, and animal study was performed according to the Guidelines for Animal Experimentation of Southwest Medical University. The ethical approval reference number of the study is 20191010.C57BL/6 SPF class-bred mice (18±2g) were obtained from the

Figure 1 Emodin significantly suppressed the fibrotic effect of TGFβ₁ in vitro. (A) HK2 cells were treated with 0, 10, 20, 40, and 60 μM emodin for 48 hours and cell viability determined with CCK8 assay. (B) HK2 cells were treated with 5 ng/mL TGFβ₁, 5 ng/mL TGFβ₁ + 10 μM emodin, or 5 ng/mL TGFβ₁ + 20 μM emodin for 48 hours. Then, collagen II, fibronectin, and αSMA expression in HK2 cells was detected by Western blot. (C) Relative expression of collagen II was quantified normalized to GAPDH. (D) Relative expression of fibronectin was quantified normalized to GAPDH. (E) Relative expression of αSMA was quantified normalized to GAPDH. ***P*<0.01 vs 0 μM group or control; ##*P*<0.01 vs TGFβ₁ group.

Laboratory Animal Center of Southwest Medical University with. Mice were housed in a 12-hour light/12-hour dark cycle at 22°C–24°C. After 1 week of adaptive feeding, mice were divided into groups for the following 3 weeks: control (n=12, treated with normal saline), unilateral ureteral obstruction (UUO; n=12, treated with normal saline) UUO plus 50 mg/kg/day emodin (n=12, orally treated with 50 mg/kg/day emodin, emodin dissolved in 2 mL normal saline), UUO plus 0.5 μg/kg/day HGF (n=12, intravenous injection of 0.5 μg/kg/day HGF, HGF dissolved in 1 mL normal saline), and UUO+ HGF + emodin (n=12, injection of 0.5 μg/kg/day HGF and orally treated with 50 mg/kg/day emodin). Standard HGF was purchased from MedChemExpress (99.36% purity, Monmouth Junction, NJ, USA). The dosages of HGF and emodin were selected based on previous research.^{22,23} After treatment, the level of creatinine (Cr) and blood urea nitrogen (BUN) were examined. Kidney tissues were extracted from each mouse and

dissected, then symptoms of renal fibrosis was detected with H&E and Masson's trichrome. The severity of fibrosis was assessed by calculating the ratio of blue-stained scarred areas to total area.

Enzyme-Linked Immunosorbent Assays

At the end of the animal study, mice were killed for collection of sera and levels of Cr and BUN measured with an ELISA kit (Jiancheng Bioengineering Institute, Nanjing, China) according to the manufacturer's instructions.

Statistical Analysis

For each group, at least three independent experiments were performed. All data are expressed as means ± SD. Comparisons among multiple groups were analyzed using one-way ANOVA followed by Tukey's test. *P*<0.05 was considered statistically significant.

Figure 2 HGF significantly decreased the fibrotic effect of TGFβ₁ in HK2 cells. HK2 cells were treated with 5 ng/mL TGFβ₁ and then with 2, 5 or 10 ng/mL HGF for 48 hours. Then, expression of (A and B) collagen II, (A and C) fibronectin and αSMA, and (A and D) in HK2 cells was detected by Western blot. Relative protein expression was quantified normalized to GAPDH (E). Collagen II, fibronectin, and αSMA expression in HK2 cells was detected by qPCR. GAPDH was used as an internal control. ***P*<0.01 vs control; #*P*<0.05 vs TGFβ₁ group; ##*P*<0.01 vs TGFβ₁.

Results

Emodin Significantly Suppressed Fibrotic Effect of TGFβ₁ In Vitro

For detection of the cytotoxicity of emodin on HK2 cells, CCK8 assays were used. As shown in Figure 1A, compared to 0 μM, 40 or 60 μM emodin significantly inhibited cell viability, while 10 or 20 μM had no significant effect. These data demonstrated that ≥40 μM emodin showed obvious cytotoxicity. Furthermore, as indicated in Figure 1B–E, the expression of collagen II in HK2 cells was notably suppressed by TGFβ₁, which was significantly rescued by 20 μM emodin. In contrast, TGFβ₁ markedly activated fibronectin and αSMA in HK2 cells, while TGFβ₁-induced increase of these two proteins was significantly reversed by 20 μM emodin. As we know, fibronectin, collagen II, and αSMA are the key markers in fibrosis.^{24,25} Emodin 10 μM affected collagen II, fibronectin, and αSMA expression to a limited. Therefore, 20 μM emodin was selected for the

following experiments. These data showed that emodin significantly suppressed the fibrotic effect of TGFβ₁ in vitro.

HGF Significantly Decreased Fibrotic Effect of TGFβ₁ in HK2 Cells

To detect protein expression, Western blot was used. As demonstrated in Figure 2A–D, TGFβ₁ obviously upregulated fibronectin and αSMA and downregulated collagen II in HK2 cells. Furthermore, 2, 5, or 10 ng/mL HGF dose-dependently activated collagen II and suppressed protein levels of fibronectin and αSMA in HK2 cells after TGFβ₁ treatment. Since 10 ng/mL HGF exhibited a better antifibrotic effect compared with the other two concentrations, 10 ng/mL HGF was selected for use in the following experiments. For the purpose of confirming this result, RT-qPCR was used. As demonstrated in Figure 2E, HGF obviously activated collagen II and fibronectin and inactivated αSMA in TGFβ₁-stimulated HK2 cells. Taken

Figure 3 Combination of emodin with HGF suppressed TGF β_1 -induced renal fibrosis via inhibition of the TGF β_1 -signaling pathway. (A) After 48 hours' incubation, α SMA, collagen II, and fibronectin expression in HK2 cells was investigated with RT-qPCR. GAPDH was used as a loading control. (B) Protein expression of collagen II, fibronectin, α SMA, p-Smad2, and Smad2 in HK2 cells was detected by Western blot. Relative protein expression of (C) collagen II, (D) fibronectin, (E) α SMA, and (F) p-Smad2 was quantified by normalizing to GAPDH. (G) Smad4 expression in HK2 cells was detected by immunofluorescence staining. ** $P<0.01$ vs control group; # $P<0.05$ vs TGF β_1 group; ## $P<0.01$ vs TGF β_1 group; ^^ $P<0.01$ vs TGF β_1 + HGF group.

together, all these data revealed that HGF significantly suppressed the fibrotic effect of TGF β_1 in HK2 cells.

Combination Treatment Exhibited Significant Suppressive Effect on Renal Fibrosis In Vitro Through Inactivation of TGF β_1 Pathway

To investigate gene expression in vitro, RT-qPCR was used. As shown in Figure 3A, the expression of collagen II in TGF β_1 -stimulated HK2 cells was notably upregulated by HGF and further increased by emodin. In contrast, HGF notably inhibited the expressions of α SMA and fibronectin in TGF β_1 -treated HK2 cells. As expected, the inhibitory effect of HGF on α SMA and fibronectin was further enhanced in the presence of emodin (Figure 3A). Additionally, protein expression of fibronectin, α SMA, and p-Smad2 in TGF β_1 -treated HK2 cells was significantly decreased by HGF and/or emodin (Figure 3B–F). In contrast, TGF β_1 -induced decrease of collagen II in HK2 cells was notably reversed by HGF and/or emodin. Consistently, the combination of HGF with emodin exhibited better effect on protein expressions compared with either treatment alone (Figure 3B–F). Also, data on Smad4 staining revealed that

TGF β_1 -induced upregulation of Smad4 expression in nuclei of HK2 cells was significantly decreased by HGF and/or emodin (Figure 3G). To sum up, combination treatment exhibited a significant suppressive effect on renal fibrosis in vitro via inactivation of the TGF β_1 pathway.

Emodin in Combination with HGF Significantly Attenuated Symptoms of UUO-Induced Mouse Renal Fibrosis In Vivo

To investigate the antifibrotic effect of emodin in combination with HGF in vivo, a UUO mice model was established. The results revealed that HGF or emodin notably suppressed renal fibrosis in UUO-treated mice. Moreover, renal fibrosis in UUO-treated mice was further attenuated by combination treatment (Figure 4A). Renal fibrosis symptoms were significantly induced by UUO treatment, which was obviously attenuated by emodin, HGF, or combination treatment (Figure 4B). Fibrosis area was significantly upregulated in UUO-treated mice and significantly downregulated by emodin or HGF alone. Consistently, combination treatment showed more significant antifibrotic effects than emodin- or HGF-alone treatment (Figure 4C). ELISA data demonstrated that HGF and/or emodin notably inhibited levels of Cr and BUN in sera of HFD-treated mice (Figure 4D and E).

Figure 4 Emodin in combination with HGF notably attenuated UUO-induced renal fibrosis of mice in vivo. **(A)** H&E staining of mouse-kidney tissue in control, UUO, UUO plus 50 mg/kg emodin, UUO plus 0.5 μg/kg HGF, and UUO + emodin + HGF. **(B)** Masson staining of mouse-kidney tissue in control, UUO, UUO plus 50 mg/kg emodin, UUO plus 0.5 μg/kg HGF, and UUO + emodin + HGF group. **(C)** Fibrosis area in mice was quantified. **(D)** Levels of creatinine in sera of mice were detected with ELISA. **(E)** Levels of BUN in sera of mice were detected with ELISA. ** $P < 0.01$ vs control group; # $P < 0.05$ vs UUO group; ## $P < 0.01$ vs UUO group; ^^ $P < 0.01$ vs UUO + 0.5 μg/kg HGF group.

Altogether, combination of emodin with HGF exhibited more significant antirenal fibrosis effect in vivo than emodin- or HGF-alone treatment.

Combination of Emodin with HGF Suppressed Progression of Renal Fibrosis via Inactivation of TGFβ₁ Pathway In Vivo

Finally, to confirm the mechanism by which emodin in combination with HGF relieved renal fibrosis symptoms, collagen II, αSMA, and p-Smad2 expression in mice was detected. As shown in Figure 5A–D, emodin or HGF significantly inactivated αSMA and p-Smad2 in tissue of UUO-treated mice. Moreover, combination treatment further decreased expression of these two protein. In contrast, protein level of collagen II in tissue of UUO-treated mice was greatly upregulated by HGF and/or emodin. Taken together, combination of emodin with HGF attenuated symptoms of renal fibrosis via inactivation of the TGFβ₁ pathway in vivo.

Discussion

In this research, we sought to detect the antifibrotic effect of combination treatment. It has been confirmed that transplantation of HGF-transgenic mesothelial cell sheets show therapeutic effects on renal fibrosis²⁶ and can be used as an antifibrotic agent.²⁷ The present study verified that HGF suppressed the progression of renal fibrosis at different concentrations. Emodin has already been confirmed to be able to be used for the treatment of multiple diseases.^{22,28} In addition, recent reports have revealed that emodin plays an inhibitory role in neuropathy and hepatopathy.^{16,29} Some natural properties show antifibrotic ability.^{30,31} Our research was consistent with these studies, further confirming that emodin can be considered an antifibrotic agent. Briefly, our findings strongly confirm that combination treatment shows more significant antifibrotic effects than HGF or emodin alone.

Combination treatment exhibited significant antifibrotic effects via inactivating fibronectin and αSMA and activating collagen II. Previous research has indicated that fibronectin plays a crucial role in fibrosis.³² Moreover, Notch3

Figure 5 Emodin in combination with HGF inhibited renal fibrosis in vivo by suppression of TGF β ₁ pathway. **(A)** Protein expression of collagen II, α SMA, Smad2, and p-Smad2 in kidney tissues of mice was measured by Western blot. Relative protein expression of **(B)** collagen II, **(C)** α SMA, and **(D)** p-Smad2 were quantified by normalizing to GAPDH. ** P <0.01 vs control group; ### P <0.01 vs UUO group; ^^ P <0.01 vs TGF β ₁ + HGF group.

relieves symptom of cardiac fibrosis via downregulation of α SMA.²⁵ Collagen II has been regarded as being downregulated during the progression of fibrosis.³³ Consistent with these data, emodin in combination with HGF suppressed the development of renal fibrosis via decreasing fibronectin and α SMA and upregulating collagen II in vitro. α SMA plays key

roles in epithelial–mesenchymal transition (EMT).^{34–36} In this study, the findings indicated that HGF can inactivate α SMA. Emodin significantly enhanced the suppressive effect of HGF on this protein. These data are similar to a recent report, suggesting that HGF and/or emodin exhibit antifibrotic effects via suppression of the EMT process.

TGF β ₁ signaling has been confirmed to be involved in fibrosis,^{37,38} and is activated persistently in fibrosis.^{39,40} Smad4 is at the core of the TGF β ₁-signaling pathway.⁴¹ In addition, it has been verified that TGF β ₁ can upregulate Smad2, which can affect this signaling pathway, resulting in fibrosis.⁴² In the current research, combination treatment further inhibited the expression of p-Smad2 in HK2 cells after TGF β ₁ treatment compared with HGF or emodin alone. Based on these data, the mechanism underlying the antifibrotic effects of emodin in combination with HGF was closely correlated with inactivation of TGF β ₁ pathways. In recent research, emodin relieved the severity of fibrosis via down-regulation of TGF β ₁-Smad signaling.⁴³ Additionally, emodin relieved fibrosis symptoms via inactivation of TGF β ₁-Smad signaling.⁴⁴ Our study was consistent with these results. Also, our study revealed that α SMA was notably activated in HK2 cells after TGF β ₁ treatment. Sisto et al revealed that activation of TGF β ₁ signaling can contribute to the upregulation of EMT in Sjögren's syndrome.⁴⁵ Consistently, our findings suggested that TGF β ₁ signaling upregulated EMT in fibrosis. To sum up, emodin has the potential ability to downregulate fibrosis by inhibiting TGF β ₁-Smad signaling. Frankly speaking, our research has paid attention only to the effect of emodin in combination with HGF on the TGF β ₁ pathway so far. We will detect the effect of combination treatment on other signaling pathways in future. We have revealed that emodin in combination with HGF shows significantly antifibrotic effects via inactivation of TGF β ₁ signaling. Therefore, the combination of emodin with HGF can be used as an effective tool in the treatment of renal fibrosis.

Funding

This research was supported by the Fund for Doctor Research Initiation from the Affiliated Hospital of Southwest Medical University (19044).

Disclosure

The authors report no conflicts of interest for this work.

References

- Hewitson TD. Renal tubulointerstitial fibrosis: common but never simple. *Am J Physiol Renal Physiol*. 2009;296(6):F1239–F1244. doi:10.1152/ajprenal.90521.2008
- Yang Q, Wang J, Zhong P, et al. The clinical prognostic value of lncRNA FAM83H-AS1 in cancer patients: a meta-analysis. *Cancer Cell Int*. 2020;20:72. doi:10.1186/s12935-020-1148-8
- Santos-Paul MA, Neves RS, Gowdak LHW, et al. Cardiovascular risk reduction with periodontal treatment in patients on the waiting list for renal transplantation. *Clin Transplant*. 2019;33(8):e13658. doi:10.1111/ctr.13658
- Elkomy N, Ibrahim I, El-Fayoumi HM, Elshazly SM. Effect of imidazoline-1 receptor agonists on renal dysfunction in rats associated with chronic, sequential fructose and ethanol administration. *Clin Exp Pharmacol Physiol*. 2019.
- Yang SZ, Ning SS, Sun YX, et al. [Effect of electroacupuncture on renal fibrosis in spontaneously hypertension rats and its related mechanisms]. *Zhen Ci Yan Jiu*. 2019;44(12):911–915. doi:10.13702/j.1000-0607.190167. Chinese.
- Matsumoto K, Funakoshi H, Takahashi H, Sakai K. HGF-met pathway in regeneration and drug discovery. *Biomedicines*. 2014;2(4):275–300. doi:10.3390/biomedicines2040275
- Atta H, El-Rehany M, Hammam O, et al. Mutant MMP-9 and HGF gene transfer enhance resolution of CCl₄-induced liver fibrosis in rats: role of ASH1 and EZH2 methyltransferases repression. *PLoS One*. 2014;9(11):e112384. doi:10.1371/journal.pone.0112384
- Du T, Zou X, Cheng J, et al. Human Wharton's jelly-derived mesenchymal stromal cells reduce renal fibrosis through induction of native and foreign hepatocyte growth factor synthesis in injured tubular epithelial cells. *Stem Cell Res Ther*. 2013;4(3):59. doi:10.1186/scrt215
- Lv W, Booz GW, Wang Y, Fan F, Roman RJ. Inflammation and renal fibrosis: recent developments on key signaling molecules as potential therapeutic targets. *Eur J Pharmacol*. 2018;820:65–76. doi:10.1016/j.ejphar.2017.12.016
- Liu S, Zhao M, Zhou Y, et al. A self-assembling peptide hydrogel-based drug co-delivery platform to improve tissue repair after ischemia-reperfusion injury. *Acta Biomater*. 2019.
- Jiang Y, Liu R, Chen J, et al. Application of multifold characteristic ion filtering combined with statistical analysis for comprehensive profiling of chemical constituents in anti-renal interstitial fibrosis I decoction by ultra-high performance liquid chromatography coupled with hybrid quadrupole-orbitrap high resolution mass spectrometry. *J Chromatogr A*. 2019;1600:197–208. doi:10.1016/j.chroma.2019.04.051
- Sargin G, Senturk T, Cildag S. Systemic sclerosis and malignancy. *Int J Rheum Dis*. 2018;21(5):1093–1097. doi:10.1111/1756-185X.13311
- Yang M, Wei Y, Ashokkumar M, et al. Effect of ultrasound on binding interaction between emodin and micellar casein and its microencapsulation at various temperatures. *Ultrason Sonochem*. 2019;104861.
- Kim YG, Park YH, Yang EY, Park WS, Park KS. Inhibition of tamoxifen's therapeutic effects by emodin in estrogen receptor-positive breast cancer cell lines. *Ann Surg Treat Res*. 2019;97(5):230–238. doi:10.4174/astr.2019.97.5.230
- Niu Y, Zhang J, Tong Y, Li J, Liu B. Physcion 8-O-beta-glucopyranoside induced ferroptosis via regulating miR-103a-3p/GLS2 axis in gastric cancer. *Life Sci*. 2019;237:116893. doi:10.1016/j.lfs.2019.116893
- Dou F, Liu Y, Liu L, et al. Aloe-emodin ameliorates renal fibrosis via inhibiting PI3K/Akt/mTOR signaling pathway in vivo and in vitro. *Rejuvenation Res*. 2019;22(3):218–229. doi:10.1089/rej.2018.2104
- Wu H, Yu Y, Huang H, et al. Progressive pulmonary fibrosis is caused by elevated mechanical tension on alveolar stem cells. *Cell*. 2020;180(1):107–121 e117. doi:10.1016/j.cell.2019.11.027
- Ishikawa K, Kohno RI, Mori K, et al. Increased expression of periostin and tenascin-C in eyes with neovascular glaucoma secondary to PDR. *Graefes Arch Clin Exp Ophthalmol*. 2019.
- Wang Y, Xing QQ, Tu JK, et al. Involvement of hydrogen sulfide in the progression of renal fibrosis. *Chin Med J (Engl)*. 2019;132(23):2872–2880. doi:10.1097/CM9.0000000000000537
- Yao W, Guo A, Han X, et al. Aerosol inhalation of a hydrogen-rich solution restored septic renal function. *Aging (Albany NY)*. 2019;11(24):12097–12113. doi:10.18632/aging.102542

21. Yasumura S, Naito Y, Okuno K, et al. Effects of HETEROZYGOUS TFR1 (Transferrin Receptor 1) deletion in pathogenesis of renal fibrosis in mice. *Hypertension*. 2020;75(2):413–421. doi:10.1161/HYPERTENSIONAHA.119.13670
22. Ma L, Li H, Zhang S, et al. Emodin ameliorates renal fibrosis in rats via TGF-beta1/Smad signaling pathway and function study of Smurf 2. *Int Urol Nephrol*. 2018;50(2):373–382. doi:10.1007/s11255-017-1757-x
23. Xu J, Yu TT, Zhang K, et al. HGF alleviates renal interstitial fibrosis via inhibiting the TGF-beta1/SMAD pathway. *Eur Rev Med Pharmacol Sci*. 2018;22(22):7621–7627. doi:10.26355/eurrev_201811_16378
24. Shin SK, Kim KO, Kim SH, et al. Exogenous 8-hydroxydeoxyguanosine ameliorates liver fibrosis through the inhibition of Rac1-NADPH oxidase signaling. *J Gastroenterol Hepatol*. 2020;35(6):1078–1087. doi:10.1111/jgh.14979
25. Feng F, Cheng P, Zhang H, et al. The protective role of tanshinone IIA in silicosis rat model via TGF-beta1/Smad signaling suppression, NOX4 Inhibition and Nrf2/ARE signaling activation. *Drug Des Devel Ther*. 2019;13:4275–4290. doi:10.2147/DDDT.S230572
26. Imafuku A, Oka M, Miyabe Y, et al. Rat mesenchymal stromal cell sheets suppress renal fibrosis via microvascular protection. *Stem Cells Transl Med*. 2019;8(12):1330–1341. doi:10.1002/sctm.19-0113
27. Kim YC, Lee J, An JN, et al. Renoprotective effects of a novel cMet agonistic antibody on kidney fibrosis. *Sci Rep*. 2019;9(1):13495. doi:10.1038/s41598-019-49756-z
28. Wang XT, Sun XJ, Li C, et al. Establishing a cell-based high-content screening assay for TCM compounds with anti-renal fibrosis effects. *Evid Based Complement Alternat Med*. 2018;2018:7942614. doi:10.1155/2018/7942614
29. Xiang Z, Sun H, Cai X, Chen D, Zheng X. The study on the material basis and the mechanism for anti-renal interstitial fibrosis efficacy of rhubarb through integration of metabonomics and network pharmacology. *Mol Biosyst*. 2015;11(4):1067–1078. doi:10.1039/C4MB00573B
30. Liu K, Zhou S, Liu J, et al. Silibinin attenuates high-fat diet-induced renal fibrosis of diabetic nephropathy. *Drug Des Devel Ther*. 2019;13:3117–3126. doi:10.2147/DDDT.S209981
31. Ma F, Li L, Wang Q, et al. Qi-dan-di-huang decoction alleviates diabetic nephropathy by inhibiting the NF-kappaB pathway. *Front Biosci (Landmark Ed)*. 2019;24:1477–1486. doi:10.2741/4792
32. Yang C, Chen Z, Yu H, Liu X. Inhibition of disruptor of telomeric silencing 1-like alleviated renal ischemia and reperfusion injury-induced fibrosis by blocking PI3K/AKT-mediated oxidative stress. *Drug Des Devel Ther*. 2019;13:4375–4387. doi:10.2147/DDDT.S224909
33. Zhuang X, Tong H, Ding Y, et al. Long noncoding RNA ABHD11-AS1 functions as a competing endogenous RNA to regulate papillary thyroid cancer progression by miR-199a-5p/SLC1A5 axis. *Cell Death Dis*. 2019;10(8):620. doi:10.1038/s41419-019-1850-4
34. Cao Q, Deji QZ, Liu YJ, et al. The role of mechanical stretch and TGF-beta2 in epithelial-mesenchymal transition of retinal pigment epithelial cells. *Int J Ophthalmol*. 2019;12(12):1832–1838. doi:10.18240/ijo.2019.12.03
35. Nam MH, Smith AJO, Pantcheva MB, et al. Aspirin inhibits TGFbeta2-induced epithelial to mesenchymal transition of lens epithelial cells: selective acetylation of K56 and K122 in histone H3. *Biochem J*. 2020;477(1):75–97. doi:10.1042/BCJ20190540
36. Geng XQ, Ma A, He JZ, et al. Ganoderic acid hinders renal fibrosis via suppressing the TGF-beta/Smad and MAPK signaling pathways. *Acta Pharmacol Sin*. 2019.
37. Li SR, Tan ZX, Chen YH, et al. Vitamin D deficiency exacerbates bleomycin-induced pulmonary fibrosis partially through aggravating TGF-beta/Smad2/3-mediated epithelial-mesenchymal transition. *Respir Res*. 2019;20(1):266. doi:10.1186/s12931-019-1232-6
38. Zhang Q, Chang X, Wang H, et al. TGF-beta1 mediated Smad signaling pathway and EMT in hepatic fibrosis induced by Nano NiO in vivo and in vitro. *Environ Toxicol*. 2019.
39. Zhou Y, Huang F, Liu Y, et al. TGF-beta1 relieves epithelial-mesenchymal transition reduction in hypospadias induced by DEHP in rats. *Pediatr Res*. 2019.
40. Hong W, Zhang G, Lu H, et al. Epithelial and interstitial Notch1 activity contributes to the myofibroblastic phenotype and fibrosis. *Cell Commun Signal*. 2019;17(1):145. doi:10.1186/s12964-019-0455-y
41. Chleil E, Pethe A, Pfeifer D, Kriegelstein K, Roussa E. TGF-beta signaling regulates SLC8A3 expression and prevents oxidative stress in developing midbrain dopaminergic and dorsal raphe serotonergic neurons. *Int J Mol Sci*. 2020;21(8):2735. doi:10.3390/ijms21082735
42. Sharma S, Goswami R, Rahaman SO. The TRPV4-TAZ mechanotransduction signaling axis in matrix stiffness- and TGFbeta1-induced epithelial-mesenchymal transition. *Cell Mol Bioeng*. 2019;12(2):139–152. doi:10.1007/s12195-018-00565-w
43. Liu F, Zhang J, Qian J, Wu G, Ma Z. Emodin alleviates CCl4-induced liver fibrosis by suppressing epithelial-mesenchymal transition and transforming growth factorbeta1 in rats. *Mol Med Rep*. 2018;18(3):3262–3270. doi:10.3892/mmr.2018.9324
44. Wang X, Niu C, Zhang X, Dong M. Emodin suppresses activation of hepatic stellate cells through p38 mitogen-activated protein kinase and Smad signaling pathways in vitro. *Phytother Res*. 2018;32(12):2436–2446. doi:10.1002/ptr.6182
45. Sisto M, Lorusso L, Ingravalle G, Ribatti D, Lisi S. TGFbeta1-Smad canonical and -Erk noncanonical pathways participate in interleukin-17-induced epithelial-mesenchymal transition in Sjogren's syndrome. *Lab Invest*. 2020;100(6):824–836. doi:10.1038/s41374-020-0373-z

Drug Design, Development and Therapy

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>

Dovepress

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.