Antimicrobial Resistance and Genotyping of Bacteria Isolated from Urinary Tract Infection in Children in an Iranian Referral Hospital

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Setareh Mamishi^{1,2}
Zohreh Shalchi³
Shima Mahmoudi¹
Reihaneh Hosseinpour Sadeghi¹
Mohammad Taghi Haghi Ashtiani⁴
Babak Pourakbari¹

¹Pediatric Infectious Diseases Research Center, Tehran University of Medical Sciences, Tehran, Iran; ²Department of Infectious Diseases, Pediatrics Center of Excellence, Children's Medical Center, Tehran University of Medical Sciences, Tehran, Iran; ³Department of Pediatrics, Hamadan University of Medical Sciences, Hamadan, Iran; ⁴Department of Pathology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Introduction: Urinary tract infection (UTI) is one of the most common bacterial infections in childhood, and the increasing rate of antibiotic resistance to the commonly prescribed antimicrobial agents against it has become a major concern. The aim of this study was to determine the antibiotic resistance and genotyping of bacteria isolated from urine cultures in patients referred to the Children's Medical Center, Tehran, Iran.

Methods: During the 1-year period, antimicrobial susceptibility profiles of strains isolated from patients with UTI were determined. Typing of the isolates causing nosocomial infections was performed by random amplified polymorphic DNA (RAPD) analysis, and the results were analyzed by Gelcompar II software.

Results: In this study, 203 children (130 girls and 73 boys) were included. The patients' age ranged from 1 day to 16 years (IQR average=4 months to 4 years). The most frequent isolated organisms were *Escherichia coli* (118 isolates, 58%), followed by *Klebsiella pneumoniae* (30 isolates, 15%). Sixty-two strains (18 strains of *E. coli*, 13 strains of *K. pneumoniae*, 11 strains of *Enterococcus faecium*, and five strains of *Burkholderia cepacia* complex) had criteria of nosocomial infection. A high resistance rate to trimethoprim-sulfamethoxazole (69%) and cefotaxime (60%) was reported in *E. coli* and *K. pneumoniae* strains, respectively. *Pseudomonas aeruginosa* strains showed high sensitivity to amikacin (100%). All *E. faecium* strains were susceptible to trimethoprim-sulfamethoxazole (100%), and 23% of the strains were resistant to vancomycin. The analysis of RAPD-typing revealed the presence of three clusters in *E. coli*, two clusters in *E. faecium*, and one clone in *K. pneumoniae*. Besides, four out of five isolates of *B. cepacia* complex had more than 90% genetic similarity.

Conclusion: The most frequent isolated pathogen was *E. coli*, and an increasing rate of antibiotic resistance to the commonly prescribed antimicrobial agents such as trimethoprim/sulfamethoxazole and cephalosporins was observed. Moreover, the results of this study showed the presence of clones with $\geq 80\%$ similarity in *E. coli*, *K. pneumoniae*, *E. faecium*, and *B. cepacia* complex isolates; therefore, the transmission of nosocomial infections from one patient to another or one ward to another is probable.

Keywords: urinary tract infection, antibiotic resistance pattern, genotyping

Correspondence: Babak Pourakbari
Pediatric Infectious Diseases Research
Center, Children Medical Center
Hospital, School of Medicine, Tehran
University of Medical Sciences, No. 62,
Gharib St., Keshavarz Blvd, Tehran, Iran
Tel/Fax +98 021 6642 8996
Email pourakbari@tums.ac.ir

Introduction

Urinary tract infections (UTIs) are one of the most common bacterial infections in children aged <2 years¹ and considered as a main cause of hospitalization in childhood.² The incidence rate of UTIs is up to 7% among febrile infants with no other recognizable causes.³ *Escherichia coli* has been known as the most common isolated organism in pediatric patients with UTI with a prevalence of 80–90%,^{4,5}

followed by others such as *Enterococcus species* (spp.), *Enterobacter spp.*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Proteus mirabilis*, and *Staphylococcus spp.*^{5,6} In the preantibiotic era, UTI caused a mortality rate as high as 20%.⁷ The early treatment of UTI decreases the rate of morbidity;² however, the increasing rate of antibiotic resistance to the commonly prescribed antimicrobial agents such as trimethoprim/sulfamethoxazole, cephalosporins, and fluoroquinolones⁸ has become a major concern.⁹ Hence, it is crucial to know the frequency of uropathogens and antimicrobial resistance in pediatrics.¹⁰ Prompt and efficient antimicrobial therapies for UTIs are vital for an appropriate outcome and prevention of long-term morbidity associated with renal scarring such as hypertension, development of chronic kidney disease, and ultimately the need of renal transplantation.¹¹ The aim of this study was to determine the antibiotic resistance pattern and genotyping of strains isolated from pediatric patients with UTI.

Materials and Methods

This study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of Tehran University of Medical Sciences, Tehran, Iran (IR.TUMS.CHMCREC.1397.47).

This study was performed over a period of 1 year, from March 2018 to February 2019 in the Children's Medical Center, an Iranian referral pediatrics hospital in Tehran, Iran. The inclusion criteria were patients having a history of fever >38°C or symptoms of UTIs including dysuria, frequency, urgency, hesitancy, small volume voids, or lower abdominal pain.⁹ UTI was defined in infants and children as growth of a single pathogen of >10⁵ colony forming units/mL by properly collected urine specimen (suprapubic aspiration, transurethral catheterization, or mid-stream urine) in febrile children or the ones who had other urinary symptoms.¹² Bacteria identification was performed using Gram staining (for bacterial morphology) and differential biochemical tests. Identification of organisms was carried out using conventional biochemical methods such as catalase, oxidase, sugar fermentation, methyl red/Voges-Proskauer (MR/VP) test, growth on Kligler's Iron Agar (KIA), growth on Bile-Esculin Agar and 6.5% NaCl media, and motility test.¹³

All isolated bacteria were checked for their susceptibility towards different antibiotics by disk diffusion method based on Clinical Laboratory Standard Institute (CLSI) guidelines, 2019.¹⁴ The list of antibiotic discs

used for gram-negative bacteria were: nitrofurantoin (300 µg), trimethoprim-sulfamethoxazole (1.25/23.75 µg), cefepime (30 µg), cefotaxime (30 µg), imipenem (10 µg), amikacin (30 µg), piperacillin/tazobactam (100–10 g), ceftazidime (30 µg), and gentamicin (10 µg). Disks used for Gram-positive bacteria were gentamicin (10 µg), penicillin (30 µg), vancomycin (30 µg), nitrofurantoin (300 µg), Trimethoprim-sulfamethoxazole (1.25/23.75 µg), piperacillin/tazobactam (100–10 g), imipenem (10 µg), meticillin (30 µg), and ampicillin (10 µg). High-level resistance to gentamicin was also determined using a 120 µg gentamicin disk. The minimal inhibitory concentration of vancomycin was determined by E-test methods. All disks were prepared by the Mast Group Ltd., Merseyside, UK. *E. coli* ATCC 25922, *Pseudomonas aeruginosa* ATCC 27853, and *Staphylococcus aureus* ATCC 25923 were used as control strains. All recovered strains which had caused nosocomial infections were typed by Random Amplified Polymorphic DNA PCR (RAPD-PCR) in a mixture consisting of 2.5 µL of the PCR buffer (10-times concentrated), 0.75 µL of MgCl₂ (final concentration 250 µM), 0.6 µL of dNTPs (Fermentas, Vilnius, Lithuania, final concentration 3 mM), 1 µL of primer (5'- AGCGGGCCAA-3'), final concentration of 0.4 mM, 0.3 µL (1.5 U) of the DFS Taq DNA polymerase (BIORON, Germany), 1 µL of the boiled DNA (final concentration 2 ng/µL), and DNase-, RNase-free deionised water (Biomedicals) to a final volume of 25 µL.¹⁵ Amplification was carried out in a thermal cycler (Applied Biosystems, USA) as follows: the primary denaturation for 5 minutes at 95°C; followed by 30 cycles of 60 seconds for 94°C, 60 seconds at annealing temperature, and 5 minutes at 72°C; and the final extension phase for 15 minutes at 72°C. The analysis of the amplified products was performed in 1% agarose (Sigma), and DNA bands were visualized by staining with gel red (Biotium), analyzed under UV light, and photographed using the GEL Doc 2000 documentation system (Bio-Rad). Comparison of RAPD-PCR banding patterns was performed using Gelcompar II, version 6.5 (Applied Maths, Sint-MatensLatem, Belgium), and the unweighed pair group method with arithmetic mean (UPGMA) and dendrograms were constructed.

Statistical Analysis

Descriptive statistics were reported by total numbers, percentages, means, and medians as appropriate.

Table 1 The Distribution of Antimicrobial Resistance Pattern in Gram Negative Bacteria Isolated from UTIs

Gram Negative Bacteria	Total	Trimethoprim-sulfamethoxazole	Nitrofurantoin	Piperacillin/Tazobactam	Amikacin	Gentamycin	Cefepime	Cefotaxime	Imipenem	Ceftazidime
	N	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)
<i>E. coli</i>	118	81 (69)	9 (8)	20 (17)	3 (2/5)	21 (18)	67 (57)	74 (63)	-	-
<i>K. pneumoniae</i>	30	11 (37)	15 (50)	14 (47)	10 (33)	7 (23)	15 (50)	18 (60)	-	-
<i>P. aeruginosa</i>	13	-	-	1 (8)	0	1 (8)	1 (8)	-	2 (15)	1 (8)
<i>E. aerogenes</i>	3	0	3 (100)	-	0	1 (33)	1 (33)	-	-	-
<i>B. cepacia</i> complex	6	0	-	2 (33)	6 (100)	6 (100)	4 (67)	-	6 (100)	0
<i>K. oxytoca</i>	3	1 (33)	0	1 (33)	1 (33)	1 (33)	1 (33)	-	-	-

Results

A total of 203 children with positive urine cultures, collected in the tertiary referral Children's Medical Hospital in Tehran, Iran, were included in our study. Among these patients, 130 (64%) subjects were female. Median age was 1 year (IQR=4 months to 4 years), and the age range was 1 day to 16 years.

E. coli (118 samples, 58%) was the most prevalent microorganism isolated from urine cultures, followed by *K. pneumoniae* (30 samples, 15%) and *P. aeruginosa* (13 samples, 6%). Among the isolated gram positive microorganisms, the most frequent isolated bacteria were *E. faecium* (13 samples, 6%) and *S. epidermidis* (5 samples, 2.5%), respectively. The other strains were identified as: *B. cepacia* complex, *E. aerogenes*, *K. oxytoca*, *E. cloacae* complex, *P. mirabilis*, *Morganella morganii*, *Acinetobacter baumannii-baumannii*, *P. vulgaris*, *Enterobacter spp.*, *E. faecalis*, *S. haemolyticus*, and *Alpha Hemolytic Streptococcus* (viridans group).

A high percentage of *E. coli* strains were isolated from female children (81%), while 70% of *K. pneumoniae* strains were isolated from male children. Most of the bacteria were isolated from the emergency ward (N=51, 25%), followed by the urology (N=33, 16%) and nephrology (N=32, 16%) wards.

The results of the antibiotic resistance pattern to the mentioned antibiotic disks for each microorganism are shown in Tables 1 and 2. Sixty percent of *K. pneumoniae* and 69% of *E. coli* strains were resistant to cefotaxime and trimethoprim-sulfamethoxazole, respectively. Twenty-three percent of *E. faecium* strains (n=3) showed resistance to vancomycin, which was lower than our previous reports.^{15,16}

Out of 203 strains isolated, 62 strains had nosocomial infection criteria which were: 18 *E. coli*, 13 *K. pneumoniae*, 11 *E. faecium*, and five *B. cepacia* complex. Molecular typing and phylogenetic analysis of these 62 isolates causing hospital-acquired infection revealed that *E. coli* strains were divided into three clusters with 70% genetic similarity (Figure 1A). Also, the presence of all *K. pneumoniae* strains in one cluster with high genetic similarity was reported (Figure 1B), most of which were isolated from the Nephrology ward and Neonatal Intensive Care Unit (NICU). Among five *B. cepacia* complex isolates, four strains had 90% genetic similarity (Figure 1C).

The dendrogram, based on analysis of *E. faecium* strains, indicated the presence of two clusters with more

Table 2 The Distribution of Antimicrobial Resistance Pattern in Gram Positive Bacteria Isolated from UTIs

Gram Positive Bacteria	Total	Trimethoprim-Sulfamethoxazole	Nitrofurantoin	Piperacillin/Tazobactam	Ampicillin	Gentamycin	Vancomycin	Penicillin	Imipenem	Meticillin
	N	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)	N (%)
<i>E. faecium</i>	13	–	3 (23)	5 of 12 (42)	5 of 12 (42)	3 of 3 (100)	3 (23)	8 (61/5)	3 (23)	–
<i>S. epidermidis</i>	5	2 (40)	0	–	–	–	0	4 (80)	–	4 (80)

than 70% genetic similarity (Figure 1D). Two strains were completely similar to each other, isolated from different wards.

Discussion

In this study, we report the antimicrobial resistance and genotyping of bacteria isolated from UTIs in children in an Iranian referral hospital. The current concern of UTIs is the prevalence of multi drug resistance and also a main problem in the treatment of UTIs.^{17,18}

In our study among 203 positive cultures, the most frequent pathogen was *E. coli* (58%), which is generally the leading uropathogen and was isolated from 56.6–84.6% of Iranian children with febrile UTI.^{19,20} Similarly, it is reported as frequent as 62% in Raman et al's²¹ study. However, in Heidary et al study *P. aeruginosa* was reported as the most common pathogen (49.65%).

In this study, *E. coli* isolates revealed a high resistance rate to trimethoprim-sulfamethoxazole (69%) which has increased compared with the percentage of this antibiotic in previous studies.²¹ According to a report by Bryce et al,²² pooled prevalence of resistance to trimethoprim-sulfamethoxazole was as high as 30% in OECD (Organization for Economic Cooperation and Development) countries and resistance to trimethoprim-sulfamethoxazole was more than twice as high in non-OECD compared with OECD countries. On the other hand, the highest percentages of susceptibility of *E. coli* in our current study were reported for amikacin (97.5%) and nitrofurantoin (91%), which was similar to our previous study.²³

A high *P. aeruginosa* susceptibility rate to amikacin (100%), ceftazidime, cefepime, gentamycin, piperacillin/tazobactam (92%), and imipenem (85%) was reported in the current study which was higher than the frequency reported by Zhang et al.²⁴ Genotyping of 18 *E. coli* strains causing nosocomial infections depicted the presence of three clusters with more than 70% genetic similarity, 12 of 18 strains were isolated from urology or surgical units. These results suggested the possibility of dissemination of these clones, especially in these two units.

More than half of the *K. pneumoniae* isolates were resistant to cefepime and cefotaxime. In addition, all *K. pneumoniae* isolates belonged to one cluster and most of the strains were isolated from the nephrology ward or NICU, suggesting a nosocomial infection between these two units. Similar to the current result, our previous study reported the presence of 93% of *K. pneumoniae* isolates causing nosocomial infections in one cluster.²⁵

Figure 1 Genotyping of *E. coli* (A), *K. pneumoniae* (B), *B. cepacia* complex (C), and *E. faecium* strains (D) that met hospital-acquired infection criteria using RAPD-PCR.

Treatment of infections caused by *B. cepacia* complex, opportunistic nosocomial pathogens, relies on ceftazidime and other extended-spectrum cephalosporins, as intrinsic resistance prevents the action of many other classes of antimicrobials.²⁶ In our study, all *B. cepacia* complex isolates were susceptible to ceftazidime and 100% resistant to amikacin and gentamycin. In a study conducted in Taiwan, 24% of *B. cepacia* complex were resistant to ceftazidime and, similar to ours, had a high susceptibility

to trimethoprim-sulfamethoxazole.²⁷ The genotyping of *B. cepacia* complex depicted that 80% of isolates had high genetic similarity, isolated from urology, nephrology units, and NICU. Therefore, the spread of one strain between these three wards is probable.

The analysis of RAPD-typing of *E. faecium* strains showed the presence of two clusters with more than 70% genetic similarity and most of the strains were isolated from urology and nephrology units.

The RAPD-PCR is a simple, rapid, easy, inexpensive, and reproducible method that has been widely used in bacterial epidemiology and finding of bacterial genetic similarity that is crucial for cross-infection assessment.^{15,25,28,31} Cross-sectional transmission, especially from the urology unit to other wards, may lead to strains transmission causing nosocomial infection through the hospital. Therefore, much attention should be paid to the basic methods of preventing infection (standard precautions).^{32,33} The length of hospitalization and the use of antibiotics in patients, particularly during long-term hospitalization and unnecessary transfer of patients between different units might have impacts on the reduction of antibiotic resistance and the occurrence of nosocomial infection.

There are several limitations for this study. First, this is a single-center retrospective study using existing data collected over years with small samples size. It is possible that we could have missed cases such as those who were treated without any culture testing, or received antibiotics before urine collection for culture. Second, there was no access to some clinical information of the patients such as the results of treatment, so determining the rate of the outcome was impossible.

In conclusion, the most frequent isolated pathogen was *E. coli* and the increasing rate of antibiotic resistance to the commonly prescribed antimicrobial agents such as trimethoprim/sulfamethoxazole and cephalosporins was observed. Moreover, the results of this study showed the presence of clones with $\geq 80\%$ similarity in *E. coli*, *K. pneumoniae*, *E. faecium*, and *B. cepacia* complex isolates; therefore, the transmission of nosocomial infections from one patient to another or one ward to another is probable.

Acknowledgments

This study was taken from Dr. Zohreh Shalchi's postgraduate thesis and was supported by a grant (number: 97-02-88-39126) from Tehran University of Medical Sciences to Dr. Babak Pourakbari.

Disclosure

The authors report no conflicts of interest in this work.

References

- Okarska-Napierała M, Wasilewska A, Kuchar E. Urinary tract infection in children: diagnosis, treatment, imaging—Comparison of current guidelines. *J Pediatr Urol.* 2017;13(6):567–573. doi:10.1016/j.jpuro.2017.07.018
- Kutasy B, Coyle D, Fossum M. Urinary tract infection in children: management in the era of antibiotic resistance—a pediatric urologist's view. *Eur Urol Focus.* 2017;3(2–3):207–211. doi:10.1016/j.euf.2017.09.013
- Wang J, He L, Sha J, et al. Etiology and antimicrobial resistance patterns in pediatric urinary tract infection. *Pediatrics Int.* 2018;60(5):418–422. doi:10.1111/ped.13526
- Awean GZ, Salameh K, Elmohamed H, Alshmayt H, Omer MRB. Prevalence of ESBL urinary tract infection in children. *J Adv Pediatr Child Health.* 2019;2(1):004–7. doi:10.29328/journal.japch.1001004
- Pouladfar G, Basiratnia M, Anvarinejad M, Abbasi P, Amirmoezi F, Zare S. The antibiotic susceptibility patterns of uropathogens among children with urinary tract infection in Shiraz. *Medicine.* 2017;96(37):37. doi:10.1097/MD.00000000000007834
- Dahle KW, Korgenski EK, Hersh AL, Srivastava R, Gesteland PH. Clinical value of an ambulatory-based antibiogram for uropathogens in children. *J Pediatric Infect Dis Soc.* 2012;1(4):333–336. doi:10.1093/jpids/pis055
- Zorc JJ, Kiddoo DA, Shaw KN. Diagnosis and management of pediatric urinary tract infections. *Clin Microbiol Rev.* 2005;18(2):417–422. doi:10.1128/CMR.18.2.417-422.2005
- Amraei S, Hashemi Karouei SM, Babakhani S, Kazemi MJ. Serotyping and antibiotic susceptibility pattern of common bacterial uropathogens in urinary tract infections in Koohdasht, Lorestan province. *Int J Enteric Pathog.* 2016;4(2):1–5.
- Nor NSM, Abu NA, Rashid MA, Ismail M, Razak R. Bacterial pathogens and antibiotic resistance patterns in children with urinary tract infection in a Malaysian tertiary hospital. *Med J Malays.* 2015;70(3):153–157.
- Doyev R, Ben-Shalom E, Megged O. The predictive utility of prior positive urine culture in children with recurrent urinary tract infections. *Eur J Pediatr.* 2020;179(3):415–421. doi:10.1007/s00431-019-03522-5
- Koçak M, Büyükkaragöz B, Çelebi Tayfur A, et al. Causative pathogens and antibiotic resistance in children hospitalized for urinary tract infection. *Pediatrics Int.* 2016;58(6):467–471. doi:10.1111/ped.12842
- Prais D, Straussberg R, Avitzur Y, Nussinovitch M, Harel L, Amir J. Bacterial susceptibility to oral antibiotics in community acquired urinary tract infection. *Arch Dis Child.* 2003;88(3):215–218. doi:10.1136/adc.88.3.215
- Mahon CR, Lehman DC, Manuselis G. *Textbook of Diagnostic Microbiology* 2007. 2008. 3rd ed. St. Louis, Mo: Saunders Elsevier.
- <melvin p. performance standards for antimicrobial susceptibility testing. 2019.pdf>
- Mamishi S, Mohammadian M, Pourakbari B, et al. Antibiotic resistance and genotyping of gram-positive bacteria causing hospital-acquired infection in patients referring to children's medical center. *Infect Drug Resist.* 2019;12:3719–3726. doi:10.2147/IDR.S220522
- Pourakbari B, Mahmoudi S, Moradzadeh M, et al. Antimicrobial Resistance Patterns of the Gram-positive Bacteria Isolated from Children with Bloodstream Infection in an Iranian Referral Hospital: A 6-year Study. *Infect Disord Drug Targets.* 2018;18(2):136–144. doi:10.2174/1871526517666170821164343
- Niranjan V, Malini A, Bahaman AR, Amran F, Khalilpour A. Antimicrobial resistance pattern in Escherichia coli causing urinary tract infection among inpatients. *Indian J Med Res.* 2014;139(6):945.
- Marialouis XA, Santhanam A. Antibiotic resistance, RAPD-PCR typing of multiple drug resistant strains of Escherichia coli from urinary tract infection (UTI). *J Clin Diagn Res.* 2016;10(3):DC05.
- Afsharpaiman S, Bairaghdar F, Torkaman M, et al. Bacterial pathogens and resistance patterns in children with community-acquired urinary tract infection: a cross sectional study. *J Comprehens Pediatr.* 2012;3(1):16–20. doi:10.5812/jcp.7078
- Mirsoleymani SR, Salimi M, Shareghi Brojeni M, Ranjbar M, Mehtarpour M. Bacterial pathogens and antimicrobial resistance patterns in pediatric urinary tract infections: a four-year surveillance study (2009–2012). *Int J Pediatr.* 2014;2014.

21. Raman G, McMullan B, Taylor P, Mallitt K-A, Kennedy SE, Multiresistant *E. coli* urine infections in children: a case-control study. *Arch Dis Child*. 2018;103(4):336–340. doi:10.1136/archdis-child-2017-312831
22. Bryce A, Hay AD, Lane IF, Thornton HV, Wootton M, Costelloe C. Global prevalence of antibiotic resistance in paediatric urinary tract infections caused by *Escherichia coli* and association with routine use of antibiotics in primary care: systematic review and meta-analysis. *BMJ*. 2016;352:i939. doi:10.1136/bmj.i939
23. Pourakbari B, Ferdosian F, Mahmoudi S, et al. Increase resistant rates and ESBL production between *E. coli* isolates causing urinary tract infection in young patients from Iran. *Br J Microbiol*. 2012;43(2):766–769. doi:10.1590/S1517-83822012000200041
24. Zhang X, Niu S, Zhang L. Antimicrobial susceptibilities and clinical characterization of *Pseudomonas aeruginosa* isolates from urinary tract infections. *Urol Int*. 2014;93(4):464–469. doi:10.1159/000358493
25. Mahmoudi S, Pourakbari B, Rahbarimanesh A, Abdosalehi MR, Ghadiri K, Mamishi S. An outbreak of ESBL-producing *Klebsiella pneumoniae* in an Iranian referral hospital: epidemiology and molecular typing. *Infect Dis Drug Targets*. 2019;19(1):46–54.
26. Rhodes KA, Schweizer HP. Antibiotic resistance in *Burkholderia* species. *Drug Resist Updates*. 2016;28:82–90. doi:10.1016/j.drug.2016.07.003
27. Tseng S-P, Tsai W-C, Liang C-Y, et al. The contribution of antibiotic resistance mechanisms in clinical *Burkholderia cepacia* complex isolates: an emphasis on efflux pump activity. *PLoS One*. 2014;9(8):8. doi:10.1371/journal.pone.0104986
28. Baker JC, Crumley RE, Eckdahl TT. Random amplified polymorphic DNA PCR in the microbiology teaching laboratory: identification of bacterial unknowns. *Biochem Mol Biol Educ*. 2002;30(6):394–397. doi:10.1002/bmb.2002.494030060135
29. Pourakbari B, Mahmoudi S, Aghdam MK, et al. Clonal spread of vancomycin resistance *Enterococcus faecalis* in an Iranian referral pediatrics center. *J Prev Med Hyg*. 2013;54(2):87–89.
30. Pourakbari B, Mahmoudi S, Habibi R, et al. An increasing threat in an Iranian referral children's hospital: multidrug-resistant *acinetobacter baumannii*. *Infect Disord Drug Targets*. 2018;18(2):129–135. doi:10.2174/1871526517666170113144541
31. Hassuna NA, Darwish MK, Sayed M, Ibrahim RA. Molecular epidemiology and mechanisms of high-level resistance to meropenem and imipenem in *pseudomonas aeruginosa*. *Infect Drug Resist*. 2020;13:285. doi:10.2147/IDR.S233808
32. Mamishi S, Pourakbari B, Teymuri M, Babamahmoodi A, Mahmoudi S. Management of hospital infection control in Iran: a need for implementation of multidisciplinary approach. *Osong Public Health Res Perspect*. 2014;5(4):179–186. doi:10.1016/j.phrp.2014.06.001
33. Movahedi Z, Pourakbari B, Mahmoudi S, et al. *Pseudomonas aeruginosa* infection among cystic fibrosis and ICU patients in the referral children medical hospital in Tehran, Iran. *J Prev Med Hyg*. 2013;54(1):24.

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

Dovepress