Magnitude of Anemia and Its Associated Factors Among Pregnant Women Attending Antenatal Care in Southern Ethiopia: A Cross-Sectional Study

This article was published in the following Dove Press journal:
Journal of Blood Medicine

Dirshaye Argaw
Robel Hussen Kabthymmer
Mahlet Birhane

Human Nutrition Unit, School of Public Health, Dilla University, Dilla, Ethiopia

Purpose: The aim of this study was to assess the magnitude of anemia and its associated factors among pregnant women attending antenatal care (ANC) at Dilla University referral Hospital, South Ethiopia.

Patients and Methods: An institution-based cross-sectional study was conducted from January to February 2019, among 373 pregnant women who attended antenatal care at Dilla University referral hospital. Socio-demographic factors, nutritional, medical and obstetric information of the study participants were collected using a structured questionnaire. Hemoglobin was measured using a hemacue machine, and fecal specimens were examined to detect intestinal parasites. Bivariate and multiple variable binary logistic regressions were used to identify predictors of anemia. A p-value less than 0.05 was used to declare statistical significance.

Results: Overall prevalence of anemia was 28.7%, of which 19.6% had mild anemia. Decreased odds of anemia were found in women with good nutritional status (MUAC ≥ 24 cm) (AOR= 0.07 95% CI: 0.03–0.1), iron supplementation (AOR=0.06 95% CI: 0.02–0.15) and birth spacing ≥ 2 yrs (AOR=0.03 95% CI: 0.009–0.45). However, increased odds of anemia were seen in pregnant women with intestinal parasites (AOR=6.11 95% CI 7.70–37.0).

Conclusion: The magnitude of anemia among pregnant women was found to be a moderate public health problem. Iron supplementation, good nutritional status (MUAC > 24 cm), and birth spacing reduce the odds of anemia. But having intestinal parasites was found to increase the likelihood of anemia during pregnancy. Counseling on birth spacing, strengthening iron supplementation, and intestinal parasite management during pregnancy should be given due emphasis.

Keywords: pregnancy, anemia, Dilla, Ethiopia, prevalence

Introduction

Anemia is a condition which exists when there is less than the normal hemoglobin (Hb) level in the body, which decreases oxygen-carrying capacity of red blood cells to tissues. The Hb levels for each class of anemia during pregnancy are 10.0–10.9 g/dl (mild), 7–9.9 g/dl (moderate) and <7 g/dl (severe).¹

Anemia affects more than 2 billion people globally, accounting for nearly 30% of the world's population. Anemia is the most common public health problem

Correspondence: Robel Hussen Kabthymmer Human Nutrition Unit, School of Public Health, Dilla University, Dilla 419, Ethiopia
Tel +251913575702
Email Robelhussen@rocketmail.com

particularly in developing countries, occurring at all stages of the life cycle. The prevalence of anemia in developing and developed countries is estimated to be 43% and 9%, respectively.² Magnitude of anemia exceeds 50% in most sub-Saharan African countries.^{3,4} As reported in 2005 Ethiopian demographic health survey (EDHS), anemia has severe public health significance in Ethiopia. In 2016, the national prevalence among pregnant women was 29% and the figure did not decline over the previous decade.^{5,6}

Anemia during pregnancy remains one of the top public health concerns in developing countries due to various socio cultural problems like illiteracy, poverty, lack of awareness, cultural and religious taboos, poor dietary habits, and high prevalence of parasitic infestation. Current estimates from the World Health Organization reports indicate that 41.8% of pregnant women are anemic, with the highest burden of share found in Africa (61.3%) and South East Asia (52.5%).^{2,7}

Pregnancy makes women more vulnerable to anemia. According to Ethiopian demographic health survey (EDHS 2016), 22% of Ethiopian women aged 15–49 years are anemic, of which 17% have mild anemia, 5% have moderate anemia, and 1% have severe anemia. Proportions of anemia during pregnancy are higher (22%) than in women who are breastfeeding (19%) and women who are neither pregnant nor breastfeeding (15%).⁵

The main risk factors of anemia include a low dietary intake of iron, poor absorption of iron from diets, and period of life where iron requirements become high (ie, growth and pregnancy).⁴ Among other causes of anemia, heavy blood loss, acute and chronic infections, cancer, tuberculosis, and HIV can also lower blood Hb concentrations. The presence of other micronutrient deficiencies like vitamin A, vitamin B12, folic acid, riboflavin and copper increases the risk of anemia.^{4,7,8}

There have been different measures to control anemia like iron/folic acid supplementation, food fortification, dietary diversification, deworming and, malaria treatment and control.² Despite these efforts, the burden of anemia remains alarmingly high. Current strategies to prevent and control anemia have to be re-evaluated to ensure that the various factors contributing to anemia have been identified and addressed properly in an integrated manner.

The availability of local information on the magnitude and related risk factors has a major role in the management and control of anemia, especially during pregnancy. Since there is inadequate scientific in the study area, the aim of

this study was to assess the magnitude and its associated factors of anemia among pregnant women attending antenatal clinic in DURH, Southern Ethiopia.

Methods

Study Area

The study was conducted in Dilla University referral hospital (DURH), which is located in Dilla city of southern Ethiopia. The hospital provides medical service for 49,849 and 3,226 clients in outpatient and inpatient settings respectively. Dilla is 365 KM away from the capital Addis Ababa. At the time of the study the total population of Dilla city was 102,624, among this 50,286 were males and 52,338 were females. In the city there is one referral and teaching hospital, and three public health centers. DURH is a referral hospital for the catchment population of more than 2,119,100. It is located in the southern part of the country along the main Addis Ababa – Nairobi international road. The ANC is run by the maternity department of the hospital. About 30 women attend the ANC clinic daily. In Ethiopia, “package of services” provided within the ANC include iron folic acid supplementation, deworming, HIV testing, nutrition counseling and intermittent preventive treatment for malaria.

Study Design and Period

Institutional-based cross-sectional study was conducted from November 2017 to January 2018.

Source Population

Source population for this study was all pregnant women attending antenatal care (ANC) in Dilla University referral hospital.

Study Population

All selected pregnant women who were attending antenatal care (ANC) in Dilla University referral hospital at the time of data collection.

Eligibility Criteria

Pregnant women who attended ANC in DURH were included in the study, while pregnant women with recent blood loss, severe dehydration were excluded.

Sample Size Determination

The sample size required for this study was determined using a single population proportion formula with the

assumption of confidence level 95%, power of 80%, margin of error 5% and prevalence of anemia 32.8% from previous institution-based cross-sectional study conducted in Arbaminch health care settings.⁹ In addition a 10% non-response rate was added.

$$n = \frac{Z^2 P (1 - p)}{d^2}$$

where,

n=sample size

Z= 95% confidence interval (1.96)

P= prevalence = 32.8%

d = margin of error = 0.05

$$n = \frac{(1.96)^2 0.328(1 - 0.328)}{(0.05)^2}$$

n = 339+10% non-response=373

The sample size was also calculated considering the significant associated factors using epi info software but 373 was found to be the largest sample size.

Sampling Procedure

First we reviewed the records of monthly flow of pregnant women for ANC utilization in DURH ANC clinic and the previous 3 months total pregnant women who visited DURH ANC were 1113. Using this data we calculated a sampling interval of 3 (K=1113/373). Hence, 373 pregnant women visiting ANC during November 2018 to January 2019 were selected by using systematic random sampling.

Operational Definition of Variables

- Anemia: defined as a condition of pregnant women in which the hemoglobin concentration measured is less than 11 g/dl. And anemia was classified as:

mild = hemoglobin concentration of 10–10.9 g/dl

moderate= 7–9.9 g/dl

severe= less than 7.0 g/dl^{1,2}

- Dietary diversity score: the total score calculated for each individual from the selected 9 food groups consumed in the last 24 hours without place and time restriction. From those nine food groups, those who consumed ≥ 6 food groups were labeled as “highly diversified food”, 4 and 5 food groups as “medium dietary diversity” and ≤ 3 food groups as “undiversified”.¹⁰
- Intestinal parasite: parasites that cause blood loss in the body were assessed by stool examination. Some

of these parasites are hook worms, schistosomiasis, *Trichuris trichiura*, *Giardia lamblia*, *Entamoeba histolytica* and *Ascaris lumbricoides*.¹¹

- Nutritional status: mid upper arm circumference (MUAC) was measured to the nearest 0.1 cm using flexible and non-stretchable measuring tapes following the standard procedures. Mothers were considered under-nourished when their MUAC value was less than 24.0 cm and those with MUAC ≥ 24 were considered well nourished.¹²

Data Collection Tools and Procedures

A structured and pretested interviewer administered questionnaire was used to obtain socio-demographic information, dietary habits, and obstetric history. To obtain dietary habits, a multiple pass 24 hour recall method was used to assess the intake of various food items for the past 24 hours. The questionnaire was developed in English and then translated into Amharic language for simplicity then back-translated to English language to check its consistency by two different language experts who speak both English and Amharic fluently. The nutritional status of the women was measured by using mid-upper arm circumference (MUAC) measuring tape.

Household Food Insecurity Level

The household food insecurity access scale was adapted from food and nutrition technical assistance (FANTA) Version 3. Each of the questions was asked with a recall period in the last four weeks (30 days). Responses were counted for occurrence, and frequency of occurrence which produced a score for each household. This score is finally classified into four categories: as “food secure”, “mildly food insecure”, “moderately food insecure”, and “severely food insecure”.¹³

Hemoglobin Measurement

Hemoglobin measurement was done from capillary blood by collecting one drop of blood carefully from the middle finger. The finger of the pregnant woman was pricked after cleaning the fingertip with sterile cotton (immersing in 70% alcohol) with a sterile disposable lancet. Calibrated HEMOCUE Hb 301, HEMOCUE AB, ANGELHOLM SWEDEN machines were used to assess the hemoglobin concentration, which was recommended for use in resource-limited settings and the results were expressed in g/dl, then categorized based on criteria of WHO cut off point.¹⁴

Stool Specimen Collection and Examination for Intestinal Parasites

Stool samples were collected for the study after explanation about the procedure and its importance. Stool samples were collected from the women using clean containers and within an hour of collection stool microscopy was done at the laboratory by Bachelor degree holder laboratory technician. Direct microscopy with Formalin-ether concentration technique was employed to identify the parasites.¹⁵

Data Quality

Two day intensive training was given to data collectors regarding study objectives, interviewing techniques, and ethical issues during data collection. A pretest was done among 5% of the total sample size at nearby Yirgacheffe district hospital. Functionality and performance of instruments were cross-checked by using quality control samples, for Hemocue it was checked against CBC machine.

Data Analysis

The collected data were checked for completeness and consistency. Data were entered into EpiData version 3.2, and then exported to SPSS version 20 for cleaning and analysis. Tables and graphs were used to summarize descriptive results

Table 1 Socio-Demographic Characteristics of Pregnant Women Attending Antenatal Care in DURH, Southern Ethiopia, 2019

Variables	Frequency (n)	Percentage (%)
Age (in years)		
<20	46	12.3
20–24	119	31.9
25–29	116	31.1
30–34	55	14.7
≥35	37	9.9
Religion		
Protestant	160	42.9
Orthodox	135	36.2
Muslim	66	17.7
Other	12	3.2
Residence		
Urban	49	13.1
Rural	324	86.9
Marital status		
Married	307	82.3
Single	40	10.7
Divorced	16	4.3
Widowed	10	2.7

of the data. Bivariate and multiple variable binary logistic regression models were used. Those variables with p-value < 0.25 at the bivariate level were included in a multivariate logistic regression model to control potential confounding factors. After adjusting, those variables with a p value < 0.05 were considered as statistically significant factors associated independently with anemia. Crude and adjusted odds ratios with 95% confidence intervals were reported.

Results

Socio-Demographic Characteristics

A total of 373 pregnant women were included in this study making a response rate of 99.9%. Majority, (324 [76.5%]) of study participants resided in urban areas and regarding marital status, 307 (82.3%) were married (Table 1).

Socio Economic Characteristics

(Table 2).

Table 2 Socio-Economic Characteristics of Pregnant Women Attending Antenatal Care in DURH, Southern Ethiopia, 2019

Variables	Frequency (n)	Percentage (%)
Educational status		
Illiterate	77	20.6
Primary education (1–8)	96	25.7
Secondary education (9–12)	117	31.4
Higher education	82	22.3
Husbands' educational status		
Illiterate	36	9.7
Primary education (1–8)	56	15.0
Secondary education (9–12)	85	22.8
Higher education	130	34.9
Employment status		
House-wife	128	34.3
Government employee	83	22.3
Self-employed	117	31.4
Other	45	12.1
Household income		
Less than 500	21	5.6
500–1499	73	19.6
1500–2499	59	15.8
2500–3499	88	23.6
Greater than 3500	132	35.4
Family size		
≤5	333	89.3
> 5	40	10.7

Dietary-Related Characteristics

Nearly three fourths (82.8%) of the respondents consumed meat once per week or less. From the total study participants, 142 (38.1%) respondents drank more than 3 cups of coffee per day. More than half, (229 [61.4%]), of the respondents had a medium dietary diversity score (4–5), and the rest, 100 (26.8%), and 44 (11.8%), had a low and high dietary diversity score respectively (Table 3).

Nutrition-Related Characteristics

Majority, 277 (74.3%), of the participants had MUAC ≥ 24 cm and 271 (72.7%) were currently taking iron with folic acid supplement (Table 4).

Chronic and Acute Illness

From the total study participants almost all (341 [91.4%]) of the respondents had no known chronic illness and only 91 (24.4%) respondents had a recent history of malarial infection.

Table 3 Dietary-Related Characteristics of Pregnant Women Attending Antenatal Care in DURH, Southern Ethiopia, 2019

Variables	Frequency (n)	Percent (%)
Frequency of animal product consumption		
Every day and Every two days	28	7.5
Every week and less	309	82.8
Consumption of coffee		
≤ 3 cups per day	175	46.9
> 3 cups per day	198	53.1
Frequency of consuming green vegetables		
Every day	44	30.6
Every two days	147	39.4
Every week and less	104	27.9
Staple food of the family		
Enset	145	38.9
Maize	71	19
Both enset and maize	8	2.1
Injera	149	39.9
Dietary diversity score (24hr)		
Low	100	26.8
Medium	229	61.4
High	44	11.8

Table 4 Nutrition-Related Characteristics of Pregnant Women Attending Antenatal Care in DURH, Southern Ethiopia, 2019

Variables	Frequency (n)	Percent (%)
Household food security status		
Food secure	149	39.9
Mildly food insecure	122	32.7
Moderately food insecure	94	25.2
Severely food insecure	8	2.1
MUAC		
< 24 cm	94	25.2
≥ 24 cm	279	74.8
Currently taking iron with folic acid		
Yes	277	74.3
No	96	25.7
Nutrition education about nutrition during pregnancy		
Yes	60	16.1
No	313	83.9

Table 5 Reproductive Health History of Pregnant Women Attending Antenatal Care in DURH, South Ethiopia, 2019

Variables	Frequency (n)	Percent (%)
Parity		
Nulliparous	117	31.4
1–2 Para	228	61.1
≥ 3 Para	28	7.5
Trimester		
1st trimester	34	9.1
2nd trimester	179	48.0
3rd trimester	160	42.9
Pregnancy interval (years)		
≤ 2	61	16.4
> 2	196	52.5
History of abortion		
Yes	78	20.9
No	271	72.1
Contraceptive use before this pregnancy		
Yes	186	49.9
No	187	50.1

Reproductive Health Characteristics

Only one third, (117 [31.4%]), of the respondents were nulliparous. From the total study participants, 160 (42.9%) were in 3rd trimester and 179 (48.0%) were in

their 2nd trimester. Majority (196 [52.5%]) of study participants spaced their latest inter pregnancy interval by greater than 2 years (Table 5).

Parasitic Infection

From the total study participants, 238 (63.8%) had no parasitic infection. More than a third of pregnant women were infected with different intestinal parasites. Of this, *Entamoeba histolytica* and *Giardialamblia* were predominant (Figure 1).

Prevalence of Anemia

The overall prevalence of anemia in this study was 28.7%, (95% CI: 24.1–33.5). Of this 68.2%, 29.9% and 1.9% were mild, moderate, and severe anemia respectively (Figure 2).

Factors Associated with Anemia Among Pregnant Women

On bivariate analysis, age, educational status, occupational status, residence, family size, household income, frequency of meat consumption, drinking coffee, presence of intestinal parasitic infection, iron supplementation, nutrition education, 24-hour dietary diversity score, latest inter pregnancy interval, and MUAC were associated with anemia with p-value > 0.25. After multivariate analysis, iron folate supplementation, presence of intestinal infection, inter pregnancy interval, and MUAC were significantly associated with anemia in pregnant women (Table 6).

Pregnant women who had good nutritional status (MUAC $\geq$ 24 cm) were 93% less likely to develop anemia as compared to women with MUAC < 24 cm [AOR= 0.07

Figure 1 Types of intestinal parasites identified from the pregnant women, DURH, South Ethiopia, 2019.

Figure 2 Severity of anemia among pregnant women attending antenatal care in DURH, Southern Ethiopia, 2019 (n=373).

(0.03, 0.18)]. In addition, pregnant women who had iron/folate supplementation were less likely to develop anemia by 94% [AOR=0.06 (0.02, 0.15)]. Pregnant women whom spaced the previous birth and their latest pregnancy by ≥ 2 yrs were 97% less likely to develop anemia as compared to their counterparts [AOR=0.03 (95% CI 0.009, 0.45)]. The odds of being anemic among pregnant women who had intestinal parasites were 6 times [AOR=6.11 (7.70, 37.0)] more likely as compared to those who had no intestinal parasites.

Discussion

In this study, the overall prevalence of anemia among pregnant women attending antenatal care was found to be 28.7% (95% CI: 24.1–33.5). Out of which 19.57% had mild anemia, 8.58% had moderate anemia, and 0.55% had severe anemia.

The magnitude of anemia among pregnant women in this study is a moderate public health problem.⁴ The prevalence found in this study is comparable with studies done in Southeast Ethiopia (27.9%), Western Ethiopia (29%), and Arbaminch town (33%).^{7,9,16} These similarities

Table 6 Factors Associated with Anemia Among Pregnant Women Attending Antenatal Care in DURH, South Ethiopia, 2019

Variables	Anemia		COR	AOR
	YES	NO		
Iron supplementation				
Yes	39 (36.4%)	238 (89.5%)	1	1
No	68(63.6%)	28(10.5%)	14.32(8.50, 25.82)	0.06(0.02, 0.15)**
MUAC				
≤ 23 cm	61 (38.5%)	33(12.4%)	1.00	1.00
≥ 24 cm	46 (43.0%)	233 (87.6%)	0.10(0.06,0.18)	0.07(0.03, 0.18)**
Inter pregnancy interval (in years)				
≤ 2	24(22.4%)	37(13.9%)	1.00	1.00
> 2	49 (45.8%)	147 (55.3%)	0.51(0.28, 0.94)	0.03 (0.09, 0.45)*
Intestinal parasites				
Yes	29 (27.1%)	209 (78.6%)	1.00	1.00
No	78 (72.9%)	57(21.4%)	9.86(5.88, 16.54)	6.11 (7.70,37.10) **

Notes: *p < 0.05; **p< 0.001.

might be due to similar socio economic characteristics and settings of the studies. But, the prevalence in this study was higher than in studies done in Tigray region (19.3%),⁷ and in Gondar town Amhara region (16.6%).¹⁷ These variations might be due to differences in cutoffs used to define anemia, geographic location, as well as the time differences (when the studies were undertaken).

In the present study, the odds of having anemia were significantly higher among pregnant who did not take iron supplementation during pregnancy as compared to those pregnant women who took iron supplementation. This finding was similar to a study done in North Western Tigray¹⁸ and Southern Ethiopia.¹⁹ The possible explanation for this association might be that pregnant women who take their iron tablets can meet their pregnancy-increased iron need, which in turn prevents the occurrence of anemia. Hence, focusing on strengthening the adherence of iron/folate supplementation will prevent anemia and its negative impacts on birth outcome.

Pregnant women who had a good nutritional status (MUAC ≥ 24 cm) were 93% less likely to develop anemia during pregnancy as compared to their counterparts. This result was supported by studies done in urban area of Eastern Ethiopia²⁰ and Oromia region.²¹ The possible explanation for the observed association could be increased overall nutritional demand during pregnancy, which results in both macro and micro nutrient deficiency, if intake is not sufficient.²² Those malnourished pregnant women might also be affected by micronutrient deficiency, hence leading to depletion of stored iron in the body and bone marrow, which results in anemia.

Birth spacing by ≥ 2 yrs reduces the likelihood of developing anemia during pregnancy by 97%. Similar finding was reported from studies done in Mekelle town⁷ and Arbaminch.⁹ The relationship between anemia and birth spacing can be explained by the fact that birth spacing will give mothers enough time to restore, replenish, and prepare their bodies for the next pregnancy. This implies family planning has a huge role in the health of mothers, children, and the family as a whole.

Having intestinal parasites increases the risk of anemia 6 times as compared to those who had no intestinal parasites. Our finding was in agreement with different studies conducted in different parts of Ethiopia and other countries that reported higher odds of anemia among pregnant women with intestinal parasites.^{16,19,23} Intestinal parasites, particularly hookworm infection has long been recognized among the major causes of anemia in poor communities.

Evidence has shown that increased hookworm infection intensity is associated with lower hemoglobin levels in pregnant women.¹⁹ Intestinal parasites reduce iron status via sucking blood from intestinal wall or by facilitating chronic blood loss, reducing appetite and nutrient intake.^{24,25}

This result implies that, other than routine prenatal iron supplementation, preventive or therapeutic deworming integrated with ANC help to reduce the burden of maternal anemia. According to the national guidelines of Ethiopia, pregnant women should be routinely dewormed in the second or third trimester of pregnancy. Different measures should be designed to increase ANC coverage to reach those who do not come to ANC clinics.

Strength and Limitation

A strength could be the high response rate. The limitation could be the fact that it was an institution-based study, with a cross-sectional design which was unadjusted for altitude as the participants come from different areas.

Conclusion

The overall prevalence of anemia among pregnant women attending antenatal care in the study area was found to be a moderate public health problem. Iron supplementation, good nutritional status (MUAC > 24 cm), and birth spacing reduce the odds of anemia. But having intestinal parasites was found to increase the likelihood of anemia during pregnancy.

Abbreviations

ANC, antenatal care; AOR, adjusted odds ratio; COR, crude odd ratio; DDS, dietary diversity score; EDHS, Ethiopian Demographic and Health Survey; Hgb, hemoglobin; HIV, human immunodeficiency virus; MUAC, mid upper arm circumference; SNNPR, Southern Nations, Nationalities, and Peoples Region; SPSS, Statistical Package for Social Sciences; WFP, World Food Program; WHO, World Health Organization.

Data Sharing Statement

The datasets used and/or analyzed during the current study are available from the corresponding author upon request.

Ethics Approval and Consent to Participate

Ethical clearance was obtained from institutional review board of Dilla University, College of health sciences and

medicine. Participants were informed about the purpose of the study, and written informed consent was obtained. Confidentiality was maintained by omitting their names on the questionnaire. We confirm this study was conducted in accordance with Declaration of Helsinki. While collecting the data, the pregnant women with Hgb <11 g/dl and women with IPIs were immediately advised to see health professionals at the ANC clinic for treatment and follow up.

Acknowledgments

We would like to thank Dilla University College of Medicine and Health Sciences for giving us permission to conduct this study. We also acknowledge our data collectors and study participants.

Author Contributions

All authors made a significant contribution to the work reported, that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

Dilla University funded the expenses for data collection.

Disclosure

The authors report no conflicts of interest in this work.

References

1. WHO. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Vitamin and mineral nutrition information system. Geneva: World Health Organization; 2011. Available from: <http://www.who.int/vmnis/indicators/haemoglobin.pdf>.
2. WHO. The global prevalence of anaemia in 2011. Geneva: World Health Organization; 2015. Available from: https://apps.who.int/iris/bitstream/9789241564960_eng.
3. Kassebaum NJ, Jasrasaria R, Naghavi M, et al. A systematic analysis of global anemia burden from 1990 to 2010. *Blood*. 2014;123(5):615–624. doi:10.1182/blood-2013-06-508325
4. De Benoist B, McLean E, Egli I, Cogswell M editors. *Worldwide Prevalence of Anaemia*. Vol. 1993–2005. Geneva: WHO Global Database on Anaemia. World Health Organization; 2008.
5. Central Statistical Agency [Ethiopia], the DHS Program. Ethiopia: demographic and health survey 2016. Addis Ababa and Maryland: CSA and ICF; 2017.
6. Central Statistical Agency [Ethiopia], the DHS Program. Ethiopia: demographic and health survey 2005. Addis Ababa and Maryland: CSA and ICF; 2006.
7. Abriha A, Yesuf ME, Wassie MM. Prevalence and associated factors of anemia among pregnant women of Mekelle town: a cross sectional study. *BMC Res Notes*. 2014;7:888. doi:10.1186/1756-0500-7-888
8. Ejeta E, Alemnew B, Fikadu A, Fikadu M, Tesfaye L, Birhanu T. Prevalence of anemia in pregnant women's and associated risk factors in Western Ethiopia. *Food Sci Qual Manag*. 2014;31:82–91.
9. Bekele A, Tilahun M, Mekuria A. Prevalence of anemia and its associated factors among pregnant women attending antenatal care in health institutions of Arba Minch Town, Southern Ethiopia. *Anemia*. 2016;2016:1–9. doi:10.1155/2016/1073192
10. FAO and FHI 360. *Minimum Dietary Diversity for Women: A Guide for Measurement*. Rome: FAO; 2016.
11. World Health Organization. Soil-transmitted helminth infections: key facts. 2018. Available from: <http://www.who.int/news-room/factsheets/detail/soil-transmitted-helminth-infections>. Accessed September 15, 2020.
12. Tang AM, Chung M, Dong K, et al. Determining a global mid-upper arm circumference cutoff to assess malnutrition in pregnant women. *Food Nutr Tech Assist*. 2016.
13. Coates J, Swindale A, Bilinsky P. Household food insecurity access scale (HFIAS) for measurement of household food access: indicator guide (v. 3). Washington, D.C: Food and Nutrition Technical Assistance Project, Academy for Educational Development; August, 2007.
14. Nkrumah B, Nguah SB, Sarpong N, et al. Hemoglobin estimation by the HemoCue® portable hemoglobin photometer in a resource poor setting. *BMC Clin Pathol*. 2011;11:5. doi:10.1186/1472-6890-11-5
15. Cheesbrough M. *District Laboratory Practice in Tropical Countries*. Cambridge: Cambridge University Press; 2006.
16. Kefiyalew F, Zemene E, Asres Y, Gedefaw L. Anemia among pregnant women in Southeast Ethiopia: prevalence, severity and associated risk factors. *BMC Res Notes*. 2014;7:771. doi:10.1186/1756-0500-7-771
17. Melku M, Addis Z, Alem M, Enawgaw B. Prevalence and predictors of maternal anemia during pregnancy in Gondar, Northwest Ethiopia: an institutional based cross-sectional study. *Anemia*. 2014. doi:10.1155/2014/108593
18. Gebre A, Mulugeta A. Prevalence of anemia and associated factors among pregnant women in North Western Zone of Tigray, Northern Ethiopia: a cross-sectional study. *J Nutr Metab*. 2015;2(4).
19. Lebso M, Anato A, Loha E. Prevalence of anemia and associated factors among pregnant women in Southern Ethiopia: a community based cross-sectional study. *PLoS One*. 2017;12(12):e0188783. doi:10.1371/journal.pone.0188783
20. Kedir H, Berhane Y, Worku A. Khat chewing and restrictive dietary behaviors are associated with anemia among pregnant women in high prevalence rural communities in eastern Ethiopia. *PLoS One*. 2013;8(11):e78601. doi:10.1371/journal.pone.0078601
21. Unicef. Prevention and control of nutritional anaemia: a South Asia priority. UNICEF Regional Office for South Asia; 2002. Available from: <https://hetv.org/pdf/anaemia-pc.pdf>. Accessed September 15, 2020.
22. Conde-Agudelo A, Rosas-Bermudez A, Castaño F, Norton MH. Effects of birth spacing on maternal, perinatal, infant, and child health: a systematic review of causal mechanisms. *Stud Fam Plann*. 2012;43(2):93–114. doi:10.1111/j.1728-4465.2012.00308.x
23. Tefera G. Determinants of anemia among pregnant women with emphasis on intestinal helminthic infection at Sher- Ethiopia Hospital, Ziway, Southern Ethiopia. *J Microbiol Immunol Infect Dis*. 2014;2(4):33–39.
24. Pullan R, Brooker S. The health impact of polyparasitism in humans: are we under-estimating the burden of parasitic diseases? *Parasitology*. 2008;135(7):783–794. doi:10.1017/S0031182008000346
25. Stephenson LS, Latham MC, Ottesen EA. Malnutrition and parasitic helminth infections. *Parasitology*. 2000;121:S23–38. doi:10.1017/S0031182000006491

Journal of Blood Medicine**Dovepress****Publish your work in this journal**

The Journal of Blood Medicine is an international, peer-reviewed, open access, online journal publishing laboratory, experimental and clinical aspects of all aspect pertaining to blood based medicine including but not limited to: Transfusion Medicine; Blood collection, Donor issues, Transmittable diseases, and Blood banking logistics; Immunohematology; Artificial and alternative blood based

therapeutics; Hematology; Biotechnology/nanotechnology of blood related medicine; Legal aspects of blood medicine; Historical perspectives. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/journal-of-blood-medicine-journal>