Medication-Induced Oral Hyperpigmentation: A Systematic Review

This article was published in the following Dove Press journal:
Patient Preference and Adherence

Nada O Binmadi

Maram Bawazir

Nada Alhindi

Hani Mawardi

Ghada Mansour

Sana Alhamed

Sarah Alfarabi

Sara Akeel

Soulafa Almazrooa

Oral Diagnostic Sciences Department,
Faculty of Dentistry, King Abdulaziz
University, Jeddah, Saudi Arabia

Background: Medication-induced oral hyperpigmentation is an oral condition that impacts patients' quality of life and has been linked to many systemic therapeutic agents. The exact pathogenesis of tissue pigmentation varies greatly and is not completely known. This systematic review aimed to present data on the causal association between medications and the development of oral/mucosal pigmentation as an adverse drug reaction.

Methods: A systematic review and analysis of literature were conducted using the following databases: PubMed, Science Direct, ProQuest, Web of Science, and Scopus. The systematic review included original articles written in English and published between January 1982 and June 2020. Following the PRISMA statement, eligible articles were systematically reviewed, and data were extracted from eligible studies and analyzed.

Results: A total of 235 articles were identified, of which 57 met the inclusion criteria and were included in this review. The mean age of included patients was 46.2 ± 16.38 years (range: 10–90 years) with a male to female ratio of 1:1.45. Oral mucosal hyperpigmentation was reported following the use of several classes of medications such as antiviral (eg, zidovudine), antibiotic (eg, minocycline), antimalarial (eg, chloroquine), anti-fungal (eg, ketoconazole), antileprotic (eg, clofazimine), antihypertensive (eg, amlodipine), chemotherapeutic, and antineoplastic drugs. The risk of developing oral pigmentation was significantly higher with antimalarial medications, antibiotics, antineoplastic and chemotherapeutic agents. Medication-induced oral hyperpigmentation was most frequent among women and in the hard palate.

Conclusion: Future research is warranted to better understand the pathogenesis and risk factors for medication-induced oral hyperpigmentation in order to reassure patients during prescription and management.

Keywords: medication-induced, oral hyperpigmentation, drugs, adverse reaction, systematic review

Introduction

The World Health Organization (WHO) defines an adverse drug reaction (ADR) as an unintended and noxious response to a drug that occurs at doses normally used in man for the prophylaxis, diagnosis or therapy of disease, or modification of physiological function.¹

An ADR in the oral cavity could present as xerostomia, lichenoid reaction/lichen planus, aphthous-like ulcers, bullous disorders, pigmentation, fibrovascular hyperplasia, epithelial keratosis, dysesthesia, osteonecrosis of the jaws, and/or angioedema.²

Medication-induced oral hyperpigmentation (MIOH) is an ADR that may develop rapidly after taking an associated drug once or for several continuous

Correspondence: Nada O Binmadi
Oral Diagnostic Sciences Department,
Faculty of Dentistry, King Abdulaziz
University, Jeddah, Saudi Arabia
Tel +966505699092
Email nmadi@kau.edu.sa

days or years.³ Clinically, MIOH presents as localized or generalized zones of blue to ill-defined black pigmentation affecting any site in the oral cavity, most commonly the gingiva, tongue, and buccal mucosa.^{4,5} Detailed patient medical history and information regarding clinical manifestations facilitate reaching a proper diagnosis.² While the exact mechanism of MIOH is still unclear, several mechanisms have been proposed, including an increase in the number of melanocytes in tissues, melanin production, and/or deposition/accumulation of medication metabolites.⁶

Management of MIOH varies and includes, but is not limited to, the discontinuation of the offending medication, switching to another drug, or laser photoablation.^{2,3} The oral pigmentation is usually involved in ADRs as common as cutaneous involvement; the dentist should therefore be informed regarding medications that cause oral ADRs to differentiate them from other etiologies and disorders.

There is currently a dearth of evidence-based literature to prove the relation between medications and oral hyperpigmentation; most published reports thus far are based on individual case studies or repeated observations. This review aimed to address this gap in the literature by systematically reviewing different aspects of MIOH and providing evidence of the causal relationship between medicinal drugs and subsequent ADRs presenting as mucosal pigmentation. We believe that the findings of this review will help familiarize dentists with this oral condition and thus provide patients with proper evaluation, diagnosis, and health management.

Materials and Methods

Protocol and Registration

The review protocol was registered with the International Prospective Register of Systematic Reviews (PROSPERO) (registration number: CRD42018087955) and followed the PRISMA statement guidelines.⁷ All study steps were conducted in accordance with the Cochrane Handbook for Systematic Review of Interventions.⁸

Inclusion and Exclusion Criteria

The inclusion criteria for this systematic review were as follows: 1) English language studies published between January 1982 and June 2020; 2) original studies on the usage of a medication or medication class associated with MIOH; and 3) observational cohort studies, cross-sectional studies, case reports, and case series. We excluded animal studies,

review articles, abstracts, and trials with outcomes irrelevant to our study or with data that were not reliable for extraction.

Literature Search

A computer-based literature search was carried out using five authentic electronic databases: Medline (PubMed), Science Direct, Scopus, ProQuest, and Web of Science. The following keywords were applied in the search strategy: ("Alkylating drugs" OR "Cyclophosphamide" OR "Antiarrhythmic drugs" OR "Amiodarone" OR "Anti-depressant drugs" OR "Imipramine" OR "Antifungal" OR "Ketoconazole" OR "Antileprotic drugs" OR "Clofazimine" OR "Antimalarial drugs" OR "Amodiaquine" OR "Chloroquine" OR "Hydroxychloroquine" OR "Quinacrine" OR "Quinidine" OR "Antimetabolites" OR "5 - Fluorouracil" OR "Antineoplastic drugs" OR "Cisplatin" OR "Hydroxyurea" OR "Antipsychotic drugs" OR "Chlorpromazine" OR "Phenothiazine" OR "Antiviral" OR "Zidovudine" OR "Contraceptives" OR "Oral contraceptives" OR "Cytotoxic antibiotic drugs" OR "Bleomycin" OR "Doxorubicin" OR "drug induced" OR "adverse drug effect" OR "adverse drug events" OR "adverse drug reaction" OR "drugs use" OR "medication use" OR "Other drugs" OR "Busulfan" OR "Desipramine" OR "Gold compounds" OR "Imatinib" OR "Minocycline" OR "Phenolphthalein" OR "Tegafur") AND ("Oral hyper-pigmentation" OR "oral ariasis" OR "oral discoloration" OR "oral melanosis" OR "oral pigmentation" OR "oral staining" OR "oral tanning").

Selection of Studies

Two authors (NB and MB) reviewed eligible studies in a two-phase stepwise manner, including screening of abstracts for eligibility and retrieval of full texts for final inclusion in our study.

Data Extraction

Data extraction from eligible studies was performed by two authors (NB and MB) independently using an online data extraction form that included study design, outcome, population, medical history relevant to patients, social history, sample size, and medication (type, dose, and duration of intake). In addition, the risk of bias assessment for each domain was conducted in order to establish transparency of the review. Cases of dispute between reviewers during data extraction were resolved upon consulting a third reviewer (SM).

Quality Assessment

We used the tool proposed by Murad et al⁹ for critical appraisal and quality assessment of the risk of bias in case reports and case series within our study. This tool uses the domains of selection, ascertainment, causality, and reporting for risk of bias assessment. Additionally, we used the Newcastle-Ottawa scale (NOS) for assessment of bias in the included cohort and cross-sectional studies.¹⁰ NOS relies on selection, comparability, and outcome domains for risk of bias revision. Each study's risk of bias was classified as low, moderate, or high.

Statistical Analysis

Statistical analysis for included studies was performed using IBM SPSS Statistics for Windows (Version 24.0. Armonk, NY: IBM Corp). Continuous variables were presented as means and standard deviations, categorical data were presented as frequencies and percentages, and *p*-values less than 0.05 were considered statistically significant.

Results

Search Results

We searched five reliable databases: PubMed, Scopus, Science Direct, ProQuest, and Web of Science. Our search using pre-specified keywords yielded a total of 235 citation

records. After removing duplicate studies, 175 unique records remained to be screened based on our intended inclusion and exclusion criteria. Title and abstract screening led to the exclusion of 15 studies because they were not written in the English language. Further screening was applied to the remaining studies. We excluded 103 full-text articles because they were either review articles or conference papers, or did not involve drug exposure or drug-induced hyperpigmentation in the oral cavity. Ultimately, 57 studies were eligible for our systematic review and statistical analysis.^{4,5,12-64} Figure 1 depicts a study flow diagram for the selection process following the PRISMA guidelines.⁷

Study Characteristics and Outcomes

We extracted data pertaining to a total of 268 patients involved in the selected studies. The mean age was 46.2 ± 16.38 years (range: 10–90 years) and the male to female ratio was 1:1.44. Data on the characteristics of the enrolled studies are shown in Table 1. Hyperpigmentation of the oral mucosa associated with antibiotic use was reported in 24 cases. The most affected sites were the gingiva, palate, and tongue. In total, 22 cases (91.7%) were associated with minocycline used for facial acne vulgaris at a dosage of 100–200 mg daily.^{4,12,13,15,17,18,23,27–30,34,37,44,65} The overall treatment duration ranged from 2 weeks to 11 years. Clinical characteristics included a bluish to grayish discoloration with

Figure 1 A flow diagram of the study selection process following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.

Notes: Liberati A, Altman D, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol.* 2009;62:10.⁷

Table 1 Summary of Characteristics of Included Patients

Characteristics		Mean	Standard Deviation
Age (years)		46.20	16.38
		Frequency	Percentage
Gender	Male	110	41.04%
	Female	158	58.95%
Race	Indian	30	11.19%
	White	19	7.09%
	Caucasian	75	27.99%
	Black	14	5.22%
	NA	115	42.9%
	Other	4	1.5%
Drug type	Antibiotic	24	8.96%
	Antimalarial	68	25.37%
	Antineoplastic	125	46.64%
	Antiviral	14	5.22%
	Chemotherapy	30	11.19%
	Antihypertensive	1	0.37%
	Antileprotic drug	1	0.37%
	Antifungal	1	0.37%
	Immunosuppressive	2	0.75%
	Hormonal therapy	1	0.37%
	Antiepileptic	1	0.37%
Pigmentation location	Hard palate	111	41.42%
	Tongue	61	22.76%
	Lip	2	0.75%
	Buccal mucosa	13	4.85%
	Gingiva	11	4.15%
	Multiple locations	70	26.12%
Pigmentation color*	Black	21	7.84%
	Brown	71	26.49%
	Blue	9	3.36%
	Gray	140	52.24%
	Melanotic Macule	14	5.22%
Cutaneous involvement	Yes	107	39.93%
	No	161	60.07%
Teeth pigmentation	Yes	7	2.61%
	No	261	97.39%
Bone pigmentation	Yes	7	2.61%
	No	261	97.39%

Note: *Each case may count for multiple pigment colors.

Abbreviations: NA, not available; SLE, systemic lupus erythematosus; RA, rheumatic arthritis; HTN, hypertension; MCL, mantle cell lymphoma.

variable size and site. Few cases reported cutaneous, tooth, or bone pigmentation. Additionally, linezolid and clindamycin use were reported in two other cases (4.1%).^{38,64}

Antimalarial medications were linked to MIOH in 68 cases affecting the palate, gingiva, and lips. Quinacrine hydrochloride,^{47,63,66} hydroxychloroquine,^{18,39} quinidine

sulfate,^{35,41} Daraclor,³² amodiaquine,⁴² and chloroquine compound^{31,33,39,45,60} were the most commonly-reported offending medications. Oral melanosis occurred with different doses, in various populations, within fluctuating periods. The lesions varied in size, color, and involvement of skin, teeth, and/or bone ([Supplementary Table 1](#)).

Oral hyperpigmentation induced by chemotherapeutic and antineoplastic agents was reported in 30 and 125 patients, respectively. The most prevalent agent was imatinib^{18,21,22,24,36,46,48–55,57,62} other agents included capecitabine,¹⁶ cyclophosphamide,^{25,40} 5-fluorouracil,⁴³ irinotecan,²⁶ and hydroxyurea.¹¹ The risk of oral pigmentation in association with imatinib therapy increased as the duration of intake increased (an average of 7 years) and occurred most commonly in the hard palate.⁵⁹ Other affected sites included the gingiva and tonsils.^{53,57}

Zidovudine, an antiviral agent, was associated with melanotic macules in the buccal mucosa and lips in 14 cases.⁵ Other reported medications included golimumab¹⁸ and cortisone⁶⁴ (immunosuppressants), ketoconazole (an anti-fungal agent),⁶⁴ amlodipine (an antihypertensive agent),¹⁴ retigabine (an antiepileptic agent),⁵⁶ and clofazimine (an antileprotic agent).⁶⁴ All baseline data of enrolled patients in the included studies are presented in [Supplementary Table 1](#).

Risk of Bias Assessment

The majority of the included studies were case reports and case series,^{4,5,11–19,21–32,35–42,44–46,48,54,58,62–66} most of which achieved a moderate quality level based on Murad et al's method.⁹ The NOS was used for the assessment of bias in the included cohort and cross-sectional studies. Included cohort studies achieved a moderate level of quality,^{5,11,28,40,60} while the cross-sectional study achieved a high level of quality.^{39,59} The quality of each included study is summarized in [Supplementary Table 2–4](#).

Discussion

In this systematic review, we aimed to provide evidence of the causal relation between medicinal drugs and subsequent ADRs presenting as oral/mucosal pigmentation and highlight the characterization of MIOH. Clinically, medication- and non-MIOH may share the same clinical features with no specific criteria for differentiation. To differentiate them, a thorough investigation should be conducted to consider the site, color, and duration of the lesion, patient's medical and social history, and medications used.⁶⁶

The current literature describes oral/mucosa hyperpigmentation as an adverse event due to the usage of several medications such as antibiotic (eg, minocycline), antiviral (eg, zidovudine), antimalarial (eg, chloroquine), chemotherapeutic, antineoplastic, anti-fungal drugs (eg, ketoconazole), antileprotic (eg, clofazimine), and antihypertensive drugs (eg, amlodipine). The pathophysiology of MIOH is still poorly understood and could differ based on the type of medication. For instance, one proposed theory of the underlying mechanisms focuses on the precipitation of minocycline metabolites in addition to a colored quinone derived from the aromatic ring.⁶⁷ However, imatinib is reported to have caused hyperpigmentation of unknown mechanisms and hypopigmentation in in-vitro studies due to the inhibition of the c-KIT pathway, which is usually involved in melanocyte development and regulation. Other agents in the same antineoplastic drug category as ipilimumab have shown yet another mechanism of reaction involving inhibition of cytotoxic T-lymphocyte antigen-4 (CTLA-4) and subsequent immune system activation.⁶⁸

The most common category of medication associated with oral pigmentation was found to be Antineoplastic. Imatinib-associated hyperpigmentation was described by several studies that reported treatment duration as a major risk factor leading to larger and darker lesions.^{59,68,69} In addition, antimalarial was the second most common category caused MIOH. A long duration (up to 15 years) of use of antimalarial drugs was observed to be associated with severe blue-black and gray diffused lesions in the hard palate with cutaneous involvement.^{31,35,42} An important development of used hydroxychloroquine in the treatment or prophylaxis of Coronavirus Disease-2019 (COVID-19) addresses the need for further analysis and follow up of MIOH in this population.⁷⁰

Minocycline, which is the most common antibiotic agent associated with MIOH, was reported to be associated with a diffuse bluish-gray discoloration affecting any site in the oral cavity within a relatively short duration around one month of use.^{4,28} Bone pigmentation was only associated with the use of minocycline, and teeth pigmentation was seen in patients who consumed either minocycline or imatinib.^{17,19,27,34,53}

Several antiviral agents were also linked to MIOH. Notably, Ficarra et al reported 14 cases of hyperpigmentation of the buccal mucosa and lips associated with zidovudine. This cohort study⁵ included human immunodeficiency virus (HIV)-seropositive patients who were administered

zidovudine. At the two-year follow-up, 14 (out of 217) HIV patients had developed pigmented lesions in the oral mucosa. Similarly, Almeida et al⁷¹ conducted a systematic review and meta-analysis comparing a group of patients on highly active antiretroviral therapy (HAART) to a non-HAART group in terms of oral lesions. They analyzed data from five studies and reported that 78 out of 724 (10.77%) patients who received HAART ultimately developed MIOH compared to 37 (8.5%) patients of the non-HAART group, with a risk ratio of 1.65 (95% Confidence Interval: 1.16–2.32).⁷¹ This could be explained by the augmented melanin production in the epithelium associated with the increased release of the melanocyte-stimulating hormone as a result of the systemic ketoconazole and zidovudine therapies.⁷² Few cases of hyperpigmentation induced by antihypertensive drugs have been reported.¹⁴

Thus far, no malignant transformation potential or serious complication with MIOH has been reported and no treatment is required. However, follow-up for the lesions is still advisable for the timely detection of any changes. Discontinuation of minocycline, imatinib, and hydroxychloroquine yielded persistence of pigmentation, not disappearance,^{17,18,39,52,55,57} while few reports did document regression of pigmentation after discontinuation of medication.^{16,18,58} In some cases, MIOH raises aesthetic concerns for patients based on the location of the lesions in the oral cavity. Few articles have reported the removal of minocycline-induced pigmentation by Q-switched Nd:YAG (neodymium-doped yttrium aluminum garnet), ruby, or alexandrite lasers.^{18,65} The best management strategies for MIOH should depend on proper diagnosis by a specialist and comprehensive patient education. Therefore, oral and maxillofacial specialists should be well-informed of all medications that can induce pigmentation in the oral mucosa in order to reassure patients about this unexpected hyperpigmentation.

This systematic review had several limitations. First, the included studies had a lower evidence level because the majority were case reports (class III evidence). Second, the heterogeneity of the study designs and outcome measures of the included studies affect the comparison of outcomes. Third, a correlation analysis between the risk factors, patient characteristics, and outcomes of the included studies was not feasible due to the lack of raw data.

Conclusion

In conclusion, there was a significant association between oral/mucosal hyperpigmentation among middle-aged

adults and prescribed medications, with most of the cases being caused by the use of antineoplastic, antimalarial and antibiotic agents. The practitioners and dentists should be aware of medications associated with hyperpigmentation to reassure the patients and monitor the potential side effects. Future research is required to better understand the risk factors, patients at risk and pathogenesis of MIOH to assist in patient management.

Disclosure

The authors report no conflicts of interest for this work.

References

- WHO. International drug monitoring: the role of national centres. *Rep WHO Meet.* 1972;1–25.
- Yuan A, Woo S-B. Adverse drug events in the oral cavity. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2015;119(1):35–47. doi:10.1016/j.oooo.2014.09.009
- Abdollahi M, Radfar M. A review of drug-induced oral reactions. *J Contemp Dent Pract.* 2003;4(1):10–31. doi:10.5005/jcdp-4-1-10
- Meyerson MA, Cohen PR, Hymes SR. Lingual hyperpigmentation associated with minocycline therapy. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 1995;79(2):180–184. doi:10.1016/S1079-2104(05)80279-3
- Ficarra G, Shillitoe EJ, Adler-Storthz K, et al. Oral melanotic macules in patients infected with human immunodeficiency virus. *Oral Surg Oral Med Oral Pathol.* 1990;70(6):748–755. doi:10.1016/0030-4220(90)90014-J
- Eisen D. Disorders of pigmentation in the oral cavity. *Clin Dermatol.* 2000;18(5):579–587. doi:10.1016/S0738-081X(00)00148-6
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol.* 2009;62(10):e1–e34. doi:10.1016/j.jclinepi.2009.06.006
- Higgins JP, Green S. *Cochrane Handbook for Systematic Reviews of Interventions.* Vol. 5. 2008. doi:10.1002/9780470712184
- Murad MH, Sultan S, Haffar S, Bazerbachi F. Methodological quality and synthesis of case series and case reports. *BMJ Evid Based Med.* 2018;23(2):60–63. doi:10.1136/bmjebm-2017-110853
- Wells GA, Shea B, O'Connell D, et al. *The Newcastle-Ottawa Scale (NOS) for Assessing the Quality of Nonrandomized Studies in Meta-Analyses.* 2016.
- Kumar B, Saraswat A, Kaur I. Mucocutaneous adverse effects of hydroxyurea: a prospective study of 30 psoriasis patients. *Clin Exp Dermatol.* 2002;27(1):8–13. doi:10.1046/j.0307-6938.2001.00947.x
- Fendrich P, Brooke RI. An unusual case of oral pigmentation. *Oral Surg Oral Med Oral Pathol.* 1984;58(3):288–289. doi:10.1016/0030-4220(84)90056-2
- Johnston S. Feeling blue? Minocycline-induced staining of the teeth, oral mucosa, sclerae and ears - A case report. *Br Dent J.* 2013;215(2):71–73. doi:10.1038/sj.bdj.2013.682
- Erbagci Z. Amlodipine associated hyperpigmentation. *Saudi Med J.* 2004;25(1):103–105.
- Treister NS, Magalnick D, Bin WS. Oral mucosal pigmentation secondary to minocycline therapy: report of two cases and a review of the literature. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004;97(6):718–725. doi:10.1016/j.tripleo.2003.11.006
- Vasudevan B. An unusual case of capecitabine hyperpigmentation: is hyperpigmentation a part of hand-foot syndrome or a separate entity. *Indian J Pharmacol.* 2010;42(5):326–328. doi:10.4103/0253-7613.70401
- Siller GM, Tod MA, Savage NW. Minocycline-induced oral pigmentation. *J Am Acad Dermatol.* 1994;30(2):350–354. doi:10.1016/S0190-9622(94)70038-9
- Tosios KI, Kalogirou EM, Sklavounou A. Drug-associated hyperpigmentation of the oral mucosa: report of four cases. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2018;125(3):e54–e66. doi:10.1016/j.oooo.2017.10.006
- Salman RA, Salman DG, Glickman RS. Minocycline induced pigmentation of the oral cavity. *J Oral Med.* 1985;40(3):154–157.
- Renald Perusse RM. Oral Pigmentation induced by premarin. *Laval Univ Sch Dent.* 1991;48:61–64.
- Lyne A, Creedon A, Bailey BMW. Mucosal pigmentation of the hard palate in a patient taking imatinib. *BMJ Case Rep.* 2015;2015(apr16 1):1–3. doi:10.1136/bcr-2015-209335
- Romeo U, Palaia G, Fantozzi PJ, Tenore G, Bosco D, Rare A. Case of melanosis of the hard palate mucosa in a patient with chronic myeloid leukemia. *Case Rep Dent.* 2015;2015:9–12. doi:10.1155/2015/817094
- Patel K, Cheshire D, Vance A. Oral and systemic effects of prolonged minocycline therapy. *Br Dent J.* 1998;185(185):560. doi:10.1038/sj.bdj.4809867
- Resende RG, Teixeira RGL, Vasconcelos FDO, Silva MEDS, Abreu MHG, Gomez RS. Imatinib-associated hyperpigmentation of the palate in post-HSCT patient. *J Craniomaxillofac Surg.* 2012;40(5):e140–e143. doi:10.1016/j.jcms.2011.07.010
- Parvaei P, Mortazavi H, Baharvand M. Severe pigmentation of oral mucosa, skin and nails due to breast cancer chemotherapy – case report. *Dent Med Probl.* 2015;52(3):363–365.
- Nestor LA, Flint S, Galvin S. Unusual case of hyperpigmentation secondary to irinotecan. *BMJ Case Rep.* 2016;2016:1–2. doi:10.1136/bcr-2016-217545
- Odell EW, Hodgson RP, Haskell R. Oral presentation of minocycline-induced black bone disease. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 1995;79(4):459–461. doi:10.1016/S1079-2104(05)80128-3
- Ozog DM, Gogstetter DS, Scott G, Gaspari AA. Minocycline-induced hyperpigmentation in patients with pemphigus and pemphigoid. *Arch Dermatol.* 2000;136(9):1133–1138. doi:10.1001/archderm.136.9.1133
- Cale AE, Freedman PD, Lumerman H. Pigmentation of the jawbones and teeth secondary to minocycline hydrochloride therapy. *J Periodontol.* 1988;59(2):112–114. doi:10.1902/jop.1988.59.2.112
- Morrow GL, Abbott RL. Minocycline-induced scleral, dental, and dermal pigmentation. *Am J Ophthalmol.* 1998;125(3):396–397. doi:10.1016/S0002-9394(99)80156-1
- de Filho MRM, da Silva CAD, da Dourado MR, de Pires MBO, Pêgo SPB, de Freitas EM. Palate hyperpigmentation caused by prolonged use of the anti-malarial chloroquine. *Head Neck Pathol.* 2012;6(1):48–50. doi:10.1007/s12105-011-0288-5
- Levy H. Chloroquine-induced pigmentation. *S Afr Med J.* 1982;62(December 1981):1981–1983.
- De Andrade BAB, Fonseca FP, Pires FR, et al. Hard palate hyperpigmentation secondary to chronic chloroquine therapy: report of five cases. *J Cutan Pathol.* 2013;40(9):833–838. doi:10.1111/cup.12182
- Beehner ME, Houston GD, Young JD. Oral pigmentation secondary to minocycline therapy. *J Oral Maxillofac Surg.* 1986;44(7):582–584. doi:10.1016/S0278-2391(86)80104-5
- Birek C, Ph D. Short communications & case reports associated with cases of oral pigmentation with quinidine therapy. *Oral Surg Oral Med Oral Pathol.* 1988;66(66):59–61. doi:10.1016/0030-4220(88)90067-9
- Bombeccari GP, Garagiola U, Pallotti F, et al. Hyperpigmentation of the hard palate mucosa in a patient with chronic myeloid leukaemia taking imatinib. *Maxillofac Plast Reconstr Surg.* 2017;39(1). doi:10.1186/s40902-017-0136-y
- Buddula A. Staining of palatal torus secondary to long term minocycline therapy. *J Indian Soc Periodontol.* 2009;13(1):48. doi:10.4103/0972-124x.51896

38. Balaji G, Maharani B, Ravichandran V, Parthasarathi T. Linezolid induced black hairy tongue. *Indian J Pharmacol.* 2014;46(6):653–655. doi:10.4103/0253-7613.144942
39. Bahloul E, Jallouli M, Garbaa S, et al. Hydroxychloroquine-induced hyperpigmentation in systemic diseases: prevalence, clinical features and risk factors: a cross-sectional study of 41 cases. *Lupus.* 2017;26(12):1304–1308. doi:10.1177/0961203317700486
40. Acharya S, Pai KM, Bhat S, Mamatha B, Bejadi V, Acharya S. Oral changes in patients undergoing chemotherapy for breast cancer. *Indian J Dent Res.* 2017;28(3):261–268. doi:10.4103/ijdr.IJDR_379_16
41. Mahler R, Sissons W, Watters K. Pigmentation induced by quinidine therapy. *Arch Dermatol.* 1986;122(9):1062–1064. doi:10.1001/archderm.1986.01660210112031
42. Watson IB, MacDonald DG. Amodioquine induced oral pigmentation - a light and electron microscopic study. *J Oral Pathol Med.* 1974;3(1):16–21. doi:10.1111/j.1600-0714.1974.tb01694.x
43. Co ML, Esteban MJ. Lingual hyperpigmentation after 5-fluorouracil chemotherapy. *BMJ Case Rep.* 2017;2017:1–2. doi:10.1136/bcr-2017-219806
44. Ayangco L, Sheridan PJ. Minocycline-induced staining of torus palatinus and alveolar bone. *J Periodontol.* 2003;74(5):669–671. doi:10.1902/jop.2003.74.5.669
45. de Andrade BAB, Padron-Alvarado NA, Muñoz-Campos EM. Hyperpigmentation of hard palate induced by chloroquine therapy. *J Clin Exp Dent.* 2017;9(12):e1487–e1491. doi:10.4317/jced.54387
46. Wong M, Sade S, Gilbert M, Klie HBE. Oral melanosis after tyrosine kinase inhibition with imatinib for chronic myelogenous leukemia: report of a case and review of the literature. *Dermatol Online J.* 2011;17(5).
47. López-Jornet P, Pons-Fuster A. Pigmented oral mucosa due to quinacrine. *N Y State Dent J.* 2011;77(6):49–51.
48. Mattsson U, Halbritter S, Möner Serikoff E, Christerson L, Warfvinge G. Oral pigmentation in the hard palate associated with imatinib mesylate therapy: a report of three cases. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2011;111(5):e12–e16. doi:10.1016/j.tripleo.2010.11.006
49. Di Tullio F, Mandel VD, Scotti R, Padalino C, Pellacani G. Imatinib-induced diffuse hyperpigmentation of the oral mucosa, the skin, and the nails in a patient affected by chronic myeloid leukemia: report of a case and review of the literature. *Int J Dermatol.* 2018;57(7):784–790. doi:10.1111/ijd.13931
50. Song HSKH, Kang HY. Imatinib mesylate-induced hyperpigmentation of the nose and palate. *Ann Dermatol.* 2014;26(4):532–533. doi:10.5021/ad.2014.26.4.532
51. Roeker LEWA, Wolanskyj AP. Imatinib-associated melanosis of the palate. *Am J Hematol.* 2014;89(5):564. doi:10.1002/ajh.23589
52. Khoo TL, Catalano A, Supple S, et al. Hyperpigmentation of the hard palate associated with imatinib therapy for chronic myeloid leukemia with a genetic variation in the proto-oncogene c-KIT. *Leuk Lymphoma.* 2013;54(1):186–188. doi:10.3109/10428194.2012.702904
53. Singh NBS, Bakhshi S. Imatinib-induced dental hyperpigmentation in childhood chronic myeloid leukemia. *J Pediatr Hematol Oncol.* 2007;29(3):208–209. doi:10.1097/MPH.0b013e318033a76c
54. Li CC, Malik SM, Blaaser BF, et al. Mucosal pigmentation caused by imatinib: report of three cases. *Head Neck Pathol.* 2012;6(2):290–295. doi:10.1007/s12105-011-0325-4
55. Lewis DM. Diffuse pigmentation of the palate. *J Okla Dent Assoc.* 2009;100:24–25.
56. Beacher NG, Brodie MJ, Goodall C. A case report: retigabine induced oral mucosal dyspigmentation of the hard palate. *BMC Oral Health.* 2015;15(1):1–5. doi:10.1186/s12903-015-0102-y
57. Steele JC, Triantafyllou A, Rajlawat BPFE, Field EA. Oral mucosal hyperpigmentation and horizontal melanonychia caused by imatinib. *Clin Exp Dermatol.* 2012;37(4):432–433. doi:10.1111/j.1365-2230.2011.04196.x
58. Dixon DR, Yassin A. Oral effects and early implant survival results after imatinib discontinuation therapy for chronic myelogenous leukemia: a case report. *J Clin Periodontol.* 2017;7(3):115–120. doi:10.1902/cap.2016.160050
59. Oliveira SR, de Azevedo Branco LG, Rocha AL, et al. Association of oral mucosa hyperpigmentation with imatinib mesylate use: a cross-sectional study and a systematic literature review. *Clin Oral Investig.* 2019;23(12):4371–4382. doi:10.1007/s00784-019-02886-0
60. Chacón-Dulcey V, López-Labady J, Villarreal-Dorrego M, et al. Oral manifestations associated with antimalarial therapy in patients with systemic lupus erythematosus. *Lupus.* 2020;29(7):761–766. doi:10.1177/0961203320922620
61. Kristensson JH, Sander BB, von Euler-chelpin M, Lynge E. Predictors of non-participation in cervical screening in Denmark. *Cancer Epidemiol.* 2014;38(2):174–180. doi:10.1016/j.canep.2013.12.007
62. Mascitti M, Luconi E, Togni L, Barlattani A, Santarelli A. Imatinib-related hyperpigmentation of oral mucosa: case report and literature review. *J Dent Sci.* 2019;14(3):335–337. doi:10.1016/j.jds.2019.02.005
63. Lerman MA, Karimbux N, Guze KA, Bin WS. Pigmentation of the hard palate. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2009;107(1):8–12. doi:10.1016/j.tripleo.2008.07.022
64. Langford A, Pohle HD, Gelderblom H, Zhang X, Reichart PA. Oral hyperpigmentation in HIV-infected patients. *Oral Surg Oral Med Oral Pathol.* 1989;67(3):301–307. doi:10.1016/0030-4220(89)90360-5
65. LaPorta VN, Nikitakis NG, Sindler AJ, Reynolds MA. Minocycline-associated intra-oral soft-tissue pigmentation: clinicopathologic correlations and review. *J Clin Periodontol.* 2005;32(2):119–122. doi:10.1111/j.1600-051X.2005.00646.x
66. Kleinegger CL, Hammond HL, Finkelstein MW. Oral mucosal hyperpigmentation secondary to antimalarial drug therapy. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2000;90(2):189–194. doi:10.1067/moe.2000.106340
67. Geria AN, Tajirian AL, Kihiczak G, Schwartz RA. Minocycline-induced skin pigmentation: an update. *Acta Dermatovenereol Croat.* 2009;17(2):123–126.
68. Dai J, Belum VR, Wu S, Sibaud V, Lacouture ME. Pigmentary changes in patients treated with targeted anticancer agents: a systematic review and meta-analysis. *J Am Acad Dermatol.* 2017;77(5):902–910.e2. doi:10.1016/j.jaad.2017.06.044
69. Mcpherson T, Sherman V, Turner R. Imatinib-associated hyperpigmentation, a side effect that should be recognized. *J Eur Acad Dermatol Venereol.* 2009;23(1):82–83. doi:10.1111/j.1468-3083.2008.02706.x
70. Eljaaly K, Alireza KH, Alshehri SA-TJ, Al-Tawfiq JA. Hydroxychloroquine safety: a meta-analysis of randomized controlled trials. *Travel Med Infect Dis.* 2020;36:101812. doi:10.1016/j.tmaid.2020.101812
71. de Almeida VL, Lima IFP, Ziegelmann PK, Paranhos LR, de Matos FR. Impact of highly active antiretroviral therapy on the prevalence of oral lesions in HIV-positive patients: a systematic review and meta-analysis. *Int J Oral Maxillofac Surg.* 2017;46(11):1497–1504. doi:10.1016/j.ijom.2017.06.008
72. Hamza OJ, Matee MI, Simon EN, et al. Oral manifestations of HIV infection in children and adults receiving highly active anti-retroviral therapy [HAART] in Dar es Salaam, Tanzania. *BMC Oral Health.* 2006;6(1):12. doi:10.1186/1472-6831-6-12

Patient Preference and Adherence**Dovepress****Publish your work in this journal**

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease

states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>