Prevalence and Risk Factors for Diabetic Peripheral Neuropathy Among Saudi Hospitalized Diabetic Patients: A Nested Case-Control Study

This article was published in the following Dove Press journal:
International Journal of General Medicine

Fahad AS Aleidan^{1,2}
Banderi A Ahmad¹
Farah A Alotaibi¹
Dalal H Aleesa¹
Nuha A Alhefdhi¹
Motasim Badri¹
Abdel Galil Abdel Gader¹

¹College of Medicine, King Saud Bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia; ²College of Medicine, King Abdulaziz Medical City, Riyadh, Saudi Arabia

Objective: To determine the prevalence and the risk factors of diabetic peripheral neuropathy (DPN) in hospitalized adult Saudi diabetics.

Methods: This is a retrospective, nested case-control study conducted at King Abdulaziz Medical City (KAMC) in Riyadh, Saudi Arabia. All diabetic patients admitted to the hospital between the January 1, and December 31, 2018 were considered for inclusion in the study. Patients with DPN were identified and three controls per case were randomly selected from the remaining diabetic patients without peripheral neuropathy (PN).

Results: A total of 2,096 adult diabetic patients were identified during the study period. Of these, 73 patients (3.5%) were confirmed to be suffering from DPN and 219 were included as controls. When comparing diabetic with the control cases, DPN cases were significantly older ($p=0.002$), had a significantly higher proportion of type 2 diabetes ($p=0.023$), chronic kidney disease ($p<0.0001$), cerebral vascular stroke ($p=0.027$), hypertension ($p=0.005$), dyslipidemia ($p=0.002$), peripheral vascular disease ($p<0.0001$), osteoarthritis ($p=0.034$), diabetic ketoacidosis ($p=0.003$), foot ulcers ($p=0.006$), gangrene ($p=0.001$), lower limb ischemia ($p=0.001$), increased duration with diabetic disease ($p=0.031$), increased BMI ($p=0.003$), higher serum creatinine ($p<0.001$) and lower serum albumin levels ($p=0.035$). In the multivariate logistic regression, only older age {odds ratio (OR) 1.02, 95% CI 1.01–1.04, $p=0.031$ }, chronic kidney disease (OR 2.39, 95% CI 1.23–4.64, $p=0.010$) and peripheral vascular disease (OR 3.14, 95% CI 1.39–7.13, $p=0.006$) were independently associated with DPN.

Conclusion: This study identified several risk factors that contributed to the development of DPN in Saudis. These must be considered in strategies and campaigns aimed at risk reduction of cardiovascular and chronic diseases, and consequently progression of DPN.

Keywords: diabetes mellitus, diabetic peripheral neuropathy, risk factors

Introduction

Diabetic peripheral neuropathy DPN is defined as, “the presence of symptoms and/or signs of peripheral nerve dysfunction in patients with diabetes after the exclusion of other causes,” and may be present despite a lack of symptoms.¹ The diagnosis relies on clinical signs as well as quantitative electrophysiological testing.¹ DPN is a common complication in patients with diabetes and its consequences can be distressing, as patients may develop neuropathic foot pain and decreased sensation that can lead to frequent falls and injuries. Patients may also develop foot ulcers. All these consequences of DPN will unavoidably end in the reduction in the quality

Correspondence: Fahad AS Aleidan
College of Medicine, King Saud Bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia
Tel +966118011111
Email faleidan@yahoo.com

of life and significant economic burden both to the patients and society.^{2–5} Although up to 50% of patients with DPN are asymptomatic, some patients may manifest burning feet pain, due to progressive sensory impairment, which responds poorly to analgesics. Rarely, this sensory impairment could lead to foot ulceration and eventually to lower limb amputation.^{6,7} diabetes is a leading cause of “non-traumatic” lower limb amputation.⁸

There is wide agreement that DPN is a consequence of diabetes longstanding hyperglycemia which is critical for peripheral nerve damage and distal-predominant nerve fiber degeneration.^{9–11} The chronic diabetes hyperglycemia is compounded by numerous metabolic aberrations and diabetic microvascular complications resulting from disturbed nutritional support especially of the most distal parts of very long nerve axons originating in the spinal cord and travel long distances to supply the lower limbs and the feet, in particular. Additionally, the sparse vascular supply in this area, resulting from diabetes, is likely to cause further hypoxic nerve damage. Other confounding metabolic aberrations resulting from chronic hypoglycemia include the excessive release of cytokines, and exaggerated oxidative stress. This topic has been reviewed in detail elsewhere.⁹ Although diabetes is considered the main underlying cause of DPN, the pre-diabetic stage with other comorbidities such as: obesity, hypertension and hyperlipidemia may increase the risk of neuropathy.^{12–15} Moreover, treatment modalities received by diabetic patients (insulin and/or oral hypoglycemic agents) play an additional important role in delaying the development of this complication among diabetic patients. A meta-analysis conducted on 1,228 patients with type 1 diabetes and 6,669 patients with type 2 diabetes revealed significant reduction of neuropathy when blood sugar is controlled within the limits of normal.¹³

Type 2 diabetes mellitus is becoming a rapidly growing health problem in the affluent oil rich Gulf Council states (Saudi Arabia, Oman, Kuwait, Bahrain, and the United Arab Emirates). Its prevalence is highest in Saudi Arabia (31.6%, 29.0%, 25.4%, 25.0%, 25.0%, and 16.7%, respectively).¹⁶ This high prevalence of type 2 diabetes was found to be significantly associated with the high Gross Domestic Product (GDP) and energy consumption. In contrast the lowest prevalence was found in the poorest Arab countries (Mauritania, 4.7% and Somalia, 3.9%).

Careful search in the literature uncovered only two small studies on painful DPN undertaken in different primary health facilities.^{17,18} The first study¹⁷ which was

undertaken in a Primary Health Centre (PHC) in Riyadh found 35% of patients with type-2 diabetes ($n = 242$) suffering from DPN. On the other hand, the second study¹⁸ which was also a PHC study found the prevalence of DPN to be 30.1% ($n = 235$). With this alarming prevalence of DPN in Saudi diabetics we felt the need to have better documentation of DPN. Thus, the aim of the current study was to determine the prevalence and risk factors associated with the development of DPN in a large population of hospitalized adult Saudi diabetics.

Methods

This is a retrospective, nested case-control study conducted at King Abdulaziz Medical City (KAMC); a tertiary health care facility in Riyadh, Saudi Arabia. All adult patients, who were admitted with diabetes mellitus at KAMC, during the study period (from January 1 to December 31, 2018) were considered for inclusion in the study. Additionally, they met the ADA diagnosis criteria (fasting blood glucose ≥ 7.0 mmol/L, glycosylated hemoglobin [HbA1C] $\geq 6.5\%$, classic symptoms of hyperglycemia, hyperglycemic crisis with a random plasma glucose ≥ 11.1 mmol/L or 2-hour plasma glucose ≥ 11.11 mmol/L). Of these, patients with confirmed DPN diagnosis were identified and served as case patients. Whereas, three controls per case were randomly selected from the remaining diabetic patients without peripheral neuropathy (PN). Patients who were under the age of 18 or did not fulfill the eligibility criteria were not considered for participation in the study.¹⁹ DPN confirmation was based on the following criteria: 1) abnormal clinical examination for pain, touch, vibration, pressure, power, and ankle reflex; and 2) abnormal nerve conduction studies (NCS) and/or electromyography.

Medical records of all enrolled patients (cases and controls) were accessed and the collected data were recorded in computerized format. These data included: demographic characteristics: age, gender, body mass index (BMI), type of diabetes, presence and duration of diabetic neuropathy, hypertension, kidney disease, dyslipidemia, and other comorbidities, life style (smoking and alcohol use), blood biochemistry indices (fasting blood glucose, HbA1C, lipid profile), medications, particularly insulin, oral hypoglycemic agents and diet, blood pressure and other clinical complications.

All methods and procedures were performed in accordance with the relevant institutional guidelines and regulations. The study was approved by the Institutional Review

Board at King Abdullah International Medical Research Center (KAIMRC). The ethical committee waived the need for written informed consent due to the retrospective nature of the study.

Statistical Analysis

Data were analyzed using the Statistical Package for the Social Sciences Program (IBM SPSS Statistics version 24). Continuous data were presented as medians (inter-quartile range), and categorical data as frequencies (%). Data were compared using Mann–Whitney *U*-test or χ^2 -test; as appropriate. Conditional logistic regression analysis was conducted to identify the factors associated with the DPN. Factors found statistically significant in logistic univariate analyses were included in the final multivariate model. Strength of the association was expressed as odds ratios (OR) with 95% confidence interval. All tests were two-sided and a *p*-value less than 0.05 was considered statistically significant.

Results

A total of 2,906 patients with diabetes were identified during the study period. Of these, 73 patients (3.5%) were found and labeled in the electronic chart by the treating clinician with DPN and 219 patients (three to one DPN case) were randomly selected as a control group. Table 1 shows the basic characteristics of the two groups. Females represented 51.8% and 63.0% of the DPN and the controls, respectively. The median (IQR) age of patients in the DPN group was significantly higher than in the controls [65 (58–75) vs 61 (45–71), *p*=0.002] and the duration of diabetes of patients with DPN was significantly higher than in the controls [8.5 (5.5–16) vs 6.5 (4–15), *p*=0.031]. Type 2 DM was diagnosed in 66 (90.4%) patients with DPN, compared with 173 (78.6%) patients in the control group, and 7 (9.6%) patients with DPN were diagnosed with type 1 DM compared with 47 (21.4%) patients in the control group. The median BMI of patients with DPN was significantly higher than controls [31 (26–35) vs 27 (22–33), *p*=0.003]. However, the two groups were similar in gender distribution (*p*=0.096), treatment modalities (*p*=0.072), smoking status (*p*=0.131) and alcohol consumption (*p*=0.413). Biochemical indices and assessment parameters were also similar in both groups (Table 1).

The median number of comorbidities in DPN patients was significantly higher than in controls [5(4–7.5) vs 4 (2–

5.75), *p*<0.001]. Hypertension was more prevalent in the DPN patients (*p*=0.005); so were peripheral vascular disease (*p*<0.001), cerebral vascular accident (*p*=0.027), chronic kidney disease (*p*<0.001) and dyslipidemia (*p*=0.002), than in control patients. Other comorbid diseases were similar in both groups. The DPN patients developed diabetic complications more than the control patients (Table 2).

Table 3 shows the results of the final multivariate logistic regression for factors associated with DPN. The independent factors associated with the likelihood of DPN were age (OR 1.02, 95% CI 1.01–1.05, *p*=0.031), peripheral vascular disease (OR 3.14, 95% CI 1.39–7.127, *p*=0.006) and chronic kidney disease (OR 2.39, 95% CI 1.23–4.64, *p*=0.010). However, the presence of hypertension, cerebral vascular disease and dyslipidemia were not significantly associated with DPN.

Discussion

DPN is a very distressing chronic complication of diabetes with an array of poor outcomes. For example, DPN leads to neuropathic pain and diminished sensation which in turn can lead to frequent falls and injuries, restriction of movement, poor quality of life and difficulties with earning a living. Other serious complications include: leg ulcers that can eventually end in leg amputation.^{1–5,20,21}

DPN is classified into numerous types and subtypes based on a mixture of phenomenological, aetiological, pathological and neurophysiological parameters.^{1,22–24} In a retrospective study, like the present one, it is difficult to assure the reliability of information on the details of these types or subtypes and be confident that such information has been documented in every patient. Accordingly, we restricted the definition of DPN, as mentioned earlier, to the presence of symptoms and/or signs of peripheral nerve dysfunction in patients with diabetes after the exclusion of other causes.¹

Interest in DPN stems from the fact that it has recently been recognized as the most common of all the neuropathy complications worldwide.²⁵ However, the reported prevalence of DPN varies widely from study to study and is heavily dependent on the type of diabetes, the population selected, and criteria used for defining cases, including the degree of control of the hyperglycemia.²⁶ The most classical citation of DPN is a study conducted in France²⁷ in which 4,400 adults with diabetes were followed for 36 years (1937–1973). Approximately 50% of these patients developed DPN at the end of the follow-up period. Pirart²⁷ found that two thirds of patients with diabetes had

Table I Clinical and Demographic Characteristics of the DPN and Control Patients

Variables	Control (n=220)	DPN (n=73)	p-value
Age, years (median, IQR)	61 (45–71)	65 (58–75)	0.002
Gender, n (%)			0.096
Male	106 (48.2)	27 (37.0)	
Female	114 (51.8)	46 (63.0)	
Duration of diabetes years, median (IQR)	6.5 (4–15)	8.5 (5.5–16)	0.031
Type of DM, n (%)			0.025
Type I	47 (21.4)	7 (9.6)	
Type II	173 (78.6)	66 (90.4)	
Type of treatment, n (%)			0.072
Insulin	158 (72.8)	62 (86.1)	
Oral hypoglycemic	53 (24.4)	9 (12.5)	
Diet	6 (2.8)	1 (1.4)	
BMI kg/cm ² , median (IQR)	27 (22–33)	31 (26–35)	0.003
Smoking status	29 (13.2)	15 (20.5)	0.131
Alcohol status	2 (0.9)	0 (0.0)	0.413
Laboratory indices			
Fasting blood glucose mmol/L (median, IQR)	9 (7–12)	10 (6–14)	0.507
HbA1C (median, IQR)	9 (7–10)	9 (7–10)	0.851
Systolic blood pressure mmHg (median, IQR)	124 (107–139)	125 (110–143)	0.254
Diastolic blood pressure mmHg (median, IQR)	69 (58–76)	66 (55–73)	0.158
Micro-albuminuria (median, IQR)	26 (2–100)	50 (50–200)	0.092
Proteinuria (median, IQR)	2 (1.25–2)	2 (1–2)	0.218
Serum Creatinine umol/L (median, IQR)	71 (61–117)	103 (80–192.5)	<0.001
Cholesterol mmol/L (median, IQR)	4 (3–5)	4 (3–4.75)	0.649
Triglycerides mmol/L (median, IQR)	1 (1–2)	1 (1–2)	0.356
Albumin g/L (median, IQR)	37 (32–40)	35 (30–39)	0.035

Abbreviations: BMI, body mass index; DPN, diabetes peripheral neuropathy; IQR, interquartile range.

objective evidence of some form of neuropathy. The most common was DPN; affecting 50% of type 2 diabetics. Outside Europe, particularly in Asia and Africa, a very wide range of prevalences were reported. For examples Saudi Arabia (19.9; n = 552);²⁸ Turkey (60%; n = 550),²⁹ China: (61.8%, n = 435);³⁰ and Tanzania (72.2%, n = 327).³¹ On the other hand, recent studies from Europe and the US have reported prevalences ranging from 6% to 51% depending on the population studied.¹ We noted that the frequency of DPN in our patients was considerably lower than expected. This maybe owing to the fact that, typically, DPN is often diagnosed clinically with little further laboratory investigations to confirm the diagnosis or poorly captured in patients' records. In a study by Day et al,³² more than 40% of diabetic patients in general

practice had no biochemical evaluation, eye or foot examination.³²

The wide geographical variations in prevalence of DPN highlight the need for a global multicenter study in which the selection criteria for diabetic patients are unified to the finest detail. Hopefully, then, the true reflection of the geographical and ethnic differences in the prevalence of DPN can be meaningful and may shed some light in the mechanism underlying the differences in the prevalences in the DPN in different ethnic groups.

This possibility proved to be true in a comparative study between South Asians and Caucasians living in Britain where the prevalence of DPN was found to be significantly lower in South Asians (38.1%) than White Caucasians (54.3%). Further observations uncovered

Table 2 Associated Comorbidities and Diabetes Complications

Variables	Control (n=220)	DPN (n=73)	p-value
Number of comorbid disease (median, IQR)	4 (2–5.75)	5 (4–7.5)	<0.001
Ischemic heart disease, n (%)	92 (42)	37 (50.7)	0.196
Congestive heart failure, n (%)	53 (24.4)	23 (31.5)	0.226
Chronic kidney disease, n (%)	32 (14.6)	27 (37)	<0.001
Cerebral vascular stroke, n (%)	32 (14.7)	19 (26)	0.027
Hypertension, n (%)	160 (73)	65 (89)	0.005
Dyslipidemia, n (%)	119 (54.6)	55 (75)	0.002
Peripheral vascular disease, n (%)	15 (7)	17 (24)	<0.001
Osteoarthritis, n (%)	15 (6.9)	11 (15)	0.034
Hypoglycemia, n (%)	10 (4.6)	2 (2.7)	0.492
Diabetes ketoacidosis, n (%)	30 (13.8)	1 (1.4)	0.003
Diabetes complications, n (%)			
Retinopathy	28 (12.8)	16 (21.9)	0.061
Foot ulcer	15 (6.9)	13 (17.8)	0.006
Gangrene	3 (1.4)	7 (9.6)	0.001
Lower limb ischemia	7 (3.2)	10 (13.7)	0.001

Abbreviations: DPN, diabetes peripheral neuropathy; IQR, interquartile range.

foot insensitivity as assessed by 10 g monofilament perception, was more common in Caucasians (43.9% vs 23.8%). South Asians have better preserved small nerve fiber integrity than equivalent Europeans.

Corneal nerve fiber length (22.0 ± 7.9 vs 19.3 ± 6.3 mm/mm²), corneal nerve branch density [(geometric mean (range): 60.0 (4.7–246.2) vs 46.0 (3.1–129.2) no./mm²], and heart rate variability [(geometric mean

Table 3 Univariate and Multivariate Conditional Logistic Regression for Risk Factors Associated with DPN

Factor	Univariate Analysis		Multivariate Analysis	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Age	1.03 (1.01–1.05)	<0.001	1.02 (1.01–1.04)	0.031
Chronic kidney disease				
Yes	3.43 (1.87–6.28)	<0.001	2.39 (1.23–4.64)	0.010
No	1		1	
Cerebral vascular stroke				
Yes	2.05 (1.08–3.89)	0.029	1.64 (0.80–3.33)	0.176
No	1		1	
Peripheral vascular disease				
Yes	4.11 (1.93–8.74)	<0.001	3.14 (1.39–7.13)	0.006
No	1		1	
Hypertension				
Yes	3.00 (1.36–6.62)	0.007	0.83 (0.27–2.54)	0.740
No	1		1	
Dyslipidemia				
Yes	2.54 (1.40–4.61)	0.002	1.56 (0.78–3.13)	0.181
No	1		1	

Abbreviations: CI, confidence interval; OR, odds ratio.

(range): 7.9 (1.4–27.7) vs 6.5 (1.5–22.0)], were significantly higher in South Asians than Europeans.^{33,34}

As mentioned earlier DPN is believed to be a consequence of the diabetes longstanding hyperglycemia which is critical for peripheral nerve damage and distal-predominant nerve fiber degeneration.^{9–11} The hyperglycemia of the chronic diabetes process is compounded by numerous metabolic aberrations and diabetic microvascular complications resulting from the disturbed nutritional support especially of the most distal parts of very long nerve axons originating in the spinal cord and travel long distances to supply the lower limbs and the feet in particular. Additionally, the sparse vascular supply in this area resulting from diabetes is likely to cause further hypoxic nerve damage. Other confounding metabolic aberrations resulting from the chronic hypoglycemia include excessive release of cytokines, and exaggerated oxidative stress. This topic has been reviewed in detail elsewhere.⁹

It is also becoming clear that the duration of diabetes and glycemic control are the most significant risk factors for DPN.³⁵ In regard to the benefit of blood glucose control, there are reports indicating that aggressive blood glucose control was associated with lower prevalence of DPN.^{36–39} In support of this notion, in the current study, we found an association between DPN and blood glucose control as reflected by the fasting blood glucose level, HbA1c and the type of treatment. Therefore, like others⁴⁰ the most effective approach to reduce the progression to neuropathy is aggressive treatment of diabetes hyperglycemia towards normoglycemia.

Our finding that multiple risk factors are associated with the development of DPN in diabetic patients should direct the attention towards the link between these risk factors and the increase in diabetes-related complications, morbidity as well as mortality.^{40,41} It was also clear that age is a significant risk factor that predisposes to DPN and this finding was observed by others.^{1,20–22} In the present study the vast majority of patients with DPN were type 2 diabetes (90.4%). Similarly, Kästenbauer et al,²³ showed a higher percentage (16% vs 37.5%) of type 2 diabetic patients with symptoms of neuropathy compared with type 1 diabetic patients.²³ Other risk factors including BMI, number of comorbid diseases such as chronic kidney disease, hypertension, dyslipidemia, cerebral vascular stroke, peripheral vascular disease and diabetic ketoacidosis were also found to be significantly associated with DPN. The close association between DPN and modifiable cardiovascular risk factors (cardiometabolic disease)

including elevated triglyceride levels (hyperlipidemia), body mass index (BMI), smoking, hypertension and cardiovascular disease was documented repeatedly in previous studies.^{1,20,21,24,42} However, no specific mechanisms linking DPN with cardiovascular disease have been suggested in any of these studies.

It also worth adding that the findings of the present study are consistent with those reported from the neighboring Gulf state of Bahrain. The Bahraini study²² found that older age, poor glycemic control, longer duration of diabetes, elevated cholesterol levels, current smoking, obesity, large waist circumference, elevated triglycerides levels and hypertension but not gender were significant risk factors for DPN. However, the vast majority of the DPN patients in the study were females and were on insulin therapy. In contrast, there were no significant differences in gender distribution or treatment modalities when compared to controls. On the other hand, many other studies have reported a significant association between female gender and the prevalence of DPN.^{1,20,21,24}

In the multivariate model, we identified three risk factors that independently correlate with the presence of DPN. These were age, chronic kidney disease and peripheral vascular disease. These risk factors were closely associated with DPN and this association has been reported before.^{1,24,26,37–43}

However, we did not find any correlation between smoking and DPN.^{24,26,44} Nonetheless, the following diseases were found to be associated with DPN: PVD, kidney disease, hypertension, dyslipidemia, foot ulcer and amputation, which is in line with many earlier studies.^{15,37,39,44,45}

Our findings concur with those reported in recent published studies on the factors associated with DPN. These studies were noticeable multi-center studies of long-term follow-up of 20–25 years.^{46,47} Most of these studies confirmed that old age, longer diabetes duration, and elevated HbA1c were the commonest risk factors for the development of DPN.^{45–48}

We acknowledge several limitations in our study. The nature of study design as a retrospective study limits the accurate assessment of the association between DPN and risk factors. Therefore, prospective studies are needed to fully characterize the sequel of DPN in diabetes patients. Poor documentation of DPN diagnosis and supportive laboratories investigations (nerve conduction studies and electromyography) reduced our ability to access important data needed for the study and to confirm how DPN

diagnosis was made. Therefore, it is likely that some DPN patients were missed.

Conclusion

This study identified several risk factors that contributed to the development of DPN in Saudis. These must be considered in strategies and campaigns aimed at risk reduction of cardiovascular and chronic diseases, and consequently progression of DPN.

Ethics

The study was approved by the Institutional Review Board, King Abdullah International Medical Research Centre, King Saud bin Abdulaziz University for Health Sciences, National Guard Health Affairs. All patient data accessed complied with relevant data protection and privacy regulations. All procedures performed in the study involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Acknowledgments

We would like to acknowledge the help of the Health Informatics Department, KAMC.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Funding

There is no funding to report.

Disclosure

The authors report no conflicts of interest for this work.

References

- Hicks CW, Selvin E. Epidemiology of peripheral neuropathy and lower extremity disease in diabetes. *Curr Diab Rep*. 2019;19:86. doi:10.1007/s11892-019-1212-8.
- Shiferaw WS, Akalu YT, Work Y, Yared Asmare Aynalem YA. Prevalence of diabetic peripheral neuropathy in Africa: a systematic review and meta-analysis. *MBC Endoc Disord*. 2020;20:49–58. doi:10.1186/s12902-020-0534-5
- Alleman CJ, Westerhout KY, Hensen M, et al. Humanistic and economic burden of painful diabetic peripheral neuropathy in Europe: a review of the literature. *Diabet Res Clin Pract*. 2015;109:215–225. doi:10.1016/j.diabres.2015.04.031
- Ebata-Kogure N, Nozawa K, Murakami A, Toyoda T. Clinical and economic burdens experienced by patients with painful diabetic peripheral neuropathy: an observational study using a Japanese claims database. *PLoS One*. 2017;12(10):e0187250. doi:10.1371/journal.pone.0187250
- John J, Stamenitis S, Clouth J, et al. The quality of life and economic burden of neuropathy in diabetic patients in Germany in 2002—results from the Diabetic Microvascular Complications (DIMICO) study. *Diabetes Res Clin Pract*. 2008;81:223–230. doi:10.1016/j.diabres.2008.03.019
- Ugwu E, Adeleye O, Ezeani I. Predictors of lower extremity amputation in patients with diabetic foot ulcer: findings from MEDFUN, a multi-center observational study. *J Foot Ankle Res*. 2019;12:34. doi:10.1186/s13047-019-0345-y
- Kuntzer T, Medlin F, Burnand B, Camain JY. Diabetic neuropathies: clinical sub-types, early detection, and asking help from neurologist. *Praxis (Bern 1994)*. 2012;101(20):1315–1319. doi:10.1024/1661-8157/a001037
- Lazzarini PA, Damien Clark D, Derhy PH. What are the major causes of lower limb amputations in a major Australian teaching hospital? The Queensland Diabetic Foot Innovation Project, 2006 – 2007. *J Foot Ankle Res*. 2011;4(Suppl1):O24. doi:10.1186/1757-1146-4-S1-O24
- Yagihashi S, Mizukami H, Sugimoto K. Mechanism of diabetic neuropathy: where are we now and where to go? A review. *J Diabetes Invest*. 2011;2:18–32. doi:10.1111/j.2040-1124.2010.00070
- Boulton A, Malik R, Arezzo J, Sosenko J. Diabetic somatic neuropathies. *Diabetes Care*. 2004;27(6):1458–1486. doi:10.2337/diacare.27.6.1458
- Won J, Kwon H, Kim C, et al. Prevalence and clinical characteristics of diabetic peripheral neuropathy in hospital patients with Type 2 diabetes in Korea. *Diabetic Med*. 2012;29(9):e290. doi:10.1111/j.1464-5491.2012.03697.x
- Boronikolos GC, Menge BA, Schenker N, et al. Upper gastrointestinal motility and symptoms in individuals with diabetes, prediabetes and normal glucose tolerance. *Diabetologia*. 2015;58(6):1175–1182. doi:10.1007/s00125-015-3538-3
- Callaghan BC, Little AA, Feldman EL, Hughes RA. Enhanced glucose control for preventing and treating diabetic neuropathy. *Cochrane Database Syst Rev*. 2012;6:CD007543.
- Tesfaye S, Stevens L, Stephenson J, et al. Prevalence of diabetic peripheral neuropathy and its relation to glycaemic control and potential risk factors: the EURODIAB IDDM Complications Study. *Diabetologia*. 1996;39(11):1377–1384. doi:10.1007/s001250050586
- Tesfaye S, Chaturvedi N, Eaton SE, et al. Vascular risk factors and diabetic neuropathy. *N Engl J Med*. 2005;352(4):341–350. doi:10.1056/NEJMoa032782
- Meo SA, Usmani AM, Qalbani E. Prevalence of type 2 diabetes in the Arab world: impact of GDP and energy consumption. *Europ Med Pharmacol Sci*. 2017;21:1303–1312.
- Algeffari MA. Painful diabetic peripheral neuropathy among Saudi diabetic patients is common but under-recognized: multicenter cross-sectional study at primary health care setting. *Family Community Med*. 2018;25(1):43–47.
- Sendi RA, Mahrus A, Saeed RM, et al. Diabetic peripheral neuropathy among Saudi diabetic patients: a multicenter cross-sectional study at primary health care setting. *J Family Med Prim Care*. 2020;9(1):197–201. doi:10.4103/jfmpc.jfmpc_927_19
- American Diabetes Association. “Diagnosis and classification of diabetes mellitus.”. *Diabetes Care*. 2010;33(supplement 1):S62–S69. doi:10.2337/dc10-S062

20. Yang H, Sloan G, Ye Y, Wang S. New perspective in diabetic neuropathy: from the periphery to the brain, a call for early detection, and precision medicine. *Front Endocrinol.* 2020;10. doi:10.3389/fendo.2019.00929
21. Dyck PJ, Kratz KM, Karnes JL, et al. The prevalence by staged severity of various types of diabetic neuropathy, retinopathy, and nephropathy in a population-based cohort: the Rochester diabetic neuropathy study. *Neurology.* 1993;43:817–824. doi:10.1212/WNL.43.4.817
22. Al-Mahroos F, Al-Roomi K. Diabetic neuropathy, foot ulceration, peripheral vascular disease and potential risk factors among patients with diabetes in Bahrain: a nationwide primary care diabetes clinic-based study. *Ann Saudi Med.* 2007;27(1):25.
23. Albers JW, Pop-Busui R. Diabetic neuropathy: mechanisms, emerging treatments, and subtypes. Review. *Curr Neurol Neurosci Rep.* 2014;14:473. doi:10.1007/s11910-014-0473-3
24. Pop-Busui R, Boulton AJM, Feldman EL, Bril V. Diabetic Neuropathy: A Position Statement by the American Diabetes Association. *Diabetes Care.* 2017;40:136–154. doi:10.2337/dc16-2042
25. Iqbal Z, Azmi S, Yadav R, et al. Diabetic peripheral neuropathy: epidemiology, diagnosis, and pharmacotherapy. *Clin Ther.* 2018;40:828–849. doi:10.1016/j.clinthera.2018.04.001
26. Ziegler D, Papanas N, Vinik AI, Shaw JE. Epidemiology of polyneuropathy in diabetes and prediabetes. *Handb Clin Neurol.* 2014;126:3–22. doi:10.1016/B978-0-444-53480-4.00001
27. Pirart J. [Diabetes mellitus and its degenerative complications: a prospective study of 4400 patients observed between 1947 and 1973 (3rd and last part) (author's transl)]. *Diabete Metab.* 1977;3:2–4. French.
28. Wang DD, Bakhotmah BA, Hu FB, Alzahrani HA. Prevalence and correlates of diabetic peripheral neuropathy in a Saudi Arabic population: a cross-sectional study. *PLoS One.* 2014;9:e106935. doi:10.1371/journal.pone.0106935
29. Börü UT, Alp R, Sargin H, et al. Prevalence of peripheral neuropathy in type 2 diabetic patients attending a diabetes center in Turkey. *Endocr J.* 2004;51:563–567.
30. Lu B, Yang Z, Wang M, et al. High prevalence of diabetic neuropathy in population-based patients diagnosed with type 2 diabetes in the Shanghai downtown. *Diabetes Res Clin Pract.* 2010;88(2010):289–294. doi:10.1016/j.diabres.2010.02.002
31. Amour AA, Chamba N, Kayandabila J, Lyaruu IA. Prevalence, patterns, and factors associated with peripheral neuropathies among diabetic patients at tertiary hospital in the Kilimanjaro region: descriptive cross-sectional study From North-Eastern Tanzania. *Int J Endocrinol.* 2019;2019:5404781. doi:10.1155/2019/5404781
32. Day JL, Humphreys H, Alban-Davies H. Problems of comprehensive shared diabetes care. *Br Med J.* 1987;294:1590–1592. doi:10.1136/bmj.294.6587.1590
33. Tahrani AA, Altaf QA, Piya MK, Barnett AH. Peripheral and autonomic neuropathy in south asians and white caucasians with type 2 diabetes mellitus: possible explanations for epidemiological differences. *J Diabetes Res.* 2017;2017:1273789. doi:10.1155/2017/1273789
34. Fadavi H, Tavakoli M, Foden P, et al. Explanations for less small fibre neuropathy in South Asian versus European subjects with type 2 diabetes in the UK. *Diabetes Metab Res Rev.* 2018;34:e3044. doi:10.1002/dmrr.3044.
35. Callaghan BC, Gao L, Li Y, et al. Diabetes and obesity are the main metabolic drivers of peripheral neuropathy. *Ann Clin Transl Neurol.* 2018;5:397–405. doi:10.1002/acn3.531.
36. Ang L, Jaiswal M, Martin C, et al. Glucose control and diabetic neuropathy: lessons from recent large clinical trials. *Curr Diab Rep.* 2014;14(528). doi:10.1007/s11892-014-0528-7
37. Harris M, Eastman R, Cowie C. Symptoms of sensory neuropathy in adults with NIDDM in the U.S. population. *Diabetes Care.* 1993;16:1446–1452. doi:10.2337/diacare.16.11.1446
38. Al-Kaabi JM, FAI M, Zoubeidi T, et al. Prevalence and determinants of peripheral neuropathy in patients with type 2 diabetes attending a tertiary care center in the United Arab Emirates. *J Diabetes Metab.* 2014;5(346):2.
39. Cameron NE, Eaton SE, Cotter MA, Tesfaye S. Vascular factors and metabolic interactions in the pathogenesis of diabetic neuropathy. *Diabetologia.* 2001;44:1973–1988. doi:10.1007/s001250100001
40. Andersen ST, Witte DR, Dalsgaard EM, et al. Risk factors for incident diabetic polyneuropathy in a cohort with screen-detected type 2 diabetes followed for 13 years: ADDITION-denmark. *Diabetes Care.* 2018;41:1068–1075. doi:10.2337/dc17-2062.
41. Abbott CA, Malik RA, van Ross ER, et al. Prevalence and characteristics of painful diabetics. Explanations for less small fibre neuropathy in a large community-based diabetic population South Asian versus European subjects with type 2 diabetes in the UK. *Diabetes Care.* 2011;34(10):2220–2224. doi:10.2337/dc11-1108
42. Yang CP, Lin CC, Li CI, et al. Cardiovascular risk factors increase the risks of diabetic peripheral neuropathy in Patients with type 2 diabetes mellitus: the Taiwan diabetes study. *Medicine (Baltimore).* 2015;94:e1783. doi:10.1097/MD.0000000000001783
43. Kiani J, Moghimbeigi A, Azizkhani H, Kosarifar S. The prevalence and associated risk factors of peripheral diabetic neuropathy in Hamedan, Iran. *Arch Iran Med.* 2013;16:17–19.
44. Kästenbauer T, Irsigler P, Sauseng S, Grimm A, Prager R. The prevalence of symptoms of sensorimotor and autonomic neuropathy in Type 1 and Type 2 diabetic subjects. *J Diabetes Complications.* 2004;18:27–31. doi:10.1016/S1056-8727(03)00071-0
45. Coppini DV, Bowtell PA, Weng C, Young PJ, Sonksen PH. Showing neuropathy is related to increased mortality in diabetic patients a survival analysis using an accelerated failure time model. *J Clin Epidemiol.* 2000;53:519–523. doi:10.1016/S0895-4356(99)00170-5
46. Braffett BH, Gubitosi-Klug RA, Albers JW, et al. Risk Factors for diabetic peripheral neuropathy and cardiovascular autonomic neuropathy in the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Study. The DCCT/EDIC and Research Group*. *Diabetes.* 2020;69:1000–1010. doi:10.2337/db19-1046
47. Pfannkuche A, Alhajjar A, Ming A, et al. Prevalence and risk factors of diabetic peripheral neuropathy in a diabetics cohort: register initiative “diabetes and nerves”. *Endoc Metab Sci.* 2020;1(1–2):100053. doi:10.1016/j.endmts.2020.100053
48. Pan Q, Li Q Q, Deng W, et al. Prevalence of and risk factors for peripheral neuropathy in Chinese patients with diabetes: a multicenter cross-sectional study. *Front Endocrinol.* 2018;9:Article 617. doi:10.3389/fendo.2018.00617

International Journal of General Medicine**Dovepress****Publish your work in this journal**

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>