Condom Utilization and Affecting Factors Among People Living with HIV/AIDS Attending ART Clinics in Addis Ababa, Ethiopia

This article was published in the following Dove Press journal:
HIV/AIDS - Research and Palliative Care

Rahel Hailu Geleta¹
Mesafint Abeje Tiruneh²

¹Nefas Silk Lafto Sub City Woreda 5
Health Center, Addis Ababa, Ethiopia;

²Bethzatha General Hospital, Addis
Ababa, Ethiopia

Background: Human immune deficiency virus or acquired immune deficiency syndrome (HIV/AIDS) has been a major public health problem globally as well as in Africa including Ethiopia. To prevent HIV effectively, condoms must be used regularly and consistently. When HIV positive individuals have sex without condom, they can easily transmit the virus to a seronegative partner. Condom utilization may be affected by various factors. Therefore, the findings from this study will give an opportunity to understand condom utilization and the barriers to condom use.

Objective: To assess condom utilization and associated factors among people living with HIV/AIDS attending anti-retro viral treatment (ART) clinics in Addis Ababa, Ethiopia.

Materials and Methods: An institution-based cross-sectional study was conducted among 677 people living with HIV/AIDS attending ART clinics in Addis Ababa from February 1, 2019 to March 30, 2019. The sampling technique was three-stage sampling and finally, the study participants were selected using a systematic random sampling technique. Data were coded, cleaned and entered in to EPIData version 3.4, and analyzed using SPSS version 23.0. The association between the dependent variable and independent variables was analyzed using binary logistic regression analysis at 5% level of significance.

Results: About 677 HIV/AIDS patients receiving ART at health centers participated in the study. From the total participants, 306 (45.2%) utilized a condom in the last six months. Sex (adjusted odds ratio (AOR)=2.02; 95%CI=1.34–3.05), marital status (AOR=0.39; 95%CI=0.19–0.77 and (AOR=0.48; 95%CI=0.24–0.94), employment status (AOR=0.27; 95%CI=0.12–0.59), type of partner (AOR=0.15; 95%CI=0.05–0.48 and AOR=0.46; 95%CI=0.27–0.79), condom access (AOR=3.97; 95%CI=2.04–7.75), disclosure of HIV status (AOR=0.21; 95%CI=0.14–0.33) and attitude towards condoms (AOR=3.76; 95%CI=2.49–5.65) had significant association with condom utilization.

Conclusion: Condom utilization was found to be low among people living with HIV/AIDS attending ART clinics in Addis Ababa. This finding indicates that high concern shall be given to minimize the gap. Sex, marital status, employment status, disclosure of HIV status, condom access, type of partner and attitude towards condoms were significantly associated with condom utilization. Hence, continuous condom provision, promotion, demonstration, and awareness creation are vital.

Keywords: condom utilization, HIV/AIDS, anti-retroviral therapy, Addis Ababa, Ethiopia

Correspondence: Rahel Hailu Geleta
Nefas Silk Lafto Sub City Woreda 5
Health Center, Addis Ababa, Ethiopia
Tel +251913551027
Email rahelhailu2002@gmail.com

Introduction

Globally, HIV epidemic threats to public health were recognized 30 years ago.¹ Seventy-six point one million people became infected with HIV and 35.0 million

people died due to HIV/AIDS since the start of the epidemic.² Thirty-six point seven million people were living with HIV, 1.8 million people became newly infected with HIV and one million people died from AIDS 2016 globally.² Young women aged 15–24 years are at particularly high risk of HIV infection among adults globally in 2015.³ Globally about 45 million HIV infections have been prevented through condom use since 1990 globally.⁴ In 2015, an estimated 1.9 million adults were newly infected with HIV, most of them were infected through sexual intercourse.⁵

Nineteen point four million people were living with HIV in Eastern and Southern Africa in 2016.² In Sub-Saharan Africa, HIV/AIDS has been a major health problem for more than 20 years. Even if Ethiopia has one of the lowest HIV prevalence rates in East Africa, still there are more than one million people living with HIV.⁶ Over the last decade free anti-retro viral treatment (ART) services scale-up in the country has been one of the achievements of the HIV/AIDS program. The scale-up of ART program services have substantially decreased deaths and contributed to reduce incidence of HIV since the initiation of the ART program in 2005.⁷

Even though there is progress, the benefits of condom utilizations are not fully being used.¹ Condoms must be used consistently and correctly to achieve the high level of protection against HIV.⁸ Epidemiologic and laboratory studies indicates a condom is effective for HIV and sexually transmitted diseases (STD) prevention.⁹ Inconsistent or incorrect use of condoms usually leads to failure to protect against STD/HIV transmission.⁸ Transmission of HIV and STDs can occur with a single sex act with an infected partner due to inconsistent or nonuse of condoms. Incorrect use of condoms minimizes the effectiveness of condoms by leading to leakage, breakage, or slippage.⁹

Many people living with HIV have partners who have not acquired HIV, sero-discordant couples. Couple and partner testing, consistent condom use and providing ART to the partner living with HIV are key prevention strategies for sero-discordant couples.¹ Condom use is important even if both partners have HIV to prevent re-infection.¹⁰ Key populations including sex workers, people who inject drugs, transgender people, prisoners, and gay men and other men who have sex with men have higher risk of HIV infection.¹¹

Knowledge of HIV prevention methods is low in Ethiopia. Only 56% of women knew that HIV could be prevented using condoms and only 65% knew that HIV

could be prevented by limiting sex to one uninfected partner in 2011. Multiple sexual partners without using a condom increases the risk of contracting HIV.⁶ If HIV-positive individuals have sex without a condom, individuals with low CD4 count or higher viral load at the initiation of ART have the potential to transmit to a sero-negative sexual partner or they may be at risk of acquiring drug-resistant viral strains.¹²

Since unprotected sexual intercourse with an HIV-infected person easily transmits infection, inconsistent and incorrect use of condoms will increase the disease incidence and prevalence. In Ethiopia condom promotion and distribution is emphasized for the general population rather for PLWHA, therefore, data regarding condom use are hardly found. Despite the fact that condoms are the most effective method to fight against HIV/AIDS there is little information regarding its magnitude and the factors affecting condom utilization among PLWHA.¹³

Data regarding condom utilization among PLWHA are rare to find in Ethiopia. Therefore, this study tries to assess condom utilization among PLWHA receiving ART and the factors that affect their utilization in Addis Ababa such as sociodemographic factors, knowledge and attitude related factors, condom access, type of partner, disclosure of HIV status, duration on ART and substance use. The government offices and nongovernmental organizations (NGOs) working in this area will gain insight from the findings that may help them identify the gaps and plan appropriate intervention methods to improve condom utilization while increasing its provision and making it acceptable among PLWHA.

Materials and Methods

Study Design and Setting

Institution-based cross-sectional study was conducted in Addis Ababa, Ethiopia from February, 1, 2019 to March 30, 2019. According to the last national census in 2007, the city has an estimated total population of 3,384,569. There are 10 sub-cities in Addis Ababa. The city accompanied 93 health centers within the 10 sub-cities which is divided by woredas and majority of them provide ART service. It has the second highest HIV/AIDS prevalence rate in the country 3(.4%), according to the 2018 Federal HIV/AIDS Prevention and Control Office report.¹⁴

Source and Study Population

The source population were PLWHA attending ART clinics of health centers in Addis Ababa and the study

population were PLWHA attending ART clinics of selected health centers in Addis Ababa. PLWHA aged above 18 years who had been on ART for at least one month prior to the data collection period were included in the study. Critically ill and those who were unable to respond were excluded from the study.

Sample Size Determination and Sampling Procedure

Double population proportion formula was used to determine the sample size by considering 95%CI, 80% power and one-to-one ratio based on a study conducted in North West Ethiopia, 2015¹³ by Epi info statistical calculation. Design effect, 1.5 and 10% nonresponse rate were considered and the final sample size was 677.

Three-stage sampling technique was used. There are 10 sub-cities in the city from these three sub-cities were selected using a simple random sampling technique. The selected three sub-cities had 17 health centers which provide ART service. Six health centers were selected from the two sub-cities (three from each) and two health centers were selected from the third sub-city using simple random sampling technique proportion to the number of health centers; a total of eight health centers were selected. A systematic random sampling technique was used to select the study participants who were interviewed after the first random start proportional to the number of patients in each health center.

Operational Definitions

Condom utilization: whether the study participants used condoms regularly and consistently in the last six months prior to the study.

Knowledge about condom use:

Knowledgeable: study participants who answered at least five of the seven knowledge-related questions about condoms.

Not knowledgeable: study participants who answered less than five of the seven knowledge-related questions about condoms.

Attitude toward condom use:

Favorable attitude: study participants who answered two of the three attitude-related questions toward condom use.

Unfavorable attitude: study participants who answered less than two of the three attitude-related questions toward condom use.

Knowledge about HIV/AIDS:

Knowledgeable: study participants who answered three of the five knowledge-related questions about HIV/AIDS transmission and prevention practice.

Not knowledgeable: study participants who answered less than three of the five knowledge-related questions about HIV/AIDS transmission and prevention practice.

Attitude towards HIV/AIDS:

Favorable attitude: study participants who answered at least one of the two attitude-related questions towards HIV/AIDS transmission and prevention practice.

Unfavorable attitude: study participants who answered none of the two attitude-related questions towards HIV/AIDS transmission and prevention practice.

Data Collection Tool, Procedure and Quality Assurance

A structured interviewer administered questionnaire was first prepared in English and then translated in to Amharic. The data was collected using the Amharic version questionnaire. The questionnaire was then back translated into English for data entry. In order to assure the quality of the data, training was given to three selected health care professional data collectors who were not working in those health centers where the study was conducted and pretest was done on 5% of the study participants who were not included in the actual study. The collected data were checked for completeness and consistency daily during the data collection period.

Data Management and Data Analysis

The data collected from each participant was coded and entered in to EpiData version 3.4 and then exported to SPSS version 23 computer software for analysis. Relevant descriptive statistics such as frequency and percent were carried out. To screen out potentially significant variables, univariate binary logistic regression analysis was done, and those with less than 25% level of significance remained in the final models. To check the adequacy of the final model Hosmer–Lemeshow goodness of fit test was used ($p=0.08$). Multicollinearity test was done and no multicollinearity identified. Sensitivity analysis done and no data bias detected. Finally, a stepwise multiple binary logistic regression analysis was done. Adjusted odds ratio and 95%CI was employed to examine associations. Variables with p -value <0.05 were considered as statistically significant.

Results

Sociodemographic Characteristics of the Study Population

The total study participants were 677 in number, with an age range of 20–65 years and the mean age was 38.5 (SD ± 5.5) years. There were more females constituting 415 (61.3%) of the total study participants. The majority of the participants 453 (66.9%) were orthodox Christians. From the total participants 232 (34.3%) were married. More than one-third of the study participants 321 (47.4%) were self-employed while one-third of the participants 226 (33.4%) had completed secondary education and 116 (17.1%) had no formal education (Table 1).

Knowledge and Attitude About HIV/AIDS, Condoms and their Utilization

The majority, 469 (69.3%) of the study participants were knowledgeable about HIV/AIDS transmission and prevention practice. About 534 (78.9%) of the total participants had a favorable attitude toward HIV/AIDS prevention practice. About 479 (70.8%) of the total respondents were knowledgeable about condoms and 358 (52.9%) respondents had a favorable attitude toward condom use (Table 2).

Condom Access and Type of Partner

About 517 (76.4%) of the total respondents had a regular sexual partner/spouse in the last six months while the rest 24 (3.5%) and 136 (20.1%) of the respondents engaged in sexual relationships with sex workers and multiple partners, respectively (Table 3).

Approximately 593 (87.6%) of the respondents had access to condoms and the rest 84 (12.4%) had not from the respondents who had no access to condoms the reasons stated were price 5 (6%), distance/nonavailability 20 (23.8%), preference/type of condom 1 (1.2%), fear of being seen when buying 20 (23.8%) and did not know where to find 38 (45.2%) (Figure 1).

Duration on ART, Substance Abuse and Disclosure About HIV Status

About 22 (3.2%) of the total respondents stayed on ART for <3 months while nearly half of them 356 (52.6%) stayed on ART for more than 36 months.

About 131 (19.4%) of the respondents had used at least one of the substances (alcohol, cigarette, or chat) while

Table 1 Sociodemographic Characteristics of Study Participants, Addis Ababa, 2019 (n=677)

Variables	Frequency	Percent (%)
Sex		
Female	415	61.3
Male	262	38.7
Age		
20–30	156	23.0
31–45	431	63.7
46–65	90	13.3
Religion		
Orthodox	453	66.9
Muslim	87	12.9
Protestant	131	19.4
Other	6	0.9
Marital status		
Single	190	28.1
Married	232	34.3
Divorced	162	23.9
Widowed	93	13.7
Educational status		
No formal education	96	14.1
Primary education	209	30.9
Secondary education	226	33.4
Tertiary education	146	21.6
Employment status		
Unemployed	143	21.1
Civil servant	142	21.0
Self-employed/business	321	47.4
Housewife	71	10.5

engaging in a sexual relationship. Regarding disclosure about HIV status, more than half 392 (57.9%) of the respondents did not inform their partners of their HIV status (Table 4).

Condom Utilization

Among the total study participants, 306 (45.2%, 95% CI=41.5–49.0%) reported that they used condoms regularly and consistently whereas the remaining 371 (54.8%, 95%CI: 51.0–58.5%) respondents did not used condoms regularly and consistently in the last six months.

Factors Associated with Condom Utilization

At a 25% level of significance univariate binary logistic regression analysis of sex, educational status, marital status,

Table 2 Knowledge and Attitude About HIV/AIDS, Condoms and Their Utilization Among Study Participants, Addis Ababa, 2019 (n=677)

Variables	Frequency	Percent (%)
Knowledge about HIV/AIDS		
Knowledgeable	469	69.3
Not knowledgeable	208	30.7
Attitude about HIV/AIDS		
Favorable attitude	534	78.9
Un favorable attitude	143	21.1
Knowledge about condoms		
Knowledgeable	479	70.8
Not knowledgeable	198	29.2
Attitude about condoms		
Favorable attitude	358	52.9
Unfavorable attitude	319	47.1

Abbreviation: HIV/AIDS, human immune deficiency virus or acquired immune deficiency syndrome.

Table 3 Condom Access and the Type of Partner Among Participants, Addis Ababa, 2019 (n=677)

Variable	Frequency	Percent (%)
Condom access		
Yes	593	87.6
No	84	12.4
Type of partner		
Regular partner/spouse	517	76.4
Sex worker	24	3.5
Multiple partners	136	20.1

employment status, condom access, knowledge about HIV, attitude toward condoms, type of partner and disclosure of HIV status were significantly associated with condom utilization. However, at a 5% level of significance multivariable binary logistic regression analysis of sex, marital status, employment status, type of partner, condom access, disclosure of HIV status and attitude toward condoms were significantly associated with condom utilization.

Male respondents were 2.02 times more likely to utilize condom compared to female respondents (AOR=2.02; 95%CI=1.34–3.05). Respondents who were married and widowed were 61% and 52% less likely to utilize condom, respectively when compared to those who were single (AOR=0.39; 95%CI=0.19–0.77 and (AOR=0.48; 95%CI=0.24–0.94)

Similarly, respondents who were self-employed/business were 73% less likely to utilize condom compared to

Figure 1 Reason for nonaccessibility of condom among respondents, Addis Ababa, 2019 (n=677).

unemployed respondents (AOR=0.27; 95%CI=0.12–0.59). Respondents who had multiple sexual partners and who had sex with sex workers were 85% and 54% less likely to utilize condoms compared to those who had regular sexual partner, respectively (AOR=0.15; 95%CI=0.05–0.48 and AOR=0.46; 95%CI=0.27–0.79).

Table 4 Duration on ART, Substance Abuse, Disclosure About HIV Status Among Participants, Addis Ababa, 2019 (n=677)

Variables	Frequency	Percent (%)
Duration on ART		
<3 months	22	3.2
3–12 months	112	16.5
12–36 months	187	27.6
>36 months	356	52.6
Substance abuse		
Yes	131	19.4
No	546	80.6
Disclosure about HIV status		
Yes	285	42.1
No	392	57.9

Abbreviations: HIV, human immune deficiency virus; ART, anti-retroviral therapy.

Respondents who had access to condoms were 3.97 times more likely to utilize them compared to respondents who had no access (AOR=3.97; 95%CI=2.04–7.75). Respondents who did not disclose their HIV status to their partners were 79% less likely to utilize condoms compared to respondents who disclosed their status (AOR=0.21; 95%CI=0.14–0.33).

Respondents who had a favorable attitude toward condoms were 3.76 times more likely to utilize

condoms than those who had an unfavorable attitude (AOR=3.76; 95%CI=2.49–5.65) (Table 5).

Discussion

The finding of this study showed that the proportion of condom utilization is 45.2%, this finding is similar to the study of condom utilization among people living with HIV/AIDS attending ART clinic in North Central Nigeria

Table 5 Factors Associated with Condom Utilization Among PLWHA Attending ART Clinics in Addis Ababa, 2019 (n=677)

Variables	Condom Utilization		COR (95%CI)	AOR (95%CI)	p-value
	Yes	No			
Sex					
Female	157	258	1.00	1.00	
Male	113	149	1.72 (1.26–2.35)	2.02 (1.34–3.05)	0.001*
Educational status					
No formal education	39	57	1.00	1.00	
Primary education	77	132	1.72 (1.02–2.90)	1.69 (0.85–3.37)	0.136
Secondary education	111	115	2.02 (1.32–3.10)	1.33 (0.78–2.28)	0.299
Tertiary education	79	67	1.22 (0.80–1.85)	1.33 (0.79–2.25)	0.280
Marital status					
Single	98	92	1.00	1.00	
Married	100	132	0.48 (0.29–0.81)	0.39 (0.19–0.77)	0.007*
Divorced	78	84	0.69 (0.42–1.14)	1.14 (0.58–2.28)	0.701
Widowed	30	63	0.61 (0.36–1.03)	0.48 (0.24–0.94)	0.031*
Employment status					
Unemployed	53	90	1.00	1.00	
Civil servant	80	62	0.74 (0.40–1.36)	0.68 (0.32–1.47)	0.328
Self-employed	150	171	0.29 (0.16–0.54)	0.27 (0.12–0.59)	0.001*
Housewife	23	48	0.32 (0.32–0.96)	0.55 (0.28–1.11)	0.096
Condom access					
Yes	292	301	4.07 (2.30–7.18)	3.97 (2.04–7.75)	<0.001*
No	14	70	1.00	1.00	
Knowledge about HIV					
Knowledgeable	217	252	1.29 (0.93–1.79)	1.39 (0.89–2.17)	0.143
Not knowledgeable	89	119	1.00	1.00	
Attitude about condom					
Favorable attitude	219	139	3.67 (2.66–5.06)	3.76 (2.49–5.65)	<0.001*
Unfavorable attitude	87	223	1.00	1.00	
Type of partner					
Spouse/regular partner	252	265	1.00	1.00	
Sex worker	18	6	0.47 (0.31–0.70)	0.46 (0.27–0.79)	0.005*
Multiple attitude	36	100	0.18 (0.07–0.46)	0.15 (0.05–0.48)	0.001*
Disclosure of HIV status					
Yes	178	107	1.00	1.00	
No	128	264	0.29 (0.21–0.40)	0.21 (0.14–0.33)	<0.001*

Note: *Statistically significant at 5% level of significance.

Abbreviations: PLWHA, people living with HIV/AIDS; ART, anti-retroviral therapy.

(45.5%).¹⁵ However, this finding is quite different from the study on consistent condom use in HIV/AIDS patients receiving ART in northwest Ethiopia (78.9%).¹³

This study showed that consistent condom use is affected by sex. Male respondents were 2.02 times more likely to utilize condom than female respondents. This finding is similar to the studies conducted in North Central Nigeria, in Northwest Ethiopia and in the Asia-Pacific Region.^{15,16} This might be due to the fact that women are less powerful to negotiate condom use and have less ability to decide on sexual life especially in developing countries. Women also involve in risky sexual behaviors due to economic problems and they might be forced not to use condoms either due to extra payment or inability to negotiate.

This study also showed that there is a significant association between disclosure of HIV status and condom utilization. Respondents who did not disclose their HIV status to their partners were 79% less likely to utilize condoms than respondents who disclosed their status. This finding is consistent with the study conducted among PLWHA in Maiduguri, Nigeria.¹⁷ This shows that individuals do not bother about knowing their partners HIV status and they may have fear that if their partners are seronegative, they might leave them and remain alone or not to be blamed by their partners if they become positive after the relationship.

Respondents who were married and widowed were 61% and 52% less likely to utilize condoms, respectively when compared to those who were single and this finding is in agreement with a study conducted in Maiduguri, Nigeria.¹⁷ This may be due to the perception that a condom is unnecessary among married couples, partners refusal of condom, the desire to have children and less empowerment for women.

There is also an association between occupational status and condom utilization. Respondents who were self-employed/business were 73% less likely to utilize condoms compared to unemployed respondents. Respondents who had multiple sexual partners and who had sex with sex workers were 85% and 54% less likely to utilize condoms compared to those who had a regular sexual partner, respectively. This indicates that despite the efforts made through counseling to decrease HIV transmission and re-infection people are still involved in risky sexual behaviors which could result in further spread of HIV infection in the society and treatment failure in PLWHA.

Respondents who had access to condoms were 3.97 times more likely to utilize them compared to respondents

who had no access. This finding is supported by a systematic review done in the Asia-Pacific region.¹⁶ This might be due to individuals do not knowing where to find condoms, being ashamed to buy condoms openly or nonavailability (the condoms from clinics may not sustain them till the next visit).

Respondents who had a favorable attitude toward condoms were 3.76 times more likely to utilize condoms than those who had an unfavorable attitude. This may be due to the misperception that condoms decrease sexual pleasure and being in ART totally prevents HIV.

Limitation of the Study

Although this study provides important information regarding condom utilization and its determinant factors, which is relevant to make interventions, it may have certain limitations of recall and socially desirable bias.

Conclusion

Condom utilization was found to be low among people living with HIV/AIDS attending ART clinics in Addis Ababa. This finding indicates that high concern shall be given to minimize the gap. Sex, marital status, employment status, disclosure of HIV status, condom access, type of partner and attitude toward condoms were significantly associated with condom utilization. Therefore, continuous condom provision, promotion, demonstration and awareness creation are vital. Community based education and sensitization on HIV/AIDS and condom use has to be a major concern of stakeholders. Health-care providers should create awareness among PLWHA that condom use is the key intervention to prevent HIV transmission and reinfection including its negative outcome and the need to use condoms even with in marriage.

Data Sharing Statement

All relevant data are available within the manuscript.

Ethics Approval and Consent to Participate

As our institutions do not have the mandate to provide ethical clearance for research, ethical clearance was obtained from GAMBY Medical and Business College and Addis Ababa Health Bureau. This study was conducted under technical support and supervision of GAMBY Medical and Business College. The findings of the study were also presented to the College and approved. The study was

conducted in accordance with the Declaration of Helsinki. The study participants were informed that participation was voluntarily and written informed consent was taken from the study participants by telling that they have the right not to respond. This study had no risk for study participants. Confidentiality of the collected data was maintained using a code number instead of their name.

Consent for Publication

Consent for publication was taken from the study participant during data collection.

Acknowledgments

Authors would like to thank all study participants, data collectors and GAMBY Medical and Business College and Addis Ababa Health Bureau.

Author Contributions

Both authors meet the ICMJE criteria for co-authorship, providing substantial intellectual contributions for the manuscript. Both authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

The authors received no specific funding for this research.

Disclosure

The authors report no conflicts of interest in this work.

References

1. World Health Organization. *Global Update on the Health Sector Response to HIV*. 2014.
2. United Nations Program for HIV/AIDS. *Global HIV Statistics Fact Sheet-World AIDS Day*. 2017.
3. United Nations Program for HIV/AIDS. *Global AIDS Update*. 2016.
4. Stover J, Avenir H. *The Contribution of Condoms to HIV Prevention. Presentation Made at Global Condom Steering Group Meeting*. 2016.
5. Lori N, Jane R, Stephen V, et al. Global estimates of the prevalence and incidence of four curable sexually transmitted infections in 2012 based on systematic review and global reporting. *PLoS One*. 2015;10(12):1–17.
6. Central Statistical Agency (CSA) [Ethiopia] and ICF. *Ethiopia Demographic and Health Survey 2011*. Addis Ababa, Ethiopia, and Rockville, Maryland, USA: CSA and ICF; 2011.
7. Federal Democratic Republic of Ethiopia. *Country Progress Report on the HIV Response*. 2014.
8. United States Aid for International Development. *Condom Fact Sheet*. 2015.
9. Center for Communicable Disease control. *Condom Fact Sheet in Brief.department of Health and Human Service*. 2013.
10. UNFPA, WHO, PATH. *Condom Programming for HIV Prevention an Operational Manual for Programme Managers*. 2005.
11. United Nations Program for HIV/AIDS. *Prevention Gap Report*. 2016.
12. Donnell D, Baeten J, Kiarie J. Partners in prevention HSV/HIV transmission study team. Heterosexual HIV-1 transmission after initiation of antiretroviral therapy: a prospective cohort analysis. *Lancet*. 2010;375(9731):2092–2098. doi:10.1016/S0140-6736(10)60705-2
13. Zewdneh S, Befikadu L, Bayew T, Akshaya S, Abyot E. Consistent condom use in HIV/AIDS patients receiving antiretroviral therapy in Northwestern Ethiopia: implication to reduce transmission and multiple infections. *HIV AIDS*. 2015;7:119–124.
14. Ethiopian Federal HIV/AIDS Prevention and Control Office. *Annual Report*. 2018.
15. Adekunle G, Salaudeen OI, Yusuf A, et al. Condom use among people living with HIV/AIDS attending Abejokolo General Hospital in Kogi State, North Central Nigeria. *Ann Afr Med*. 2014;13(3):99–103. doi:10.4103/1596-3519.134378
16. Keshab D, Verena K, Sushil K, Anna M. Condom use behaviour among people living with HIV: a seven-country community-based participatory research in the Asia-Pacific region. *BMJ*. 2017;1–8.
17. Ballah A, Sasilu A, David B, Umaru I, Babajide B. Evaluation of condom use and associated factors adult HIV clients in Maiduguri, North Eastern Nigeria: a comparative cross sectional study. *World J AIDS*. 2014;4(02):169–177. doi:10.4236/wja.2014.42022

HIV/AIDS - Research and Palliative Care

Dovepress

Publish your work in this journal

HIV/AIDS - Research and Palliative Care is an international, peer-reviewed open-access journal focusing on advances in research in HIV, its clinical progression and management options including antiviral treatment, palliative care and public healthcare policies to

control viral spread. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/hivaid—research-and-palliative-care-journal>