Validity of Anthropometric Cut-Offs for Early Diagnosis of Dyslipidemia Among Ethiopian Adults

This article was published in the following Dove Press journal:
Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Sileshi Kenate¹
Temamen Tesfaye²
Solomon Berhanu Mogas³
Belay Zawdie⁴
Yonas Tesfaye²
Lelisa Sena Dadi³
Mulualem Tadesse⁵
Ayantu Kebede³
Esayas Kebede Gudina⁶
Dessalegn Tamiru⁴

¹Jimma Town Health Office, Jimma, Ethiopia; ²School of Nursing, Jimma University, Jimma, Ethiopia; ³Department of Epidemiology, Jimma University, Jimma, Ethiopia; ⁴Department of Nutrition and Dietetics, Jimma University, Jimma, Ethiopia; ⁵Department of Medical Laboratory Sciences, Jimma University, Jimma, Ethiopia; ⁶Department of Internal Medicine, Jimma University, Jimma, Ethiopia

Introduction: Lack of regional- and local-based cut-off points of lipid profile and/or anthropometric measurements remains one of the challenges in prevention, early detection and control of non-communicable diseases. This study aimed to validate anthropometric-based screening of lipid profiles to develop locally appropriate optimal cut-off points for metabolic syndrome screening.

Methods: A community-based cross-sectional study was conducted among randomly selected 977 adults in Jimma Town, Ethiopia. Data were collected using structured questionnaire, anthropometric and biochemical measurements. Data were analyzed using SPSS windows version 21 and Kappa statistic was used to validate the agreement between anthropometric measurement and lipid profile. A p-value of <0.05 was considered statistically significant.

Results: Body mass index (BMI) at ≥ 24.5 was used as screening of dyslipidemia (TG ≥ 150 mg/dl) with slight Kappa coefficient of 0.138 (P<0.001) among females while it was ≥ 22.2 among males with fair (0.275) Kappa coefficient (P<0.001). Waist circumference-based screening of dyslipidemia (TG ≥ 150 mg/dl) at ≥ 78.0 cm had negative (-0.005) Kappa coefficient (Pp<0.001) among females (sensitivity: 72.6% and specificity: 26.7%). Yet, waist circumference at ≥ 83.7 cm had slight Kappa coefficient of 0.13 (P<0.005) among males (sensitivity: 38% and specificity: 74.9%). Waist hip ratio-based screening of dyslipidemia (TG ≥ 150 mg/dl) at ≥ 0.82 among females had negative (-0.001) Kappa coefficient (p=0.763), whereas among males at ≥ 0.88 , there was a slight (0.105) Kappa coefficient of (p=0.002) (sensitivity: 77.5% and specificity: 36.8%). This study showed that anthropometric-based high-density lipoprotein measurement was not applicable.

Conclusion: This study indicated that BMI-based screening of triglyceride was more applicable for both sexes than other anthropometric measurements. Waist circumference and waist-to-hip ratio-based screening of triglyceride were slightly applicable only for males. However, anthropometric-based screening of high-density lipoprotein measurement was not applicable for both sexes. In conclusion, researchers and policy makers need to consider locally validated cut-off points to be used for screening metabolic syndrome in the community.

Keywords: metabolic syndrome, lipid profile, anthropometric measurements, Jimma

Correspondence: Dessalegn Tamiru
Department of Nutrition and Dietetics,
Jimma University, P. O. Box-378, Jimma,
Oromia Region, Ethiopia
Tel +251912373397
Email dessalegn97@gmail.com

Introduction

Metabolic syndrome is a subset of non-communicable diseases (NCDs), which is interconnected by physiological, biochemical, clinical and metabolic factors that directly increases the risk of cardiovascular disease and type 2 diabetes mellitus.

Insulin resistance, visceral adiposity, atherogenic dyslipidemia, endothelial dysfunction, genetic susceptibility, elevated blood pressure, hypercoagulable state and chronic stress are among several factors which constitute the syndrome.¹ Individuals with metabolic syndrome exhibit a characteristic pattern of abnormalities in serum lipid levels consisting of a low level of high density lipoprotein and an elevated level of triglyceride.²

Different studies show that the burdens of NCDs and their impact on the health of the population and health services are increasing in developing countries. Cardiovascular diseases, cancer, chronic respiratory diseases and diabetes are responsible for 82% of deaths. Currently, low income countries are witnessing epidemiological transition from infectious communicable diseases to chronic non-communicable diseases due to changes in lifestyles, rapid urbanization and diminishing levels of physical activity.^{3,4} Consequently, non-communicable diseases are predicted to increase globally by 17% and by 27% in the African region in the next 10 years, and projected to overtake communicable diseases as the major cause of morbidity in sub-Saharan Africa by the year 2030.³

The 2017 WHO report progress monitor also showed that NCDs kill 41 million people each year, which is 71% of all global deaths. In Ethiopia, the proportion of death from NCDs has been reported at 42% of all causes of deaths, which accounts for 27% of premature deaths before the age 70 years.⁵ The increment of NCDs on top of the preexisting highly prevalent communicable diseases, maternal, prenatal and nutritional conditions constrains the already meager health resources and hinders economic development of the country.⁶

However, it is only relatively recently that NCDs have received attention in low-income countries. Consequently, unrecognized/undiagnosed NCDs are found to be major problems as the individuals are unaware of their status in this regard. It has been indicated that the international anthropometric cutoff for detecting obesity is not appropriate for Ethiopians.⁷ There is a need to reorient the national health system to ensure recognition of the NCDs burden and improve organization and delivery of NCDs services at primary health care (PHC) level.^{8,9}

Lack of local based cut-off points remains one of the challenges in management, prevention, and control of NCDs, and their modifiable risk factors. Studies show that the international cut-off points for different anthropometric measurements and body composition are not

appropriate for the Ethiopian population; it leads to underestimation and misclassifies the risk of metabolic syndrome. Therefore, developing standardized local cut-off points for early detection of metabolic syndrome is crucial for prevention and management of NCDs. Thus, this study aimed to validate anthropometric cut-off points among Ethiopian adults⁷ through household level screening using waist circumference, waist-to-hip ratio and BMI measurement for estimation and classification of the risk of metabolic syndrome in Jimma Town, Oromia Regional State, Southwest Ethiopia.

Methods

Study Setting

A community-based cross-sectional study was conducted in Jimma Town from June to July 2019 among adults aged ≥ 18 years. Jimma is found in Southwest Ethiopia, located 352 km from the capital city of Addis Ababa. Sample size was calculated using sensitivity estimation formula¹⁰ taking highest prevalence of high density lipoprotein (23.4%) based on a study done in Addis Ababa among Ethiopian adults,¹¹ 5% margin of error, 95% confidence level and a 90% anticipated sensitivity.⁷ Finally, sample size of 977 was calculated by considering 10% non-response rate and 1.5 design effects. Study participants were selected using two-stage cluster sampling technique. Six kebeles (smallest administrative unit) out of 17 were randomly selected. Study participants were selected from each kebele proportionally based on the number of households in each selected kebele using systematic sampling technique.

Inclusion and Exclusion Criteria

All residents who lived in the town for six months prior to the data collection were included into the study. However, adults who had physical deformity (kyphosis and scoliosis), pregnancy, known chronic illness and serious illness were excluded.

Data Collection Procedure

Data were collected using WHO stepwise questionnaire and adapted to the local context. The survey tools included socio-demographic characteristics, anthropometric measurements and laboratory analyses of lipid profile. Height of the study participants was measured to the nearest 0.1 cm using a stadiometer (Seca GmbH & Co. KG, Germany) with the subjects positioned at the Frankfurt Plane and the four points (heel, calf, buttocks and shoulder) touching the

vertical stand and their shoes taken off. Before starting the measurements, the stadiometer was checked using calibration rods. Weight was measured using digital weight scale to the nearest 0.1 kg with the subjects wearing light clothes and shoes taken off. The validity of the scale was checked using an object of a known weight of 1kg.

Waist circumference was measured at midpoint between the inferior margin of the last rib and the iliac crest just at wider area using a stretch tape. Just before taking the measurement, participants were requested to stand with their feet together, place their arms at the side of their body with the palms of their hands facing inwards, and breathe out gently. Hip circumference was measured at the level of the greater trochanter of the femur with the subjects wearing pants. All anthropometric measurements were done in triplicate and the average value was used for further analysis. Standardization of anthropometric measurements were done to reduce inter observer error.

The laboratory parameter was determined according to the standard operating procedures. Five 5 mL venous blood samples were collected to determine participant's lipid profiles in mg/dL. Serum was carried out in ABX Pentra 400 Automated Chemistry Machine (Horiba ABX SAS, 34184) Montpellier¹² at Jimma medium Clinical chemistry Core laboratory to determine lipid profile including total cholesterol, high-density lipoprotein (HDL) and triglyceride (TG). Low density lipoprotein (LDL) level was calculated by using Freidwald formula.¹³

The optimal cut-off values of waist circumference (WC) was 85.3 cm for male and 78.0 cm for females and the values greater than this indicates risk factor for over-nutrition. The optimal cut-off values of waist-to-hip ratio (WHR) were calculated by dividing waist-to-hip in centimeters. The cut-off values were 0.88 cm for male and 0.82 cm for females; values greater than the cut-off value was considered at risk for over-nutrition. Similarly the optimal cut-off of values body mass index (BMI) was 24.5kg/m^2 for female and 22.2kg/m^2 for male; values greater than the cut off value was considered as at risk for over-nutrition. A lipid level was classified according to WHO classification and triglyceride was considered as normal when it was $<150\text{ mg/dl}$ and HDL-C was considered as normal if it is $>40\text{ mg/dl}$ male and $>50\text{mg/dl}$ for female.¹⁴

Data Quality Management

Data collectors were trained for four days before the actual data collection on interviewing approach, anthropometric measurement and data recording. To ensure the quality of

the data, a properly designed data collection instrument was used. Calibration of instruments was done to increase the reliability of the measurements. A questionnaire for the survey was first prepared in English language and translated into local languages (Afan-Oromo and Amharic), then back-translated into English to check for consistency. The questionnaire was pretested on 5% of sample size in Agaro Town and based on the pretest, additional adjustments were made based on identified gaps.

Data Processing and Analysis

Data were checked for completeness. The data were cleaned and checked for outliers and missing values. Epi Data version 3.1 and SPSS version 21 were used for data entry and analysis, respectively. The data were checked for normality using QQ-plot. Descriptive analysis was used to describe the study subjects and presented using tables and figures. Validity measures including sensitivity, specificity, positive predictive value, negative predictive value and kappa agreement were used to validate anthropometric based screening of lipid profile. The Kappa coefficient was used to measure the agreement between local anthropometric cut-off points and lipid profiles; and the kappa coefficient were classified as perfect ($k=0.81-1.00$), substantial ($k=0.61-0.80$), moderate ($k=0.41-0.60$), fair ($k=0.21-0.40$), slight ($k=0.01-0.20$) and no or poor agreement ($k=0$).^{7,15}

Results

A total of 915 individuals were included into this study with a 93.7% response rate. The mean age of study participants was 38.35 ± 13.51 years. A large proportion of the respondents (71.5%) were married. The majority of participants were Oromo ethnic (66.0%) and Muslim followers (56.0%). Close to three-fourths (64.5%) of the study participants attended at least primary school (Table 1).

BMI-based screening of triglyceride in both sexes had slight Kappa agreement among female and fair agreement among males. Using BMI at ≥ 24 for screening of hyper triglyceride ($\text{TG} \geq 150\text{mg/dl}$) among females showed slight (0.12) Kappa agreement ($P=0.024$) with 72.6% sensitivity and 65.0% specificity. Similarly, in case of male, BMI at $\geq 22.2\text{kg/m}^2$ based screening of dyslipidemia ($\text{TG} \geq 150\text{mg/dl}$) had fair (0.275) Kappa agreement ($P<0.001$) with 64.1% sensitivity and 66.8% specificity. Body mass index-based screening of high-density lipoprotein ($\text{HDL-C} < 50\text{mg/dl}$) among females at $\geq 24.5\text{kg/m}^2$ had negative (-0.003) Kappa coefficient with 58.0%

Table 1 The Socio-Demographic Characteristics of Adults in Jimma Town, Ethiopia

Variables	Categories	Number (n)	Percent (%)
Sex	Male	476	52.0
	Female	439	48.0
Age (years)	≤30	323	35.3
	31–40	301	32.9
	41–50	126	13.8
	≥51	165	18.0
Educational status	Degree and Above	79	8.6
	Diploma	96	10.5
	Illiterate	150	16.4
	Primary School	377	41.2
	Secondary School	213	23.3
Marital Status	Single	130	14.2
	Married	654	71.5
	Divorced	79	8.6
	Widowed	40	4.4
	Separated	12	1.3
Religion	Orthodox	290	31.7
	Muslim	512	56.0
	Protestant	111	12.1
	Catholic and	2 (0.2)	2 (0.2)
	Wakefata		
Ethnic Group	Oromo	604	66.0
	Amhara	81	8.9
	Dawuro	97	10.6
	Yemi	30	3.3
	Gurage	33	3.6
	Others*	70	7.7
Occupation	Unemployed	117	12.8
	Student	41	4.5
	Housewife	158	17.3
	Daily laborer	124	13.6
	Merchant	81	8.9
	Government	183	20.0
	Employee		
	Private employee	133	14.5
	Others ⁺	78	8.5

Notes: *Kafa, Tigre, Hadiya, Silte; ⁺Carpenter, Driver, Car Washer, No Specific Job.

sensitivity and 41.7% specificity, which is not statistically significant ($p=0.949$). Similarly, BMI at $\geq 22.2 \text{ kg/m}^2$ had a slight (0.032) Kappa coefficient ($P=0.449$) with 58.1% sensitivity and 45.45% specificity (Table 2).

Waist circumference based screening of dyslipidemia ($\text{TG} \geq 150 \text{ mg/dl}$) at ≥ 78.0 among females had -0.005 Kappa coefficient with 72.6% sensitivity and 26.7% specificity, which was not statistically significant ($p=0.877$).

On the other hand, for males, waist circumference at $\geq 83.7 \text{ cm}$ based screening of dyslipidemia ($\text{TG} \geq 150 \text{ mg/dl}$) had slight Kappa agreement of 0.13 ($p=0.005$) with 38.0% sensitivity and 74.9%, specificity.

Waist circumference based screening of high-density lipoprotein cholesterol ($< 50 \text{ mg/dl}$) at $\geq 78.0 \text{ cm}$ had slight Kappa coefficient (0.04) with 27.5% sensitivity and 77.3%, specificity which was not statistically significant ($p=0.263$). Similarly, for male waist circumference $\geq 83.7 \text{ cm}$ based screening of HDL-C had slight Kappa coefficient of 0.02, but not statistically significant (Table 3).

Waist circumference to hip ratio based screening of triglyceride was valid only for male at ≥ 0.88 cm with slight (0.105) Kappa coefficient ($p=0.002$) and 77.46% sensitivity and 36.83% specificity. However, in case females, WHR at $\geq 0.82 \text{ cm}$ was not statistically significant. WHR based screening of high-density lipoprotein cholesterol was not statistically significant for both female ($p=0.263$), and male ($p=0.661$) (Table 4).

Discussion

Developing locally appropriate optimal cut-off points using obesity based anthropometric measurement is important for early detection of metabolic syndrome to prevent farther complication and disabled from unrecognized NCD.⁷ According to the present study, BMI-based screening of triglyceride had fair and slight agreement of Kappa coefficient among males and females, respectively. This finding indicate that though the cut-off points for BMI is not the same for males ($\geq 22.2 \text{ kg/m}^2$) and females ($\geq 24 \text{ kg/m}^2$) like the international cut-off points, it can be used to early detect dyslipidemia among Ethiopian adults. A similar study done in Brazil also revealed that anthropometric indices had slight differences in screening of triglyceride among males and females. This study also revealed that BMI-based screening of triglyceride was applicable when compared with other anthropometric measurements.¹⁶ Likewise, other hospital-based studies done in India, Canada and others indicated presence of correlation between BMI and lipid profiles.^{17–19} However, in this study no significant association was observed between local BMI cut-off points and other lipid profiles like high and low density lipoprotein.

Waist circumference based screening of triglyceride among males showed slight agreement ($k=0.13$). However, similar studies from Canada and India found that waist circumference was a better predictor of dyslipidemia (TG)

Table 2 Comparison of BMI-Based Screening of Triglyceride and HDL with Laboratory Results Among Jimma Town Adults, Ethiopia

BMI-Based Screening				Laboratory Results									
Marker	Sex	Cut-Off Point	True Positive (a)	False Positive (b)	False Negative (c)	True Negative (d)	Total (P)	Sensitivity (%)	Specificity (%)	PPV	NPV	Kappa	P-value
TG≥150	F	BMI ≥24.5	53	114	60	212	439	72.6	65	32	83	0.12	<0.001
	M	BMI ≥22.2	86	101	56	233	476	64.1	66.8	45	81	0.275	<0.001
HDL<50	F	BMI ≥24.5	160	95	116	68	439	58	41.7	63	37	−0.003	0.949
HDL<40	M	BMI ≥22.2	104	162	75	135	476	58.1	45.5	39	64	0.032	0.449

Notes: Sensitivity=a/a+c; specificity=d/b+d; PPV=a/a+b; NPV=d/c+d; Kappa agreement (0=no/poor), (0.01–0.20: slight), (0.21–0.40=fair), (0.41–0.60=moderate), (0.61–0.80=substantial), and (0.81–1.00=perfect).

Table 3 Comparison of Waist Circumference-Based Screening of Triglyceride and HDL with Laboratory Results Among Adults, Jimma Town, Ethiopia

WC-Based Screening			Laboratory Results										
Markers	Sex	Cut-Off Value	True Positive (a)	False Positive (b)	False Negative (c)	True Negative (d)	Total (P)	Sensitivity (%)	Specificity (%)	PPV	NPV	Kappa	P-value
TG≥150	F	WC ≥78	82	239	31	87	439	72.6	26.7	25.5	73.7	−.005	0.877
	M	WC ≥83.7	54	84	88	250	476	38.0	74.9	39.1	73.9	0.13	0.005
HDL<50	F	WC ≥78	76	37	200	126	439	27.5	77.3	67.3	38.7	0.04	0.263
HDL>40	M	WC≥83.7	54	84	125	213	476	30.2	71.7	39.1	63.0	0.02	0.661

Notes: Sensitivity=a/a+c; specificity=d/b+d; PPV=a/a+b; NPV=d/c+d; Kappa agreement (0=no/poor), (0.01–0.20: slight), (0.21–0.40=fair), (0.41–0.60=moderate), (0.61–0.80=substantial), and (0.81–1.00=perfect).

Table 4 Comparison of WHR-Based Screening of Triglyceride and HDL with Laboratory Results Among Adults, Jimma Town, Ethiopia

WHR-Based Screening			Laboratory Results										
I	Sex	Cut-Off Value	True Positive (a)	False Positive (b)	False Negative (c)	True Negative (d)	Total (P)	Sensitivity (%)	Specificity (%)	PPV	NPV	Kappa	P-value
TG≥150	F	WHR≥0.82	112	324	1	2	439	99.12	0.613	25.7	66.7	−0.001	0.763
	M	WHR≥0.88	110	211	32	123	476	77.46	36.83	34.3	79.4	0.105	0.002
HDL<50	F	WHR≥0.82	274	162	2	1	439	99.28	0.613	62.84	33.3	−.001	0.891
HDL<40	M	WHR≥0.88	137	223	42	74	476	76.5	24.9	38.1	638	0.012	0.721

Notes: Sensitivity=a/a+c; specificity=d/b+d; PPV=a/a+b; NPV=d/c+d; Kappa agreement (0=no/poor), (0.01–0.20: slight), (0.21–0.40=fair), (0.41–0.60=moderate), (0.61–0.80=substantial), and (0.81–1.00=perfect).

in both sexes.^{18,20,21} Similar to waist circumference, the screening of triglyceride based on waist to hip ratio among males had slight Kappa agreement ($k=0.105$), which is statistically significant ($p=0.002$). Study done in India among adult population (≤ 60 years old) and aged group of population in Brazil (60 years old and above) revealed that anthropometric indices of waist to hip ratio was the best indicators for screening dyslipidemia (TG).^{16,22} This discrepancy might be due to the difference in sample size, age difference or ethnic differences, which can be addressed by a large scale and multi-centered study involving different populations.

Limitations of the Study

Anthropometric measurements are prone to errors. However, this study was conducted with highest precautions in training the anthropometric data collectors, equipment calibration and standardization of procedure to minimize the possibility of error occurrence.

Conclusion

Anthropometric indices can be used as a simple screening for dyslipidemia at primary health care unit. BMI cut-off points at ≥ 24.5 kg/m² for females and ≥ 22.2 kg/m² for males

can be used for screening of dyslipidemia ($TG \geq 150 \text{ mg/dl}$). Similarly, waist circumference at $\geq 83.7 \text{ cm}$ and waist to hip ratio at ≥ 0.88 can be considered as screening of dyslipidemia ($TG \geq 150 \text{ mg/dl}$) among males. Generally, with a further investigation healthcare providers and researchers can use locally validated anthropometric indices as alternative to investigate lipid profiles to be used in the prevention and management of metabolic syndrome and other non-communicable diseases.

Abbreviations

TG, triglycerides; HDL, high-density lipoprotein; LDL, low-density lipoprotein; WC, waist circumference; WHR, waist-to-hip ratio; BMI, body mass index.

Data Sharing Statement

The deidentified datasets used for this study are available from the corresponding author on reasonable request.

Ethical Approval

Ethical approval was obtained from Jimma University Ethical Review Board (Ref. IHRPGD/339/2019) and the study was conducted in accordance with the principles of the Declaration of Helsinki. The verbal consent was taken from study subjects based on the ethical approval letter granted by Jimma University. Prior to the first interview each respondent was informed about objective of the study and privacy during the interview. The privacy of study participants was respected and results regarding the participants made confidential.

Acknowledgments

The authors would like to express their gratitude to Jimma University for facilitating this study. We would also like to thank study participants who gave valuable information.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed on the journal to which the article will be submitted; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Funding

The authors received no financial support for the publication of the article.

Disclosure

The authors declare that they have no competing interests.

References

1. Kaur J. A comprehensive review on metabolic syndrome. *Cardiol Res Pract.* 2014;2014:943162. doi:10.1155/2014/943162
2. Brinck J, Thomas A, Lauer E, et al. Diabetes mellitus is associated with reduced high-density lipoprotein sphingosine-1-phosphate content and impaired. *Arterioscler Thromb Vasc Biol.* 2016;36(5):817–824. doi:10.1161/ATVBAHA.115.307049
3. World Health Organization. Global report on diabetes; 2016. Available from: https://apps.who.int/iris/bitstream/10665/204871/1/9789241565257_eng.pdf. Accessed January 11, 2020.
4. Kavishe B, Biraro S, Baisley K, et al. High prevalence of hypertension and of risk factors for non-communicable diseases (NCDs): a population based cross-sectional survey of NCDs and HIV infection in Northwestern Tanzania and Southern Uganda. *BMC Med.* 2015;13(1):126. doi:10.1186/s12916-015-0357-9
5. Shiferaw F, Letebo M, Misganaw A, et al. Non-communicable diseases in Ethiopia: disease burden, gaps in health care delivery and strategic directions. *Ethiop J Health Dev.* 2018;32(3).
6. Misganaw A, Haregu T, Deribe K, Tessema G, Deribew A. National mortality burden due to communicable, non-communicable, and other diseases in Ethiopia, 1990 – 2015: findings from the Global Burden of Disease Study 2015. *Popul Health Metr.* 2017;15(1):29. doi:10.1186/s12963-017-0145-1
7. Sinaga M, Worku M, Yemane T, et al. Optimal cut-off for obesity and markers of metabolic syndrome for Ethiopian adults. *Nutr J.* 2018;17(1):109. doi:10.1186/s12937-018-0416-0
8. Federal Democratic Republic of Ethiopia. The Ethiopia Noncommunicable Diseases and Injuries (NCDI) commission report summary: addressing the impact of noncommunicable diseases and injuries in Ethiopia. Available from: <https://static1.squarespace.com/static/55d4de6de4b011a1673a40a6/t/5bfc17ab21c67c558106119e/1543247788342/Ethiopia+NCDI+Commission+Report+Summary+FINAL.pdf>. Accessed March 11, 2019.
9. Asfaw D, Teshome E. Tobacco control and prevention efforts in Ethiopia pre- and post-ratification of WHO FCTC: current challenges and future directions. *Tob Induc Dis.* 2019;17:13. doi:10.18332/tid/102286
10. Sorana D. Medical diagnostic tests: a review of test anatomy, phases, and statistical treatment of data. *Comput Math Methods Med.* 2019;2019:1891569. doi:10.1155/2019/1891569
11. Gebreyes Y, Goshu D, Geletew T, et al. Prevalence of high blood-pressure, hyperglycemia, dyslipidemia, metabolic syndrome and their determinants in Ethiopia. *PLoS One.* 2018;13(5):e0194819. doi:10.1371/journal.pone.0194819
12. Green R, Wachsmann-Hogiu S. Development, history, and future of automated cell counters. *Clin Lab Med.* 2015;35(1):1–10. doi:10.1016/j.cll.2014.11.003
13. Hajian-Tilaki K. Sample size estimation in diagnostic test studies of biomedical informatics. *J Biomed Inform.* 2014;48:193–204. doi:10.1016/j.jbi.2014.02.013
14. Sinaga M, Yemane T, Tegene E, Lidstrom D, Belachew T. Performance of newly developed body mass index cut-off for diagnosing obesity among Ethiopian adults. *J Physiol Anthropol.* 2019;38(1):14. doi:10.1186/s40101-019-0205-2
15. Cohen W. A coefficient of agreement for nominal scales. *Educ Psychol Meas.* 1960;20(1):37–46. doi:10.1177/001316446002000104
16. Silva R, Fares D, Rodrigues A, Dél T, Matos W, Henrique M. Anthropometric indicators as predictors of serum triglycerides and hypertriglyceridemia in older adults. *MedicalExpress.* 2014;1(4):202–205. doi:10.5935/MedicalExpress.2014.04.08

17. Manjareeka M, Nanda S, Mishra J, Mishra S. Correlation between anthropometry and lipid profile in healthy subjects of Eastern India. *J Midlife Health*. 2015;6(4):164–168. doi:10.4103/0976-7800.172302
18. Brenner D, Tepylo K, Eny K, Cahill L, El-Sohehy A. Comparison of body mass index and waist circumference as predictors of cardiometabolic health in a population of young Canadian adults. *Diabetol Metab Syndr*. 2010;2(1):28. doi:10.1186/1758-5996-2-28
19. Ramoteme L, Martinique S, Sarah J, Makama A. The association between dyslipidemia and anthropometric indicators in black and white adolescents residing in tlokwe municipality, North-West Province, South Africa: the PAHL study. *Afr Health Sci*. 2011;14(4):929–938. doi:10.4314/ahs.v14i4.23
20. Rao S, Parab-Waingankar P. Performance of waist circumference relative to BMI in predicting risk of obesity and hypertension among affluent Indian adults. *Health (Irvine Calif)*. 2013;05(08):16–22. doi:10.4236/health.2013.58A3003
21. Silvana C, Miralles W, Wollinger LM, et al. Waist-to-height ratio (WHtR) and triglyceride to HDL-c ratio (TG/HDL-c) as predictors of cardiometabolic risk. *Nutr Hosp*. 2015;31(5):2115–2121. doi:10.3305/nh.2015.31.5.7773
22. Rocha F, Menezes TN, Melo RL, Pedraza DF. Correlation between indicators of abdominal obesity and serum lipids in the elderly. *Rev Assoc Med Bras*. 2013;59(1):48–55.

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion

and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>